

ALMOST  $Q$ -RINGS

C. JAYARAM

ABSTRACT. In this paper we establish some new characterizations for  $Q$ -rings and Noetherian  $Q$ -rings.

## 1. INTRODUCTION

Throughout this paper  $R$  is assumed to be a commutative ring with identity.  $L(R)$  denotes the lattice of all ideals of  $R$ .  $R$  is said to be a  $Q$ -ring [4], if every ideal is a finite product of primary ideals. It is well known that if  $R$  is a  $Q$ -ring, then  $R_M$  is a  $Q$ -ring for every maximal ideal  $M$  of  $R$  [4, Lemma 4]. But in general the converse need not be true. For example, if  $R$  is an almost Dedekind domain which is not a Dedekind domain, then  $R_M$  is a  $Q$ -ring, for every maximal ideal  $M$  of  $R$ , but  $R$  is not a  $Q$ -ring. We call a ring  $R$  an almost  $Q$ -ring if  $R_M$  is a  $Q$ -ring, for every maximal ideal  $M$  of  $R$ . The goal of this paper is to characterize those almost  $Q$ -rings which are also  $Q$ -rings. We prove that  $R$  is an almost  $Q$ -ring if and only if every non-maximal prime ideal is locally principal (see Theorem 1). Using this result, we characterize  $Q$ -rings in terms of almost  $Q$ -rings (see Theorem 2). Finally, we establish some equivalent conditions for Noetherian  $Q$ -rings (see Theorem 3).

For any  $A, B \in L(R)$ , we denote  $A \setminus B = \{x \in A \mid x \notin B\}$ . We use  $\subset$  for proper set containment. For any  $x \in R$ , the principal ideal generated by  $x$  is denoted by  $(x)$ . For any ideal  $I \in L(R)$ , we denote  $\theta(I) = \sum\{(I_1 : I) \mid I_1 \subseteq I \text{ and } I_1 \text{ is a finitely generated ideal}\}$ . Recall that an ideal  $I$  of  $R$  is called a *multiplication ideal* if for every ideal  $J \subseteq I$ , there exists an ideal  $K$  with  $J = KI$ . If  $I$  is a multiplication ideal, then  $I$  is locally principal [1, Theorem 1 and Page 761]. An ideal  $M$  of  $R$  is called a *quasi-principal ideal* [9, Exercise 10, Page 147] (or a principal element of  $L(R)$  [11]) if it satisfies the following identities (i)  $(A \cap (B : M))M = AM \cap B$  and (ii)  $(A + BM) : M = (A : M) + B$ , for all  $A, B \in L(R)$ . Obviously, every quasi-principal ideal is a multiplication ideal. It should be mentioned that every quasi-principal ideal is finitely generated and also a finite product of quasi-principal ideals of  $R$  is again a quasi-principal ideal [9, Exercise 10, Page 147]. In fact, an ideal  $I$  of  $R$  is quasi-principal if and only if it is finitely generated and locally

---

2000 *Mathematics Subject Classification*: 13A15, 13F20, 13G05.

*Key words and phrases*:  $Q$ -ring, almost  $Q$ -ring, Noetherian  $Q$ -ring.

Received July 11, 2002.

principal (see [6, Theorem 4]) or [11, Theorem 2]). A  $B_w$ -prime of  $I$  is a prime ideal  $P$  such that  $P$  is minimal over  $(I : x)$  for some  $x \in R$ .  $R$  is said to be a *Laskerian ring* [8], if every ideal is a finite intersection of primary ideals. It is well known that  $R$  is a  $Q$ -ring if and only if  $R$  is a Laskerian ring in which every non-maximal prime ideal is quasi-principal [4, Theorem 13].  $R$  is a  $\pi$ -ring if every principal ideal is a finite product of prime ideals. We say that  $R$  has Noetherian spectrum, if  $R$  satisfies the ascending chain condition for radical ideals [12]. It is well known that  $R$  has Noetherian spectrum if and only if every prime ideal is the radical of a finitely generated ideal [12, Corollary 2.4]. Also it is well known that if  $R$  has Noetherian spectrum, then every ideal has only finitely many minimal primes.

For general background and terminology, the reader is referred to [9].

We shall begin with the following definition.

**Definition 1.** A quasi-local ring  $R$  with maximal ideal  $M$  is said to satisfy the condition  $(*)$  if for each non-maximal prime ideal  $P$  with  $P = PM$ , there exists  $t \in M$  such that  $P + (t)$  is finitely generated.

Note that valuation rings (i.e., any two ideals are comparable), quasi-local rings in which the maximal ideals are principal and one dimensional quasi-local domains are examples of quasi-local rings satisfying the condition  $(*)$ .

**Lemma 1.** *Let  $R$  be a quasi-local  $Q$ -ring with maximal ideal  $M$ . Then  $R$  satisfies the condition  $(*)$ .*

**Proof.** The proof of the lemma follows from [4, Lemma 5]. □

**Lemma 2.** *Let  $R$  be a quasi-local ring with maximal ideal  $M$  satisfying the condition  $(*)$ . Suppose every principal ideal is a finite product of primary ideals. If  $P$  is a non-maximal prime ideal with  $P = PM$ , then  $P = (0)$ .*

**Proof.** Suppose  $P$  is a non-maximal prime ideal with  $P = PM$ . By hypothesis, there exists  $a \in M$  such that  $P + (a)$  is finitely generated. If  $a \in P$ , then  $P = P + (a)$  is finitely generated, so by Nakayama's lemma,  $P = 0$ . Suppose  $a \notin P$ . Since  $P + (a)$  is finitely generated, it follows that  $P + (a) = P_1 + (a)$  for some finitely generated ideal  $P_1 \subseteq P$ . Since  $P = PM$ , we have  $(P + (a))M = PM + (a)M = P_1M + (a)M$ , so  $P + (a)M = P_1M + (a)M$  and hence  $P + (a) = P_1M + (a)$ . Again since  $P_1 \subseteq P + (a) = (a) + P_1M$  and  $P_1$  is finitely generated, by Nakayama's lemma, it follows that  $P_1 \subseteq (a)$ . Therefore  $P \subseteq (a)$ . Let  $x \in P$ . By hypothesis  $(x) = QA$  for some primary ideal  $Q \subseteq P$  and  $A \in L(R)$ . Since  $Q \subseteq (a)$ , it follows that  $Q = (a)Q$ . Therefore  $(x) = QA = Q(a)A = (x)(a)$  and hence by Nakayama's lemma,  $(x) = (0)$ . This shows that  $P = (0)$ . □

**Lemma 3.** *Let  $R$  be a quasi-local ring with maximal ideal  $M$ . Suppose every ideal generated by two elements is a finite product of primary ideals. If  $P$  is a non-maximal prime ideal with  $P \neq PM$ , then  $P$  is principal.*

**Proof.** Let  $P$  be a non-maximal prime ideal with  $P \neq PM$ . Choose any element  $a \in P$  such that  $a \notin PM$ . Let  $t \in M$  be any element such that  $t \notin P$ . Suppose

$x \in P$ . Then by hypothesis,  $(a) + (xt)$  is a finite product of primary ideals. Since  $a \notin PM$ , it follows that  $(a) + (xt)$  is primary. Again since  $(xt) \subseteq (a) + (xt)$  and  $t \notin \sqrt{(a) + (xt)} \subseteq P$ , it follows that  $x \in (a) + (xt)$ , so by Nakayama's lemma  $(x) \subseteq (a)$ . Therefore  $P = (a)$ .  $\square$

**Lemma 4.** *Let  $R$  be a quasi-local ring with maximal ideal  $M$  satisfying the condition (\*). Suppose every ideal generated by two elements is a finite product of primary ideals. Then the non-maximal prime ideals are principal. Hence  $\dim R \leq 2$ .*

**Proof.** By Lemma 2 and Lemma 3, every non-maximal prime ideal is principal. Again as shown in the last paragraph of the proof of Lemma 5 of [4],  $\dim R \leq 2$ . This completes the proof of the lemma.  $\square$

**Lemma 5.** *Suppose  $I$  is an ideal of  $R$  such that every prime minimal over  $I$  is finitely generated. Then  $I$  contains a finite product of prime ideals minimal over  $I$ . Further  $I$  has only finitely many minimal primes.*

**Proof.** Suppose  $I$  does not contain a finite product of prime ideals minimal over  $I$ . Let  $\mathfrak{S} = \{J \in L(R) \mid I \subseteq J \text{ and } J \text{ does not contain a finite product of prime ideals minimal over } I\}$ . By Zorn's lemma,  $\mathfrak{S}$  has a maximal element, say  $P$ . It can be easily shown that  $P$  is a prime ideal. Again note that  $P$  contains a prime ideal  $P_0$  which is minimal over  $I$ , a contradiction. Therefore  $I$  contains a finite product of prime ideals minimal over  $I$ . Consequently,  $I$  has only finitely many minimal primes.  $\square$

**Lemma 6.** *Suppose  $R$  is a quasi-local ring. Then the following statements are equivalent:*

- (i)  $R$  is a  $Q$ -ring.
- (ii)  $R$  satisfies the condition (\*) and every ideal generated by two elements is a finite product of primary ideals.
- (iii) Every non-maximal prime ideal is principal.

**Proof.** (i) $\Rightarrow$ (ii) follows from Lemma 1.

(ii) $\Rightarrow$ (iii) follows from Lemma 4.

(iii) $\Rightarrow$ (i). Suppose (iii) holds. Then every ideal  $I$  is either  $M$ -primary ( $M$  is a maximal ideal of  $R$ ) or by Lemma 5,  $I$  has only finitely many minimal primes. Again by the last paragraph of the proof of [4, Lemma 5],  $R$  is Laskerian. Now the result follows from [4, Theorem 10].  $\square$

**Lemma 7.** *Let  $R$  be an almost  $Q$ -ring. Suppose every principal ideal is a finite product of primary ideals. Then every non-maximal prime ideal of  $R$  is a multiplication ideal.*

**Proof.** Using Lemma 6 and by imitating the proof of [4, Lemma 7], we can get the result.  $\square$

**Lemma 8.** *Let  $\dim R \leq 2$  and let every ideal generated by two elements has only finitely many minimal primes. Then  $R$  has Noetherian spectrum.*

**Proof.** First we show that every minimal prime ideal is the radical of a finitely generated ideal. By hypothesis,  $R$  has only finitely many minimal primes. Let  $P_1, P_2, \dots, P_n$  be the distinct minimal prime ideals. If  $n = 1$ , then  $P_1$  is the radical of the zero ideal. Suppose  $n > 1$ . Then  $P_1 \not\subseteq \bigcup_{i=2}^n P_i$ . Choose any  $x \in P_1$  such that  $x \notin \bigcup_{i=2}^n P_i$ . Let  $Q_1, Q_2, \dots, Q_m$  be the distinct primes minimal over  $(x)$ . Then  $P_1 = Q_j$  for some  $j$ , say  $P_1 = Q_1$ . If  $m = 1$ , then  $P_1$  is the radical of  $(x)$ . Suppose  $m > 1$ . Then  $P_1 \not\subseteq \bigcup_{i=2}^m Q_i$ . Choose any  $y \in P_1$  such that  $y \notin \bigcup_{i=2}^m Q_i$ . By hypothesis,  $(x) + (y)$  has only finitely many minimal primes. Let  $Q'_1, Q'_2, \dots, Q'_k$  be the distinct primes minimal over  $(x) + (y)$ . Note that  $P_1 = Q'_j$  for some  $j$ , say  $P_1 = Q'_1$ . If  $k = 1$ , then  $P_1$  is the radical of  $(x) + (y)$ . Suppose  $k > 1$ . Observe that any  $Q'_j$  different from  $P_1$  contains  $Q_i$  properly, for some  $i \neq 1$ , and each  $Q_i$  different from  $P_1$ , is non-minimal. So each  $Q'_j$  is maximal, for  $j = 2, 3, \dots, k$ . Choose any element  $z \in P_1$  such that  $z \notin \bigcup_{i=2}^k Q'_i$ . Now it can be easily shown that  $P_1$  is the radical of  $(x) + (y) + (z)$ . Thus we have shown that every minimal prime ideal is the radical of a finitely generated ideal.

Next we show that every non-minimal prime ideal is the radical of a finitely generated ideal. Let  $P$  be a non-minimal prime ideal. Then  $P \not\subseteq \bigcup_{i=1}^n P_i$ . Choose any  $x \in P$  such that  $x \notin \bigcup_{i=1}^n P_i$ . Let  $Q_1, Q_2, \dots, Q_m$  be the distinct primes minimal over  $(x)$ . Then  $P \supseteq Q_j$  for some  $j$ , say  $P \supseteq Q_1$ . If  $m = 1$  and  $P = Q_1$ , then  $P$  is the radical of  $(x)$  and so we are through. Suppose  $m \geq 1$  and  $Q_1 \subset P$ . Then  $P \not\subseteq \bigcup_{i=1}^m Q_i$ . Choose any  $y \in P$  such that  $y \notin \bigcup_{i=1}^m Q_i$ . Then  $(x) + (y)$  has only finitely many minimal primes and every prime minimal over  $(x) + (y)$  is a maximal ideal. Therefore there exists a finitely generated ideal  $I$  such that  $P$  is the radical of  $I$ . Finally assume that  $m > 1$  and  $P = Q_1$ . Then  $P \not\subseteq \bigcup_{i=2}^m Q_i$ . Choose any  $y \in P$  such that  $y \notin \bigcup_{i=2}^m Q_i$ . Let  $Q'_1, Q'_2, \dots, Q'_k$  be the distinct primes minimal over  $(x) + (y)$ . Note that  $P_1 \supseteq Q'_j$  for some  $j$ , say  $P_1 \supseteq Q'_1$ . Since  $x \in Q'_1$  and  $Q_1 = P \supseteq Q'_1$ , it follows that  $P = Q_1 = Q'_1$ . If  $k = 1$ , then  $P_1$  is the radical of  $(x) + (y)$ . Suppose  $k > 1$ . Then  $P \not\subseteq \bigcup_{i=2}^k Q'_i$  and each  $Q'_i$  different from  $P$ , is maximal. Choose any element  $z \in P$  such that  $z \notin \bigcup_{i=2}^k Q'_i$ . Then  $P$  is the radical of  $(x) + (y) + (z)$ . Thus every prime ideal is the radical of a finitely generated ideal and hence  $R$  has Noetherian spectrum.  $\square$

For any  $I \in L(R)$  and for any prime ideal  $P$  minimal over  $I$ , we denote  $P_I = \bigcap \{Q \in L(R) \mid Q \text{ is a } P\text{-primary ideal containing } I\}$ . It can be easily seen that  $P_I$  is the smallest  $P$ -primary ideal containing  $I$ . For any  $x \in R$ , and for any prime ideal  $P$  minimal over  $(x)$ , we denote  $P_x = \bigcap \{Q \in L(R) \mid Q \text{ is a } P\text{-primary ideal containing } (x)\}$ .

For any  $x \in R$ , we denote  $(x)^* = \bigcap \{P_x \mid P \text{ is a prime ideal minimal over } (x)\}$ .

**Lemma 9.** *Let  $P$  be a prime minimal over an ideal  $I$  of  $R$  and let  $P_1$  be a prime properly containing  $P$ . Then the following statements hold:*

- (i) *If  $P$  is a multiplication ideal, then  $P \subset ((I + PP_I) : P_I)$ .*
- (ii) *If  $P_1$  is a  $B_w$ -prime of  $I$ , then  $(P_I)_M \neq I_M$  (in  $R_M$ ) for all maximal ideals  $M$  containing  $P_1$ .*

**Proof.** (i) Consider the ideal  $((I + PP_I) : P_I)$ . Note that  $P \subseteq ((I + PP_I) : P_I)$ . Suppose  $P = ((I + PP_I) : P_I)$ . We claim that  $I + PP_I$  is  $P$ -primary. Let  $yz \in I + PP_I$  and  $z \notin P$ . Then  $yz \in P_I$ , so  $y \in P_I$ . Since  $P$  is a multiplication ideal, by [3, Lemma 1] and [2, Corollary],  $P_I$  is a multiplication ideal. As  $P_I$  is a multiplication ideal, it follows that  $(y) = P_I C$  for some ideal  $C$  of  $R$ . If  $C \subseteq P$ , then we are through. Suppose  $C \not\subseteq P$ . Then  $(yz) = (z)P_I C \subseteq I + PP_I$ , so  $zC \subseteq ((I + PP_I) : P_I) = P$ , a contradiction. Therefore  $I + PP_I$  is  $P$ -primary and hence  $P_I = I + PP_I$ . Consequently,  $1 \in ((I + PP_I) : P_I) = P$ , a contradiction. Therefore  $P \subset ((I + PP_I) : P_I)$ .

(ii) Suppose  $P_1$  is a  $B_w$ -prime of  $I$ . Then  $P_1$  is minimal over  $(I : r)$  for some  $r \in R$ . Since  $(I : r)r \subseteq I \subseteq P_I$ ,  $(I : r) \not\subseteq P$  and  $P_I$  is  $P$ -primary, it follows that  $r \in P_I$ . If  $(P_I)_M = I_M$  for some maximal ideal  $M$  containing  $P_1$ , then  $rs \in I$  for some  $s \notin M$ . So  $s \in (I : r) \subseteq M$ , a contradiction. Therefore the result is true.  $\square$

**Lemma 10.** *Let every non-maximal prime ideal of  $R$  be a multiplication ideal. Suppose  $P$  is a non-maximal minimal prime and minimal over an ideal  $I$  of  $R$ . Then the following statements hold:*

- (i) *Any  $B_w$ -prime of  $I$  which contains  $P$  properly, is a rank one maximal ideal and minimal over  $((I + PP_I) : P_I)$ .*
- (ii) *If the maximal ideals of  $R$  are finitely generated, then the ideal  $((I + PP_I) : P_I)$  has only finitely many minimal primes.*

**Proof.** (i) Let  $P_1$  be any  $B_w$ -prime of  $I$  which contains  $P$  properly. If  $M$  is a maximal ideal containing  $P_1$ , then by Lemma 9(ii),  $R_M$  is not a domain. Note that by Lemma 6,  $R$  is an almost  $Q$ -ring. As  $R$  is an almost  $Q$ -ring, by [4, Corollary 6], it follows that  $\text{rank } M = 1$ , so  $P_1$  is a rank one maximal ideal. If  $((I + PP_I) : P_I) \not\subseteq P_1$ , then  $(P_I)_{P_1} \subseteq I_{P_1} + (PP_I)_{P_1}$ . As  $P$  is a multiplication ideal, it follows that  $P_I$  is a multiplication ideal, so  $P_I$  is locally principal, and hence by Nakayama's lemma, it follows that  $(P_I)_{P_1} = I_{P_1}$ . But this contradicts the statement of Lemma 9(ii). Therefore  $((I + PP_I) : P_I) \subseteq P_1$  and hence by Lemma 9(i),  $P_1$  is minimal over  $((I + PP_I) : P_I)$ .

(ii) Note that by hypothesis,  $R$  is an almost  $Q$ -ring and so  $\dim R \leq 2$ . By Lemma 9(i), every prime minimal over  $((I + PP_I) : P_I)$  is either a non-minimal maximal ideal or a rank one non-maximal prime. As every non-maximal prime is a multiplication ideal, by [2, Theorem 3], the rank one non-maximal primes are quasi-principal. By hypothesis, the minimal primes over  $((I + PP_I) : P_I)$  are finitely generated and so by Lemma 5, the ideal  $((I + PP_I) : P_I)$  has only finitely many minimal primes.  $\square$

**Lemma 11.** *Suppose every ideal (of  $R$ ) generated by two elements has only finitely many minimal primes and the non-maximal prime ideals are multiplication ideals. Then the non-maximal prime ideals are quasi-principal.*

**Proof.** Let  $P$  be a non-maximal prime ideal. As  $\dim R \leq 2$ , it follows that  $P$  is either minimal or a rank one prime. If  $P$  is non-minimal, then  $P$  is quasi-principal [2, Theorem 3]. Suppose  $P$  is minimal. By Lemma 8,  $P = \sqrt{I}$  for some finitely generated ideal  $I$  of  $R$ . Note that every  $B_w$ -prime of  $I$  contains  $P$ , and by Lemma 8, the ideal  $((I + PP_I) : P_I)$  has only finitely many minimal primes. Therefore by Lemma 10(i),  $I$  has only finitely many  $B_w$ -primes. Again note that by Lemma 10(i), for every finitely generated ideal  $I_0$  with  $I \subseteq I_0 \subseteq P$ ,  $I_0$  has only finitely many  $B_w$ -primes. As  $\dim R \leq 2$ , by [7, Theorem 1.3],  $P$  is finitely generated and hence quasi-principal.  $\square$

**Lemma 12.** *Suppose every non-maximal prime ideal of  $R$  is a multiplication ideal, the maximal ideals of  $R$  are finitely generated and every principal ideal has only finitely many minimal primes. Then every principal ideal is a finite intersection of primary ideals.*

**Proof.** Note that by hypothesis,  $R$  is an almost  $Q$ -ring, so by Lemma 4,  $\dim R \leq 2$ . Let  $x \in R$ . Then by hypothesis,  $(x)^*$  is a finite intersection of primary ideals. Suppose  $(x)$  is not contained in any minimal prime. We show that  $(x) = (x)^*$ . Let  $M$  be a maximal ideal. If  $x \notin M$ , then  $(x)_M = (x)^*_M$ . Suppose  $x \in M$ . If  $M$  is minimal over  $(x)$ , then  $(x)_M = (x)^*_M$ . Suppose  $M$  is not minimal over  $(x)$ . Then  $\text{rank } M = 2$ , so by [4, Corollary 6],  $R_M$  is a  $\pi$ -domain. Therefore  $(x)_M = (x)^*_M$  (see the proof of [10, Theorem 1.2] or [5, Theorem 3]). This shows that  $(x)_M = (x)^*_M$  for all maximal ideals containing  $x$  and hence  $(x) = (x)^*$ .

Now assume that  $P_1, P_2, \dots, P_m$  be the primes minimal over  $(x)$ . Let  $P_1, P_2, \dots, P_t$  be the non-maximal minimal primes and let  $P_{t+1}, P_{t+2}, \dots, P_m$  be the primes which are either maximal or rank one non-maximal primes. By Lemma 10(ii), the ideals  $((x) + P_i(P_i)_x : (P_i)_x)$  for  $i = 1, 2, \dots, t$  have only finitely many minimal primes, say  $M_1, M_2, \dots, M_n$ . Again by the proof of Lemma 10(ii), these are either non-minimal maximal ideals or rank one non-maximal prime ideals. Without loss of generality, assume that  $M_1, M_2, \dots, M_k$  are the rank one maximal prime ideals and  $M_{k+1}, M_{k+2}, \dots, M_n$  are either rank two maximal ideals or rank one non-maximal prime ideals. Let  $M$  be any maximal ideal different from  $M_1, M_2, \dots, M_k$ . We claim that  $(x)_M = (x)^*_M$ . Obviously, if  $x \notin M$ , then  $(x)_M = (x)^*_M$ . Suppose  $x \in M$ . If either  $M$  is minimal over  $(x)$  or  $\text{rank } M = 2$ , then  $(x)_M = (x)^*_M$ . Suppose  $M$  is not minimal over  $(x)$  and  $\text{rank } M = 1$ . Then  $M$  is different from  $M_1, M_2, \dots, M_n$ , so  $((x) + P_i(P_i)_x : (P_i)_x) \not\subseteq M$  for  $i = 1, 2, \dots, t$  and hence  $((P_i)_x)_M = (x)_M$  for  $i = 1, 2, \dots, t$ . Consequently,  $(x)_M = (x)^*_M$ . If  $(x)_{M_i} = (x)^*_{M_i}$  for  $i = 1, 2, \dots, k$ , then  $(x)_M = (x)^*_M$  for all maximal ideals, so  $(x) = (x)^*$ . Suppose  $(x)_{M_i} \neq (x)^*_{M_i}$  for  $i = 1, 2, \dots, l$  ( $1 \leq l \leq k$ ). As  $R_{M_i}$  is a Laskerian ring, it follows that there exist  $M_i$ -primary  $Q_i$  such that  $(x)_{M_i} = ((x)^*_{M_i} \cap (Q_i)_{M_i})$  for  $i = 1, 2, \dots, l$ . Then  $(x)_M = ((x)^* \cap Q_1 \cap Q_2 \cap \dots \cap Q_l)_M$  for all maximal ideals  $M$  of  $R$ . Therefore  $(x) = (x)^* \cap Q_1 \cap Q_2 \cap \dots \cap Q_l$  and

hence  $(x)$  is a finite intersection of primary ideals. This completes the proof of the lemma.  $\square$

**Lemma 13.** *Suppose  $R$  is a quasi-local ring in which the maximal ideal  $M$  is finitely generated. If every ideal generated by two elements is a finite product of primary ideals, then  $R$  is a Noetherian  $Q$ -ring.*

**Proof.** If  $M$  is minimal, then we are through. Suppose  $M$  is non-minimal. By Lemma 6, it is enough if we show that  $R$  satisfies the condition  $(*)$ . Let  $P$  be a non-maximal prime ideal with  $P = PM$ . Let  $\Psi = \{P_\alpha \mid P \subseteq P_\alpha, P_\alpha \text{ is prime and } P_\alpha = P_\alpha M\}$ . Clearly  $\Psi \neq \emptyset$  and by Zorn's lemma,  $\Psi$  has a maximal element, say  $P_0$ . Note that  $P_0 \neq M$ . If  $P_0 \subset P_1 \subset M$  for some prime ideal  $P_1$ , then  $P_1 \neq P_1 M$ , so by Lemma 3,  $P_1$  is principal and hence  $P$  is contained in a principal ideal. Now assume that  $M$  covers  $P_0$ . Choose any  $x \in M$  such that  $x \notin P_0$ . Then  $P_0 + (x)$  is  $M$ -primary. As  $M$  is finitely generated, it follows that  $M^k \subseteq P_0 + (x)$  for some positive integer  $k$ . Again since  $P_0 = P_0 M$ , it follows that  $P_0 \subseteq M^n$  for all positive integers  $n$ . Therefore  $M^k \subseteq P_0 + (x) \subseteq (x) + M^{k+1} = (x) + M^k M$  and hence by Nakayama's lemma  $P_0 \subset M^k \subseteq (x)$ . This shows that  $P$  is properly contained in  $(x)$  and hence  $R$  satisfies the condition  $(*)$ .  $\square$

**Lemma 14.** *Suppose every finitely generated ideal of  $R$  is a finite product of primary ideals. Suppose  $I$  is an ideal of  $R$  such that  $I$  is locally finitely generated and every prime minimal over  $I$  is a maximal ideal. Then  $I$  is finitely generated.*

**Proof.** We claim that  $\theta(I) = R$ . Suppose  $\theta(I) \neq R$ . Then  $\theta(I) \subseteq M$  for some maximal ideal  $M$  of  $R$ . Since  $I$  is locally finitely generated, it follows that  $I_M = (I_1)_M$  for some finitely generated ideal  $I_1$  contained in  $I$ . By hypothesis, there exist primary ideals  $Q_1, Q_2, \dots, Q_n$  such that  $I_1 = Q_1 Q_2 \dots Q_n$ . Without loss of generality, assume that  $Q_i \subseteq M$  for  $i = 1, 2, \dots, k$  and  $Q_j \not\subseteq M$  for  $j = k+1, k+2, \dots, n$ . Then  $I_M = (I_1)_M = (Q_1)_M (Q_2)_M \dots (Q_k)_M$ . Since  $I_M \subseteq (Q_i)_M$ , it follows that  $I \subseteq Q_i$  for  $i = 1, 2, \dots, k$ . Since  $M$  is minimal over  $I$ , it follows that each  $Q_i$  is  $M$ -primary and hence  $Q_1 Q_2 \dots Q_k$  is  $M$ -primary. Therefore  $I \subseteq Q_1 Q_2 \dots Q_k$ . Choose elements  $x_j \in Q_j$  such that  $x_j \notin M$  for  $j = k+1, k+2, \dots, n$ . Let  $z = x_{k+1} x_{k+2} \dots x_n$ . Since  $I \subseteq Q_1 Q_2 \dots Q_k$  and  $z \in Q_{k+1} Q_{k+2} \dots Q_n$ , it follows that  $Iz \subseteq Q_1 Q_2 \dots Q_n = I_1$ , so  $z \in (I_1 : I) \subseteq \theta(I) \subseteq M$ , which is a contradiction. Therefore  $\theta(I) = R$  and hence  $R = \sum_{i=1}^n (I_i : I)$ , where  $I_i$ 's are finitely generated ideals contained in  $I$ . So  $I = \sum_{i=1}^n I_i$ . This shows that  $I$  is a finitely generated ideal.  $\square$

**Theorem 1.**  *$R$  is an almost  $Q$ -ring if and only if every non-maximal prime ideal is locally principal.*

**Proof.** The result follows from Lemma 6.  $\square$

**Corollary 1.** *Suppose every principal ideal is a finite product of primary ideals. Then  $R$  is an almost  $Q$ -ring if and only if every non-maximal prime ideal is a multiplication ideal.*

**Proof.** The proof of the corollary follows from Theorem 1 and Lemma 7.  $\square$

**Corollary 2.** *Suppose every principal ideal is a finite intersection of primary ideals. Then  $R$  is an almost  $Q$ -ring if and only if every non-maximal prime ideal is quasi-principal.*

**Proof.** The proof of the corollary follows from Theorem 1 and [4, Theorem 12].  $\square$

**Theorem 2.** *The following statements on  $R$  are equivalent:*

- (i)  $R$  is a  $Q$ -ring.
- (ii)  $R$  is an almost  $Q$ -ring in which every ideal generated by two elements is a finite intersection of primary ideals.
- (iii)  $R$  is an almost  $Q$ -ring in which every ideal generated by two elements is a finite product of primary ideals.
- (iv) Every ideal generated by two elements is a finite product of primary ideals and for every maximal ideal  $M$  of  $R$ ,  $R_M$  satisfies the condition (\*).
- (v) Every non-maximal prime ideal is a multiplication ideal and every ideal generated by two elements has only finitely many minimal primes.

**Proof.** (i) $\Rightarrow$ (ii) and (i) $\Rightarrow$ (iii) follow from [4, Lemma 4 and Theorem 10].

(ii) $\Rightarrow$ (v) follows from Corollary 2.

(iii) $\Rightarrow$ (iv) follows from Lemma 1.

(iv) $\Rightarrow$ (v) follows from Lemma 6 and Corollary 1.

(v) $\Rightarrow$ (i). Suppose (v) holds. By Lemma 4 and Lemma 6,  $\dim R \leq 2$ . By Lemma 8,  $R$  has Noetherian spectrum. Also by Lemma 11, every non-maximal prime ideal is quasi-principal. Therefore by [4, Lemma 1], every primary ideal whose radical is non-maximal is a power of its radical and hence quasi-principal. Consequently, every primary ideal whose radical is non-maximal is finitely generated. Again by [8, Corollary 2.3],  $R$  is Laskerian and hence by [4, Theorem 13],  $R$  is a  $Q$ -ring.  $\square$

The following theorem gives some new equivalent conditions for Noetherian  $Q$ -rings.

**Theorem 3.** *The following statements on  $R$  are equivalent:*

- (i)  $R$  is a Noetherian  $Q$ -ring.
- (ii) The maximal ideals are locally finitely generated and every ideal generated by two elements is a finite product of primary ideals.
- (iii)  $R$  is an almost  $Q$ -ring in which the maximal ideals are finitely generated and every principal ideal is a finite product of primary ideals.

**Proof.** (i) $\Rightarrow$ (ii) is obvious.

(ii) $\Rightarrow$ (iii). Suppose (ii) holds. By Lemma 13,  $R$  is locally Noetherian and an almost  $Q$ -ring. By Theorem 2,  $R$  is a  $Q$ -ring and so by Lemma 14, the maximal ideals are finitely generated. Therefore (iii) holds.

(iii) $\Rightarrow$ (i). Suppose (iii) holds. By Corollary 1, Corollary 2 and Lemma 12,  $R$  is a Noetherian ring and hence by Theorem 2,  $R$  is a Noetherian  $Q$ -ring. This completes the proof of the theorem.  $\square$



## REFERENCES

- [1] Anderson, D. D., *Multiplication ideals, Multiplication rings and the ring  $R(X)$* , Canad. J. Math. XXVIII (1976), 760–768.
- [2] Anderson, D. D., *Some remarks on multiplication ideals*, Math. Japon. **25** (1980), 463–469.
- [3] Anderson, D. D., *Noetherian rings in which every ideal is a product of primary ideals*, Canad. Math. Bull. **23** (4), (1980), 457–459.
- [4] Anderson, D. D. and Mahaney, L. A., *Commutative rings in which every ideal is a product of primary ideals*, J. Algebra **106** (1987), 528–535.
- [5] Anderson, D. D. and Mahaney, L. A., *On primary factorizations*, J. Pure Appl. Algebra **54** (1988), 141–154.
- [6] Becerra, L. and Johnson, J. A., *A note on quasi-principal ideals*, Tamkang J. Math. (1982), 77–82.
- [7] Heinzer, W. and Ohm, J., *Locally Noetherian commutative rings*, Trans. Amer. Math. Soc. **158** (1971), 273–284.
- [8] Heinzer, W. and Lantz, D., *The Laskerian property in commutative rings*, J. Algebra **72** (1981), 101–114.
- [9] Larsen, M. D. and McCarthy, P. J., *Multiplicative theory of ideals* Academic Press, New York 1971.
- [10] Levitz, K. B., *A characterization of general ZPI-rings*, Proc. Amer. Math. Soc. **32** (1972), 376–380.
- [11] McCarthy, P. J., *Principal elements of lattices of ideals*, Proc. Amer. Math. Soc. **30** (1971), 43–45.
- [12] Ohm, J. and Pendleton, R. L., *Rings with Noetherian spectrum*, Duke Math. J. **35** (1968), 631–639.

UNIVERSITY OF BOTSWANA, DEPARTMENT OF MATHEMATICS  
P/ BAG 00704, GABORONE, BOTSWANA  
E-mail: chillumu@mopipi.ub.bw