

RIEMANNIAN SYMMETRIES IN FLAG MANIFOLDS

PAOLA PIU AND ELISABETH REMM

ABSTRACT. Flag manifolds are in general not symmetric spaces. But they are provided with a structure of \mathbb{Z}_2^k -symmetric space. We describe the Riemannian metrics adapted to this structure and some properties of reducibility. The conditions for a metric adapted to the \mathbb{Z}_2^2 -symmetric structure to be naturally reductive are detailed for the flag manifold $SO(5)/SO(2) \times SO(2) \times SO(1)$.

1. INTRODUCTION

In this work we call flag manifold any homogeneous space

$$G/H = \frac{SO(n)}{SO(p_1) \times \cdots \times SO(p_k)}$$

with $p_1 + \cdots + p_k = n$. We suppose also that $p_1 \geq p_2 \geq \cdots \geq p_k$. When $k = 2$ and $p_2 = 1$, this space is isomorphic to the sphere S^{n-1} and it is a symmetric space. When $k = 2$ and $p_1 \neq 1$, the homogeneous space G/H is isomorphic to the Grassmannian manifold and it is also a symmetric space. But if $k > 2$ the homogeneous space G/H is reductive but not symmetric. In [2] and [4] we have shown that we can define on G/H a structure of \mathbb{Z}_2^k -symmetric space, that is, the Lie algebra \mathfrak{g} of G admits a \mathbb{Z}_2^k -grading. A Riemannian metric on G/H is called associated with the \mathbb{Z}_2^k -symmetric structure if the natural symmetries defining the \mathbb{Z}_2^k -symmetric structure are isometries. The Riemannian geometry is more complicated than the Riemannian geometry of symmetric spaces (or \mathbb{Z}_2 -symmetric spaces); in particular, it is not always naturally reductive. In the paper we investigate the \mathbb{Z}_2^k -symmetric Riemannian tensor on flag manifolds and develop all the computations when $G/H = SO(5)/SO(2) \times SO(2) \times SO(1)$.

2. RIEMANNIAN \mathbb{Z}_2^k -SYMMETRIC SPACES

This notion has been introduced in [9]. Let G/H be an homogeneous space with a connected Lie group G . We denote by \mathfrak{g} and \mathfrak{h} respectively the Lie algebras of G and H .

2010 *Mathematics Subject Classification*: primary 53C30.

Key words and phrases: \mathbb{Z}_2^k -symmetric space, flag manifolds, Riemannian metrics.

The first author was supported by a Visiting Professor fellowship at Université de Haute Alsace – Mulhouse on February 2012 and by GNSAGA(Italy).

DOI: 10.5817/AM2012-5-387

Definition 1. The homogeneous space G/H is \mathbb{Z}_2^k -symmetric if the Lie algebra \mathfrak{g} admits a \mathbb{Z}_2^k -grading, that is,

$$\mathfrak{g} = \bigoplus_{\gamma \in \mathbb{Z}_2^k} \mathfrak{g}_\gamma, \quad [\mathfrak{g}_{\gamma_1}, \mathfrak{g}_{\gamma_2}] \subset \mathfrak{g}_{\gamma_1 \gamma_2},$$

with $\mathfrak{g}_e = \mathfrak{h}$ where e is the identity of \mathbb{Z}_2^k .

If $\mathfrak{m} = \bigoplus_{\gamma \in \mathbb{Z}_2^k, \gamma \neq e} \mathfrak{g}_\gamma$ then $\mathfrak{g} = \mathfrak{g}_e \oplus \mathfrak{m}$ and the \mathbb{Z}_2^k -grading implies $[\mathfrak{g}_e, \mathfrak{m}] \subset \mathfrak{m}$. But if $k \geq 2$, we have not $[\mathfrak{m}, \mathfrak{m}] \subset \mathfrak{g}_e$. Thus the decomposition $\mathfrak{g} = \mathfrak{g}_e \oplus \mathfrak{m}$ is reductive, not symmetric when $k \geq 2$.

Consequence. A \mathbb{Z}_2^k -symmetric homogeneous space G/H is reductive.

Example 2. From [2], [4] and [7], it is possible to give a classification of the \mathbb{Z}_2^2 -grading of complex simple Lie algebras. In the following list we give the pairs $(\mathfrak{g}, \mathfrak{g}_e = \mathfrak{h})$ which correspond to the (local) classification of \mathbb{Z}_2^2 -symmetric structures when G is simple complex or G is simple compact and real.

\mathfrak{g}	$\mathfrak{g}_e = \mathfrak{h}$	\mathfrak{g}	$\mathfrak{g}_e = \mathfrak{h}$
$so(k_1 + k_2 + k_3 + k_4),$ $k_1 \geq k_2 \geq k_3 \geq k_4, k_3 \neq 0$	$\bigoplus_{i=1}^4 so(k_i)$	E_6	$so(6) \oplus \mathbb{C}$
$sp(k_1 + k_2 + k_3 + k_4),$ $k_1 \geq k_2 \geq k_3 \geq k_4, k_3 \neq 0$	$\bigoplus_{i=1}^4 sp(k_i)$		$sp(2) \oplus sp(2)$
$so(2m)$	$gl(m)$		$sp(3) \oplus sp(1)$
$so(2k_1 + 2k_2)$	$gl(k_1) \oplus gl(k_2)$		$su(3) \oplus su(3) \oplus \mathbb{C}^2$
$sp(2m)$	$gl(m)$		$su(4) \oplus sp(1) \oplus sp(1) \oplus \mathbb{C}$
$sp(2k_1 + 2k_2)$	$gl(k_1) \oplus gl(k_2)$		$su(5) \oplus \mathbb{C}^2$
$so(2m)$	$so(m)$		$so(8) \oplus \mathbb{C}^2$
$so(4m)$	$sp(2m)$	E_7	$so(9)$
$sp(4m)$	$sp(2m)$		$so(8)$
$sp(2m)$	$so(m)$		$su(4) \oplus su(4) \oplus \mathbb{C}$
$sl(2n)$	$sl(n)$		$sp(4)$
$sl(k_1 + k_2)$	$\bigoplus_{i=1}^2 sl(k_i) \oplus \mathbb{C}$		$su(6) \oplus sp(1) \oplus \mathbb{C}$
$sl(k_1 + k_2 + k_3)$	$\bigoplus_{i=1}^3 sl(k_i) \oplus \mathbb{C}^2$		$so(8) \oplus so(4) \oplus sp(1)$
$sl(k_1 + k_2 + k_3 + k_4)$	$\bigoplus_{i=1}^4 sl(k_i) \oplus \mathbb{C}^3$		$u(6) \oplus \mathbb{C}$
$so(8)$	$gl(3) \oplus gl(1)$		$so(10) \oplus \mathbb{C}^2$
			F_4
\mathfrak{g}	$\mathfrak{g}_e = \mathfrak{h}$	\mathfrak{g}	$\mathfrak{g}_e = \mathfrak{h}$
E_8	$so(8) \oplus so(8)$	F_4	$u(3) \oplus \mathbb{C}$
	$su(8) \oplus \mathbb{C}$		$sp(2) \oplus sp(1) \oplus sp(1)$
	$so(12) \oplus sp(1) \oplus sp(1)$		$so(8)$
	$E_6 \oplus \mathbb{C}^2$	G_2	\mathbb{C}^2

If $M = G/H$ is a \mathbb{Z}_2^k -symmetric space, the grading

$$\mathfrak{m} = \bigoplus_{\gamma \in \mathbb{Z}_2^k, \gamma \neq e} \mathfrak{g}_\gamma$$

is associated with a spectral decomposition of \mathfrak{g} defined by a family $\{\sigma_\gamma, \gamma \in \mathbb{Z}_2^k\}$ of automorphisms of \mathfrak{g} satisfying

$$\begin{cases} \sigma_\gamma^2 = \text{Id}, \\ \sigma_{\gamma_1} \circ \sigma_{\gamma_2} = \sigma_{\gamma_2} \circ \sigma_{\gamma_1}, \end{cases}$$

for any $\gamma, \gamma_1, \gamma_2 \in \mathbb{Z}_2^k$. Any $\sigma_\gamma \in \text{Aut}(\mathfrak{g})$ defines an automorphism s_γ of G and H , if it is connected, corresponds to the identity component of the group $\{A \in G/\forall \sigma \in \mathbb{Z}_2^k, s_\gamma(A) = A\}$. The family $\{s_\gamma, \gamma \in \mathbb{Z}_2^k\}$ is a subgroup of $\text{Aut}(G)$ isomorphic to \mathbb{Z}_2^k . For any $x \in G/H$, it determines a subgroup $\{s_{\gamma,x}, \gamma \in \mathbb{Z}_2^k\}$ of $\text{Diff}(M)$ isomorphic to \mathbb{Z}_2^k . The diffeomorphisms $s_{\gamma,x}$ are called the symmetries of the \mathbb{Z}_2^k -symmetric space G/H . By extension, we will also call symmetries, the automorphisms s_γ of G .

Remark. A Γ -symmetric space is usually called generalized symmetric space when Γ is a cyclic group. In this case the grading of the Lie algebra is defined in the complex field and corresponds to the roots of the unity. For a general presentation, see [8].

Definition 3 ([3]). A Riemannian metric g on the (reductive) \mathbb{Z}_2^k -symmetric space G/H is called adapted to the \mathbb{Z}_2^k -symmetric structure if the symmetries $s_{\gamma,x}$ are isometries for any γ in \mathbb{Z}_2^k and x in G/H .

In this case we will say that G/H is a Riemannian \mathbb{Z}_2^k -symmetric space.

As G/H is a reductive homogeneous space, we consider only G -invariant Riemannian metrics on G/H which are positive-definite $\text{ad}(H)$ -invariant symmetric bilinear form B on \mathfrak{m} , with the following correspondence

$$B(X, Y) = g(X, Y)_0$$

for $X, Y \in \mathfrak{m}$. Since in this paper we consider H connected, the invariance of B writes as

$$B([Z, X], Y) + B(X, [Z, Y]) = 0,$$

for X, Y in \mathfrak{m} and $Z \in \mathfrak{h}$.

Proposition 4. Let G/H be a \mathbb{Z}_2^k -symmetric structure with G and H connected. Any Riemannian metric g adapted to the \mathbb{Z}_2^k -symmetric structure is in one-to-one correspondence with the $\text{ad}(H)$ -invariant positive-definite bilinear form B on \mathfrak{m} such that $B(\mathfrak{g}_\gamma, \mathfrak{g}_{\gamma'}) = 0$ for $\gamma \neq \gamma'$ in \mathbb{Z}_2^k where $\mathfrak{g} = \bigoplus_{\gamma \in \mathbb{Z}_2^k} \mathfrak{g}_\gamma$ is the \mathbb{Z}_2^k -grading corresponding to the \mathbb{Z}_2^k -symmetric structure of G/H . Let $U(X, Y)$ be the symmetric bilinear mapping of $\mathfrak{m} \times \mathfrak{m}$ in \mathfrak{m} defined by

$$2B(U(X, Y), Z) = B(X, [Z, Y]_{\mathfrak{m}}) + B([Z, X]_{\mathfrak{m}}, Y),$$

for all $X, Y, Z \in \mathfrak{m}$, where $[\]_{\mathfrak{m}}$ denotes the projection on \mathfrak{m} of the bracket of \mathfrak{g} .

Thus the Riemannian connection for g is given by

$$\nabla_X Y = U(X, Y) + \frac{1}{2}[X, Y]_{\mathfrak{m}}$$

and the curvature tensor satisfies

$$R(X, Y)(Z) = \nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X, Y]_{\mathfrak{m}}} Z - [[X, Y]_{\mathfrak{h}}, Z],$$

for $X, Y, Z \in \mathfrak{m}$ and where the term $[[X, Y]_{\mathfrak{h}}, Z]$ corresponds to the linear isotropy representation of H into G/H .

3. THE RIEMANNIAN \mathbb{Z}_2^2 -SYMMETRIC SPACE $SO(5)/SO(2) \times SO(2) \times SO(1)$

Any Riemannian symmetric space is naturally reductive. It is not usually the case for Riemannian \mathbb{Z}_2^k -symmetric spaces as soon as $k \geq 2$. In this section we describe the Riemannian and Ricci tensors for any \mathbb{Z}^2 -symmetric metric on the flag manifold $SO(5)/SO(2) \times SO(2) \times SO(1)$. In particular we study some properties of these metrics when they are not naturally reductive. A \mathbb{Z}_2^2 -grading of the Lie algebra $so(5)$ is given by the decomposition

$$\mathfrak{g} = \mathfrak{g}_e \oplus \mathfrak{g}_a \oplus \mathfrak{g}_b \oplus \mathfrak{g}_c,$$

corresponding to the multiplication of \mathbb{Z}_2^2 : $a^2 = b^2 = c^2 = e, ab = c, ac = b, bc = a, e$ being the identity. To describe the components of the grading, we consider

$$so(5) = \left\{ \left(\begin{array}{ccccc} 0 & x_1 & a_1 & a_2 & b_1 \\ -x_1 & 0 & a_3 & a_4 & b_2 \\ -a_1 & -a_3 & 0 & x_2 & c_1 \\ -a_2 & -a_4 & -x_2 & 0 & c_2 \\ -b_1 & -b_2 & -c_1 & -c_2 & 0 \end{array} \right), x_i, a_i, b_i, c_i \in \mathbb{R} \right\}.$$

Thus

$$\begin{aligned} \mathfrak{g}_e &= \{X \in so(5) / a_i = b_i = c_i = 0\}, \\ \mathfrak{g}_a &= \{X \in so(5) / x_i = b_i = c_i = 0\}, \\ \mathfrak{g}_b &= \{X \in so(5) / x_i = a_i = c_i = 0\}, \\ \mathfrak{g}_c &= \{X \in so(5) / x_i = a_i = b_i = 0\}. \end{aligned}$$

We have $\mathfrak{g}_e = so(2) \oplus so(2) \oplus so(1)$ and this grading induces the \mathbb{Z}_2^2 -symmetric structure on $SO(5)/SO(2) \times SO(2) \times SO(1)$. Moreover, from [2], this grading is unique up to an equivalence of \mathbb{Z}_2^2 -gradings. We denote by $\{\{X_1, X_2\}, \{A_1, A_2, A_3, A_4\}, \{B_1, B_2\}, \{C_1, C_2\}\}$ the basis of $so(5)$ where each big letter corresponds to the matrix of $so(5)$ with the small letter equal to 1 and other coefficients zero. This basis is adapted to the grading. Let us denote by $\{\omega_1, \omega_2, \alpha_1, \alpha_2, \alpha_3, \alpha_4, \beta_1, \beta_2, \gamma_1, \gamma_2\}$ the dual basis.

Theorem 5. *Any \mathbb{Z}_2^2 -symmetric Riemannian metric g on $SO(5)/SO(2) \times SO(2) \times SO(1)$ is given by an $ad(H)$ -invariant symmetric bilinear form B on $\mathfrak{m} = \mathfrak{g}_a \oplus \mathfrak{g}_b \oplus \mathfrak{g}_c$*

$$B = t^2(\alpha_1^2 + \alpha_2^2 + \alpha_3^2 + \alpha_4^2) + u(\alpha_1\alpha_4 - \alpha_2\alpha_3) + v^2(\beta_1^2 + \beta_2^2) + w^2(\gamma_1^2 + \gamma_2^2)$$

with $tvw \neq 0$ and $u \in]-4t^2, 4t^2[$.

Proof. It is explain in detail in [3] and [4]. □

The Riemannian connection is given by

$$\nabla_X(Y) = \frac{1}{2}[X, Y]_{\mathfrak{m}} + U(X, Y),$$

for any $X, Y \in \mathfrak{m}$, where U is the symmetric bilinear mapping on $\mathfrak{m} \times \mathfrak{m}$ into \mathfrak{m} defined by

$$2B(U(X, Y), Z) = B(X, [Z, Y]_{\mathfrak{m}}) + B([Z, X]_{\mathfrak{m}}, Y),$$

for any $X, Y, Z \in \mathfrak{m}$. The bilinear mapping U is reduced to

$$B = \sum_{i=1}^4 \tilde{\alpha}_i^2 + \sum_{i=1}^2 \tilde{\beta}_i^2 + \sum_{i=1}^2 \tilde{\gamma}_i^2$$

with

$$\begin{aligned} \tilde{\alpha}_1 &= t\alpha_1 + \frac{u}{2t}\alpha_4, & \tilde{\alpha}_2 &= t\alpha_2 - \frac{u}{2t}\alpha_3, & \tilde{\alpha}_3 &= K\alpha_3, & \tilde{\alpha}_4 &= K\alpha_4, \\ \tilde{\beta}_1 &= v\beta_1, & \tilde{\beta}_2 &= v\beta_2, & \tilde{\gamma}_1 &= w\gamma_1, & \tilde{\gamma}_2 &= w\gamma_2. \end{aligned}$$

with $K = \sqrt{t^2 - \frac{u^2}{4t^2}}$. If $\{\tilde{A}_j, \tilde{B}_i, \tilde{C}_i\}$ is the dual basis, this basis is orthonormal and the brackets $[\ , \]_{\mathfrak{m}}$ are given by

	\tilde{A}_1	\tilde{A}_2	\tilde{A}_3	\tilde{A}_4	\tilde{B}_1	\tilde{B}_2	\tilde{C}_1	\tilde{C}_2
\tilde{A}_1	0	0	0	0	$-\frac{w}{tv}\tilde{C}_1$	0	$\frac{v}{tw}\tilde{B}_1$	0
\tilde{A}_2		0	0	0	$-\frac{w}{tv}\tilde{C}_2$	0	0	$\frac{v}{tw}\tilde{B}_1$
\tilde{A}_3			0	0	$-\frac{uw}{2t^2vK}\tilde{C}_2$	$-\frac{w}{Kv}\tilde{C}_1$	$\frac{v}{Kw}\tilde{B}_2$	$\frac{uv}{2t^2wK}\tilde{B}_1$
\tilde{A}_4				0	$\frac{uw}{2t^2vK}\tilde{C}_1$	$-\frac{w}{Kv}\tilde{C}_2$	$-\frac{uv}{2t^2wK}\tilde{B}_1$	$\frac{v}{Kw}\tilde{B}_2$
\tilde{B}_1					0	0	$-\frac{t}{vw}\tilde{A}_1$	$-\frac{t}{vw}\tilde{A}_2$
\tilde{B}_2						0	$\frac{u}{2vwt}\tilde{A}_2 - \frac{K}{vw}\tilde{A}_3$	$-\frac{u}{2vwt}\tilde{A}_1 - \frac{K}{vw}\tilde{A}_4$
\tilde{C}_1							0	0
\tilde{C}_2								0

Recall that the Riemannian connection ∇ is given by

$$\nabla_X(Y) = U(X, Y) + \frac{1}{2}[X, Y]_{\mathfrak{m}},$$

and the Riemannian curvature $R(X, Y)$ is the matrix given by

$$R(X, Y)(Z) = \nabla_X\nabla_Y Z - \nabla_Y\nabla_X Z - \nabla_{[X, Y]_{\mathfrak{m}}} Z - [[X, Y]_{\mathfrak{h}}, Z],$$

for $X, Y, Z \in \mathfrak{m}$ and where the term $[[X, Y]_{\mathfrak{h}}, Z]$ corresponds to the linear isotropy representation of H into G/H .

The symmetric mapping U is given by:

$U(X, Y)$	\widetilde{A}_1	\widetilde{A}_2	\widetilde{A}_3	\widetilde{A}_4	\widetilde{B}_1	\widetilde{B}_2	\widetilde{C}_1	\widetilde{C}_2
\widetilde{A}_1	0	0	0	0	$\frac{t^2-v^2}{2tvw} \widetilde{C}_1$	$\frac{u}{4vwt} \widetilde{C}_2$	$-\frac{t^2+w^2}{2tvw} \widetilde{B}_1$	$-\frac{u}{4vwt} \widetilde{B}_2$
\widetilde{A}_2		0	0	0	$\frac{t^2-v^2}{2tvw} \widetilde{C}_2$	$-\frac{u}{4vwt} \widetilde{C}_1$	$\frac{u}{4vwt} \widetilde{B}_2$	$-\frac{t^2+w^2}{2tvw} \widetilde{B}_1$
\widetilde{A}_3			0	0	$\frac{-uv}{4t^2wK} \widetilde{C}_2$	$\frac{K^2-v^2}{2Kvw} \widetilde{C}_1$	$-\frac{K^2+w^2}{2Kvw} \widetilde{B}_2$	$\frac{uw}{4t^2vK} \widetilde{B}_1$
\widetilde{A}_4				0	$\frac{uv}{4t^2wK} \widetilde{C}_1$	$\frac{K^2-v^2}{2Kvw} \widetilde{C}_2$	$-\frac{uw}{4t^2vK} \widetilde{B}_1$	$-\frac{K^2+w^2}{2Kvw} \widetilde{B}_2$
\widetilde{B}_1					0	0	$\frac{v^2-w^2}{2vwt} \widetilde{A}_1 + \frac{u(v^2-w^2)}{4t^2vwK} \widetilde{A}_4$	$\frac{v^2-w^2}{2vwt} \widetilde{A}_2 - \frac{u(v^2-w^2)}{4t^2vwK} \widetilde{A}_3$
\widetilde{B}_2						0	$\frac{v^2-w^2}{2vwK} \widetilde{A}_3$	$\frac{v^2-w^2}{2vwK} \widetilde{A}_4$
\widetilde{C}_1							0	0
\widetilde{C}_2								0

Infinitesimal isometries are given by the vectors $X \in \mathfrak{m}$ satisfying

$$B(A_X Y, Z) = -B(Y, A_X Z),$$

for any $Y, Z \in \mathfrak{m}$, where $A_X Y = [X, Y] - \nabla_X Y$. Since $\nabla_X Y = U(X, Y) + \frac{1}{2}[X, Y]$, we have $A_X Y = -U(X, Y) + \frac{1}{2}[X, Y]$. Thus $X \in \mathfrak{m}$ is an infinitesimal isometry if

$$B([X, Y], Z) + B(Y, [X, Z]) = 0,$$

for any $Y, Z \in \mathfrak{m}$.

Proposition 6. *If $X \in \mathfrak{m}$ is an infinitesimal isometry, thus*

(1) *If $u = 0$ and*

(a) *If $t^2 = v^2$, $X = c_1 \widetilde{C}_1 + c_2 \widetilde{C}_2$,*

(b) *If $t^2 = w^2$, $X = c_1 \widetilde{B}_1 + c_2 \widetilde{B}_2$,*

(c) *If $v^2 = w^2$, $X = a_1 \widetilde{A}_1 + a_2 \widetilde{A}_2 + a_3 \widetilde{A}_3 + a_4 \widetilde{A}_4$.*

(2) *If $u \neq 0$ and*

(a) *If $v^2 = w^2$, $X = a_1 \widetilde{A}_1 + a_2 \widetilde{A}_2 + a_3 \widetilde{A}_3 + a_4 \widetilde{A}_4$,*

(b) *If $v^2 \neq w^2$, $X = 0$.*

In particular, if $u = 0$ and $t^2 = v^2 = w^2$, the bilinear form B is $\text{ad}(\mathfrak{m})$ -invariant.

4. ON THE (NON)NATURALLY REDUCTIVITY

We consider on the \mathbb{Z}_2^2 -symmetric space $SO(5)/SO(2) \times SO(2) \times SO(1)$ the \mathbb{Z}_2^2 -symmetric Riemannian metric associated with the bilinear form

$$B = t^2(\alpha_1^2 + \alpha_2^2 + \alpha_3^2 + \alpha_4^2) + u(\alpha_1\alpha_4 - \alpha_2\alpha_3) + v^2(\beta_1^2 + \beta_2^2) + w^2(\gamma_1^2 + \gamma_2^2)$$

with $tvw \neq 0$ and $u \in]-4t^2, 4t^2[$.

Proposition 7. *The \mathbb{Z}_2^2 -symmetric space $SO(5)/SO(2) \times SO(2) \times SO(1)$ is naturally reductive if and only if*

$$u = 0 \quad \text{and} \quad t^2 = v^2 = w^2.$$

Proof. Indeed, naturally reductivity means that

$$B(X, [Z, Y]_{\mathfrak{m}}) + B([Z, X]_{\mathfrak{m}}, Y) = 0,$$

that is, $B(U(X, Y), Z) = 0$ for any $X, Y, Z \in \mathfrak{m}$. From the expression of U we deduce $u = 0$ and $t^2 = v^2 = w^2 = K^2$. But $u = 0$ implies $K^2 = t^2$. \square

Now, we assume that we have not the naturally reductivity property. In this case, we can study if such a Riemannian space is a d'Atri space, that is, the geodesic symmetries preserve the volume. Let us recall that naturally reductive homogeneous spaces are d'Atri spaces, but these two notions are not equivalent. We know some examples of d'Atri spaces which are not naturally reductive. The condition of being a d'Atri space is often difficult to compute because it is necessary to know the equations of geodesics. But the D'Atri definition is equivalent to the Ledger (infinite) system whose first equation writes

$$L(X, Y, Z) = (\nabla_X \rho)(Y, Z) + (\nabla_Y \rho)(Z, X) + (\nabla_Z \rho)(X, Y) = 0,$$

for any $X, Y, Z \in \mathfrak{m}$, where ρ is the Ricci tensor, that is, the trace of the linear operator $V \rightarrow R(V, X)Y$. In the orthonormal basis previously defined the Ricci tensor is the symmetric matrix

$$\begin{pmatrix} \rho_{11} & 0 & 0 & \rho_{14} & 0 & 0 & 0 & 0 \\ 0 & \rho_{22} & \rho_{23} & 0 & 0 & 0 & 0 & 0 \\ 0 & \rho_{23} & \rho_{33} & 0 & 0 & 0 & 0 & 0 \\ \rho_{14} & 0 & 0 & \rho_{44} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \rho_{55} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \rho_{66} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \rho_{77} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \rho_{88} \end{pmatrix}$$

with

$$\begin{aligned} \rho_{11} = \rho_{22} &= \frac{4t^4 + u^2 - 4(v^4 - 6v^2w^2 + w^4)}{8t^2v^2w^2}, \\ \rho_{14} = -\rho_{23} &= \frac{u(K^2t^2 + v^4 - 6v^2w^2 + w^4)}{4Kt^3v^2w^2}, \\ \rho_{33} = \rho_{44} &= \frac{(4t^4 - u^2)^2 - 4(v^4 - 6v^2w^2 + w^4)(u^2 + 4t^4)}{8t^2(4t^4 - u^2)v^2w^2}, \\ \rho_{55} = \rho_{66} &= \frac{-4t^6 + 12t^4w^2 + t^2(u^2 + 4v^4 - 4w^4) - 3u^2w^2}{(4t^4 - u^2)v^2w^2}, \\ \rho_{77} = \rho_{88} &= \frac{-4t^6 + 12t^4v^2 + t^2(u^2 - 4v^4 + 4w^4) - 3u^2v^2}{(4t^4 - u^2)v^2w^2}. \end{aligned}$$

The non trivial Ledger equations (for the first condition) are

$$\begin{cases} L(\widetilde{A}_1, \widetilde{B}_1, \widetilde{C}_1) = L(\widetilde{A}_2, \widetilde{B}_1, \widetilde{C}_2), \\ L(\widetilde{A}_1, \widetilde{B}_2, \widetilde{C}_2) = L(\widetilde{A}_2, \widetilde{B}_2, \widetilde{C}_1), \\ L(\widetilde{A}_3, \widetilde{B}_1, \widetilde{C}_2) = L(\widetilde{A}_4, \widetilde{B}_1, \widetilde{C}_1), \\ L(\widetilde{A}_3, \widetilde{B}_2, \widetilde{C}_1) = L(\widetilde{A}_4, \widetilde{B}_2, \widetilde{C}_2). \end{cases}$$

Thus we obtain

$$(*) \quad \begin{cases} (v^2 - w^2)\rho_{11} + (w^2 - t^2)\rho_{55} + (t^2 - v^2)\rho_{77} + \frac{u(w^2 - v^2)}{2tK}\rho_{14} = 0, \\ -\frac{u}{2t}\rho_{55} + \frac{u}{2t}\rho_{77} + \frac{v^2 - w^2}{K}\rho_{14} = 0, \\ \frac{u(v^2 - w^2)}{2tvwK}\rho_{33} + \frac{uw}{2tvK}\rho_{55} - \frac{v^2 - w^2}{vw}\rho_{14} - \frac{uv}{2twK}\rho_{77} = 0, \\ (v^2 - w^2)\rho_{33} + (w^2 - K^2)\rho_{55} + (K^2 - v^2)\rho_{77} = 0. \end{cases}$$

- If $u = 0$, thus $\rho_{11} = \rho_{33}$, $\rho_{14} = 0$ and $(*)$ is equivalent to

$$(v^2 - w^2)\rho_{33} + (w^2 - t^2)\rho_{55} + (t^2 - v^2)\rho_{77} = 0,$$

that is,

$$(v^2 - w^2)(9t^4 + v^4 + 10v^2w^2 + w^4 - 10t^2v^2 - 10t^2w^2) = 0.$$

We obtain

$$\begin{cases} v^2 = w^2 \\ \text{or } 9t^4 + v^4 + 10v^2w^2 + w^4 - 10t^2v^2 - 10t^2w^2 = 0. \end{cases}$$

Let us study the second equation. For this, since $t \neq 0$, we can consider the change of variables

$$V = \frac{v^2}{t^2}, \quad W = \frac{w^2}{t^2}.$$

The equation becomes

$$9 - 10(V + W) + (V + W)^2 + 8VW = 0.$$

Now we put $S = V + W$, $P = VW$ and we obtain

$$P = \frac{-S^2 + 10S - 9}{8}.$$

Since V and W are strictly positive, $P > 0$, which implies that $-S^2 + 10S - 9 > 0$, that is, $S \in]1, 9[$. With these conditions, V and W are the roots of $X^2 - SX + P = 0$. In fact they always exist since $X^2 - SX + P = X^2 - SX + \frac{-S^2+10S-9}{8}$ has as discriminant

$$\Delta = S^2 - 4P = S^2 - \frac{-S^2 + 10S - 9}{2} = \frac{3S^2 - 10S + 9}{2}$$

which is always positive. The roots of $X^2 - SX + P$ are

$$\begin{cases} X_1 = \frac{S - \sqrt{\frac{3S^2 - 10S + 9}{2}}}{2}, \\ X_2 = \frac{S + \sqrt{\frac{3S^2 - 10S + 9}{2}}}{2}. \end{cases}$$

Since $S \in]1, 9[$, $P > 0$ and $X_1 > 0$, $X_2 > 0$. We obtain

$$\left(\frac{v^2}{t^2}, \frac{w^2}{t^2}\right) = (X_1, X_2) \quad \text{or} \quad (X_2, X_1).$$

Proposition 8. *Assume that the Riemannian \mathbb{Z}_2^3 -symmetric metric on $SO(5)/SO(2) \times SO(2) \times SO(1)$ associated with a bilinear form B with $u = 0$ satisfies the first Ledger condition. Then B is one of the following bilinear form*

$$\begin{aligned} B_1 &= t^2(\alpha_1^2 + \alpha_2^2 + \alpha_3^2 + \alpha_4^2) + v^2(\beta_1^2 + \beta_2^2 + \gamma_1^2 + \gamma_2^2), \\ B_2(S) &= t^2(\alpha_1^2 + \alpha_2^2 + \alpha_3^2 + \alpha_4^2) + \frac{S - \sqrt{\frac{3S^2 - 10S + 9}{2}}}{2}(\beta_1^2 + \beta_2^2) \\ &\quad + \frac{S + \sqrt{\frac{3S^2 - 10S + 9}{2}}}{2}(\gamma_1^2 + \gamma_2^2), \\ B_3(S) &= t^2(\alpha_1^2 + \alpha_2^2 + \alpha_3^2 + \alpha_4^2) + \frac{S + \sqrt{\frac{3S^2 - 10S + 9}{2}}}{2}(\beta_1^2 + \beta_2^2) \\ &\quad + \frac{S - \sqrt{\frac{3S^2 - 10S + 9}{2}}}{2}(\gamma_1^2 + \gamma_2^2), \end{aligned}$$

with $S \in]1, 9[$. The metrics associated with $B_2(S)$ and $B_3(S)$ are not naturally reductive and the matrix associated with B_1 is not naturally reductive as soon as $t^2 \neq v^2$.

- Assume that $u \neq 0$. The first Ledger condition writes

$$\begin{cases} -t^2(v^2 - w^2)(36t^6 - 40t^4(v^2 + w^2) + 6u^2(v^2 + w^2) \\ \quad + t^2(-9u^2 + 4(v^4 + 10v^2w^2 + w^4))) = 0, \\ t^2u(v^2 - w^2)(28t^4 - 7u^2 - 16t^2(v^2 + w^2) + 4(v^4 - 6v^2w^2 + w^4)) = 0, \\ (v^2 - w^2)(144t^8 - 160t^6(v^2 + w^2) + 40t^2u^2(v^2 + w^2) \\ \quad - 16t^4(4u^2 - v^4 - 10v^2w^2 - w^4) + u^2(7u^2 - 4(v^4 - 6v^2w^2 + w^4))) = 0, \\ t^2u(v^2 - w^2)(4t^6 - 12t^4(v^2 + w^2) + 3u^2(v^2 + w^2) - t^2(u^2 - 32v^2w^2)) = 0. \end{cases}$$

We obtain $v^2 = w^2$ or

$$\begin{cases} 36t^6 - 40t^4(v^2 + w^2) + 6u^2(v^2 + w^2) + t^2(-9u^2 + 4(v^4 + 10v^2w^2 + w^4)) = 0, \\ 28t^4 - 7u^2 - 16t^2(v^2 + w^2) + 4(v^4 - 6v^2w^2 + w^4) = 0, \\ 144t^8 - 160t^6(v^2 + w^2) + 40t^2u^2(v^2 + w^2) \\ \quad - 16t^4(4u^2 - v^4 - 10v^2w^2 - w^4) + u^2(7u^2 - 4(v^4 - 6v^2w^2 + w^4)) = 0, \\ 4t^6 - 12t^4(v^2 + w^2) + 3u^2(v^2 + w^2) - t^2(u^2 - 32v^2w^2) = 0. \end{cases}$$

The change of variables: $U = \frac{u}{t^2}$, $V = \frac{v^2}{t^2}$, $W = \frac{w^2}{t^2}$, $S = V + W$ and $P = VW$ shows that the first three equations of the previous system are equivalent. So the system reduces to the two equations

$$(1) \quad 64P - 24PS + 4S - 13S^2 + 3S^3 = 0,$$

$$(2) \quad 7U^2 = 28 - 16S + 4(S^2 - 8P).$$

Equation (1) gives

$$P = \frac{-4S + 13S^2 - 3S^3}{8(8 - 3S)} = \frac{-S(S - 4)(3S - 1)}{8(8 - 3S)},$$

because $S = \frac{8}{3}$ is not a solution. Thus V and W are roots of $X^2 - SX + P = 0$ if there are two real solutions of this equation. But

$$X^2 - SX + P = X^2 - SX + \frac{-S(S - 4)(3S - 1)}{8(8 - 3S)}$$

and its discriminant is

$$\Delta = S^2 - 4P = S^2 + \frac{S(S - 4)(3S - 1)}{2(8 - 3S)} = \frac{S(-3S^2 + 3S + 4)}{2(8 - 3S)}.$$

We obtain a condition on S to the existence of V and W which is

$$S \in]0, \frac{3 + \sqrt{57}}{6} [\cup] \frac{8}{3}, +\infty [.$$

Moreover we want V and W to be positive solutions so $P > 0$ and $S > 0$; we have to take $S \in]\frac{1}{3}, \frac{8}{3}[\cup]4, +\infty[$. Finally

$$S \in]\frac{1}{3}, \frac{3 + \sqrt{57}}{6}[\cup]4, +\infty[.$$

Then Equation (2) gives conditions on S to the existence of U . In fact Equation (2) is equivalent to

$$U^2 = 4 \frac{8 - 7S + S^2}{8 - 3S}.$$

Then $S \in]0, \frac{7 - \sqrt{17}}{2}[\cup]\frac{8}{3}, \frac{7 + \sqrt{17}}{2}[$. But we have also that $u \in]-4t^2, 4t^2[$ so $U^2 < 16$ which reduces to $S^2 + 5S - 24 < 0$ and $S \in]0, \frac{7 - \sqrt{17}}{2}[\cup]\frac{8}{3}, 3[$.

With the conditions coming from (1) we finally have that $S \in]\frac{1}{3}, \frac{7 - \sqrt{17}}{2}[$.

Proposition 9. *Assume that the \mathbb{Z}_2^2 -symmetric metric on $SO(5)/SO(2) \times SO(2) \times SO(1)$ associated with a bilinear form B with $u \neq 0$ satisfies the first Ledger condition. Then S belongs to $]\frac{1}{3}, \frac{7 - \sqrt{17}}{2}[$ and $B = t^2(\alpha_1^2 + \alpha_2^2 + \alpha_3^2 + \alpha_4^2) + u(\alpha_1\alpha_4 - \alpha_2\alpha_4) + v^2(\beta_1^2 + \beta_2^2) + w^2(\gamma_1^2 + \gamma_2^2)$, with*

(1)

$$v^2 = \frac{-16S + 6S^2 - \sqrt{2S(32 + 12S - 33S^2 + 9S^3)}}{-32 + 12S} t^2$$

and

$$w^2 = \frac{-3S(S - 4)(S - \frac{1}{3})}{8(8 - 3S)V} t^2$$

(2)

$$w^2 = \frac{-16S + 6S^2 - \sqrt{2S(32 + 12S - 33S^2 + 9S^3)}}{-32 + 12S} t^2$$

and

$$v^2 = \frac{-3S(S - 4)(S - \frac{1}{3})}{8(8 - 3S)V} t^2$$

and in each case $u^2 = 4 \frac{8 - 7S + S^2}{8 - 3S} t^4$. None of these metrics is naturally reductive.

REFERENCES

[1] Arias-Marco, T., Kowalski, O., *Classification of 4-dimensional homogeneous D'Atri spaces*, Czechoslovak Math. J. **58** (1) (2008), 203–239.
 [2] Bahturin, Y., Goze, M., *\mathbb{Z}_2^2 -symmetric spaces*, Pacific J. Math. **236** (1) (2008), 1–21.
 [3] Bouyakoub, A., Goze, M., Remm, E., *On Riemannian non symmetric spaces and flag manifolds*, arXiv:math/0609790.
 [4] Goze, M., Remm, E., *Γ -symmetric spaces*, Differential geometry, World Sci. Publ., Hackensack, NJ, 2009, pp. 195–206.
 [5] Kobayashi, S., Nomizu, K., *Foundations of differential geometry*, Volumes I, II, Interscience Tracts in Pure and Applied Mathematics, No. 15, Interscience Publishers John Wiley and Sons, Inc., New York–London–Sydney, 1969.

- [6] Kobayashi, S., Nomizu, K., *Foundations of differential geometry*, volume I, Interscience Publishers, John Wiley and Sons, New York–London, 1963.
- [7] Kollross, A., *Exceptional $Z_2 \times Z_2$ -symmetric spaces*, Pacific J. Math. **242** (1) (2009), 113–130.
- [8] Kowalski, O., *Generalized symmetric spaces*, volume II, Lecture Notes in Math. **805**, Springer–Verlag, Berlin–New York, 1980.
- [9] Lutz, R., *Sur la géométrie des espaces Γ -symétriques*, C. R. Acad. Sci. Paris Sér. I Math. **293** (1) (1981), 55–58.

DIPARTIMENTO DI MATEMATICA E INFORMATICA, UNIVERSITÀ DEGLI STUDI DI CAGLIARI,
VIA OSPEDALE 72, 09124 CAGLIARI, ITALIA
E-mail: piu@unica.it

LABORATOIRE DE MATHÉMATIQUES ET APPLICATIONS,
UNIVERSITÉ DE HAUTE ALSACE, FACULTÉ DES SCIENCES ET TECHNIQUES,
4, RUE DES FRÈRES LUMIÈRE, 68 093 MULHOUSE CEDEX, FRANCE
E-mail: Elisabeth.Remm@uha.fr