

## CONVERGENCE OF BANACH LATTICE VALUED STOCHASTIC PROCESSES WITHOUT THE RADON-NIKODYM PROPERTY

V. MARRAFFA

ABSTRACT. We obtain almost sure convergence theorems for stochastic processes consisting of Bochner integrable functions taking values in a Banach lattice without assuming the Radon-Nikodym property. It is shown that if the limit exists in a weak sense then the almost sure convergence follows.

### 1. INTRODUCTION

For Banach lattice valued subpramarts the Radon-Nikodym property is equivalent to the convergence a. e. (see [4], [11] and [6]). If the Radon-Nikodym property is not assumed it is natural to ask how small can be the class  $T$  of functionals  $f$  such that the a.s. convergence of  $fX_n$  to  $fX$  for  $f \in T$  implies the convergence of  $X_n$  to  $X$  in some stronger sense. In case of Banach valued processes it was established that  $T$  can be a total set. In particular in [8] it was proved that an amart  $(X_n)$  converges scalarly almost surely to a random variable  $X$  if  $fX_n$  converges to  $fX$  a.s for each  $f$  in a total subset of the dual. In [3], under the same assumption, the strong a.s. convergence for martingales follows. Analogous results has been obtained also for weak amarts and uniform amarts in [1].

In §3 we obtain similar results for subpramarts taking values in a Banach lattice (see Theorem 2).

In §4, under a suitable covering condition (Vitali condition  $V$ ), we generalize the subpramarts result to directed sets.

### 2. DEFINITIONS AND NOTATIONS

Throughout this note  $(\Omega, \mathcal{F}, P)$  is a probability space and  $(\mathcal{F}_n)_{n \in \mathbb{N}}$  a family of sub- $\sigma$ -algebras of  $\mathcal{F}$  such that  $\mathcal{F}_m \subset \mathcal{F}_n$  if  $m < n$ . Moreover, without loss of generality, we will assume that  $\mathcal{F}$  is the completion of  $\sigma(\cup_n \mathcal{F}_n)$ . From now on  $E$  will denote a Banach lattice with norm  $\|\cdot\|$  and  $E^*$  its dual. A subset  $T$  of  $E^*$  is called a *total set* over  $E$  if  $f(x) = 0$  for each  $f \in T$  implies  $x = 0$ . For an element  $x \in E$

---

Received December 22, 2003.

2000 *Mathematics Subject Classification*. Primary 60G48, 28B05.

*Key words and phrases*. Random variable, stopping time, subpramart, scalar convergence.

Supported by MURST of Italy.

we denote by  $x^+$  the least upper bound between  $x$  and 0. The Banach lattice  $E$  is said to *have the order continuous norm* or, briefly, to be *order continuous*, if for every downward directed set  $\{x_\alpha\}_\alpha$  in  $E$  with  $\bigwedge_\alpha x_\alpha = 0$ , then  $\lim_\alpha \|x_\alpha\| = 0$ . The norm on  $E$  has the *Kadec-Klee property with respect to a set  $D \subset E^*$*  if whenever  $\lim_n f(x_n) = f(x)$  for every  $f \in D$  and  $\lim_n \|x_n\| = \|x\|$ , then  $\lim_n x_n = x$  strongly. If  $D = E^*$  we say that the norm has the *Kadec-Klee property*. It was proved in [2] the following renorming theorem for Banach lattices.

**Theorem 1.** *A Banach lattice  $E$  is order continuous if and only if there is an equivalent lattice norm on  $E$  with the Kadec-Klee property.*

It is obvious that if  $E$  is separable, the equivalent norm has the Kadec-Klee property with respect to a countable set of functionals.

A *stopping time* is a map  $\tau : \Omega \rightarrow \mathbb{N} \cup \{\infty\}$  such that, for each  $n \in \mathbb{N}$ ,  $\{\tau \leq n\} = \{\omega \in \Omega : \tau(\omega) \leq n\} \in \mathcal{F}_n$ . We denote by  $\Gamma$  the collection of all simple stopping times (i.e. taking finitely many values and not taking the value  $\infty$ ). Then  $\Gamma$  is a set filtering to the right.

We recall that a stochastic process  $(X_n, \mathcal{F}_n)$  is called

- (i) a *submartingale* if  $X_n \leq E(X_{n+1} | \mathcal{F}_n)$  a.s. for each  $n \in \mathbb{N}$ , or equivalently if

$$\int_A X_n \leq \int_A X_{n+1},$$

for each  $A \in \mathcal{F}_n$  and for each  $n \in \mathbb{N}$ ;

- (ii) a *subpramart* if for each  $\varepsilon > 0$  there exists  $\tau_0 \in \Gamma$  such that for all  $\tau$  and  $\sigma$  in  $\Gamma$ ,  $\tau > \sigma > \tau_0$  then

$$P(\{\|(X_\sigma - E(X_\tau | \mathcal{F}_\sigma))^+\| > \varepsilon\}) \leq \varepsilon.$$

We remind that if  $(X_n, \mathcal{F}_n)$  is a positive subpramart (i.e.  $X_n(\omega) \geq 0$  for each  $n \in \mathbb{N}$  and  $\omega \in \Omega$ ), then for each  $f \in (E^*)^+$ , where  $(E^*)^+$  denotes the nonnegative cone in  $E^*$ ,  $(fX_n, \mathcal{F}_n)$  and  $(\|X_n\|, \mathcal{F}_n)$  are real valued positive subpramarts [5, Lemma viii.1.12].

### 3. CONVERGENCE THEOREMS FOR PROCESSES INDEXED BY $\mathbb{N}$

We will need the following Propositions.

**Proposition 1.** [5, p. 303] *Let  $E$  be a Banach space and let  $(X_n, \mathcal{F}_n)$  be a  $L^1$ -bounded stochastic process. Then there exists a subsequence  $(n_k)_k$  in  $\mathbb{N}$  such that for every  $k \in \mathbb{N}$*

$$X_{n_k} = Y_{n_k} + Z_{n_k}$$

where  $Y_{n_k}$  and  $Z_{n_k}$  are  $\mathcal{F}_{n_k}$ -measurable,  $(Y_{n_k})_k$  is uniformly integrable and  $\lim_k Z_{n_k} = 0$  a.s..

**Proposition 2.** [5, p. 298] *Let  $(X_n^m, \mathcal{F}_n)$  be a sequence of real valued positive subpramarts for which for each  $\varepsilon > 0$  there exists  $\tau_0 \in \Gamma$  such that for all  $\tau$  and  $\sigma$  in  $\Gamma$ ,  $\tau > \sigma > \tau_0$  then*

$$P(\{\sup_m (X_\sigma^m - E(X_\tau^m | \mathcal{F}_\sigma)) \leq \varepsilon\}) \geq 1 - \varepsilon.$$

Suppose, moreover, that there is a subsequence  $(n_k)_k$  such that

$$\sup_k \int \sup_m X_{n_k}^m < \infty.$$

Then each subpramart  $(X_n^m, \mathcal{F}_n)_n$  converges a.s. to an integrable function  $X^m$  and we have

$$\lim_n (\sup_m X_n^m) = \sup_m X^m \text{ a.s..}$$

We are able to prove the following theorem.

**Theorem 2.** [9, Theorem 3.8] *Let  $E$  be an order continuous Banach lattice, which is weakly sequentially complete and let  $T$  be a total subset of  $E^*$ . Let  $(X_n, \mathcal{F}_n)$  be a positive subpramart with an  $L^1$ -bounded subsequence and let  $X$  be a strongly measurable random variable. Assume that, for each  $f \in T$ ,  $fX_n$  converges to  $fX$  a.s. (the null depends on  $f$ ). Then  $X_n$  converges to  $X$  strongly, a.s..*

*Proof.* Since  $(X_n)$  and  $X$  are strongly measurable it is possible to assume that  $E$  is separable. Using Proposition 1 and the fact that a subsequence of  $(X_n)_n$ , still denoted by  $(X_n)_n$ , is  $L^1$ -bounded we can also assume that

$$X_{n_k} = Y_{n_k} + Z_{n_k}$$

where  $Y_{n_k}$  and  $Z_{n_k}$  are  $\mathcal{F}_{n_k}$ -measurable,  $(Y_{n_k})_k$  is uniformly integrable and

$$\lim_k Z_{n_k} = 0 \text{ a.s..}$$

For each  $f \in (E^*)^+$ ,  $(fX_n)_n$  is a real valued subpramart with a  $L^1$ -bounded subsequence, then it converges a.s. to a real random variable  $X_f$ . Also  $fY_{n_k}$  converges to  $X_f$  a.s. and in  $L^1$ . In particular for each  $f \in T$ ,  $\lim_k fY_{n_k} = fX$ . So for  $A \in \mathcal{F}$

$$\lim_k \int_A fY_{n_k}$$

exists in  $\mathbb{R}$ . Hence  $(\int_A Y_{n_k})_k$  is weakly Cauchy. Since the Banach lattice  $E$  is weakly sequentially complete, let for every  $A \in \mathcal{F}$

$$\mu(A) = w - \lim_k \int_A Y_{n_k}.$$

Then  $\mu$  is a measure of bounded variation and it is absolutely continuous with respect to  $P$ . For each  $f \in T$  we have

$$f(\mu(A)) = \lim_k \int_A fY_{n_k} = \int_A fX.$$

Let  $A_n = \{\|X\| \leq n\}$ , then  $XI_{A_n}$  is Bochner integrable and

$$f(\mu(A_n)) = \int_{A_n} fX = f \int_{A_n} X.$$

Since  $T$  is a total set it follows that

$$\mu(A_n) = \int_{A_n} X.$$

Moreover the uniform integrability of  $(Y_{n_k})_k$  implies that

$$(1) \quad \int_{A_n} \|X\| = \|\mu\|(A_n) \leq \sup_k \int_{\Omega} Y_{n_k},$$

and since  $X$  is strongly measurable,  $P(\cup_n(\|X\| \leq n)) = 1$ . Letting  $n \rightarrow \infty$  in (1), we get that  $X$  is Bochner integrable and for each  $A \in \mathcal{F}$

$$\mu(A) = \int_A X.$$

It follows that

$$\int_A fX = f(\mu(A)) = \lim_k \int_A fY_{n_k} = \int_A X_f,$$

for each  $f \in (E^*)^+$  and  $A \in \cup \mathcal{F}$ . Hence  $fX = X_f$  a.s. and for each  $f \in (E^*)^+$ ,  $fX_n$  converges to  $fX$  a.s.. Let  $\|\cdot\|$  denote the Kadec-Klee norm equivalent to  $\|\cdot\|$ , as in Theorem 1, and let  $D \in (E^*)^+$  be a countable norming subset. Applying Proposition 2 to the sequence  $\{(fX_n, \mathcal{F}_n), n \in \mathbb{N}, f \in D\}$  it follows that  $\lim_n \|X_n\| = \|X\|$ , a.s.. Now invoking again Theorem 1 we get the strong convergence of  $X_n$  to  $X$  and the assertion follows.  $\square$

The following corollary holds.

**Corollary 1.** *Let  $E$  be a Banach lattice not containing  $c_0$  as an isomorphic copy and let  $T$  be a total subset of  $E^*$ . Let  $(X_n, \mathcal{F}_n)$  be a positive subpramart with a  $L^1$ -bounded subsequence and let  $X$  be a strongly measurable random variable. Assume that, for each  $f \in T$ ,  $fX_n$  converges to  $fX$  a.s. (the null set depends on  $f$ ). Then  $X_n$  converges to  $X$  strongly a.s..*

*Proof.* If  $E$  does not contain  $c_0$ ,  $E$  is an order continuous Banach lattice which is weakly sequentially complete [7, p. 34] and the assertion follows from Theorem 2.  $\square$

Since a submartingale is a subpramart we get

**Corollary 2.** [3, Proposition 11] *Let  $E$  be a Banach lattice not containing  $c_0$  as an isomorphic copy and let  $T$  be a total subset of  $E^*$ . Let  $(X_n, \mathcal{F}_n)$  be a  $L^1$ -bounded positive submartingale and let  $X$  be a strongly measurable random variable. Assume that, for each  $f \in T$ ,  $fX_n$  converges to  $fX$  a.s. (the null set depends on  $f$ ). Then  $X_n$  converges to  $X$  strongly a.s..*

#### 4. A CONVERGENCE THEOREM FOR SUBPRAMARTS INDEXED BY A DIRECTED SET

In this section we will consider stochastic processes indexed by a directed set. Let  $J$  be a directed set filtering to the right. Throughout this section we assume that there is an increasing cofinal sequence  $(t_n)$  in  $J$ . Let  $(\mathcal{F}_t)$  be a filtration, that is an increasing family of sub- $\sigma$ -algebras of  $\mathcal{F}$ . A filtration  $(\mathcal{F}_t)$  is said to satisfy the *Vitali condition V* if for every adapted family of sets  $(A_t)$  and for every  $\varepsilon > 0$  there exists a simple stopping time  $\tau \in \Gamma$  such that  $P(\limsup_J A_t \setminus A_\tau) < \varepsilon$ . Even in the real-valued case the Vitali condition on the filtration is necessary for the

convergence of classes of random variables. Under the condition  $V$ , the analogue of Theorem 2 holds for subpramarts indexed by directed sets.

**Theorem 3.** *Let the filtration satisfy the condition  $V$  and let  $E$  be a separable order continuous Banach lattice, which is weakly sequentially complete. Let  $(X_t, \mathcal{F}_t)$  be a  $L^1$ -bounded positive subpramart and let  $X$  be a strongly measurable random variable. Let  $T$  be a total subset of  $E^*$  and assume that, for each  $f \in T$ ,  $fX_t$  converges to  $fX$  a.s.. Then  $X_t$  converges to  $X$  strongly a.s..*

*Proof.* Let  $(t_n)$  be an increasing cofinal sequence in  $J$ . Set  $X_{t_n} = Y_n$  and  $\mathcal{F}_{t_n} = \mathcal{G}_n$ . We first show that  $(Y_n, \mathcal{G}_n)$  is a subpramart sequence. Since  $(X_t)$  is a subpramart, for every  $\varepsilon > 0$  there exists  $\tau_o \in \Gamma$  such that if  $\tau > \sigma > \tau_o$  then

$$P(\{\|(X_\sigma - E(X_\tau|\mathcal{F}_\sigma))^+\| > \varepsilon\}) \leq \varepsilon.$$

Now if  $\sigma$  is a stopping time for  $\mathcal{G}$  then  $t_\sigma$  is a stopping time for  $\mathcal{F}_t$ . Thus choose  $\sigma_o$  such that  $t_{\sigma_o} \geq \tau_o$ . Now for each  $\tau > \sigma > \sigma_o$  it follows

$$P(\{\|(Y_\sigma - E(Y_\tau|\mathcal{G}_\sigma))^+\| > \varepsilon\}) = P(\{\|(X_{t_\sigma} - E(X_{t_\tau}|\mathcal{F}_{t_\sigma}))^+\| > \varepsilon\}) \leq \varepsilon.$$

Then  $Y_n$  is a subpramart sequence. For each  $f \in T$ ,  $fY_n$  converges to  $fX$  a.s.. Therefore by Theorem 2,  $Y_n$  converges to  $X$  a.s. and also scalarly. As  $E$  is a separable Banach lattice there exists a countable norming subset  $D$  of  $(E^*)^+$  (i.e.  $\|x\| = \sup\{|x^*(x)| : x^* \in D \cap \mathcal{B}(X^*)\}$ ). Now, for each  $f \in D$ ,  $fX_t$  is a  $L^1$ -bounded real valued subpramart and since the filtration satisfies  $V$ , by [10] Theorem 4.3,  $fX_t$  converges to  $fX$  a.s.. Since  $fX_{t_n}$  converges to  $fX$ , it follows that  $fX = X_f$ . As in Theorem 1, we denote by  $\|\cdot\|$  the Kadec-Klee norm equivalent to  $\|\cdot\|$ . Applying [6] Lemma 2.3 to the sequence  $\{(fX_t, \mathcal{F}_t), t \in T, f \in D\}$  it follows that  $\lim_t \|X_t\| = \|X\|$ , a.s.. Now invoking again Theorem 1 we get the strong convergence of  $X_t$  to  $X$  and the assertion follows.  $\square$

#### REFERENCES

1. Bouzar N., *On almost sure convergence without the Radon-Nikodym property*, Acta Math. Univ. Comenianae, **LXX**(2) (2001), 167–175.
2. Davis W. J., Ghossoub N. and Lindenstrauss J., *A lattice renorming theorem and applications to vector-valued processes*, Trans. Amer. Math. Soc., **263**(2) (1981), 531–540.
3. Davis W. J., Ghossoub N., Johnson W. B., Kwapien S. and Maurey B., *Weak convergence of vector valued martingales*, Probability in Banach spaces, **6** (1990), 41–50.
4. Egghe L., *Strong convergence of positive subpramarts in Banach lattices*, Bull. Polish Acad. Sci. Math., **31**(9–12) (1984), 415–426.
5. Egghe L., *Stopping Time Techniques for Analysts and Probabilist*, London Mathematical Society, Lecture Notes, **100**, Cambridge University Press, Cambridge 1984.
6. Frangos N. K., *On convergence of vector valued pramarts and subpramarts*, Can. J. Math. **XXXVII**(2) (1985), 260–270.
7. Lindenstrauss J. and Tzafriri L., *Classical Banach Spaces II. Function Spaces*, Ergebnisse der Math. und ihrer Grensgeb. 97, Springer Verlag, Berlin 1979.
8. Marraffa V., *On almost sure convergence of amarts and martingales without the Radon-Nikodym property*, J. of Theoretical Prob. **1**(3) (1988), 255–261.
9. Marraffa V., *Convergenza di processi stocastici senza la proprietà di Radon-Nikodym*, Ph.D. Univ. of Palermo 1988.

10. Millet A. and Sucheston L., *Convergence of classes of amarts indexed by directed sets*, Can. J. Math., **32**(1) (1980), 86–125.
11. Slaby M., *Strong convergence of vector valued pramarts and subpramarts*, Probab. Math. Statist. **5**(2) (1985), 187–196.

V. Marraffa, Department of Mathematics, Via Archirafi 34, 90123 Palermo, Italy,  
*e-mail*: [marraffa@math.unipa.it](mailto:marraffa@math.unipa.it)