

**ADDITIVE STRUCTURE OF THE GROUP OF UNITS MOD p^k ,
WITH CORE AND CARRY CONCEPTS
FOR EXTENSION TO INTEGERS**

N. F. BENSCHOP

ABSTRACT. The additive structure of multiplicative semigroup $Z_{p^k} = Z(\cdot) \bmod p^k$ is analysed for prime $p > 2$. Order $(p-1)p^{k-1}$ of cyclic group G_k of units mod p^k implies product $G_k \cong A_k B_k$, with cyclic 'core' A_k of order $p-1$ so $n^p \equiv n$ for core elements, and 'extension subgroup' B_k of order p^{k-1} consisting of all units $n \equiv 1 \pmod p$, generated by $p+1$. The p -th power residues $n^p \bmod p^k$ in G_k form an order $|G_k|/p$ subgroup F_k , with $|F_k|/|A_k| = p^{k-2}$, so F_k properly contains core A_k for $k \geq 3$.

The additive structure of subgroups A_k , F_k and G_k is derived by successor function $S(n) = n+1$, and by considering the two arithmetic symmetries $C(n) = -n$ and $I(n) = n^{-1}$ as functions, with commuting $IC = CI$, where S does not commute with I nor C . The four distinct compositions SCI , CIS , CSI , ISC all have period 3 upon iteration. This yields a *triplet* structure in G_k of three inverse pairs (n_i, n_i^{-1}) with $n_i + 1 \equiv -(n_{i+1})^{-1}$ for $i = 0, 1, 2$ where $n_0 \cdot n_1 \cdot n_2 \equiv 1 \pmod p^k$, generalizing the cubic root solution $n + 1 \equiv -n^{-1} \equiv -n^2 \pmod p^k$ ($p \equiv 1 \pmod 6$).

Any solution *in core*: $(x+y)^p \equiv x+y \equiv x^p + y^p \pmod p^{k>1}$ has exponent p distributing over a sum, shown to imply the known *FLT* inequality for integers. In such equivalence mod p^k (*FLT case₁*) the three terms can be interpreted as naturals $n < p^k$, so $n^p < p^{kp}$, and the $(p-1)k$ produced carries cause *FLT* inequality. In fact, inequivalence mod p^{3k+1} is derived for the cubic roots of 1 mod p^k ($p \equiv 1 \pmod 6$).

INTRODUCTION

The commutative semigroup $Z_{p^k}(\cdot)$ of multiplication mod p^k (prime $p > 2$) has for all $k > 0$ just two idempotents: $1^2 \equiv 1$ and $0^2 \equiv 0$, and is the disjoint union of the corresponding maximal subsemigroups (Archimedean components [4], [8]). Namely the group G_k of units ($n^i \equiv 1 \pmod p^k$ for some $i > 0$) which are all relative prime to p , and maximal ideal N_k as nilpotent subsemigroup of all p^{k-1} multiples of p ($n^i \equiv 0 \pmod p^k$ for some $i > 0$). Notice that, since the analysis holds for any odd prime p , the index p in G_k and N_k is omitted for brevity of notation. Order $|G_k| = (p-1)p^{k-1}$ has two coprime factors, so that $G_k \cong A_k B_k$, with 'core' A_k

Received June 17, 2003.

2000 *Mathematics Subject Classification.* Primary 11D41, 11P99, 11A15.

Key words and phrases. Residue arithmetic, ring, group of units, multiplicative semigroup, additive structure, triplet, cubic roots of unity, carry, Hensel, Fermat, *FST*, *FLT*.

and 'extension group' B_k of orders $p - 1$ and p^{k-1} respectively. Residues of n^p form a subgroup $F_k \subset G_k$ of order $|F_k| = |G_k|/p$, to be analysed for its additive structure. Each $n \in A_k$ has $n^p \equiv n \pmod{p^k}$ denoted as FST_k , since this is related to Fermat's Small Theorem where $k = 1$.

Notation: Base p number representation is used, which is useful for computer experiments, as reported in Tables 1 and 2. This models residue arithmetic mod p^k by considering only the k less significant digits, and ignoring the more significant digits. Congruence class $[n] \pmod{p^k}$ is represented by natural number $n < p^k$, encoded by k digits (base p). Class $[n]$ consists of all integers with the same least significant k digits as n . As usual, concatenation of operands indicates multiplication.

Define the 0 -extension of residue $n \pmod{p^k}$ as the natural number $n < p^k$ with the same k -digit representation (base p), and all more significant digits (at p^m , $m \geq k$) set to 0.

Signed residue $-n$ is only a convenient notation for the complement $p^k - n$ of n , which are both positive. $C[n]$ is a cyclic group of order n , such as $Z_{p^k}(+) \cong C[p^k]$. Units mod p form a cyclic group $G_1 = C[p - 1]$, and G_k of order $(p - 1)p^{k-1}$ is also cyclic for $k > 1$ [1]. Finite *semigroup structure* is applied, and *digit analysis* of prime-base residue arithmetic, to study the combination of $(+)$ and $(\cdot) \pmod{p^k}$, especially the additive properties of multiplicative subgroups of ring $Z_{p^k}(+, \cdot)$.

Elementary residue arithmetic, cyclic groups, and (associative) function composition will be used, starting at the known cyclic (one generator) nature [1] of the group G_k of units mod p^k . The direct product structure of G_k (Lemma 1.1 and Corollary 1.2) on the p^{k-2} extensions of $n^p \pmod{p^2}$ to cover all p -th power residues mod p^k for $k > 2$ are known, but they are derived for completeness. Results beyond Section 1 are believed to be new.

The two symmetries of residue arithmetic mod p^k , defined as automorphisms of order 2, are complement $-n$ under $(+)$ and inverse n^{-1} under (\cdot) . Their role as functions $C(n) = -n$ and $I(n) = n^{-1}$, in the *triplet* additive structure of $Z(\cdot) \pmod{p^k}$ (Lemma 3.1 and Theorem 3.1) is essential.

Symbols	and Definitions (odd prime p)
$Z_{p^k}(\cdot)$	multiplicative semigroup mod p^k (k -digit arithmetic base p)
$C[m]$	cyclic group of order m : e.g. $Z_{p^k}(+) \cong C[p^k]$
$x \in Z_{p^k}(\cdot)$	unique product $x = g^i p^{k-j} \pmod{p^k}$ ($g^i \in G_j$ coprime to p)
0-extension X	of residue $x \pmod{p^k}$: the smallest non-negative integer $X \equiv x \pmod{p^k}$
(finite) extension U	of $x \pmod{p^k}$: any integer $U \equiv x \pmod{p^k}$
$G_k \equiv A_k \cdot B_k$	group of units $n: n^i \equiv 1 \pmod{p^k}$ (some $i > 0$), $ G_k \equiv (p - 1)p^{k-1}$
A_k	core of G_k , $ A_k = p - 1$ ($n^p \equiv n \pmod{p^k}$ for $n \in A_k$)
$B_k \equiv (p + 1)^*$	extension group of all $n \equiv 1 \pmod{p}$, $ B_k = p^{k-1}$
F_k	subgroup of all p -th power residues in G_k , $ F_k = G_k /p$
$A_k \subset F_k \subset G_k$	proper inclusions only for $k \geq 3$ ($A_2 \equiv F_2 \subset G_2$)

Symbols	and Definitions (odd prime p)
$d(n)$	core increment $A(n+1) - A(n)$ of core func'n $A(n) \equiv n^q$, $q = B_k $
FST_k	core A_k ($p-1$ residues) extends FST ($n^p \equiv n \pmod p$) to $\pmod{p^{k>1}}$
solution in core	$x^p + y^p \equiv z^p \pmod{p^k}$ with x, y, z in core A_k .
period of $n \in G_k$	order $ n^* $ of subgroup generated by n in $G_k(\cdot)$
normation	divide $x^p + y^p \equiv z^p \pmod{p^k}$ by one term (in F_k) to yield one term ± 1
complement $-n$	unique in $Z_{p^k}(+)$: $-n + n \equiv 0 \pmod{p^k}$
inverse n^{-1}	unique in $G_k(\cdot)$: $n^{-1} \cdot n \equiv 1 \pmod{p^k}$
1-complement pair	pair $\{m, n\}$ in $Z_{p^k}(+)$: $m + n \equiv -1 \pmod{p^k}$
inverse-pair	pair $\{a, a^{-1}\}$ of inverses in G_k
triplet	3 inv. pairs: $a + b^{-1} \equiv b + c^{-1} \equiv c + a^{-1} \equiv -1$, ($abc \equiv 1 \pmod{p^k}$)
triplet ^{p}	a triplet of p -th power residues in subgroup F_k
symmetry mod p^k	$-n$ and n^{-1} : order 2 automorphism of $Z_{p^k}(+)$ resp. $G_k(\cdot)$
EDS property	Exponent Distributes over a Sum: $(a + b)^p \equiv a^p + b^p \pmod{p^k}$

1. STRUCTURE OF THE GROUP G_k OF UNITS

Lemma 1.1. $G_k \cong A'_k \times B'_k \cong C[p-1] \cdot C[p^{k-1}]$ and $Z(\cdot) \pmod{p^k}$ has a sub-semigroup isomorphic to $Z(\cdot) \pmod p$.

Proof. Cyclic group G_k of units n ($n^i \equiv 1$ for some $i > 0$) has order $(p-1)p^{k-1}$, namely p^k minus p^{k-1} multiples of p . Then $G_k = A'_k \times B'_k$, the direct product of two relative prime cycles, with corresponding subgroups A_k and B_k , so that $G_k \cong A_k B_k$ where:

extension group $B_k = C[p^{k-1}]$ consists of all p^{k-1} residues mod p^k that are $1 \pmod p$, and

core $A_k = C[p-1]$, so $Z_{p^k}(\cdot)$ contains sub-semigroup $A_k \cup 0 \cong Z_p(\cdot)$ □

Core A_k , as $p-1$ cycle mod p^k , is Fermat's Small Theorem $n^p \equiv n \pmod p$ extended to $k > 1$ for p residues (including 0), to be denoted as FST_k .

Recall that $n^{p-1} \equiv 1 \pmod p$ for $n \not\equiv 0 \pmod p$ (FST), then Lemma 1.1 implies:

Corollary 1.1. With $|B| = p^{k-1} = q$ and $|A| = p-1$, core $A_k = \{n^q\} \pmod{p^k}$ ($n = 1, \dots, p-1$) extends FST for $k > 1$, and $B_k = \{n^{p-1}\} \pmod{p^k}$ consists of all p^{k-1} residues $1 \pmod p$ in G_k .

Subgroup $F_k \equiv \{n^p\} \pmod{p^k}$ of all p -th power residues in G_k , with $F_k \supseteq A_k$ (only $F_2 \equiv A_2$) and order $|F_k| = |G_k|/p = (p-1)p^{k-2}$, consists of all p^{k-2} extensions mod p^k of the $p-1$ p -th power residues in G_2 , which has order $(p-1)p$. Consequently:

Corollary 1.2. Each extension of $n^p \pmod{p^2}$ (in F_2) is a p -th power residue (in F_k).

Core generation: The $p - 1$ residues $n^q \bmod p^k$ ($q = p^{k-1}$) define core A_k for $0 < n < p$. Cores A_k for successive k are produced as the p -th power of each $n_0 < p$ recursively

$$(n_0)^p \equiv n_1, (n_1)^p \equiv n_2, (n_2)^p \equiv n_3, \dots$$

where n_i has $i + 1$ digits (base p). In more detail:

Lemma 1.2. *For non-negative digits $a_i < p$ the $p-1$ naturals $a_0 < p$ define core*

$$A_k(a_0) \equiv (a_0)^{p^{k-1}} \equiv a_0 + \sum_{i=1}^{k-1} a_i p^i \pmod{p^k},$$

and

$$A_{k+1}(a_0) \equiv [A_k(a_0)]^p \pmod{p^{k+1}}.$$

Proof. Let $a = a_0 + mp < p^2$ be in core A_2 , so $a^p \equiv a \pmod{p^2}$. Then

$$a^p = (mp + a_0)^p \equiv a_0^{p-1} mp^2 + a_0^p \equiv mp^2 + a_0^p \pmod{p^3},$$

by *FST*. Core digit a_1 of weight p is not found in this way as function of a_0 , requiring actual computation, except for $a \equiv p \pm 1$ as in (1) and (1'). It depends on the *carries* produced in computing the p -th power of a_0 . Similarly, the *next* more significant digit in core $A_{k+1}(n)$ is found by computing, with $k+1$ digit precision, the p -th power a^p of 0-extension $a < p^k$ in core A_k , leaving core A_k fixed, because $a^p \equiv a \pmod{p^k}$. \square

Notice $(p^2 \pm 1)^p \equiv p^3 \pm 1 \pmod{p^5}$, and $(p + 1)^p \equiv p^2 + 1 \pmod{p^3}$ yields by induction on m :

$$(1) \quad (p + 1)^{p^m} \equiv p^{m+1} + 1 \pmod{p^{m+2}}$$

$$(1') \quad (p - 1)^{p^m} \equiv p^{m+1} - 1 \pmod{p^{m+2}}$$

Lemma 1.3. *Extension group B_k is generated by $p + 1 \pmod{p^k}$, with $|B_k| = p^{k-1}$, and each subgroup $S \subseteq B_k$, $|S| = |B_k|/p^s$ has sum*

$$\sum S \equiv |S| \pmod{p^k} \not\equiv 0 \pmod{p^k}.$$

Proof. For the smallest x with $(p + 1)^x \equiv 1 \pmod{p^k}$, the *period* of $p + 1$, (1) implies $m + 1 = k$. So $m = k - 1$, thus period p^{k-1} . No smaller x generates $1 \pmod{p^k}$ since $|B_k|$ has only divisors p^s .

B_k consists of all p^{k-1} residues which are $1 \pmod{p}$. The order of each subgroup $S \subset B_k$ must divide $|B_k|$, so that $|S| = |B_k|/p^s$ ($0 \leq s < k$) and $S = \{1 + m \cdot p^{s+1}\}$ ($m = 0, \dots, |S| - 1$). Then $\sum S = |S| + p^{s+1} \cdot |S|(|S| - 1)/2 \pmod{p^k}$, where $p^{s+1} \cdot |S| = p \cdot |B_k| = p^k$, so that $\sum S = |S| = p^{k-1-s} \pmod{p^k}$. Hence no subgroup of B_k sums to $0 \pmod{p^k}$. \square

Corollary 1.3. *For core $A_k \equiv g^*$, each unit $n \in G_k \equiv A_k B_k$ has the form:*

$$n \equiv g^i (p + 1)^j \pmod{p^k}$$

for a unique pair of non-negative exponents $i < |A_k|$ and $j < |B_k|$.

Pair (i, j) are the exponents in the core- and extension- component of unit n . In case $p = 2$, the most interesting prime for computer engineering purposes, the next binary number representation is readily verified [3], [7]:

Lemma 1.4. *For $p = 2$: $p + 1 = 3$ is a semi-primitive root of 1 mod 2^k for $k > 2$.*

In other words, for base $p = 2$ and precision $k > 2$: each odd residue mod 2^k is a unique signed power of 3. Hence an efficient k -bit binary number code is

$$n = \pm 3^i \cdot 2^j \pmod{2^k},$$

for all integers $0 \leq n < 2^k$, with unique non-negative index pair $i < 2^{k-2}$ and $j \leq k$.

Clearly, this allows a dual-base (2, 3) binary logarithmic code, which reduces multiplication to addition of the two indices, and XOR (add mod 2) of the involved signs (see US-patent [7]).

Theorem 1.1. *Each subgroup $S \supset 1$ of core A_k sums to 0 mod p^k ($k > 0$).*

Proof. For even $|S|$: -1 in S implies pairwise zero-sums. In general: $c \cdot S = S$ for all c in S , and $c \sum S = \sum S$, so $S \cdot x = x$, writing x for $\sum S$. Now for any g in G_k : $|S \cdot g| = |S|$ so that $|S \cdot x|=1$ implies x not in G_k , hence $x = g \cdot p^e$ for some g in G_k and $0 < e < k$ or $x = 0$ ($e = k$). Then $S \cdot x = S(g \cdot p^e) = (S \cdot g)p^e$ with $|S \cdot g| = |S|$ if $e < k$. So $|S \cdot x|=1$ yields $e=k$ and $x = \sum S=0$. \square

Consider the normation of an additive equivalence $a + b \equiv c \pmod{p^k}$ in units group G_k , by multiplying all terms with the inverse of one of these terms, to yield rhs -1 as right hand side:

(2) 1-complement form: $a + b \equiv -1 \pmod{p^k}$ in G_k ,
 (digitwise sum $p - 1$, no carry).

For instance the well known p -th power residue equivalence: $x^p + y^p \equiv z^p$ in F_k yields:

(2') normal form: $a^p + b^p \equiv -1 \pmod{p^k}$ in G_k ,
 with a special case in core A_k , considered next.

2. THE CUBIC ROOT SOLUTION IN CORE, AND CORE SYMMETRIES

Lemma 2.1. *Cubic roots $a^3 \equiv 1 \pmod{p^k}$ ($p \equiv 1 \pmod{6}$, $k > 1$) are p -th power residues in core A_k , and $a + a^{-1} \equiv -1 \pmod{p^k}$ ($a \neq -1$) has no corresponding integers as p -th powers $< p^{kp}$.*

Proof. If $p \equiv 1 \pmod{6}$ then 3 divides $p - 1$, implying a core subgroup $S = \{a, a^2, 1\}$ of three p -th powers: the cubic roots $a^3 \equiv 1$ in G_k , with sum 0 mod p^k (Theorem 1.1). Now $a^3 - 1 = (a - 1)(a^2 + a + 1)$, so for $a \neq 1$: $a^2 + a + 1 \equiv 0$, hence $a + a^{-1} \equiv -1$ solves the normed (2'), being a root-pair of inverses with $a^2 \equiv a^{-1}$. Subgroup $S \subset A_k$ consists of p -th power residues with $n^p \equiv n \pmod{p^k}$.




Figure 1. Core $A_2 \pmod{7^2}$ (6-cycle), Cubic roots $\{42, 24, 01\}$ (3-cycle) in core.

Write b for a^{-1} , then $a^p + b^p \equiv -1$ and $a + b \equiv -1$, hence $a^p + b^p \equiv (a + b)^p \pmod{p^k}$. The “exponent p distributes over a sum” (*EDS*) property implies $A^p + B^p < (A + B)^p$ for the corresponding 0-extensions $A, B, A + B$ of residues $a, b, a + b \pmod{p^k}$. \square

1. Successive powers g^i of generator g of G_k produce $|G_k|$ points (k -digit residues) counter clockwise on a unit circle (Figures 1, 2). Inverse pairs (a, a^{-1}) are connected *vertically*, complements $(a, -a)$ *diagonally*, and pairs $(a, -a^{-1})$ *horizontally*, representing functions I, C and $IC = CI$ respectively (Theorem 3.1).
2. Scaling any equation, such as $a + 1 \equiv -b^{-1}$, by a factor $s \equiv g^i \in G_k \equiv g^*$, yields $s(a + 1) \equiv -s/b \pmod{p^k}$, represented by a rotation counter clockwise over i positions.

2.1. Another derivation of the cubic roots of 1 mod p^k

The cubic root solution was derived, for 3 dividing $p - 1$, via subgroup $S \subset A_k$ of order 3 (Theorem 1.1). For completeness a derivation using elementary arithmetic follows.

Notice $a + b \equiv -1$ to yield $a^2 + b^2 \equiv (a + b)^2 - 2ab \equiv 1 - 2ab$, and:

$$a^3 + b^3 \equiv (a + b)^3 - 3(a + b)ab \equiv -1 + 3ab.$$

The combined sum is $ab - 1$:

$$\sum_{i=1}^3 (a^i + b^i) \equiv \sum_{i=1}^3 a^i + \sum_{i=1}^3 b^i \equiv ab - 1 \pmod{p^k}.$$

Find a, b for $ab \equiv 1 \pmod{p^k}$. Now

$$n^2 + n + 1 = (n^3 - 1)/(n - 1) = 0 \quad \text{for} \quad n^3 \equiv 1 \quad (n \neq 1),$$

hence $ab \equiv 1 \pmod{p^k}$, ($k > 0$) if $a^3 \equiv b^3 \equiv 1 \pmod{p^k}$, with 3 dividing $p - 1$ ($p \equiv 1 \pmod{6}$). Cubic roots $a^3 \equiv 1 \pmod{p^k}$ exist for any prime $p \equiv 1 \pmod{6}$ at any precision $k > 0$.

In the next section other solutions of $\sum_{i=1}^3 a^i + \sum_{i=1}^3 b^i \equiv 0 \pmod{p^k}$ will be shown, depending not only on p but also on k , with $ab \equiv 1 \pmod{p^2}$ but $ab \not\equiv 1 \pmod{p^3}$, for some primes $p \geq 59$.

2.2. Core increment symmetry mod p^{2k+1} and asymmetry mod p^{3k+1}

Consider:

- core function* $A_k(n) = n^q$ ($q = |B_k| = p^{k-1}$) as natural monomial,
- core increment* $d_k(n) = A_k(n + 1) - A_k(n) = (n + 1)^q - n^q$ (even degree $q - 1$),
- natural core* $C_k(n) < p^k$ with $A_k(n) \equiv C_k(n) \pmod{p^k}$,
- integer core increment* $D_{k+1}(n) = [C_k(n + 1)]^p - [C_k(n)]^p$, with absolute value less than p^{kp} .

Recall: for natural $n < p$ the p -th power residues $[A_k(n)]^p \pmod{p^{k+1}}$ form core A_{k+1} (Lemma 1.2). For any core element $a \in C_k$: $a^{p-1} \equiv 1 \pmod{p^k}$. By FST: $C_k(n) \equiv n \pmod{p}$, so $D_k(n) \equiv 1 \pmod{p}$, and $D_k(n)$ is called *core increment*, although in general $D_k(n) \not\equiv 1 \pmod{p^k}$ for $k > 2$. Core naturals $C_k(n) < p^k$ are considered in order to study natural p -th power sums.

For example consider $p = 7$ (Figure 1). The cubic roots in core A_2 are $\{42, 24, 01\} \pmod{7^2}$, with 7-th powers $\{642, 024, 001\}$ in core A_3 . In full 14 digits (base 7):

$$42^7 + 24^7 = 0\ 14\ 24\ 06\ 25\ 00\ 66\ 6 \quad (k=2) \quad \text{versus} \quad 66^7 = 6\ 02\ 62\ 04\ 64\ 00\ 66\ 6$$

which are equivalent mod $7^{2k+1} = 7^5$, but differ mod 7^6 hence also mod $7^{3 \cdot 2 + 1} = 7^7$. Cubic roots $\{3642, 3024\}$ in core A_4 , as 7-th powers of cubic roots in A_3 ($k=3$), have increment $1 \pmod{7^7}$ with increment symmetry mod $7^{2k+1} = 7^7$, and asymmetry mod $p^{3k+1} = 7^{10}$. See also Table 1. This core- and carry effect is generalized for integers as follows.

Lemma 2.2 (Core increment symmetry and asymmetry). *For $q = |B_k| = p^{k-1}$ ($k \geq 1$) and natural $m, n < p$:*

- (a) *Core residues $A_k(n) \equiv n^q \pmod{p^k}$ and increments $d_k(n) \equiv A_k(n + 1) - A_k(n) \pmod{p^k}$ have period p in n .*
- (b) *If $m + n = p$ then $A_k(p - n) \equiv A_k(-n) \equiv -A_k(n) \pmod{p^k}$ (odd symm.).*
- (c) *If $m + n = p - 1$ then $D_{k+1}(m) \equiv D_{k+1}(n) \pmod{p^{2k+1}}$ (even symm.).*
- (d) *If $m + n = p - 1$ and natural cubic roots $C_k(m) + C_k(n) = p^k - 1$ then $D_{k+1}(m) \not\equiv D_{k+1}(n) \pmod{p^{3k+1}}$ (asymmetry)*

Proof. (a) Core function $A_k(n) \equiv n^q \pmod{p^k}$ ($q = p^{k-1}$, $n \not\equiv 0 \pmod{p}$) has just $p - 1$ distinct residues with $(n^q)^p \equiv n^q \pmod{p^k}$, and $A_k(n) \equiv n \pmod{p}$ (FST). Include non-core $A_k(0) \equiv 0$ then $A_k(n) \pmod{p^k}$ is periodic in n with period p , so $A_k(n + p) \equiv A_k(n) \pmod{p^k}$. Hence difference $d_k(n) \pmod{p^k}$ of two functions of period p also has period p .

n	core C_k	core C_{k+1}==(C_k)^p	Core_incr.	p=7 (base 7) mod p^3
	C_1	C_2		
1.	0 0 0 0 0 0 0 1	0 0 0 0 0 0 0 1	0 0 0 0 0 2 4 1	
2.	0 0 0 0 0 0 0 2	0 0 0 0 0 2 4 2	0 0 0 0 6 0 0 1	∧
3.	0 0 0 0 0 0 0 3	0 0 0 0 6 2 4 3	0 0 0 5 6 2 5 1	sym
4.	0 0 0 0 0 0 0 4	0 0 0 6 5 5 2 4	0 0 3 4 5 0 0 1	∨
5.	0 0 0 0 0 0 0 5	0 0 4 4 3 5 2 5	0 1 5 0 0 2 4 1	
6.	0 0 0 0 0 0 0 6	0 2 2 4 4 0 6 6	6 4 4 2 2 6 0 1	
	C_2	C_3		mod p^5
1.	0 0 0 0 0 0 0 1	0 0 0 0 0 0 0 1	4 6 6 3 4 6 4 1	
2.	0 0 0 0 0 0 0 2	4 6 6 3 4 6 4 2	5 4 3 0 0 0 0 1	∧
3.	0 0 0 0 0 0 0 3	3 4 2 3 4 6 4 3	4 5 2 6 5 0 5 1	sym
4.	0 0 0 0 0 0 0 4	1 2 5 3 3 0 2 4	6 0 0 0 0 0 0 1	∨
5.	0 0 0 0 0 0 0 5	0 2 5 3 3 0 2 5	4 3 5 3 4 6 4 1	
6.	0 0 0 0 0 0 0 6	4 6 4 0 0 6 6 6	2 0 2 6 6 0 0 1	
	C_3	C_4		mod p^7
1.	0 0 0 0 0 0 0 1	0 0 0 0 0 0 0 1	6 4 1 4 3 6 4 1	
2.	0 0 0 0 0 0 0 2	6 4 1 4 3 6 4 2	136 0 0 0 0 0 0 1	∧
3.	0 0 0 0 0 0 0 3	5 4 1 4 3 6 4 3	2 5 3 5 6 0 5 1	sym
4.	0 0 0 0 0 0 0 4	1 2 5 3 3 0 2 4	666 0 0 0 0 0 0 1	∨
5.	0 0 0 0 0 0 0 5	0 2 5 3 3 0 2 5	3 4 1 4 3 6 4 1	
6.	0 0 0 0 0 0 0 6	4 0 0 0 6 6 6 6	2 6 6 6 0 0 0 1	

Table 1. Cores $C_1..C_3$, increment symmetry mod $p^{[2k+1]}$ of $C_2..C_4$. For cubic roots of 1 mod p^k : asymmetry mod $p^{[3k+1]}$ in $C_2..C_4$.

(b) $(-n)^q = -n^q$, odd $q = p^{k-1}$, yields *odd symmetry*

$$A_k(p - n) \equiv A_k(-n) \equiv -A_k(n) \pmod{p^k}$$

(c) Difference polynomial $d_k(n)$ has leading term $q n^{q-1}$. Even degree $q - 1$ results in *even symmetry*

$$d_k(n - 1) = n^q - (n - 1)^q = -(-n)^q + (-n + 1)^q = d_k(-n).$$

Now $C_k(n) = p^k - C_k(p - n) < p^k$, hence for $m + n = p - 1$, $C_k(m + 1) = p^k - C_k(n)$, so

$$D_{k+1}(m) = [p^k - C_k(n)]^p - [C_k(m)]^p \quad \text{and} \quad D_{k+1}(n) = [p^k - C_k(m)]^p - [C_k(n)]^p.$$

Briefly denote naturals $C_k(m) = a$, $C_k(n) = b$, and $h = (p - 1)/2$ then

$$\begin{aligned}
 D_{k+1}(m) - D_{k+1}(n) &= [(p^k - b)^p + b^p] - [(p^k - a)^p + a^p] \\
 (*) \quad &\equiv -h [b^{p-2} - a^{p-2}] p^{2k+1} + [b^{p-1} - a^{p-1}] p^{k+1} \pmod{p^{3k+1}} \\
 &\equiv 0 \pmod{p^{2k+1}},
 \end{aligned}$$

because by FST: $a^{p-1} \equiv b^{p-1} \equiv 1 \pmod{p^k}$.

(d) Carry difference $(b^{p-1} - a^{p-1})/p^k \not\equiv h(b^{p-2} - a^{p-2}) \pmod{p^k}$ is required, to avoid cancellation in (*). It suffices to show this for $k = 1$ and 0-extensions $1 < a, b < p$ of cubic roots of 1 mod p . Using $b \equiv a^2 \equiv a^{-1}$, $b^{p-2} - a^{p-2} \equiv -(b - a) \pmod{p}$, and $h = (p - 1)/2 \equiv -1/2 \pmod{p}$ the *carry difference* must satisfy (cd)

$$(cd) \quad \frac{(b^{p-1} - a^{p-1})}{p} \not\equiv \frac{(b - a)}{2} \pmod{p}.$$

Let $a^3 \equiv cp + 1 \pmod{p^2}$ with some carry c , then for $m > 0$: $a^{3m} \equiv mcp + 1 \pmod{p^2}$. So $a^{p-1} \equiv [(p-1)/3]cp + 1 \pmod{p^2}$, and similarly for cubic root power b^3 . In other words, in extension group $B_2 \equiv \{xp+1\} \equiv (p+1)^x \pmod{p^2}$ the coefficient of p is proportional to the exponent. For a^{p-1} versus a^3 the ratio is $(p-1)/3$. However in (cd), adapted for third powers a^3, b^3 it is $(p-1)/(3/2) = 2(p-1)/3$, hence the (cd) inequivalence holds.

So for the cubic roots of $1 \pmod{p^k}$, with $a + b = C_k(m) + C_k(n) = p^k - 1$ core increment has asymmetry

$$D_{k+1}(m) \not\equiv D_{k+1}(n) \pmod{p^{3k+1}}. \quad \square$$

Corollary 2.1. *Let prime $p \equiv 1 \pmod{6}$, and any precision $k > 0$. For $x^3 \equiv y^3 \equiv 1 \pmod{p^k}$ (cubic roots $x, y \not\equiv 1$) 0-extensions $X, Y < p^k$ of x, y have $X^p, Y^p \pmod{p^{k+1}}$ in core A_{k+1} with $X^p + Y^p \equiv -1 \pmod{p^{k+1}}$ and $X^p + Y^p \not\equiv (p^k - 1)^p \pmod{p^{3k+1}}$.*

3. SYMMETRIES AS FUNCTIONS YIELD 'TRIPLETS'

Any solution of (2'): $a^p + b^p = -1 \pmod{p^k}$ has at least one term (-1) in core, and at most all three terms in core A_k . To characterize such solution by the number of terms in core A_k , quadratic analysis $(\pmod{p^3})$ is essential since proper inclusion $A_k \subset F_k$ requires $k \geq 3$. The cubic root solution, involving one inverse pair (Lemma 2.1) has all three terms in core A_k ($k > 1$). However, a computer search (Table 2) reveals another type of solution of (2') $\pmod{p^2}$ for some $p \geq 59$, namely three inverse pairs of p -th power residues, denoted triplet ^{p} , in core A_2 .

Lemma 3.1. *A triplet ^{p} of three inverse-pairs of p -th power residues in F_k satisfies*

- (3a) $a + b^{-1} \equiv -1 \pmod{p^k}$
- (3b) $b + c^{-1} \equiv -1 \pmod{p^k}$
- (3c) $c + a^{-1} \equiv -1 \pmod{p^k}$ with $abc \equiv 1 \pmod{p^k}$.

Proof. Multiplying by b, c, a resp. maps (3a) to (3b) if $ab \equiv c^{-1}$, and (3b) to (3c) if $bc \equiv a^{-1}$, and (3c) to (3a) if $ac \equiv b^{-1}$. All three conditions imply $abc \equiv 1 \pmod{p^k}$. □

Table 2 shows all normed solutions of (2') $\pmod{p^2}$ for $p < 200$, with a triplet ^{p} at $p = 59, 79, 83, 179, 193$. The cubic roots, indicated by C_3 , occur only at $p \equiv 1 \pmod{6}$, while a triplet ^{p} can occur for either prime type $\pm 1 \pmod{6}$. More than one triplet ^{p} can occur per prime: two at $p = 59$, three at 1093 (dec) = [1111111] base 3 (one of the two known Wieferich primes [9], [6], and four at 36847, each the first occurrence of such multiple triplet ^{p}). There are primes for which both root forms occur, e.g. $p = 79$ has a cubic root solution as well as a triplet ^{p} .

Such loop of inverse-pairs in residue ring $Z \pmod{p^k}$ cannot have a length beyond 3, seen as follows. Consider the successor $S(n) = n+1$ and the two symmetries: complement $C(n) = -n$ and inverse $I(n) = n^{-1}$, as functions which compose associatively.

```

Find a+b = -1 mod p^2 (in A=F < G): Core A={n^p=n}, F={n^p}=A if k=2.
G(p^2)=g*, log-code: log(a)=i, log(b)=j; a.b=1 --> i+j=0 (mod p-1)
TRIPLET^p: a+ 1/b= b+ 1/c= c+ 1/a=-1; a.b.c=1; (p= 59 79 83 179 193 ...
~~~~~
Root-Pair: a+ 1/a=-1; a^3=1 ('C3') <--> p=6m+1 (Cubic rootpair of 1)
~~~~~
p:6m+1 g=generator; p < 2000: two triplets at p= 59, 701, 1811
5:- 2 three triplets at p= 1093
7:+ 3 C3 11:- 2
13:+ 2 C3 17:- 3
19:+ 2 C3 23:- 5 29:- 2
31:+ 3 C3
37:+ 2 C3 41:- 6
43:+ 3 C3 47:- 5
53:- 2 log lin mod p^2
59:- 2 -----
-2,-25( 40 15, 18 43) 25, 23( 35 11, 23 47) -23, 2( 53 54, 5 4)
27, 19( 18 44, 40 14) -19, 8( 13 38, 45 20) -8,-27( 5 3, 53 55)
61:+ 2 C3
67:+ 2 C3 71:- 7
73:+ 5 C3
79:+ 3 C3
30, 20( 40 46, 38 32) -20, 10( 36 42, 42 36) -10,-30( 77 11, 1 67)
83:- 2
21, 3( 9 74, 73 8) -3, 18( 54 52, 28 30) -18,-21( 13 36, 69 46)
89:- 3
97:+ 5 C3 101:- 2
103:+ 5 C3 107:- 2
109:+ 6 C3 113:- 3
127:+ 3 C3 131:- 2 137:- 3
139:+ 2 C3 149:- 2
151:+ 6 C3
157:+ 5 C3
163:+ 2 C3 167:- 5 173:- 2
179:- 2
19, 1( 78 176,100 2) -1, 18( 64 90,114 88) -18,-19( 88 59, 90 119)
181:+ 2 C3 191:- 19
193:+ 5 C3
-81, 58( 64 106,128 86) -58, 53( 4 101,188 91) -53, 81(188 70, 4 122)
197:- 2
199:+ 3 C3

```

Table 2. FLT₂ root: inv-pair (C3) & triplet^p (for p < 200).

Theorem 3.1 (Two basic solution types). *Each normed solution of (2') is (an extension of) a triplet^p or an inverse-pair.*

Proof. Assume that r equations $1 - n_i^{-1} \equiv n_{i+1}$ form a loop of length r (indices mod r). Consider function $ICS(n) \equiv 1 - n^{-1}$, composed of the three elementary functions: Inverse, Complement and Successor, in that sequence. Let $E(n) \equiv n$ be the identity function, and $n \neq 0, 1, -1$ to prevent division by zero, then under function composition the third iteration $[ICS]_3 = E$, since $[ICS]_2(n) \equiv -1/(n-1) \rightarrow [ICS]_3(n) \equiv n$ (repeat substituting $1 - n^{-1}$ for n). Since C and I commute, $IC=CI$, the $3! = 6$ permutations of $\{I, C, S\}$ yield only four distinct dual-folded-successor “dfs” functions:

$$\begin{aligned}
 ICS(n) &= 1 - n^{-1}, & SCI(n) &= -(1 + n)^{-1}, \\
 CSI(n) &= (1 - n)^{-1}, & ISC(n) &= -(1 + n^{-1}).
 \end{aligned}$$

Starting at $n_0 \in G_k$ six triplet residues are generated upon iteration of e.g. $SCI(n)$: $n_{i+1} \equiv -(n_i + 1)^{-1}$ (indices mod 3), or another *dfs* function to prevent a non-invertable residue. Less than 6 residues are involved if 3 or 4 divides $p - 1$

If $3|(p - 1)$ then a cubic root of 1 ($a^3 \equiv 1$, $a \not\equiv 1$) generates just 3 residues: $a + 1 \equiv -a^{-1}$ – together with its complement this yields a subgroup $(a + 1)^* \equiv C_6$ (Figure 1, $p = 7$).

If 4 divides $p - 1$ then an x on the vertical axis has $x^2 \equiv -1$ so $x \equiv -x^{-1}$, so the three inverse pairs involve then only five residues (Figure 2, $p = 5$).

1. It is no coincidence that the period 3 of each *dfs* composition exceeds by one the number of symmetries of finite ring $Z(+, \cdot) \pmod{p^k}$.
2. No duplet occurs: multiply $a + b^{-1} \equiv -1$, $b + a^{-1} \equiv -1$ by b resp. a . Then $ab + 1 \equiv -b$ and $ab + 1 \equiv -a$, so that $-b \equiv -a$ and $a \equiv b$.
3. Basic triplet mod 3^2 : $G_2 \equiv 2^* \equiv \{2, 4, 8, 7, 5, 1\}$ is a 6-cycle of residues mod 9. Iterate $SCI(1)^* : -(1 + 1)^{-1} \equiv 4$, $-(4 + 1)^{-1} \equiv 7$, $-(7 + 1)^{-1} \equiv 1$, and $abc \equiv 1 \cdot 4 \cdot 7 \equiv 1 \pmod{9}$.

3.2. The *EDS* argument extended to non-core triplets

The *EDS* argument for the cubic root solution *CR* (Lemma 2.1), with all three terms in core, also holds for any triplet ^{p} mod p^2 . Because $A_2 \equiv F_2 \pmod{p^2}$, so all three terms are in core for some linear transform (5). Then for each of the three equivalences (3a) – (3c) holds the *EDS* property: $(x + y)^p \equiv x^p + y^p$, and thus no finite (equality preserving) extension exists, yielding inequality for the corresponding integers for all $k > 1$, to be shown next. A cubic root solution is a special triplet ^{p} for $p \equiv 1 \pmod{6}$, with $a \equiv b \equiv c$ in (3a) – (3c).

Denote the $p - 1$ core elements as residues of integer function $A_k(n) = n^{|B_k|}$ ($0 < n < p$), then for any $k > 2$ consider core increment form:

$$(4) \quad A_k(n + 1) - A_k(n) \equiv (r_n)^p \pmod{p^k}, \quad \text{where } (r_n)^p \equiv 1 \pmod{p^2}.$$

This triplet ^{p} rootform with two terms in core, and $(r_n)^p \not\equiv 1 \pmod{p^3}$, is useful for the additive analysis of subgroup F_k of p -th power residues mod p^k , in essence: the known Fermat's Last Theorem *FLT* case₁ for residues coprime to p , discussed in the next section.

Any assumed *FLT* case₁ solution (5) for integers less than p^{kp} can be transformed to (4), in two equality preserving steps. Namely first a multiplicative scaling by an integer p -th power factor s^p that is $1 \pmod{p^2}$ (so $s \equiv 1 \pmod{p}$), to yield as one lefthand term the core residue $A_k(n + 1) \pmod{p^k}$. And secondly an additive translation by integer term t which is $0 \pmod{p^2}$ applied to both sides, resulting in the other lefthand term $-A_k(n) \pmod{p^k}$, while preserving integer equality. Assuming, without loss, the normed form with $z^p \equiv 1 \pmod{p^2}$, such linear transformation (s, t) yields:

$$(5) \quad x^p + y^p = z^p \longleftrightarrow (sx)^p + (sy)^p + t = (sz)^p + t \quad [\text{integers}],$$

with $s^p \equiv A_k(n + 1)/x^p$, $(sy)^p + t \equiv -A_k(n) \pmod{p^k}$, so:

$$(5') \quad A_k(n + 1) - A_k(n) \equiv (sz)^p + t \pmod{p^k}, \quad \text{equivalent to } 1 \pmod{p^2}.$$

With $s^p \equiv z^p \equiv 1$, $t \equiv 0 \pmod{p^2}$ this yields an equivalence which is $1 \pmod{p^2}$, hence a p -th power residue, and (5') has two of the three terms in core, for $k > 2$. All three terms of a triplet ^{p} mod p^2 are in core (Corollary 1.2). In core increment form (4) for $k > 2$ this holds apparently only if the righthand side $(r_n)^p \equiv 1 \pmod{p^k}$, yielding:

Corollary 3.2 (For precision $k > 2$ (base p)). *Core increment form (4) with all three terms in core A_k is the cubic root solution, and an FLT equivalence mod p^k with three terms in core is a (scaled) cubic root solution.*

Lemma 3.2. *The p -th powers of 0-extended terms of a triplet ^{p} (mod p^k) yield integer inequality.*

Proof. In a triplet ^{p} for some odd prime p the core increment form (4) holds for three distinct values of $n < p$. Consider each triplet ^{p} equivalence separately. To simplify notation let r be any of the three r_n , and core residues $A_k(n+1) \equiv x^p \equiv x$, $-A_k(n) \equiv y^p \equiv y \pmod{p^k}$. Then $x^p + y^p \equiv x + y \equiv r^p \pmod{p^k}$, where $r^p \equiv 1 \pmod{p^2}$, has both summands in core, but $r^p \not\equiv 1 \pmod{p^k}$ for $k > 2$ is not in core: deviation $d \equiv r - r^p \not\equiv 0 \pmod{p^k}$.

Hence $r \equiv r^p + d \equiv (x + y) + d \pmod{p^k}$ (with $d \equiv 0 \pmod{p^k}$ in the cubic root case), and $x^p + y^p \equiv x + y \equiv (x + y + d)^p \pmod{p^k}$. The corresponding 0-extensions yield integer p -th power inequality: $X^p + Y^p < (X + Y + D)^p$. □

In the case of cubic roots in core A_k , less than full pk digit precision (base p), namely mod p^{3k+1} suffices to yield the FLT inequality (Corollary 2.1). For any triplet ^{p} mod p^2 , necessarily in core A_2 (Corollary 1.2), and for cubic roots of 1 mod p^k (any $k > 0$), there holds $(x + y)^p \equiv x + y \equiv x^p + y^p$, where exponent p distributes over a sum. By binomial expansion the sum of mixed terms yields integer $(X + Y)^p - (X^p + Y^p) \not\equiv 0$ of precision kp , which is $0 \pmod{p^2}$ for any triplet ^{p} .

For any triplet ^{p} mod p^k ($k > 2$), say in core increment form (5'), it is conjectured that there is a least precision $m(k)$ (base p), not exceeding that for cubic roots, which implies inequivalence $X^p - Y^p \not\equiv Z^p \pmod{p^m}$ ($Z^p \equiv 1 \pmod{p^2}$) for successive core 0-extensions $X, Y < p^k$.

Conjecture. *The 0-extensions $X, Y, Z < p^k$ of terms in any triplet ^{p} mod p^k equivalence in core increment form (5') with $X - Y = Z \equiv 1 \pmod{p^2}$ yield: $X^p - Y^p \not\equiv Z^p \pmod{p^{3k+1}}$.*

4. RELATION TO FERMAT'S SMALL AND LAST THEOREM

Core A_k as FST extension mod p^k ($k > 1$), the additive zero-sum property of its subgroups (Theorem 1.1), and the triplet structure of units group G_k (Theorem 3.1), allow a direct approach to Fermat's Last Theorem:

- (6) $x^p + y^p = z^p$ (prime $p > 2$) has no solution for positive integers x, y, z
 with case₁: $xyz \not\equiv 0 \pmod{p}$, and case₂: p divides one of x, y, z .

Usually (6) mentions exponent $n > 2$, but it suffices to show inequality for primes $p > 2$, because composite exponent $m = p \cdot q$ yields $a^{pq} = (a^p)^q = (a^q)^p$. In case₂: p divides just one term, because if p divides two terms then it also divides the third, and all terms can be divided by p^p .

A finite integer *FLT* solution of (6) has three p -th powers, each less than p^m for some finite fixed $m = kp$, with $x, y, z < p^k$, so (6) holds mod p^m , yet with no carry beyond p^{m-1} , 0-extending all terms.

The present approach needs only a simple form of Hensel's lemma [5] (in the general p -adic number theory), which is a direct consequence of Corollary 1.2, extend digit-wise the normed 1-complement form (2') such that the i -th digit of weight p^i in a^p and b^p sum to $p-1$ ($0 \leq i < k$), with p choices per extra digit. Thus to each normed solution of (2') mod p^2 there correspond p^{k-2} solutions mod p^k .

Corollary 4.1 (1-complement extension). *For $k > 2$, a normed FLT_k root is an extended FLT_2 root.*

4.1. Proof of the FLT inequality

Regarding *FLT* case₁, cubic root of 1 and triplet ^{p} are the only (normed) FLT_k roots (Theorem 3.1). Any assumed integer case₁ solution has a corresponding equivalent core increment form (4) with two terms in core, which by Lemma 3.2 has no integer extension, contradicting the assumption, as follows :

Theorem 4.1 (*FLT* case₁). *For prime $p > 2$ and integers $x, y, z > 0$ coprime to p equation $x^p + y^p = z^p$ has no solution.*

Proof. An FLT_k ($k > 1$) solution is a linear transformed extension of an FLT_2 root in core $A_2 = F_2$ (Corollary 4.1). By Lemma 3.2 it has no finite p -th power extension, yielding the theorem. \square

In *FLT* case₂ just one of x, y, z is a multiple of p , hence p^p divides one of the three p -th powers in $x^p + y^p = z^p$. Again, any assumed case₂ equality can be transformed to an equivalence mod p^p with two terms in core A_p , having no integer extension, contra the assumption.

Theorem 4.2 (*FLT* case₂). *For prime $p > 2$ and positive integers x, y, z , if p divides only one of x, y, z then $x^p + y^p = z^p$ has no solution.*

Proof. In a case₂ solution p divides a lefthand term, $x = cp$ or $y = cp$ ($c > 0$), or the right hand side $z = cp$. Bring the multiple of p to the right hand side, for instance if $y = cp$ then $z^p - x^p = (cp)^p$, while otherwise $x^p + y^p = (cp)^p$. So the sum or difference of two p -th powers coprime to p must be shown not to yield a p -th power $(cp)^p$ for any $c > 0$:

$$(7) \quad x^p \pm y^p = (cp)^p \text{ has no solution for integers } x, y, c > 0.$$

Notice that core increment form (4) does not apply here. However, by *FST* the two lefthand terms, coprime to p , are either complementary or equivalent mod p , depending on their sum or difference being $(cp)^p$. Scaling by s^p for some $s \equiv 1 \pmod{p}$, so $s^p \equiv 1 \pmod{p^2}$, transforms one lefthand term into a core residue

$A_p(n) \bmod p^p$, with $n \equiv x \bmod p$. And translation by adding $t \equiv 0 \bmod p^2$ yields the other term $A_p(n)$ or $-A_p(n) \bmod p^p$, respectively. The right hand side then becomes $s^p(cp)^p + t$, equivalent to $t \bmod p^p$. So the assumed equality (7) yields, by two equality preserving transformations, the next equivalence (8), where $A_p(n) \equiv u \equiv u^p \bmod p^p$ (u in core A_p for $0 < n < p$ with $x \equiv n \bmod p$) and $s \equiv 1, t \equiv 0 \bmod p^2$

$$(8) \quad \begin{aligned} u^p \pm u^p &\equiv u \pm u \equiv t \pmod{p^p} \quad (u \in A_p), \text{ with } u \equiv (sx)^p, \\ \pm u &\equiv \pm (sy)^p + t \pmod{p^p}. \end{aligned}$$

Equivalence (8) does not extend to integers, because $U^p + U^p > U + U$, and $U^p - U^p = 0 \neq T$, where U, T are the 0-extensions of $u, t \bmod p^p$, respectively. But this contradicts assumed equalities (7), which consequently must be false. \square

Note. From a practical point of view the *FLT* integer inequality with terms less than p^{pk} of a 0-extended FLT_k root (case₁) is caused by the *carries* beyond p^k , amounting to a multiple of the modulus p^k , produced in the arithmetic (base p). In the expansion of $(a + b)^p$, the mixed terms *can* vanish mod p^k for some a, b, p . Ignoring the carries yields $(a + b)^p \equiv a^p + b^p \bmod p^k$, and the *EDS'* property is as it were the *syntactical* expression of ignoring the carry (*overflow*) in residue arithmetic. In other words, in terms of p -adic number theory, this means 'breaking the Hensel lift': the residue equivalence of an FLT_k root mod p^k , although it holds for all $k > 0$, *does* imply inequality for integer p -th powers less than p^{pk} due to its special triplet structure, where exponent p distributes over a sum.

Conclusions and Remarks

1. The two symmetries $-n, n^{-1}$ determine FLT_k roots, which are necessary for an *FLT* integer solution. However, these symmetries (automorphisms) do not exist for positive integers.
2. Another proof of *FLT* case₁ might use product $1 \bmod p^k$ of FLT_k root terms: $ab \equiv 1$ or $abc \equiv 1$, which is impossible for integers > 1 . The p -th power of a k -digit natural requires upto pk digits. Arithmetic mod p^k ignores carries of weight p^k and beyond. Interpreting a given FLT_k equivalence in naturals less than p^k , their p -th powers produce for $p > 2$ carries that cause inequality.
3. Core $A_k \subset G_k$ as extension of *FST* to mod p^k $k > 1$, and the zero-sum of its subgroups (Theorem 1.1) yielding the cubic *FLT* root (Lemma 2.1), initiated this work. The triplets were found by analysing a computer listing (Table 2) of the *FLT* roots mod p^2 for primes $p < 200$.
4. Linear analysis (mod p^2) suffices for root existence (Hensel, Corollary 4.1), but triplet ^{p} core increment form (4) with two successor terms in core requires *quadratic* analysis (mod p^3). Similarly, *FLT* case₁ inequivalence mod p^{3k+1} holds for increments of $C_{k+1} \equiv (C_k)^p$ for 0-extended core A_k .
5. "*FLT* eqn(1) has no finite solution" and "[*ICS*]³ has no finite fixed point" are equivalent (Theorem 3.1), yet each $n \in G_k$ is a fixed point of [*ICS*]³

mod p^k (re: FLT_2 roots imply all roots for $k > 2$, yet no 0-extension to integers).

6. Crucial in finding the arithmetic triplet structure were extensive computer experiments, and the application of *associative function composition*, the essence of semi-groups, to the three elementary functions (Theorem 3.1): successor $S(n) = n+1$, complement $C(n) = -n$ and inverse $I(n) = n^{-1}$, with period 3 for $SCI(n) = -(n+1)^{-1}$ and the other three such compositions. In this sense FLT is not a purely arithmetic problem, but essentially requires non-commutative and associative function composition for its proof.

REFERENCES

1. Apostol T., *Introduction to Analytical Number Theory*, Springer Verlag, 1976.
2. Benschop N., *The semigroup of multiplication mod p^k , an extension of Fermat's Small Theorem, and its additive structure*, International conference Semigroups and their Applications, Prague, July 1996.
3. Benschop N., *Powersums representing residues mod p^k , from Fermat to Waring*, Computers and Mathematics with Applications, **39**(7-8) (2000), 253-261.
4. Clifford A. and Preston G., *The Algebraic Theory of Semigroups*, AMS survey 7, **1** (1961), 130-135.
5. Hardy G. and Wright E., *An Introduction to the Theory of Numbers*, Oxford University Press 1979.
6. Mohit S and Ram Murty, *Wieferich Primes and Hall's Conjecture*, Comptes Rendus de l'Academie. Science, **20**(1) (1998), 29-32.
7. Patent US-5923888 (July 13, 1999) on a Logarithmic Binary Multiplier, with dual bases 2 and 3 (using 3 as semi-primitive root of 1 mod 2^k).
8. Schwarz S., *The Role of Semigroups in the Elementary Theory of Numbers*, Mathematica Slovaca, **31**(4) (1981), 369-395.
9. Wieferich A., *Zum letzten Fermat'schen Theorem*, J. Reine Angew. Math, **136** (1909), 293-302.

N. F. Benschop, Schoutstraat 4, 5663EZ Geldrop, The Netherlands,
e-mail: n.benschop@chello.nl – Amspade Research