

# NONAUTONOMOUS LIVŠIĆ COLLIGATIONS AND HYPONORMAL OPERATORS

I. V. VOROBYOV

ABSTRACT. In this work, the model of a nonautonomous Livšić colligation for operators that are close to the normal is studied. The connection between the determining function for operator and the characteristic function for colligation is obtained.

In this paper the model of nonautonomous Livšić colligation is studied relying on the singular integral model for operators that are close to normal, and also its characteristic function is introduced. By means of the Plemel formula (extended to vector-valued functions) [4], [1], the Riemann-Hilbert problem that connects the boundary values of the determining function with the characteristic function of colligation is obtained.

## 1. MAIN NOTATION AND DEFINITIONS

Let  $\delta$  be a bounded closed set on the real axis,  $\Lambda$  be a set of all Borel sets that are contained in  $\delta$ ,  $m$  be the Lebesgue measure on  $\delta$ ,  $\nu$  be a singular measure on  $\Lambda$  (i. e. there is a set  $Q \in \Lambda$  such that  $m(Q) = 0$  and  $\nu(\delta \setminus Q) = 0$ ), and  $\mu = m + \nu$  be  $\sigma$ -finite measure,  $\Omega = (\delta, \Lambda, \mu)$  be a space with measure,  $D$  be a complex separable Hilbert space,  $\langle a, b \rangle_D$  be a scalar product of vectors  $a$  and  $b$  in  $D$ ,  $L(D)$  be the algebra of all linear operators in  $D$ ,  $f(x) \in D$  be a vector valued ( $D$ -valued)

---

Received August 8, 2007; revised October 17, 2008.

2000 *Mathematics Subject Classification.* Primary 53A04.

*Key words and phrases.* singular integral model; hyponormal operator.




Go back

Full Screen

Close

Quit



measurable function defined for all  $x \in \delta$ ,

$$L^2(\Omega, D) = \left\{ f \in D : \int_{\delta} \|f(x)\|_D^2 d\mu(x) < \infty \right\}$$

be the Hilbert space of measurable vector valued quadratically summable functions with the scalar product

$$\langle f, g \rangle_{L^2(\Omega, D)} = \int_{\delta} \langle f(x), g(x) \rangle_D d\mu(x), \quad \forall f, g \in L^2(\Omega, D).$$

We identify functions  $f(\cdot)$  and  $g(\cdot)$  from  $L^2(\Omega, D)$  if they differ only on a set of zero  $\mu$ -measure.

Let  $\{0\} = D_0 \subset D_1 \subset \dots \subset D_n \subset \dots$  be a sequence of spaces of finite dimension in  $D$ ,  $F_0 \subset F_1 \subset \dots \subset F_n \subset \dots$  be a sequence of sets in  $\Lambda$ . An  $L(D)$ -valued measurable function  $R(x)$ ,  $x \in \delta$ , is said to be the standard projector valued function [4] if

1.  $R(x) = 0$ ,  $x \in F_0$ ;
2.  $R(x)$  is an orthoprojector from  $D$  into  $D_n$  if  $x \in F_n \setminus F_{n-1}$ ,  $n = 1, 2, \dots$ ;
3.  $R(x) = I$ ,  $x \notin \bigcup_{n \geq 0} F_n$ .

If  $\mu(\delta) = 1$ , then the vector  $a \in D$ . One can identify with the vector valued function with the constant value of  $a$ . From this point of view,  $D$  is a subspace of  $L^2(\Omega, D)$ , and their scalar products coincide. In the general case, one can define the orthoprojector from  $L^2(\Omega, D)$  into  $D$  as follows:

$$\tilde{P}_0 : f(\cdot) \rightarrow \frac{1}{\mu(\delta)} \int_{\delta} f(x) d\mu(x).$$




Go back

Full Screen

Close

Quit



Hereinafter, we will use the following operator:

$$(1) \quad P_0 = \mu(\delta) \tilde{P}_0.$$

Introduce into consideration the following space:

$$(2) \quad H = L^2(\Omega, D, R(\cdot)) = \{f \in L^2(\Omega, D) : R(\cdot) f(\cdot) = f(\cdot)\}.$$

It, as a subspace of  $L^2(\Omega, D)$ , also is a Hilbert space.

Let  $F$  be the vector valued Fourier transform of a function  $f(\cdot)$  defined in  $L^2(D)$  on the real axis, i. e. the next strong limit

$$(Ff)(x) = s - \lim_{N \rightarrow \infty} \frac{1}{\sqrt{2\pi}} \int_{-N}^N f(t) e^{itx} dt.$$

It is well-known that  $F$  is a unitary operator in  $L^2(D)$  on the axis. Define the following operator

$$P = F^* P_{[0, +\infty)} F$$

where  $P_{[0, +\infty)}$  is the operator of multiplication by the characteristic function of semiaxis  $[0, +\infty)$ . By the Plemel formula, one can explicitly express the operator  $P$  [4], [1]:

$$(3) \quad (Pf)(x) = s - \lim_{\varepsilon \rightarrow -0} \frac{1}{2\pi i} \int_{-\infty}^{+\infty} \frac{f(t) dt}{(x + i\varepsilon) - t} = \frac{f(x)}{2} + \frac{1}{2\pi i} v.p. \int_{-\infty}^{+\infty} \frac{f(t) dt}{x - t}$$

where the symbol  $v.p.$  signifies the main value of singular integral.




Go back

Full Screen

Close

Quit



## 2. CONSTRUCTION OF SINGULAR INTEGRAL MODEL

Let  $\alpha(\cdot)$  and  $\beta(\cdot) = \beta^*(\cdot)$  be uniformly bounded measurable functions defined in  $\Omega$ , with the values in  $L(D)$ , such that

$$(4) \quad \begin{aligned} \alpha(\cdot) R(\cdot) &= R(\cdot) \alpha(\cdot) = \alpha(\cdot), \\ \beta(\cdot) R(\cdot) &= R(\cdot) \beta(\cdot) = \beta(\cdot), \\ \alpha(x) &= 0, \quad x \in Q, \end{aligned}$$

where  $Q$  is a set in  $\delta$ , for which  $m(Q) = 0$  and  $\nu(\delta \setminus Q) = 0$ .

Define the self-adjoint operator  $X$  in the Hilbert space  $H$  (2) in the following way:

$$(5) \quad (Xf)(x) = xf(x), \quad \forall f \in H.$$

Let  $\sigma$  be a self-adjoint operator in  $D$ . It is easy to check that the following operator

$$(6) \quad (Yf)(x) = \beta(x)f(x) + \alpha^*(x)\sigma P(\alpha(\cdot)f(\cdot)), \quad \forall f \in H,$$

where  $P$  is defined in (3), is self-adjoint in  $H$  (here we assume that  $\alpha f \equiv 0$  outside  $\delta$ , i. e. the singular integral we take by  $\delta$ ). Let the operators  $X$  and  $Y$  be respectively the real and the imaginary part of operator  $T$ , i. e.

$$(7) \quad T = X + iY.$$

Define the bounded linear operator from  $D$  in  $H$ :

$$(8) \quad Ka = \frac{1}{\sqrt{\pi}}\alpha^*(\cdot)a, \quad \forall a \in D.$$

The adjoint operator  $K^*$  as an operator from  $H$  in  $D$  is given by

$$(9) \quad K^*f = \frac{1}{\sqrt{\pi}}P_0(\alpha(\cdot)f(\cdot)), \quad \forall f \in H,$$




Go back

Full Screen

Close

Quit



where  $P_0$  is defined in (1). Using the definitions of self-adjoint operators, one can calculate the self-commutator of operator  $T$  (7):

$$(10) \quad \langle [T^*, T] f, f \rangle_H = \frac{1}{\pi} \langle \sigma P_0 (\alpha(\cdot) f(\cdot)), P_0 (\alpha(\cdot) f(\cdot)) \rangle_D, \quad \forall f \in H,$$

Besides, the properties of scalar product yield

$$(11) \quad ([T^*, T] f)(x) = \frac{1}{\pi} \alpha^*(x) \sigma P_0 (\alpha(\cdot) f(\cdot)), \quad \forall f \in H.$$

In the terms of operator  $K$  from (8) and (9), the self-commutator of  $T$  is given by

$$(12) \quad [T^*, T] = 2i[X, Y] = K\sigma K^*,$$

this implies that when  $\sigma \geq 0$  the operator  $T$  is hyponormal.

In the book by Xia [4] it is proved that every hyponormal operator is unitarily equivalent to the operator  $T$  (5)-(7) with  $\sigma = I_D$  and  $\alpha(x) = \alpha^*(x) \geq 0$ . Thus, the singular integral operator model is the functional model of the hyponormal operator.

**Definition 2.1.** An arbitrary operator  $A$  is said to be the completely non-normal (see [4]) if there is no subspace reducing  $A$ , on which  $A$  induces a normal operator.

**Lemma 2.2.** If a singular integral operator  $T$  (7) is completely non-normal, then the singular measure  $\nu \equiv 0$ .

*Proof.* From the contrary. Suppose that  $\nu \neq 0$  is concentrated in the set  $Q \in \Lambda$  and  $m(Q) = 0$ . Consider the following subspace  $H$ :

$$N = \{f \in H : f(x) = 0, \forall x \notin Q\}.$$




Go back

Full Screen

Close

Quit



Then from (4)  $\alpha f \equiv 0$  for all  $f \in N$ , therefore

$$\begin{aligned} (Tf)(x) &= xf(x) + i\beta(x)f(x), & \forall f \in N, \\ (T^*f)(x) &= xf(x) - i\beta(x)f(x), & \forall f \in N. \end{aligned}$$

Consequently,  $N$  is a nontrivial normal subspace that reduces the operator  $T$ , it contradicts the complete non-normality of  $T$ . The lemma is proved.

### 3. NONAUTONOMOUS LIVŠIĆ COLLIGATION

Introduce the open system  $\{\mathfrak{R}_\Delta, S_\Delta\}$  :

$$(13) \quad \begin{cases} \mathfrak{R}_\Delta : i \frac{d}{dx} h(x) + \beta(x)h(x) = \alpha^*(x)\sigma u(x), & h(0) = h_0, \\ S_\Delta : v(x) = u(x) + \alpha(x)h(x) \end{cases}$$

where  $\alpha(x)$  and  $\beta(x)$  are from the singular integral model for  $T$ .

The operators in this open system form the objects  $(\beta(x), D, \beta(x), D, \sigma)$  for a Livšić node  $\mathcal{C}_x$  [2] depending on the real parameter  $x$ . The associated input-state-output system (13) is nonautonomous [2].

By direct computation, it is easy to see that the following lemma is true.

**Lemma 3.1.** *The following energy conservation law is true:*

$$(14) \quad \begin{aligned} \frac{d}{dx} \|h(x)\|_D^2 &= \langle i\sigma v(x), u(x) \rangle_D - \langle i\sigma u(x), v(x) \rangle_D \\ &= \left\langle \begin{pmatrix} 0 & i\sigma \\ -i\sigma & 0 \end{pmatrix} \begin{pmatrix} u(x) \\ v(x) \end{pmatrix}, \begin{pmatrix} u(x) \\ v(x) \end{pmatrix} \right\rangle_{D \oplus D}. \end{aligned}$$




Go back

Full Screen

Close

Quit



Similar to [2], for all the non-real  $z$  we can introduce into consideration the following  $D$ -valued function

$$(15) \quad S_{\Delta}(x, z) = I_D + \alpha(x) (\beta(x) - zI_D)^{-1} \alpha^*(x) \sigma,$$

which we will call the characteristic function of the colligation  $\mathcal{C}_x$ . Directly from definition (15), it is easy to see that the following relations are correct:

$$(16) \quad S_{\Delta}(x, z) \alpha(x) = \alpha(x) (\beta(x) - zI_D)^{-1} (\beta(x) + \alpha^*(x) \sigma \alpha(x) - zI_D),$$

$$(17) \quad \alpha^*(x) \sigma S_{\Delta}(x, z) = (\beta(x) + \alpha^*(x) \sigma \alpha(x) - zI_D) (\beta(x) - zI_D)^{-1} \alpha^*(x) \sigma,$$

$$(18) \quad S_{\Delta}(x, z)^{-1} = I_D - \alpha(x) (\beta(x) + \alpha^*(x) \sigma \alpha(x) - zI_D)^{-1} \alpha^*(x) \sigma;$$

$$(19) \quad \frac{1}{\bar{w} - z} [S_{\Delta}^*(x, w) \sigma - \sigma S_{\Delta}(x, z)] = \kappa^*(x, w) \kappa(x, z),$$

$$\kappa(x, z) = (\beta(x) - zI_D)^{-1} \alpha^*(x) \sigma.$$

Let the operator  $\sigma$  be invertible. For all the non-real  $z$  and  $w$  define the following operator valued function  $E(z, w) : D \rightarrow D$ :

$$(20) \quad E(z, w) = \sigma^{-1} + \frac{1}{21} K^* (X - zI_H)^{-1} (Y - wI_H)^{-1} K,$$




Go back

Full Screen

Close

Quit



which we will call the determining function of the operator  $T$ . It is easy to see that the determining function satisfies the following relations:

$$(21) \quad K\sigma E(z, w) = \{(X - zI_H), (Y - wI_H)\} K,$$

$$(22) \quad E(z, w) \sigma K^* = K^* \left\{ (X - zI_H)^{-1}, (Y - wI_H)^{-1} \right\},$$

$$(23) \quad E(z, w)^{-1} = \sigma - \frac{1}{2i} \sigma K^* (Y - wI_H)^{-1} (X - zI_H)^{-1} K \sigma,$$

where  $\{A, B\} = ABA^{-1}B^{-1}$  is a multiplicative commutator of operators  $A$  and  $B$ .

Denote  $f(x, w; a) = (Y - wI_H)^{-1} \alpha^*(\cdot) a$ . It follows from (6) that for each  $a \in D$

$$(Y - \beta(\cdot)) f(\cdot, w; a) = \alpha^*(\cdot) \sigma P_- [\alpha(\cdot) f(\cdot, w; a)],$$

where  $P_- = P$  is the singular integral operator (3). Since

$$(Y - wI_H) f(\cdot, w; a) = \alpha^*(x) a,$$

then it is obvious that

$$(\beta(\cdot) - wI_D) f(\cdot, w; a) + \alpha^*(\cdot) \sigma P_- [\alpha(\cdot) f(\cdot, w; a)] = \alpha^*(\cdot) a.$$

Multiply both sides from the left by  $\alpha(\cdot) (\beta(\cdot) - wI_D)^{-1}$  and denote  $P_+ = I_H - P_-$ . Then

$$\begin{aligned} (P_+ + P_-) [\alpha(\cdot) f(\cdot, w; a)] + \alpha(\cdot) (\beta(\cdot) - wI_D)^{-1} \alpha^*(\cdot) \sigma \\ \times P_- [\alpha(\cdot) f(\cdot, w; a)] = \alpha(\cdot) (\beta(\cdot) - wI_D)^{-1} \alpha^*(\cdot) \sigma \sigma^{-1} a. \end{aligned}$$

Add  $\sigma^{-1}a$  to the both sides and take (15) into account. Hence

$$(24) \quad \sigma^{-1}a + P_+ [\alpha(\cdot) f(\cdot, w; a)] = S(\cdot, w) [\sigma^{-1}a - P_- [\alpha(\cdot) f(\cdot, w; a)]] .$$




Go back

Full Screen

Close

Quit





Using (5), (8) and (9), one can write the determining function as an integral

$$(25) \quad E(z, w) a = \sigma^{-1} a + \frac{1}{2\pi i} \int_{\delta} \frac{\alpha(x) f(x, w; a)}{x - z} dx,$$

whence by the Plemel formulae [1] one obtains that for all real  $x$

$$(26) \quad E(x \pm i0, w) a = \sigma^{-1} a \pm P_{\pm} [\alpha(\cdot) f(\cdot, w; a)].$$

Thus, from (24) and (25) we finally obtain that the determining function satisfies the following Riemann-Hilbert problem :

$$(27) \quad E(x + i0, w) = S(x, w) E(x - i0, w).$$

1. Aleshkov Yu. Z. and Smishlyayev P. P., *Theory of functions of complex variable and its applications.*, The Leningad University Publishing House 1986, (in Russian).
2. Vinnikov V., *Commuting operators and functions on a Riemann surface*, Holomorphic spaces (Ed. S. Axler, J. E. McCarthy and D. Sarason), Mathematical Science Research Institute Publications **33**, Cambridge 1998, 445–476.
3. Livšic M. S. and Yantsevich A. A., *Theory of operator colligations in Hilbert spaces*, Kharkov University Publishing House, Kharkov. 1971, (in Russian).
4. Xia Daoxing, *Spectral theory of hyponormal operators*, Operator Theory: Advances and Applications, Basel; Boston; Stuttgart: Birkhauser-Verlag 1983.
5. Zolotarev V. A., *Method of open systems. Triangular and functional models of commutative system of two operators*, Manuscript. Dep. in Ukr. NIINTI RZh. Mat. 9-630. Dep. 1984, (in Russian).

I. V. Vorobyov, Department of Mechanics and Mathematics, V. N. Karazin Kharkov National University, 4 Svobody Sq., Kharkov, Ukraine, e-mail: [ivvorobyov@yahoo.com](mailto:ivvorobyov@yahoo.com)



Go back

Full Screen

Close

Quit