



ON INTEGERS EXPRESSIBLE BY SOME SPECIAL LINEAR FORM

A. DUBICKAS AND A. NOVIKAS

ABSTRACT. Let $E(4)$ be the set of positive integers expressible by the form $4M - d$, where M is a multiple of the product ab and d is a divisor of the sum $a + b$ of two positive integers a, b . We show that the set $E(4)$ does not contain perfect squares and three exceptional positive integers 288, 336, 4545 and verify that $E(4)$ contains all other positive integers up to $2 \cdot 10^9$. We conjecture that there are no other exceptional integers. This would imply the Erdős-Straus conjecture asserting that each number of the form $4/n$, where $n \geq 2$ is a positive integer, is the sum of three unit fractions $1/x + 1/y + 1/z$. We also discuss similar problems for sets $E(t)$, where $t \geq 3$, consisting of positive integers expressible by the form $tM - d$. The set $E(5)$ is related to a conjecture of Sierpiński, whereas the set $E(t)$, where t is any integer greater than or equal to 4, is related to the most general in this context conjecture of Schinzel.

1. INTRODUCTION

Let t be a fixed positive integer. In this paper we consider the set of positive integers

$$E(t) := \{n : n = tM - d\},$$

where M is a positive multiple of the product and d is a positive divisor of the sum of two positive integers, namely,

$$ab|M \quad \text{and} \quad d|(a + b)$$

Received November 10, 2011; revised June 21, 2012.

2010 *Mathematics Subject Classification*. Primary 11D68, 11D09, 11Y50.

Key words and phrases. Egyptian fractions; Erdős-Straus conjecture; Sierpiński conjecture; Schinzel's conjecture.



Go back

Full Screen

Close

Quit



for some $a, b \in \mathbb{N}$. Evidently,

$$E(t') \subseteq E(t) \quad \text{whenever} \quad t|t'.$$

It is easy to see that

$$(1) \quad E(1) = E(2) = \mathbb{N}.$$

Indeed, suppose first that $t = 1$. Then, for each $n \in \mathbb{N}$ selecting $a = 2n + 1$, $b = 1$, $M = ab = 2n + 1$ and $d = (a + b)/2 = n + 1$, we find that

$$n = 2n + 1 - (n + 1) = M - d,$$

giving $E(1) = \mathbb{N}$. In case $t = 2$, for each $n \in \mathbb{N}$ we may choose $a = n + 1$, $b = 1$, $M = ab = n + 1$ and $d = a + b = n + 2$. Then $2M - d = 2(n + 1) - (n + 2) = n$, so that $E(2) = \mathbb{N}$.

Apart from (1) the situation with $t \geq 3$ is not clear. In this context, the sets $E(4)$ and $E(5)$ are of special interest because an integer n belongs to the set $E(t)$ if and only if

$$n = tM - d = tuab - (a + b)/v$$

with some $a, b, u, v \in \mathbb{N}$. Therefore, $n \in E(t)$ yields the representation

$$\frac{t}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}$$

with positive integers

$$x := uab, \quad y := uvna, \quad z := uvnb.$$

Thus if $n \in E(t)$, then the fraction t/n is expressible by the sum of three unit fractions. In particular, if every prime number p belongs to the set $E(4)$, then the Erdős-Straus conjecture (asserting that for each integer $n \geq 2$, the fraction $4/n$ is expressible by the sum $1/x + 1/y + 1/z$ with $x, y, z \in \mathbb{N}$) is true, whereas if every prime number p belongs to $E(5)$, then the corresponding conjecture of Sierpiński (asserting that for each $n \geq 4$, the fraction $5/n$ is expressible by the sum $1/x + 1/y + 1/z$) is true [10]. In this context, the most general Schinzel's conjecture asserts that



Go back

Full Screen

Close

Quit



the fraction t/n for each $n \geq n(t)$ is expressible by the sum $1/x + 1/y + 1/z$. This clearly holds for $t \leq 3$, but is open for each fixed $t \geq 4$. Conjecture 5 given in Section 3 implies that there is an integer $C(t)$ such that each prime number $p > C(t)$ belongs to $E(t)$. This would imply Schinzel's conjecture as well.

Yamamoto [12], [13] and Mordell [8] observed that it is sufficient to prove the Erdős-Straus conjecture for those prime numbers p which modulo 840 are 1, 121, 169, 289, 361 or 529. Vaughan [11] showed that the Erdős-Straus conjecture is true for almost all positive integers n . See also the list of references in D11 for the literature concerning the conjectures of Erdős-Straus, Sierpiński and Schinzel on Egyptian fractions. More references on the Erdős-Straus (including recent ones) can be found in a paper of Elsholtz and Tao [4] on the average number of solutions of the equation $4/p = 1/x + 1/y + 1/z$ with prime numbers p . At the computational side the calculations of Swett <http://math.uindy.edu/swett/esc.htm> show that the Erdős-Straus conjecture holds for integers n up to 10^{14} .

In this note we observe that the following holds

Theorem 1. *The set $E(4)$ does not contain perfect squares and the numbers 288, 336, 4545.*

Suppose $k^2 \in E(4)$, i.e., there exist $u, v, a, b, k \in \mathbb{N}$ such that

$$(2) \quad v(4uab - k^2) = a + b.$$

To show that $k^2 \notin E(4)$, we shall use the following fact

Lemma 2. *The equation (2) has no solutions in positive integers u, v, a, b, k .*

Lemma 2 implies that $-d$ is a quadratic nonresidue modulo $4ab$ if $d|(a+b)$. Indeed, if the number $-d$ were a quadratic residue modulo $4ab$, then by selecting the positive integer $v := (a + b)/d$, we would see that the equation $k^2 = -d + 4uab$ with $u \in \mathbb{N}$ has a solution $k \in \mathbb{N}$, which is impossible in view of Lemma 2. Note that the set of divisors of $a + b$, when $a < b$ both run through the set $\{1, 2, \dots, n\}$, contains the set $\{1, 2, \dots, 2n - 1\}$. Thus, by Lemma 2, it holds



Go back

Full Screen

Close

Quit



Corollary 3. For each positive integer n the $2n - 1$ consecutive integers

$$4n! - 2n + 1, 4n! - 2n + 2, \dots, 4n! - 1$$

are quadratic nonresidues modulo $4n!$.

Corollary 3 gives the example of at least $(2 - \varepsilon) \log m / \log \log m$ consecutive quadratic nonresidues modulo $m = 4n!$ (by a completely elementary method). In this direction, the most interesting problem is to determine how many consecutive quadratic residues and consecutive quadratic nonresidues modulo m may occur for prime numbers m . See, e.g., [3], [5], where it is shown that we have at least $c_1 \log m \log \log \log m$ consecutive quadratic residues modulo m for infinitely many primes m , and [7], where the factor $\log \log \log m$ is replaced by $\log \log m$ under assumption of the generalized Riemann hypothesis.

A set of positive integers which is a subset of $\cup_{q=0}^{\infty} E(4q + 3)$ was recently considered in [1]. For $M = ab$ and $d = a + b$, where a, b are positive integers and $b \equiv 3 \pmod{4}$, put

$$E^*(t) := \{n : n = tab - a - b\}.$$

Evidently, $E^*(t) \subseteq E(t)$. In [1] it was shown that the set $E := \cup_{q=0}^{\infty} E^*(4q + 3)$ does not contain perfect squares and that all prime numbers of the form $4s + 1$ less than 10^{10} belong to E .

2. PROOF OF THEOREM 1

Lemma 2 was apparently first proved by Yamamoto [13]. See also [9, Lemma 2] and [4, Proposition 1.6]. Here is a short proof.

Since $a = vd - b$, equality (2) yields

$$k^2 = 4u(vd - b)b - d = (4bu v - 1)d - 4b^2u.$$



Go back

Full Screen

Close

Quit



So if (2) has a solution in positive integers, then the Jacobi symbol $\left(\frac{-4b^2u}{4buv-1}\right) = \left(\frac{k^2}{4buv-1}\right)$ must be equal to 1. Indeed, since $-4b^2u$ and $4buv - 1$ are relatively prime, we have $\left(\frac{-4b^2u}{4buv-1}\right) \neq 0$, and so $\left(\frac{k^2}{4buv-1}\right) = 1$. We will show, however, that the Jacobi symbol $\left(\frac{-4b^2u}{4buv-1}\right)$ is equal to -1 . Indeed, write $u \in \mathbb{N}$ in the form $u = 2^r u_0$, where $r \geq 0$ is an integer and $u_0 \geq 1$ is an odd integer. Using $\left(\frac{-1}{4buv-1}\right) = -1$ and also $\left(\frac{2}{4buv-1}\right) = 1$ in case u is even, i.e., $r \geq 1$, we find that

$$\left(\frac{-4b^2u}{4buv-1}\right) = \left(\frac{-2^{r+2}b^2u_0}{4buv-1}\right) = -\left(\frac{2^r u_0}{4buv-1}\right) = -\left(\frac{u_0}{4buv-1}\right).$$

Further, by the quadratic reciprocity law, in view of $u_0|u$ we conclude that

$$-\left(\frac{u_0}{4buv-1}\right) = -(-1)^{(u_0-1)/2} \left(\frac{4buv-1}{u_0}\right) = -(-1)^{(u_0-1)/2} \left(\frac{-1}{u_0}\right) = -1.$$

Lemma 2 implies that $k^2 \notin E(4)$. To complete the proof of Theorem 1 we need to show that 288, 336, 4545 $\notin E(4)$.

The case $n = 288$ can be easily checked ‘by hand’. Observe that $288 = 4M - d$ implies that $d = 4s$ and $M = (288 + 4s)/4 = 72 + s$. Furthermore, from

$$72 + s = M \geq ab \geq a + b - 1 \geq d - 1 = 4s - 1$$

we find that $1 \leq s \leq 24$. So for each $s = 1, 2, \dots, 24$, it remains to check that there are no positive integers a, b for which $4s|(a+b)$ and $ab|(72+s)$.

Note first that for $s \geq 11$ we must have $a+b = 4s$ and $ab = 72+s$. Indeed, if $a+b > 4s$, then $a+b \geq 8s$ and so

$$72 + s = M \geq ab \geq a + b - 1 \geq 8s - 1,$$




Go back

Full Screen

Close

Quit



which is impossible, because $s \geq 11$. If $ab < 72 + s$, then $2ab \leq 72 + s$, so that

$$72 + s \geq 2ab \geq 2(a + b - 1) \geq 2(d - 1) = 2(4s - 1) = 8s - 2,$$

which is a contradiction again. However, from $a + b = 4s$ and $ab = 72 + s$ it follows that

$$(4s)^2 - 4(72 + s) = 4(4s^2 - s - 72)$$

is a perfect square. So $4s^2 - s - 72$ must be a perfect square. It remains to check the values of s between 11 and 24 which modulo 4 are 0 or 3, namely, $s = 11, 12, 15, 16, 19, 20, 23, 24$. For none of these values, $4s^2 - s - 72$ is a perfect square.

The values of s between 1 and 10 can also be excluded, because there are no a, b with $ab | (72 + s)$ for which $4s$ divides $a + b$; see the Table 1 below.

To complete the proof of the theorem, observe that if $n = tM - d$, then

$$n \geq tab - a - b \geq ta^2 - 2a$$

in case $a \leq b$. Hence $(at - 1)^2 \leq nt + 1$ and $b \leq (a + n)/(ta - 1)$. Therefore, all values from 1 to 10000 which do not belong to $E(4)$ can be found with Maple as follows:

```
for n from 1 to 10000 do s := true;
  for a from 1 by 1 while (s and (at - 1)^2 <= tn + 1) do
    B := (a + n)/(ta - 1);
    for b from a by 1 while (s and b <= B) do
      if a + b (mod tab - n (mod tab)) = 0 then s := false endif;
    endfor;
  endfor;
  if s then k := k + 1;
    print(n);
  endif;
endfor;
print(k):
```




Go back

Full Screen

Close

Quit



For every particular value of n from 1 to 10000, we check all the pairs (a, b) which satisfy the above inequalities for the existence of an appropriate value of d , i.e., for the divisibility of $a + b$ by some positive integer of the form $d = tuab - n$. However, $d \leq a + b \leq tab$ which means that there is only one possible such integer d . Take a unique integer in the interval $[1, tab]$ which equals $-n$ modulo tab . It can be expressed as $tab - n \pmod{tab}$, as in the pseudocode describing our algorithm above. To obtain a code for Maple, one only needs to change both 'endfor' and 'endif' to 'end'.

s	1	2	3	4	5	6	7	8	9	10
$4s$	4	8	12	16	20	24	28	32	36	40
$72 + s$	73	$2 \cdot 37$	$3 \cdot 5^2$	$2^2 \cdot 19$	$7 \cdot 11$	$2 \cdot 3 \cdot 13$	79	$2^4 \cdot 5$	3^4	$2 \cdot 41$


Table 1. .

As a result (in less than three seconds), we got that only 100 perfect squares and three exceptional numbers 288, 336, 4545 less than 10000 do not lie in $E(4)$. This completes the proof of Theorem 1.

The calculation to the bound 10^6 with Maple took us almost 40 minutes, so all the calculations of the next section to the bound $2 \cdot 10^9$ have been performed with C++.

3. SOME SPECULATIONS CONCERNING THE SETS $E(t)$

As we already observed in (1), the sets $\mathbb{N} \setminus E(1)$ and $\mathbb{N} \setminus E(2)$ are empty. By Lemma 2, the equation $v(4uab - k^2) = a + b$ has no solutions in positive integers u, v, a, b, k . In particular, if t is a positive integer divisible by 4 and $s \in \mathbb{N}$ is such that $4s|t$, then the equation $vs(4(t/4s)uab - k^2) =$



Go back

Full Screen

Close

Quit



$a + b$ has no solutions in positive integers u, v, a, b, k . The latter is equivalent to the equation $v(tuab - sk^2) = a + b$. Consequently, we obtain the following corollary

Corollary 4. *The set $E(t)$, where $4|t$, does not contain the numbers of the form sk^2 , where $s \in \mathbb{N}$ satisfies $4s|t$ and $k \in \mathbb{N}$.*

In particular, this implies that the set $\mathbb{N} \setminus E(t)$ is infinite when $4|t$. We conjecture that all other sets, namely, $\mathbb{N} \setminus E(t)$ with $t \in \mathbb{N}$ which is not a multiple of 4 are finite. More precisely, we get next conjecture

Conjecture 5. *There exists an integer $C(t) \in \mathbb{N} \cup \{0\}$ such that the set $E(t)$ contains all integers greater than or equal to $C(t) + 1$ if 4 does not divide t and all integers greater than or equal to $C(t) + 1$ except for sk^2 , where $4s|t$ and $k \in \mathbb{N}$, if $4|t$.*

By (1), we have $C(1) = C(2) = 0$. It is known that the total number of representations of t/n by the sum $1/x + 1/y + 1/z$ does not exceed $c(\varepsilon)(n/t)^{2/3}n^\varepsilon$, where $\varepsilon > 0$ (see [2]). We know that if $n \in E(t)$ then t/n is expressible by the sum of three unit fractions, so this bound also holds for the number of representations of n in the form $tM - d$. On the other hand, by the above mentioned result of Vaughan [11], almost all positive integers are expressible by the sum of three unit fractions. It is easy to see that for each fixed $t \geq 3$, almost all positive integers belong to the set $E(t)$.

In fact, one can easily show a much stronger statement that almost all positive integers can be written in the form $pa - 1$ with some prime number $p \equiv -1 \pmod{t}$ and some $a \in \mathbb{N}$. If $n \in \mathbb{N}$ can be written in this way, then

$$n = pa - 1 = (p + 1)a - a - 1 = tM - d \in E(t)$$

with $b = 1$, $d = a + 1$ and $M = (p + 1)a/t$. By the above, it suffices to show that the density of positive integers n that have no prime divisors of the form $p \equiv -1 \pmod{t}$ is zero. This can be



Go back

Full Screen

Close

Quit



easily done using a standard sieve argument. Let $p_1 < p_2 < p_3 < \dots$ denote consecutive primes in the arithmetic progression $kt - 1$, $k = 1, 2, 3, \dots$. By Dirichlet's theorem, the sum $\sum_{j=1}^{\infty} 1/p_j$ diverges. Thus for each $\varepsilon > 0$, we can pick $s \in \mathbb{N}$ for which $\prod_{j=1}^s (1 - 1/p_j) < \varepsilon/2$. Further, for each $N \geq P := p_1 p_2 \dots p_s$, select a unique $k \in \mathbb{N}$ for which $kP \leq N < (k+1)P$. The number of positive integers $n \leq N$ without prime divisors in the set $\{p_1, \dots, p_s\}$ does not exceed the number of such positive integers in the interval $[1, (k+1)P]$. The latter, by the inclusion-exclusion principle, is equal to

$$(k+1)P \prod_{j=1}^s \left(1 - \frac{1}{p_j}\right) \leq \frac{(k+1)P\varepsilon}{2} \leq \frac{(1+1/k)N\varepsilon}{2} \leq \frac{2N\varepsilon}{2} = N\varepsilon.$$

This implies the claim.

Coming back to Conjecture 5, by calculation with C++, in the range $[1, 2 \cdot 10^9]$ we found only three exceptional integers 6, 36, 3600 which do not belong to the set $E(3)$. So we conjecture that

$$E(3) = \mathbb{N} \setminus \{6, 36, 3600\} \quad \text{and} \quad C(3) = 3600.$$

For $t = 4$, we have

$$288, 336, 4545, \mathbb{N}^2 \in \mathbb{N} \setminus E(4),$$

and we conjecture that $C(4) = 4545$.

There are much more integers which do not lie in $E(5)$. In the range $[1, 2 \cdot 10^9]$ there are 48 such integers:

1, 2, 5, 6, 10, 12, 20, 21, 30, 32, 45, 46, 50, 60, 92, 102, 105, 126, 141, 182,
192, 210, 282, 320, 330, 366, 406, 600, 650, 726, 732, 842, 846, 920, 992, 1020,
1446, 1452, 1905, 1920, 2100, 2250, 2262, 3962, 7320, 9050, 11520, 40500.

We conjecture that this list is full, i.e., $C(5) = 40500$. The list of integers in $[1, 2 \cdot 10^9]$ which do not lie in $E(6)$ contains 108 numbers, the largest one being 684450. We are more cautious to claim




Go back

Full Screen

Close

Quit



that $C(6) = 684450$ since this number is quite large compared to the computation bound $2 \cdot 10^9$. Here is a result of our calculations with C++ for $3 \leq t \leq 9$.

t	computation bound	number of exceptions	largest exception
3	$2 \cdot 10^9$	3	3600
4	$2 \cdot 10^9$	3	4545
5	$2 \cdot 10^9$	48	40500
6	$2 \cdot 10^9$	108	684450
7	10^9	270	9673776
8	10^9	335	3701376
9	10^9	932	18481050

Table 2. .

In Table 2 for $t = 4$, all squares k^2 are excluded, whereas for $t = 8$, all squares k^2 and all numbers of the form $2k^2$ are excluded (see Corollary 4 and Conjecture 5).

Acknowledgment. We thank the referee for some useful comments.

1. Bello-Hernández M., Benito M. and Fernández E., *On Egyptian fractions*, preprint at [arXiv:1010.2035v1](https://arxiv.org/abs/1010.2035v1), 2010.
2. Browning T.D. and Elsholtz C., *The number of representations of rationals as a sum of unit fractions*, Illinois J. Math. (to appear).
3. Buell D. A. and Hudson R. H., *On runs of consecutive quadratic residues and quadratic nonresidues*, BIT **24** (1984), 243–247.
4. Elsholtz C. and Tao T., *Counting the number of solutions to the Erdős-Straus equation on unit fractions*, preprint at [arXiv:1107.1010v3](https://arxiv.org/abs/1107.1010v3), 2011.




Go back

Full Screen

Close

Quit



5. Graham S. W. and Ringrose C. J., *Lower bounds for least quadratic non-residues*. In: Analytic number theory, Proc. Conf. in Honor of Paul T. Bateman, Urbana, IL, USA, 1989, Prog Math. 85, 1990, 269–309.
6. Guy R., *Unsolved problems in number theory*, 3rd. ed, Springer, New-York, 2004.
7. Montgomery H. L., *Topics in multiplicative number theory*, Lecture Notes in Mathematics 227, Springer, New York 1971.
8. Mordell L.J., *Diophantine equations*, Academic Press, London, New-York, 1969.
9. Schinzel A., *On sums of three unit fractions with polynomial denominators*, Funct. Approx. Comment. Math. **28** (2000), 187–194.
10. Sierpiński W., *Sur les décompositions de nombres rationnelle en fractions primaires*, Mathesis **65** (1956), 16–32.
11. Vaughan R. C., *On a problem of Erdős, Straus and Schinzel*, Mathematika **65** (1970), 193–198.
12. Yamamoto K., *On a conjecture of Erdős*, Mem. Fac. Sci. Kyuchu Univ. Ser. A **18** (1964), 166–167.
13. Yamamoto K., *On the Diophantine equation $\frac{4}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}$* , Mem. Fac. Sci. Kyuchu Univ. Ser. A **19** (1965), 37–47.

A. Dubickas, Department of Mathematics and Informatics, Vilnius University, Naugarduko 24, Vilnius LT-03225 Lithuania, e-mail: aivaras.novikas@mif.vu.lt

A. Novikas, Department of Mathematics and Informatics, Vilnius University, Naugarduko 24, Vilnius LT-03225 Lithuania, e-mail: arturas.dubickas@mif.vu.lt



Go back

Full Screen

Close

Quit