

ON CONDITIONS FOR UNIVALENCE OF SOME INTEGRAL OPERATORS

DANIEL BREAZ AND VIRGIL PESCAR

ABSTRACT. In this paper, we obtain new univalence conditions for the integral operators $F_{[\delta]}(z)$ and $G_{[\delta]}(z)$ of analytic functions defined in the open unit disk.

2000 *Mathematics Subject Classification:* 30C45.

Keywords: Integral operator, univalence, integer part, modulus.

1. INTRODUCTION

Let \mathcal{A} denote the class of functions of the form $f(z) = z + a_2z^2 + \dots$ which are analytic in the open unit disk $\mathcal{U} = \{z : |z| < 1\}$. Further, by \mathcal{S} we shall denote the class of all functions in \mathcal{A} which are univalent in \mathcal{U} .

Pescar [7], has obtained the following univalence criteria

Theorem 1.1.[7]. Let $\gamma \in \mathbb{C}$, $f \in \mathcal{S}$, $f(z) = z + a_2z^2 + \dots$.

If

$$\left| \frac{zf'(z) - f(z)}{zf(z)} \right| \leq 1, \quad \forall z \in \mathcal{U}$$

and

$$|\gamma| \leq \frac{1}{\max_{|z| \leq 1} \left[(1 - |z|^2) \cdot |z| \cdot \frac{|z| + |a_2|}{1 + |a_2||z|} \right]},$$

then

$$F_\gamma(z) = \int_0^z \left(\frac{f(t)}{t} \right)^\gamma dt$$

is in the class \mathcal{S} .

Theorem 1.2.[7]. Let $\alpha, \beta, \gamma \in \mathbb{C}$, $f \in \mathcal{S}$, $f(z) = z + a_2z^2 + \dots$

If

$$\left| \frac{zf'(z) - f(z)}{zf(z)} \right| \leq 1, \quad \forall z \in \mathcal{U},$$

$$\operatorname{Re}\beta \geq \operatorname{Re}\alpha > 0$$

and

$$|\gamma| \leq \frac{1}{\max_{|z| \leq 1} \left[\frac{1-|z|^{2\operatorname{Re}\alpha}}{\operatorname{Re}\alpha} \cdot |z| \cdot \frac{|z|+|a_2|}{1+|a_2||z|} \right]},$$

then

$$G_{\beta,\gamma}(z) = \left[\beta \int_0^z t^{\beta-1} \left(\frac{f(t)}{t} \right)^\gamma dt \right]^{\frac{1}{\beta}}$$

is in the class \mathcal{S} .

We define the next two integral operators

$$F_{[\delta]}(z) = \int_0^z \left(\frac{f_1(t)}{t} \right)^{\alpha_1} \cdot \dots \cdot \left(\frac{f_{[\delta]}(t)}{t} \right)^{\alpha_{[\delta]}} dt,$$

where $\delta \in \mathbb{C}$, $|\delta| \notin [0, 1)$, $\alpha_i \in \mathbb{C}$, $f_i \in \mathcal{A}$, $i = \overline{1, [\delta]}$, $\alpha_1 \cdot \dots \cdot \alpha_{[\delta]} = \delta$ and

$$G_{[\gamma]}(z) = \left[\gamma \int_0^z t^{\gamma-1} \left(\frac{f_1(t)}{t} \right)^{\alpha_1} \cdot \dots \cdot \left(\frac{f_{[\gamma]}(t)}{t} \right)^{\alpha_{[\gamma]}} dt \right]^{\frac{1}{\gamma}},$$

$\gamma \in \mathbb{C}$, $|\gamma| \notin [0, 1)$, $\alpha_i \in \mathbb{C}$, $f_i \in \mathcal{A}$, $i = \overline{1, [\gamma]}$, $\alpha_1 \cdot \dots \cdot \alpha_{[\gamma]} = \gamma$.

In this paper, we obtain new univalence conditions for the integral operators $F_{[\delta]}(z)$ and $G_{[\delta]}(z)$.

2. PRELIMINARY RESULTS

In order to derive our main results, we have to recall here the following lemmas:

Lemma 2.1.[2]. If the function f is regular in unit disk \mathcal{U} , $f(z) = z + a_2z^2 + \dots$ and

$$(1 - |z|^2) \left| \frac{zf''(z)}{f'(z)} \right| \leq 1,$$

for all $z \in \mathcal{U}$, then the function f is univalent in \mathcal{U} .

Lemma 2.2. [5]. Let α be a complex number, $\operatorname{Re}\alpha > 0$ and $f(z) = z + a_2z^2 + \dots$ be a regular function in \mathcal{U} . If

$$\frac{1 - |z|^{2\operatorname{Re}\alpha}}{\operatorname{Re}\alpha} \left| \frac{zf''(z)}{f'(z)} \right| \leq 1,$$

for all $z \in \mathcal{U}$, then for any complex number β , $\operatorname{Re}\beta \geq \operatorname{Re}\alpha$, the function

$$F_\beta(z) = \left[\beta \int_0^z t^{\beta-1} f'(t) dt \right]^{\frac{1}{\beta}}$$

is in the class \mathcal{S} .

Lemma 2.3. [3]. If the function g is regular in \mathcal{U} and $|g(z)| < 1$ in \mathcal{U} , then for all $\xi \in \mathcal{U}$, the following inequalities hold

$$\left| \frac{g(\xi) - g(z)}{1 - \overline{g(z)}g(\xi)} \right| \leq \left| \frac{\xi - z}{1 - \bar{z}\xi} \right| \quad (2.1)$$

and

$$|g'(z)| \leq \frac{1 - |g(z)|^2}{1 - |z|^2},$$

the equalities hold in the case $g(z) = \epsilon \frac{z+u}{1+uz}$, where $|\epsilon| = 1$ and $|u| < 1$.

Remark 2.4. [3]. For $z = 0$, from inequality (2.1) we obtain for every $\xi \in \mathcal{U}$,

$$\left| \frac{g(\xi) - g(0)}{1 - \overline{g(0)}g(\xi)} \right| \leq |\xi|$$

and hence,

$$|g(\xi)| \leq \frac{|\xi| + |g(0)|}{1 + \overline{g(0)}g(\xi)}.$$

Considering $g(0) = a$ and $\xi = z$, then

$$|g(z)| \leq \frac{|z| + |a|}{1 + |a||z|},$$

for all $z \in \mathcal{U}$.

3. MAIN RESULTS

Theorem 3.1. Let $M > 1$, $\delta \in \mathbb{C}$, $|\delta| \notin [0, 1)$, $\alpha_i \in \mathbb{C}$, for $i = \overline{1, [\lceil \delta \rceil]}$ and $\alpha_1 \cdots \alpha_{[\lceil \delta \rceil]} = \delta$. If $f_i \in \mathcal{A}$, $f_i(z) = z + a_2^i z^2 + \dots$, for $i = \overline{1, [\lceil \delta \rceil]}$ and

$$\left| \frac{zf'_i(z) - f_i(z)}{zf_i(z)} \right| \leq 1, \quad \forall i = \overline{1, [\lceil \delta \rceil]}, \quad z \in \mathcal{U}, \quad (3.1)$$

$$\frac{|\alpha_1| + \dots + |\alpha_{[\lceil \delta \rceil]}|}{|\alpha_1 \cdots \alpha_{[\lceil \delta \rceil]}|} \leq M, \quad (3.2)$$

$$|\alpha_1 \cdots \alpha_{[\lceil \delta \rceil]}| \leq \frac{1}{M \max_{|z| \leq 1} \left[(1 - |z|^2) \cdot |z| \cdot \frac{|z| + |c|}{1 + |c||z|} \right]}, \quad (3.3)$$

where

$$|c| = \frac{\left| \alpha_1 a_2^1 + \dots + \alpha_{[\lceil \delta \rceil]} a_2^{[\lceil \delta \rceil]} \right|}{M |\alpha_1 \cdots \alpha_{[\lceil \delta \rceil]}|},$$

then

$$F_{[\lceil \delta \rceil]}(z) = \int_0^z \left(\frac{f_1(t)}{t} \right)^{\alpha_1} \cdots \left(\frac{f_{[\lceil \delta \rceil]}(t)}{t} \right)^{\alpha_{[\lceil \delta \rceil]}} dt$$

is in the class \mathcal{S} .

Proof. We have $f_i \in \mathcal{A}$, for all $i = \overline{1, [\lceil \delta \rceil]}$ and $\frac{f_i(z)}{z} \neq 0$, for all $i = \overline{1, [\lceil \delta \rceil]}$.

Let g be the function $g(z) = \left(\frac{f_1(z)}{z} \right)^{\alpha_1} \cdots \left(\frac{f_{[\lceil \delta \rceil]}(z)}{z} \right)^{\alpha_{[\lceil \delta \rceil]}}$, $z \in \mathcal{U}$. We have $g(0) = 1$.

Consider the function

$$h(z) = \frac{1}{M |\alpha_1 \cdots \alpha_{[\lceil \delta \rceil]}|} \cdot \frac{F''_{[\lceil \delta \rceil]}(z)}{F'_{[\lceil \delta \rceil]}(z)}, \quad z \in \mathcal{U}.$$

The function $h(z)$ has the form:

$$h(z) = \frac{1}{M |\alpha_1 \cdots \alpha_{[\lceil \delta \rceil]}|} \sum_{i=1}^{[\lceil \delta \rceil]} \alpha_i \frac{zf'_i(z) - f_i(z)}{zf_i(z)}.$$

Also,

$$h(0) = \frac{1}{M |\alpha_1 \cdots \alpha_{[\lceil \delta \rceil]}|} \sum_{i=1}^{[\lceil \delta \rceil]} \alpha_i a_2^i.$$

By using the relations (3.1) and (3.2) we obtain that $|h(z)| < 1$ and

$$|h(0)| = \frac{\left| \alpha_1 a_2^1 + \dots + \alpha_{[\delta]} a_2^{[\delta]} \right|}{M |\alpha_1 \cdot \dots \cdot \alpha_{[\delta]}|} = |c|.$$

Applying Remark 2.4 for the function h we obtain

$$\frac{1}{M |\alpha_1 \cdot \dots \cdot \alpha_{[\delta]}|} \cdot \left| \frac{F''_{[\delta]}(z)}{F'_{[\delta]}(z)} \right| \leq \frac{|z| + |c|}{1 + |c| |z|}, \forall z \in \mathcal{U}$$

and

$$\left| \left(1 - |z|^2\right) \cdot z \cdot \frac{F''_{[\delta]}(z)}{F'_{[\delta]}(z)} \right| \leq M |\alpha_1 \cdot \dots \cdot \alpha_{[\delta]}| \left(1 - |z|^2\right) \cdot |z| \cdot \frac{|z| + |c|}{1 + |c| |z|}, \forall z \in \mathcal{U}. \quad (3.4)$$

Consider the function $H : [0, 1] \rightarrow \mathbb{R}$ defined by

$$H(x) = (1 - x^2)x \frac{x + |c|}{1 + |c| x}; \quad x = |z|.$$

We have

$$H\left(\frac{1}{2}\right) = \frac{3}{8} \cdot \frac{1 + 2|c|}{2 + |c|} > 0 \Rightarrow \max_{x \in [0, 1]} H(x) > 0.$$

Using this result and from (3.4) we have:

$$\left| \left(1 - |z|^2\right) \cdot z \cdot \frac{F''_{[\delta]}(z)}{F'_{[\delta]}(z)} \right| \leq M |\alpha_1 \cdot \dots \cdot \alpha_{[\delta]}| \max_{|z| < 1} \left[\left(1 - |z|^2\right) \cdot |z| \cdot \frac{|z| + |c|}{1 + |c| |z|} \right], \forall z \in \mathcal{U}. \quad (3.5)$$

Applying the condition (3.3) in the form (3.5) we obtain that

$$\left(1 - |z|^2\right) \cdot \left| z \cdot \frac{F''_{[\delta]}(z)}{F'_{[\delta]}(z)} \right| \leq 1, \forall z \in \mathcal{U},$$

and from Lemma 2.1 we obtain that $F_{[\delta]} \in \mathcal{S}$.

Theorem 3.2. Let $M > 1$, $\gamma, \delta \in \mathbb{C}$, $|\gamma| \notin [0, 1]$, $\alpha_i \in \mathbb{C}$, for $i = \overline{1, [\gamma]}$, $\alpha_1 \cdot \dots \cdot \alpha_n = [\gamma]$. If $f_i \in \mathcal{A}$, $f_i(z) = z + a_2^i z^2 + \dots$, for $i = \overline{1, [\gamma]}$ and

$$\left| \frac{zf'_i(z) - f_i(z)}{zf_i(z)} \right| \leq 1, \quad \forall i = \overline{1, [\gamma]}, \quad z \in \mathcal{U}, \quad (3.6)$$

$$\frac{|\alpha_1| + \dots + |\alpha_{[\gamma]}|}{|\alpha_1 \cdot \dots \cdot \alpha_{[\gamma]}|} \leq M, \quad (3.7)$$

$$\operatorname{Re}\gamma \geq \operatorname{Re}\delta > 0,$$

$$|\alpha_1 \cdot \dots \cdot \alpha_{[\gamma]}| \leq \frac{1}{M \max_{|z| \leq 1} \left[(1 - |z|^2) \cdot |z| \cdot \frac{|z| + |c|}{1 + |c||z|} \right]}, \quad (3.8)$$

where

$$|c| = \frac{\left| \alpha_1 a_2^1 + \dots + \alpha_{[\gamma]} a_2^{[\gamma]} \right|}{M |\alpha_1 \cdot \dots \cdot \alpha_{[\gamma]}|},$$

then

$$G_{[\gamma]}(z) = \left[\gamma \int_0^z t^{\gamma-1} \left(\frac{f_1(t)}{t} \right)^{\alpha_1} \cdot \dots \cdot \left(\frac{f_{[\gamma]}(t)}{t} \right)^{\alpha_{[\gamma]}} dt \right]^{\frac{1}{\gamma}}$$

is in the class \mathcal{S} .

Proof. We consider the function

$$h(z) = \int_0^z \left(\frac{f_1(t)}{t} \right)^{\alpha_1} \cdot \dots \cdot \left(\frac{f_{[\gamma]}(t)}{t} \right)^{\alpha_{[\gamma]}} dt.$$

Define the function

$$p(z) = \frac{1}{M |\alpha_1 \cdot \dots \cdot \alpha_{[\gamma]}|} \cdot \frac{h''(z)}{h'(z)}, \quad z \in \mathcal{U}.$$

The function $p(z)$ has the form:

$$p(z) = \frac{1}{M |\alpha_1 \cdot \dots \cdot \alpha_{[\gamma]}|} \sum_{i=1}^{[\gamma]} \alpha_i \frac{zf'_i(z) - f_i(z)}{zf_i(z)}.$$

By using the relations (3.6) and (3.7) we obtain $|p(z)| < 1$ and

$$|p(0)| = \frac{\left| \alpha_1 a_2^1 + \dots + \alpha_{[\gamma]} a_2^{[\gamma]} \right|}{M |\alpha_1 \cdot \dots \cdot \alpha_{[\gamma]}|} = |c|.$$

Applying Remark 2.4 for the function h we obtain

$$\frac{1}{M |\alpha_1 \cdot \dots \cdot \alpha_{[\gamma]}|} \cdot \left| \frac{h''(z)}{h'(z)} \right| \leq \frac{|z| + |c|}{1 + |c||z|}, \quad \forall z \in \mathcal{U}$$

and

$$\left| \frac{1 - |z|^{2\operatorname{Re}\delta}}{\operatorname{Re}\delta} \cdot z \cdot \frac{h''(z)}{h'(z)} \right| \leq M |\alpha_1 \cdot \dots \cdot \alpha_{[\gamma]}| \frac{1 - |z|^{2\operatorname{Re}\delta}}{\operatorname{Re}\delta} \cdot |z| \cdot \frac{|z| + |c|}{1 + |c||z|}, \forall z \in \mathcal{U}. \quad (3.9)$$

Consider the function $Q : [0, 1] \rightarrow \mathbb{R}$ defined by

$$Q(x) = \frac{1 - x^{2\operatorname{Re}\delta}}{\operatorname{Re}\delta} \cdot x \cdot \frac{x + |c|}{1 + |c|x}; \quad x = |z|.$$

We have $Q\left(\frac{1}{2}\right) > 0 \Rightarrow \max_{x \in [0,1]} Q(x) > 0$.

Using this result in (3.9), we have:

$$\frac{1 - |z|^{2\operatorname{Re}\delta}}{\operatorname{Re}\delta} \left| \frac{zh''(z)}{h'(z)} \right| \leq M |\alpha_1 \cdot \dots \cdot \alpha_{[\gamma]}| \cdot \max_{|z| < 1} \left[\frac{1 - |z|^{2\operatorname{Re}\delta}}{\operatorname{Re}\delta} \cdot |z| \cdot \frac{|z| + |c|}{1 + |c||z|} \right], \forall z \in \mathcal{U}. \quad (3.10)$$

Applying the condition (3.8) in the relation (3.10), we obtain that

$$\frac{1 - |z|^{2\operatorname{Re}\delta}}{\operatorname{Re}\delta} \left| \frac{zh''(z)}{h'(z)} \right| \leq 1, \forall z \in \mathcal{U}$$

and from Lemma 2.2, we obtain that $G_{[\gamma]} \in \mathcal{S}$.

REFERENCES

- [1] D. Breaz, N. Breaz, *Two integral operators*, Studia Universitatis "Babeş-Bolyai", Mathematica, Cluj-Napoca, No.3, 2002, 13-21.
- [2] J. Becker, *Lownersche Differentialgleichung und quasikonform fortsetzbare schlichte Functionen*, J. Reine Angew. Math. 255(1972), 23-43.
- [3] G. M. Goluzin, *Geometricheskaya teoria funktsii kompleksnogo peremenogo*, Moscow, 1966.
- [4] Y.J. Kim, E.P. Merkes, *On an integral of powers of a spirallike function*, Kyungpook Math., J., 12 (1972), 2, 249-253.
- [5] N.N. Pascu, *An improvement of Becker's Univalence Criterion*, Proceedings of the Commemorative Session Simion Stoilow, University of Braşov, 1987, 43-48.
- [6] N.N. Pascu, V.Pescar, *On the integral operators of Kim-Merkes and Pfaltzgraff*, Mathematica, Univ. "Babeş-Bolyai", Cluj-Napoca, 32 (55), 2(1990), 185-192.

[7] V. Pescar, *On some integral operations which preserve the univalence*, Journal of Mathematics, Vol. XXX(1997), pp. 1-10, Punjab University.

Daniel Breaz
Department of Mathematics,
”1 Decembrie 1918” University of Alba Iulia
510009, Alba Iulia, Romania
email: dbreaz@uab.ro

Virgil Pescar
Department of Mathematics
”Transilvania” University of Brașov
Faculty of Mathematics and Computer Science
500091 Brașov, Romania
email: virgilpescar@unitbv.ro