

**STRONG CONSISTENCY RATES AND ASYMPTOTIC  
NORMALITY OF CONDITIONAL HAZARD FUNCTION FOR  
FUNCTIONAL DATA**

N. BELKHIR, A. RABHI, S. SOLTANI

**ABSTRACT.** We consider the problem of nonparametric estimation of the conditional hazard function for functional mixing data. More precisely, given a strictly stationary random variables  $Z_i = (X_i, Y_i)_{i \in \mathbb{N}}$ , we investigate a kernel estimate of the conditional hazard function of univariate response variable  $Y_i$  given the functional variable  $X_i$ . The principal aim of this paper is to give the mean squared convergence rate and to prove the asymptotic normality of the proposed estimator.

2010 *Mathematics Subject Classification*: 62E20, 62F12, 62H12, 62N02.

*Keywords*: Asymptotic normality, Functional data, Kernel conditional hazard function, Nonparametric estimation, Small ball probability, Strong mixing process.

1. INTRODUCTION

The statistical problems involved in the modelization of functional data have received an increasing interest in the literature. The infatuation for this topic is linked with many fields of applications in which the data are collected in the functional order. Under this supposition, the statistical analysis focuses on a framework of infinite dimension for the data under study. This field of modern statistics has received much attention in the last 20 years, and it has been popularized in the book of Ramsay and Silverman [23]. This type of data appears in many fields of applied statistics: environmetrics [8], chemometrics [2], meteorological sciences [3], etc..

In this paper, we are interested in the nonparametric estimation of the conditional hazard function when the covariates are of functional nature.

The nonparametric estimation of the hazard and/or the conditional hazard function is quite important in a variety of fields including medicine, reliability, survival analysis or in seismology. The literature on this model in multivariate statistics is abundant. Historically, the hazard estimate was introduced by Watson and Leadbetter [30], since, several results have been added, see for example, Roussas [26] (for previous works).

From a theoretical point of view, a sample of functional data can be involved in many different statistical problems, such as for example: classification and principal components analysis (PCA)[4, 24] or longitudinal studies, regression and prediction [2, 7].

The recent monograph by Ferraty and Vieu [13] summarizes many of their contributions to the non-parametric estimation with functional data; among other properties, consistency of the conditional density, conditional distribution and regression estimates are established in the i.i.d. case as well as under dependence conditions (strong mixing). Almost complete rates of convergence are also obtained, and the different techniques are applied to several examples of functional data samples. Related work can be seen in the paper of Masry [19], where the asymptotic normality of the functional non-parametric regression estimate is proven, considering strong mixing dependence conditions for the sample data. For automatic smoothing parameter selection in the regression setting, see Rachdi and Vieu [22].

The literature is strictly not limited in the case where the data is of functional nature (a curve). The first result in this context, was given by Ferraty *et al.* [12]. They established the almost complete convergence of the kernel estimate of the conditional hazard function in the i.i.d. case and under  $\alpha$ -mixing condition. Recently, Rabhi *et al.* [21] studied the mean quadratic convergence in the i.i.d. case of this estimate. More recently Mahiddine *et al.* [18] established the pointwise almost complete convergence and the uniform almost complete convergence (with the rate) of the kernel estimate in the single-index models in the i.i.d. case.

The estimation of the hazard function is a problem of considerable interest, especially to inventory theorists, medical researchers, logistics planners, reliability engineers and seismologists. The non-parametric estimation of the hazard function has been extensively discussed in the literature. Beginning with Watson and Leadbetter [30], there are many papers on these topics: Ahmad [1], Singpurwalla and Wong [27], etc. We can cite Quintela [20] for a survey.

When hazard rate estimation is performed with multiple variables, the result is an estimate of the conditional hazard rate for the first variable, given the levels of the remaining variables. Many references, practical examples and simulations in the case of non-parametric estimation using local linear approximations can be found in Spierdijk [28].

The main aim of this paper, is to study, under general conditions, the asymptotic properties of the functional data kernel estimate of the conditional hazard function introduced by Ferraty *et al.* [12]. More precisely, we treat the  $L^2$ -convergence rate by giving the exact expression involved in the leading terms of the quadratic error. In addition, we establish the asymptotic normality of the construct estimator. We point out that our asymptotic results are useful in some statistical problems such as the

choice of the smoothing parameters, the determination of confidence intervals and in risk analysis. The present work extended to dependent case the result of Rabhi *et al.* [21] given in i.i.d. case functional. We note that, one of the main difficulty, when dealing with functional variables, relies on the difficulty for choosing some appropriate measure of reference in infinite dimensional spaces. The main feature of our approach is to build estimates and to derive their asymptotic properties without any notion of density for the functional variable  $X$ . This approach allows us to avoid the use of a reference measure in such functional spaces. In each of the above described sections, we will give general asymptotic results without assuming existence of such a density, and each of these results will be discussed in relation with earlier literature existing in the usual finite dimensional case.

Our paper presents some asymptotic properties related with the non-parametric estimation of the conditional hazard function. In a functional data setting, the conditioning variable is allowed to take its values in some abstract semi-metric space. In this case, Ferraty *et al.* [29] define non-parametric estimators of the conditional density and the conditional distribution. They give the rates of convergence (in an almost complete sense) to the corresponding functions, in an  $\alpha$ -mixing) context. In Rabhi *et al.* [21], the same properties are shown in an i.i.d. context in the data sample. We extend their results to dependent case by calculating the bias and variance of these estimates, and establishing their asymptotic normality, considering a particular type of kernel for the functional part of the estimates. Because the hazard function estimator is naturally constructed using these two last estimators, the same type of properties is easily derived for it. Our results are valid in a real (one- and multi-dimensional) context.

The paper is organized as follows: the next section we present our model. Section 3 is dedicated to fixing notations and hypotheses. We state our main results in Section 4. The Section 5 is devoted to some discuss on the applicability of our asymptotic result in some statistical problems such as the choice of the smoothing parameters, the determination of confidence intervals and in seismology analysis.

## 2. THE MODEL

Consider  $Z_i = (X_i, Y_i)$ ,  $i \in \mathbb{N}$  be a  $\mathcal{F} \times \mathbb{R}$ -valued measurable strictly stationary process, defined on a probability space  $(\Omega, \mathcal{A}, \mathbb{P})$ , where  $(\mathcal{F}, d)$  is a semi-metric space.

In the following  $x$  will be a fixed point in  $\mathcal{F}$  and  $N_x$  will denote a fixed neighborhood of  $x$ . We assume that the regular version of the conditional probability of  $Y$  given  $X$  exists. Moreover, we suppose that, for all  $z \in N_x$  the conditional distribution function of  $Y$  given  $X = z$ ,  $F^z(\cdot)$ , is 3-times continuously differentiable and we denote by  $f^z$  its conditional density with respect to (w.r.t.) Lebesgue's measure

over  $\mathbb{R}$ . In this paper, we consider the problem of the nonparametric estimation of the conditional hazard function defined, for all  $y \in \mathbb{R}$  such that  $F^x(y) < 1$ , by

$$h^x(y) = \frac{f^x(y)}{1 - F^x(y)}.$$

In our spatial context, we estimate this function by

$$\widehat{h}^x(y) = \frac{\widehat{f}^x(y)}{1 - \widehat{F}^x(y)}$$

where

$$\widehat{F}^x(y) = \frac{\sum_{i=1}^n K(h_K^{-1}d(x, X_i))H(h_H^{-1}(y - Y_i))}{\sum_{i=1}^n K(h_K^{-1}d(x, X_i))}, \quad \forall y \in \mathbb{R}$$

and

$$\widehat{f}^x(y) = \frac{h_H^{-1} \sum_{i=1}^n K(h_K^{-1}d(x, X_i))H'(h_H^{-1}(y - Y_i))}{\sum_{i=1}^n K(h_K^{-1}d(x, X_i))}, \quad \forall y \in \mathbb{R}$$

with  $K$  is the kernel,  $H$  is a given continuously differentiable distribution function,  $h_K = h_{K,n}$  (resp.  $h_H = h_{H,n}$ ) is a sequence of positive real numbers and  $H'$  is the derivative of  $H$ . Furthermore, the estimator  $\widehat{h}^x(y)$  can be written as

$$\widehat{h}^x(y) = \frac{\widehat{f}_N^x(y)}{\widehat{F}_D^x - \widehat{F}_N^x(y)} \tag{1}$$

where

$$\begin{aligned} \widehat{F}_D^x &:= \frac{1}{n\mathbb{E}[K_1]} \sum_{i=1}^n K(h_K^{-1}d(x, X_i)), \quad K_1 = K(h_K^{-1}d(x, X_1)) \\ \widehat{F}_N^x(y) &:= \frac{1}{n\mathbb{E}[K_1]} \sum_{i=1}^n K(h_K^{-1}d(x, X_i))H(h_H^{-1}(y - Y_i)) \\ \widehat{f}_N^x(y) &:= \frac{1}{nh_H\mathbb{E}[K_1]} \sum_{i=1}^n K(h_K^{-1}d(x, X_i))H'(h_H^{-1}(y - Y_i)). \end{aligned}$$

Our main purpose is to study the  $L^2$ -consistency and the asymptotic normality of the nonparametric estimate  $\widehat{h}^x$  of  $h^x$  when the random filed  $(Z_i, i \in \mathbb{N})$  satisfies the following mixing condition.

### 3. NOTATIONS AND HYPOTHESES

All along the paper, when no confusion is possible, we will denote by  $C$  and  $C'$  some strictly positive generic constants. In order to establish our asymptotic results we need the following hypotheses:

(H0)  $\forall r > 0, \mathbb{P}(X \in B(x, r)) =: \phi_x(r) > 0$ , where  $B(x, r) = \{x' \in \mathcal{F} / d(x, x') < r\}$ .

(H1)  $(X_i, Y_i)_{i \in \mathbb{N}}$  is an  $\alpha$ -mixing sequence whose the coefficients of mixture verify:

$$\exists a > 0, \exists c > 0 : \forall n \in \mathbb{N}, \alpha(n) \leq cn^{-a}.$$

(H2)  $0 < \sup_{i \neq j} \mathbb{P}((X_i, X_j) \in B(x, h) \times B(x, h)) = \mathcal{O}\left(\frac{(\phi_x(h))^{(a+1)/a}}{n^{1/a}}\right)$ .

Note that (H0) can be interpreted as a concentration hypothesis acting on the distribution of the *f.r.v.*  $X$ , whereas (H2) concerns the behavior of the joint distribution of the pairs  $(X_i, X_j)$ . In fact, this hypothesis is equivalent to assume that, for  $n$  large enough

$$\sup_{i \neq j} \frac{\mathbb{P}((X_i, X_j) \in B(x, h) \times B(x, h))}{\mathbb{P}(X \in B(x, h))} \leq C \left(\frac{\phi_x(h)}{n}\right)^{1/a}.$$

(H3) For  $l \in \{0, 2\}$ , the functions  $\Psi_l(s) = \mathbb{E}\left[\frac{\partial^l F^X(y)}{\partial y^l} - \frac{\partial^l F^x(y)}{\partial y^l} \middle| d(x, X) = s\right]$ ,  $\Phi_l(s) = \mathbb{E}\left[\frac{\partial^l f^X(y)}{\partial y^l} - \frac{\partial^l f^x(y)}{\partial y^l} \middle| d(x, X) = s\right]$  and  $\Phi'_l(s) = \mathbb{E}\left[\frac{\partial^l f'^X(y)}{\partial y^l} - \frac{\partial^l f'^x(y)}{\partial y^l} \middle| d(x, X) = s\right]$  are derivable at  $s = 0$ .

(H4) The bandwidth  $h_K$  satisfies for  $j = 0, 1$ :

$$h_K \downarrow 0, \quad \forall t \in [0, 1] \quad \lim_{h_K \rightarrow 0} \frac{\phi_x(th_K)}{\phi_x(h_K)} = \beta_x(t) \quad \text{and} \quad nh_K^{2j+1} \phi_x(h_K) \rightarrow \infty \quad \text{as } n \rightarrow \infty,$$

(H5) The kernel  $K$  from  $\mathbb{R}$  into  $\mathbb{R}^+$  is a differentiable function supported on  $[0, 1]$ . Its derivative  $K'$  exists and is such that there exist two constants  $C$  and  $C'$  with  $-\infty < C < K'(t) < C' < 0$  for  $0 \leq t \leq 1$ .

(H6)  $H$  has even bounded derivative function supported on  $[0, 1]$  that verifies

$$\int_{\mathbb{R}} t^2 H'(t) dt < \infty \quad \text{and} \quad \int_{\mathbb{R}} |t|^{b_2} (H^{(2)})^2(t) dt < \infty.$$

(H7) There exist sequences of integers  $(u_n)$  and  $(v_n)$  increasing to infinity such that  $(u_n + v_n) \leq n$ , satisfying for  $j = 0, 1$

- (i)  $v_n = o((nh_H^{2j+1}\phi_x(h_K))^{1/2})$  and  $\left(\frac{n}{h_H^{2j+1}\phi_x(h_K)}\right)^{1/2} \alpha(v_n) \rightarrow 0$  as  $n \rightarrow 0$ ,
- (ii)  $q_n v_n = o((nh_H^{2j+1}\phi_x(h_K))^{1/2})$  and  $q_n \left(\frac{n}{h_H^{2j+1}\phi_x(h_K)}\right)^{1/2} \alpha(v_n) \rightarrow 0$  as  $n \rightarrow 0$

where  $q_n$  is the largest integer such that  $q_n(u_n + v_n) \leq n$ .

### 3.1. Remarks on the assumptions

**Remark 1.** *Assumption (H0) plays an important role in our methodology. It is known as (for small  $h$ ) the "concentration hypothesis acting on the distribution of  $X$ " in infinite-dimensional spaces. This assumption is not at all restrictive and overcomes the problem of the non-existence of the probability density function. In many examples, around zero the small ball probability  $\phi_x(h)$  can be written approximately as the product of two independent functions  $\psi(z)$  and  $\varphi(h)$  as  $\phi_x(h) = \psi(z)\varphi(h) + o(\varphi(h))$ . This idea was adopted by Masry [19] who reformulated the Gasser et al. [14] one. The increasing property of  $\phi_x(\cdot)$  implies that  $\zeta_h^x(\cdot)$  is bounded and then integrable (all the more so  $\zeta_0^x(\cdot)$  is integrable).*

*Without the differentiability of  $\phi_x(\cdot)$ , this assumption has been used by many authors where  $\psi(\cdot)$  is interpreted as a probability density, while  $\varphi(\cdot)$  may be interpreted as a volume parameter. In the case of finite-dimensional spaces, that is  $\mathcal{S} = \mathbb{R}^d$ , it can be seen that  $\phi_x(h) = C(d)h^d\psi(x) + oh^d$ , where  $C(d)$  is the volume of the unit ball in  $\mathbb{R}^d$ . Furthermore, in infinite dimensions, there exist many examples fulfilling the decomposition mentioned above. We quote the following (which can be found in Ferraty et al. [10]):*

1.  $\phi_x(h) \approx \psi(h)h^\gamma$  for some  $\gamma > 0$ .
2.  $\phi_x(h) \approx \psi(h)h^\gamma \exp\{C/h^p\}$  for some  $\gamma > 0$  and  $p > 0$ .
3.  $\phi_x(h) \approx \psi(h)/|\ln h|$ .

*The function  $\beta_h^x(\cdot)$  which intervenes in Assumption (H4) is increasing for all fixed  $h$ . Its pointwise limit  $\beta_0^x(\cdot)$  also plays a determinant role. It intervenes in all asymptotic properties, in particular in the asymptotic variance term. With simple algebra, it is possible to specify this function (with  $\beta_0(u) := \beta_0^x(u)$  in the above examples by:*

1.  $\beta_0(u) = u^\gamma$ ,
2.  $\beta_0(u) = \delta_1(u)$  where  $\delta_1(\cdot)$  is Dirac function,

3.  $\beta_0(u) = \mathbf{1}_{]0,1]}(u)$ .

Assumption (H2) is classical and permits to make the variance term negligible.

**Remark 2.** Assumptions (H3) is a regularity condition which characterize the functional space of our model and is needed to evaluate the bias.

**Remark 3.** Assumptions (H5) and (H6) are classical in functional estimation for finite or infinite dimension spaces.

### 3.2. Main results

### 3.3. Mean squared convergence

The first result concerns the  $L^2$ -consistency of  $\widehat{h}^x(y)$ .

**Theorem 1.** Under assumptions (H0)-(H6), we have

$$\mathbb{E} \left[ \widehat{h}^x(y) - h^x(y) \right]^2 = B_n^2(x, y) + \frac{\sigma_h^2(x, y)}{n h_H \phi_x(h_K)} + o(h_H^4) + o(h_K) + o\left(\frac{1}{n h_H \phi_x(h_K)}\right),$$

and

$$\int_{\mathbb{R}} \mathbb{E} \left[ \widehat{h}^x(y) - h^x(y) \right]^2 dx = \int_{\mathbb{R}} B_n^2(x, y) dx + \int_{\mathbb{R}} \frac{\sigma_h^2(x, y)}{n h_H \phi_x(h_K)} dx + o\left(\frac{1}{n h_H \phi_x(h_K)}\right),$$

where

$$B_n(x, y) = \frac{(B_H^f - h^x(y)B_H^F)h_H^2 + (B_K^f - h^x(y)B_K^F)h_K}{1 - F^x(y)}$$

with

$$\begin{aligned} B_H^f(x, y) &= \frac{1}{2} \frac{\partial^2 f^x(y)}{\partial y^2} \int t^2 H'(t) dt \\ B_K^f(x, y) &= h_K \Phi'_0(0) \frac{\left( K(1) - \int_0^1 (sK(s))' \beta_x(s) ds \right)}{\left( K(1) - \int_0^1 K'(s) \beta_x(s) ds \right)} \\ B_H^F(x, y) &= \frac{1}{2} \frac{\partial^2 F^x(y)}{\partial y^2} \int t^2 H'(t) dt \\ B_K^F(x, y) &= h_K \Psi'_0(0) \frac{\left( K(1) - \int_0^1 (sK(s))' \beta_x(s) ds \right)}{\left( K(1) - \int_0^1 K'(s) \beta_x(s) ds \right)}. \end{aligned}$$

and

$$\sigma_h^2(x, y) = \frac{\beta_2 h^x(y)}{(\beta_1^2 (1 - F^x(y)))} \quad (\text{with } \beta_j = K^j(1) - \int_0^1 (K^j)'(s) \beta_x(s) ds, \text{ for } j = 1, 2),$$

*Proof.* By using the same decomposition used in ( Theorem 3.1 Rabhi *et al.* [21], P.408), we show that the proof of Theorem 1 can be deduced from the following intermediates results:

**Lemma 2.** *Under the hypotheses of Theorem (1), we have*

$$\mathbb{E} \left[ \widehat{f}_N^x(y) \right] - f^x(y) = B_H^f(x, y)h_H^2 + B_K^f(x, y)h_K + o(h_H^2) + o(h_K)$$

and

$$\mathbb{E} \left[ \widehat{F}_N^x(y) \right] - F^x(y) = B_H^F(x, y)h_H^2 + B_K^F(x, y)h_K + o(h_H^2) + o(h_K).$$

**Remark 4.** *Observe that, the result of this lemma permits to write*

$$\left[ \mathbb{E} \widehat{F}_N^x(y) - F^x(y) \right] = \mathcal{O}(h_H^2) + \mathcal{O}(h_K)$$

and

$$\left[ \mathbb{E} \widehat{f}_N^x(y) - f^x(y) \right] = \mathcal{O}(h_H^2 + h_K).$$

**Lemma 3.** *Under the hypotheses of Theorem (1), we have*

$$\text{Var} \left[ \widehat{f}_N^x(y) \right] = \frac{\sigma_f^2(x, y)}{nh_H\phi_x(h_K)} + o\left(\frac{1}{nh_H\phi_x(h_K)}\right),$$

$$\text{Var} \left[ \widehat{F}_N^x(y) \right] = o\left(\frac{1}{nh_H\phi_x(h_K)}\right)$$

and

$$\text{Var} \left[ \widehat{F}_D^x \right] = o\left(\frac{1}{nh_H\phi_x(h_K)}\right).$$

where  $\sigma_f^2(x, y) := f^x(y) \int H^2(t)dt$ .

**Lemma 4.** *Under the hypotheses of Theorem (1), we have*

$$\text{Cov}(\widehat{f}_N^x(y), \widehat{F}_D^x) = o\left(\frac{1}{nh_H\phi_x(h_K)}\right),$$

$$\text{Cov}(\widehat{f}_N^x(y), \widehat{F}_N^x(y)) = o\left(\frac{1}{nh_H\phi_x(h_K)}\right)$$

and

$$\text{Cov}(\widehat{f}_D^x, \widehat{F}_N^x(y)) = o\left(\frac{1}{nh_H\phi_x(h_K)}\right).$$

**Remark 5.** *It is clear that, the results of Lemmas (3 and 4) allows to write*

$$\text{Var} \left[ \widehat{F}_D^x - \widehat{F}_N^x \right] = o\left(\frac{1}{nh_H\phi_x(h_K)}\right)$$



### 3.4. Asymptotic normality

This section contains results on the asymptotic normality of  $\widehat{h}^x(y)$  and  $\widehat{h}'^x(y)$ . Let us assume that  $h^Z$  is sufficiently smooth ( at least of class  $\mathcal{C}^2$ )

We can write an estimator of the first derivative of the hazard function through the first derivative of the estimator.

It is therefore natural to try to construct an estimator of the derivative of the function  $h^X$  on the basis of these ideas. To estimate the conditional distribution function and the conditional density function in the presence of functional conditional random variable  $X$ .

The kernel estimator of the derivative of the function conditional random functional  $h^Z$  can therefore be constructed as follows:

$$\widehat{h}'^X(y) = \frac{\widehat{f}'^X(y)}{1 - \widehat{F}^Y(y)} + (\widehat{h}^X(y))^2, \quad (2)$$

the estimator of the derivative of the conditional density is given in the following formula:

$$\widehat{f}'^X(y) = \frac{\sum_{i=1}^n h_H^{-2} K(h_K^{-1}d(X, X_i))H''(h_H^{-1}(y - Y_i))}{\sum_{i=1}^n K(h_K^{-1}d(X, X_i))} \quad (3)$$

Later, we need assumptions on the parameters of the estimator, ie on  $K, H, H', h_H$  and  $h_K$  are little restrictive. Indeed, on one hand, they are not specific to the problem estimate of  $h^X$  (but inherent problems of  $F^X, f^X$  and  $f'^X$  estimation), and secondly they consist with the assumptions usually made under functional variables.

To obtain the asymptotic normality of the conditional estimates, we have to add the following assumptions:

(H8)  $H'$  is twice differentiable.

(H9) The bandwidth  $h_H$  and  $h_K$ , small ball probability  $\phi_z(h)$  and arithmetical  $\alpha$  mixing coefficient with order  $a > 3$  satisfying

$$\left\{ \begin{array}{l} \text{(H9a)} \exists C > 0, h_H^{2j+1} \phi_z(h_K) \geq \frac{C}{n^{2/(a+1)}}, \text{ for } j = 0, 1 \\ \text{(H9b)} \left( \frac{\phi_z(h_K)}{n} \right)^{1/a} + \phi_z(h_K) = o\left( \frac{1}{n^{2/(a+1)}} \right), \text{ for } j = 0, 1 \\ \text{(H9c)} \lim_{n \rightarrow \infty} h_K = 0, \lim_{n \rightarrow \infty} h_H = 0, \text{ and } \lim_{n \rightarrow \infty} \frac{\log n}{nh_H^{2j+1} \phi_x(h_K)} = 0, j = 0, 1; \end{array} \right.$$

**Theorem 5.** Assume that (H0)-(H7) hold, and if the following inequalities

$$\exists \eta > 0, C, C' > 0 \text{ such that } C n^{\frac{3-a}{a+1}+\eta} \leq h_H \phi_x(h_K) \text{ and } \phi_x(h_K) \leq C' n^{\frac{1}{1-a}} \quad (4)$$

are verified with  $a > (5 + \sqrt{17})/2$ , then we have for any  $x \in \mathcal{A}$ ,

$$\left( \frac{nh_H \phi_x(h_K)}{\sigma_h^2(x, y)} \right)^{1/2} \left( \widehat{h}^x(y) - h^x(y) - B_n(x, y) \right) \xrightarrow{\mathcal{D}} \mathcal{N}(0, 1) \quad \text{as } n \rightarrow \infty.$$

where

$$\mathcal{A} = \{x \in \mathcal{F}, f^x(y)(1 - F^x(y)) \neq 0\}$$

and  $\xrightarrow{\mathcal{D}}$  means the convergence in distribution.

Obviously, if one imposes some additional assumptions on the function  $\phi_x(\cdot)$  and the bandwidth parameters ( $h_K$  and  $h_H$ ) we can improved our asymptotic normality by removing the bias term  $B_n(x, y)$ .

**Corollary 6.** Under the hypotheses of Theorem 5 and if the bandwidth parameters ( $h_K$  and  $h_H$ ) and if the function  $\phi_x(h_K)$  satisfies:

$$\lim_{n \rightarrow \infty} (h_H^2 + h_K) \sqrt{n \phi_x(h_K)} = 0$$

we have

$$\left( \frac{nh_H \phi_x(h_K)}{\sigma_h^2(x, y)} \right)^{1/2} \left( \widehat{h}^x(y) - h^x(y) \right) \xrightarrow{\mathcal{D}} \mathcal{N}(0, 1) \quad \text{as } n \rightarrow \infty.$$

*Proof. Proof of Theorem 5 and Corollary 6* We consider the decomposition

$$\begin{aligned} \widehat{h}^x(y) - h^x(y) &= \frac{1}{\widehat{F}_D^x - \widehat{F}_N^x(y)} \left( \widehat{f}_N^x(y) - E \widehat{f}_N^x(y) \right) \\ &+ \frac{1}{\widehat{F}_D^x - \widehat{F}_N^x(y)} \left\{ h^x(y) \left[ \mathbb{E} \widehat{F}_N^x(y) - F^x(y) \right] + \left( \mathbb{E} \widehat{f}_N^x(y) - f^x(y) \right) \right\} \\ &+ \frac{h^x(y)}{\widehat{F}_D^x - \widehat{F}_N^x(y)} \left\{ 1 - \mathbb{E} \widehat{F}_N^x(y) - \left( \widehat{F}_D^x - \widehat{F}_N^x(y) \right) \right\} \end{aligned} \quad (5)$$

Therefore, Theorem 5 and Corollary 6 are a consequence of Lemma 2, remark (4) and the following results.

**Lemma 7.** Under the hypotheses of Theorem 5

$$\left( \frac{nh_H \phi_x(h_K)}{\sigma_f^2(x, y)} \right)^{1/2} \left( \widehat{f}_N^x(y) - \mathbb{E} \left[ \widehat{f}_N^x(y) \right] \right) \rightarrow \mathcal{N}(0, 1).$$

**Lemma 8.** *Under the hypotheses of Theorem 5*

$$\widehat{F}_D^x - \widehat{F}_N^x(y) \rightarrow 1 - F^x(y) \quad \text{in probability}$$

and

$$\left( \frac{nh_H \phi_x(h_K)}{\sigma_h^2(x, y)} \right)^{1/2} \left( \widehat{F}_D^x - \widehat{F}_N^x(y) - 1 + \mathbb{E}[\widehat{F}_N^x(y)] \right) = o_{\mathbb{P}}(1).$$

**Theorem 9.** *Under conditions (H0)-(H9), then we have for any  $x \in \mathcal{A}$ ,*

$$(nh_H^3 \phi_x(h_K))^{1/2} \left( \widehat{h}'^X(y) - h'^X(y) \right) \xrightarrow{\mathcal{D}} N(0, \sigma_{h'}^2(y))$$

$$a_l^z = K^l(1) - \int_0^1 \left( K^l(u) \right)' \beta_0^z(u) du \quad \text{for } l = 1, 2$$

and

$$\sigma_{h'}^2(y) = \frac{a_2^z h^X(y)}{(a_1^z)^2 (1 - F^X(y))} \int (H''(t))^2 dt.$$

*Proof.* Let

$$\widehat{h}'^X(y) = \frac{\widehat{f}'^X(y)}{1 - \widehat{F}^X(y)} + (\widehat{h}^X(y))^2, \quad (6)$$

with

$$\widehat{h}'^X(y) - h'^X(y) = \left\{ (\widehat{h}^X(y))^2 - (h^X(y))^2 \right\} + \left\{ \frac{\widehat{f}'^X(y)}{1 - \widehat{F}^X(y)} - \frac{f'^X(y)}{1 - F^X(y)} \right\} \quad (7)$$

Using again (7), and the fact that

$$\frac{(1 - F^X(y))}{(1 - \widehat{F}^X(y))(1 - F^X(y))} \rightarrow \frac{1}{1 - F^X(y)}$$

and

$$\frac{\widehat{f}'^X(y)}{(1 - \widehat{F}^X(y))(1 - F^X(y))} \rightarrow \frac{f'^X(y)}{(1 - F^X(y))^2}$$

The asymptotic normality of  $(nh_H^3 \phi_x(h_K))^{1/2} \left( \widehat{h}'^X(y) - h'^X(y) \right)$  can be deduced from both following lemmas and Corollary 6.

**Lemma 10.** *Under Assumptions (H0)-(H6) and (H8), we have*

$$(n\phi_x(h_K))^{1/2} \left( \widehat{F}^X(y) - F^X(y) \right) \xrightarrow{\mathcal{D}} N(0, \sigma_{F^X}^2(y)) \quad (8)$$

where

$$\sigma_{F^X}^2(y) = \frac{a_2^z F^X(y) (1 - F^X(y))}{(a_1^z)^2}$$

**Lemma 11.** *Under Assumptions of Theorem 9, we have*

$$(nh_H^3 \phi_x(h_K))^{1/2} \left( \widehat{f}'^X(y) - f'^X(y) \right) \xrightarrow{\mathcal{D}} N(0, \sigma_{f'^X}^2(y)) \quad (9)$$

where

$$\sigma_{f'^X}^2(y) = \frac{a_2^z f^X(y)}{(a_1^z)^2} \int_{\mathbb{R}} (H''(t))^2 dt$$

The proofs of Lemma 10 can be seen in Ezzahrioui and Ould-Saïd [9].

#### 4. DISCUSSIONS AND APPLICATIONS

In this section we emphasize the potential impact of our work by studying its practical interest in some important statistical problems. Moreover, in order to show the easily implementation of our approach on a concrete cases, we discuss in the second part of this section the practical utilization of our model in risk analysis.

##### 4.1. Some derivatives

- *On the choices of the bandwidths parameters:* As all smoothing by a kernel method, the choice of bandwidths parameters has crucial role in determining the performance of the estimators. The mean quadratic error given in Theorem (1) is a basic ingredient to solve this problem. Usually, the ideal theoretical choices are obtained by minimizing this error. Here, we have explicated its leading term which is

$$B_n^2(x, y) + \frac{\sigma_h^2(x, y)}{nh_H \phi_x(h_K)}.$$

Then, the smoothing parameters minimizing this leading term is asymptotically optimal with respect the  $L^2$ -error. However, the practical utilization of this criterium requires some additional computational efforts. More precisely, it requires the estimation of the unknown quantities  $\Psi'_0$ ,  $\Phi'_0$ ,  $f^x(y)$  and

$F^x(y)$ . Clearly, all these estimations can be obtained by using a pilots estimators of the conditional distribution function  $F^x(y)$  and of the conditional density  $f^x(y)$ . Such estimations are possible by using the kernel methods, with a separate choice of the bandwidths parameters between both models. More preciously, for the conditional density, we propose to adopt, to the functional case, the bandwidths selectors studied by Bouraine *et al.* [6] by considering the following criterion

$$CVPDF = \frac{1}{n} \sum_{i=1} W_1(X_i) \int \hat{f}^{X_i^{-i^2}}(y) W_2(y) dy - \frac{2}{n} \sum_{i=1} \hat{f}^{X_i^{-i}}(Y_i) W_1(X_i) W_2(Y_i) \quad (10)$$

while, for the the conditional distribution function we can use the cross-validation rule proposed by De Gooijer and Gannoun (2000) (in vectorial case)

$$CVCDF = \frac{1}{n} \sum_{k,l \in I_n} \left[ \mathbf{1}_{Y_k \leq Y_l} - \hat{F}^{X_k^{-k}}(Y_l) \right]^2 W(X_k)$$

where  $W_1$ ,  $W_2$  and  $W$  are some suitable trimming functions and

$$\hat{F}^{X_k^{-k}}(Y_l) = \frac{\sum_{i \in I_{n,\varsigma_n}^{k,l}} K(h_K^{-1}d(X_k, X_i)) H(h_H^{-1}(Y_l - Y_i))}{\sum_{i \in I_{n,\varsigma_n}^{k,l}} K(h_K^{-1}d(X_k, X_i))}$$

and

$$\hat{f}^{X_i^{-i}}(y) = \frac{h_H^{-1} \sum_{j \in I_{n,\varsigma_n}^i} K(h_K^{-1}d(X_i, X_j)) H'(h_H^{-1}(y - Y_j))}{\sum_{j \in I_{n,\varsigma_n}^i} K(h_K^{-1}d(X_i, X_j))}$$

with

$$\begin{cases} I_{n,\varsigma_n}^{k,l} = \{i \text{ such that } |i - k| \geq \varsigma_n \text{ and } |i - l| \geq \varsigma_n\}, \\ I_{n,\varsigma_n}^i = \{j \text{ such that } |j - i| \geq \varsigma_n\}. \end{cases}$$

Of course, we can also adopt another selection methods, such that, the parametric bootstrap method, proposed by Hall *et al.* [15] and Hyndman *et al.* [16] for, respectively, the conditional cumulative distribution function and the conditional density in the finite dimensional case. Nevertheless, a data-driven method allows to overcome this additional computation is very important in practice and is one of the natural prospects of the present work.

- *Confidence intervals:* The main application of Theorem 5 is to build confidence band for the true value of  $h^x(y)$ . Similarly to the previous application, the practical utilization of our result in this topic requires the estimation of the quantity  $\sigma_h^2(x, y)$ . A plug-in estimate for the asymptotic standard deviation

$\sigma_h^2(x, y)$  can be obtained by using the estimators  $\widehat{f}^x(y)$  and  $\widehat{F}^x(y)$  of  $f^x(y)$  and  $F^x(y)$ . Then we get

$$\widehat{\sigma}_h^2(x, y) := \frac{\widehat{\zeta}_2 \widehat{f}^x(y)}{\left(\widehat{\zeta}_1^2 (1 - \widehat{F}^x(y))^2\right)}$$

where

$$\widehat{\zeta}_1 = \frac{1}{n\phi_x(h_K)} \sum_{i=1}^n K(h_K^{-1}d(x, X_i)) \quad \text{and} \quad \widehat{\zeta}_2 = \frac{1}{n\phi_x(h_K)} \sum_{i=1}^n K^2(h_K^{-1}d(x, X_i)).$$

Clearly, the function  $\phi_x(\cdot)$  does not appear in the calculation of the confidence interval by simplification. More precisely, we obtain the following approximate  $(1 - \zeta)$  confidence band for  $h^x(y)$

$$\widehat{h}^x(y) \pm t_{1-\zeta/2} \times \left( \frac{\widehat{\sigma}_h^2(x, y)}{nh_H\phi_x(h_K)} \right)^{1/2}$$

where  $t_{1-\zeta/2}$  denotes the  $1 - \zeta/2$  quantile of the standard normal distribution.

## 5. APPENDIX

In the following, we will denote  $\forall i$

$$K_i = K(h_H^{-1}d(x, X_i)), \quad H_i = H(h_H^{-1}(y - Y_i)) \quad \text{and} \quad H'_i = H'(h_H^{-1}(y - Y_i)).$$

*Proof of Lemma 2.* Firstly, for  $\mathbb{E}[\widehat{f}_N^x(y)]$ , we start by writing

$$\mathbb{E}[\widehat{f}_N^x(y)] = \frac{1}{\mathbb{E}[K_1]} \mathbb{E} [K_1 \mathbb{E}[h_H^{-1}H'_1|X]] \quad \text{with} \quad h_H^{-1} \mathbb{E} [H'_1|X] = \int_{\mathbb{R}} H'(t) f^X(y - h_H t) dt.$$

The latter can be re-written, by using a Taylor expansion under (H3), as follows

$$h_H^{-1} \mathbb{E}[H'_1|X] = f^X(y) + \frac{h_H^2}{2} \left( \int t^2 H'(t) dt \right) \frac{\partial^2 f^X(y)}{\partial^2 y} + o(h_H^2).$$

Thus, we get

$$\mathbb{E} \left[ \widehat{f}_N^x(y) \right] = \frac{1}{\mathbb{E}[K_1]} \left( \mathbb{E} [K_1 f^X(y)] + \left( \int t^2 H'(t) dt \right) \mathbb{E} \left[ K_1 \frac{\partial^2 f^X(y)}{\partial^2 y} \right] + o(h_H^2) \right).$$

Let  $\psi_l(\cdot, y) := \frac{\partial^l f(y)}{\partial^l y}$ : for  $l \in \{0, 2\}$ , since  $\Phi_l(0) = 0$ , we have

$$\begin{aligned}\mathbb{E}[K_1 \psi_l(X, y)] &= \psi_l(x, y) \mathbb{E}[K_1] + \mathbb{E}[K_1 (\psi_l(X, y) - \psi_l(x, y))] \\ &= \psi_l(x, y) \mathbb{E}[K_1] + \mathbb{E}[K_1 (\Phi_l(d(x, X)))] \\ &= \psi_l(x, y) \mathbb{E}[K_1] + \Phi'_l(0) \mathbb{E}[d(x, X) K_1] + o(\mathbb{E}[d(x, X) K_1]).\end{aligned}$$

So,

$$\begin{aligned}\mathbb{E}[\widehat{f}_N^x(y)] &= f^x(y) + \frac{h_H^2}{2} \frac{\partial^2 f^x(y)}{\partial y^2} \int t^2 H'(t) dt + o\left(h_H^2 \frac{\mathbb{E}[d(x, X) K_1]}{\mathbb{E}[K_1]}\right) \\ &\quad + \Phi'_0(0) \frac{\mathbb{E}[d(x, X) K_1]}{\mathbb{E}[K_1]} + o\left(\frac{\mathbb{E}[d(x, X) K_1]}{\mathbb{E}[K_1]}\right).\end{aligned}$$

Similarly to Ferraty *et al.* (2007) we show that

$$\frac{1}{\phi_x(h_K)} \mathbb{E}[d(x, X) K_1] = h_K \left( K(1) - \int_0^1 (sK(s))' \beta_x(s) ds + o(1) \right)$$

and

$$\frac{1}{\phi_x(h_K)} \mathbb{E}[K_1] = K(1) - \int_0^1 K'(s) \beta_x(s) ds + o(1).$$

Hence,

$$\begin{aligned}\mathbb{E}[\widehat{f}_N^x(y)] &= f^x(y) + \frac{h_H^2}{2} \frac{\partial^2 f^x(y)}{\partial y^2} \int t^2 H'(t) dt \\ &\quad + h_K \Phi'_0(0) \frac{\left( K(1) - \int_0^1 (sK(s))' \beta_x(s) ds \right)}{\left( K(1) - \int_0^1 K'(s) \beta_x(s) ds \right)} + o(h_H^2) + o(h_K).\end{aligned}$$

Secondly, concerning  $\mathbb{E}[\widehat{F}_N^x(y)]$ , we write by an integration by part

$$\mathbb{E}[\widehat{F}_N^x(y)] = \frac{1}{\mathbb{E}[K_1]} \mathbb{E}[K_1 E[H_1|X]] \quad \text{with} \quad \mathbb{E}[H_1|X] = \int_{\mathbb{R}} H'(t) F^X(y - h_H t) dt.$$

The same steps used to studying  $\mathbb{E}[\widehat{f}_N^x(y)]$  can be followed to prove that

$$\begin{aligned}\mathbb{E}[\widehat{F}_N^x(y)] &= F^x(y) + \frac{h_H^2}{2} \frac{\partial^2 F^x(y)}{\partial y^2} \int t^2 H'(t) dt \\ &\quad + h_K \Psi'_0(0) \frac{\left( K(1) - \int_0^1 (sK(s))' \beta_x(s) ds \right)}{\left( K(1) - \int_0^1 K'(s) \beta_x(s) ds \right)} + o(h_H^2) + o(h_K).\end{aligned}$$

*Proof of Lemma 3.* For the first quantity  $Var[\widehat{f}_N^x(y)]$ , we have

$$s_n^2 = Var[\widehat{f}_N^x(y)] = \frac{1}{(nh_H \mathbb{E}[K_1(x)])^2} Var \left[ \sum_{i=1}^n \Gamma_i(x) \right]$$

where

$$\Gamma_i(x) = K_i(x)H'_i(y) - \mathbb{E}[K_i(x)H'_i(y)].$$

Thus

$$\begin{aligned} Var[\widehat{f}_N^x(y)] &= \frac{1}{(nh_H \mathbb{E}[K_1])^2} \underbrace{\sum_{i \neq j} Cov(\Gamma_i(x), \Gamma_j(x))}_{s_n^{cov}} + \underbrace{\sum_{i=1}^n Var(\Gamma_i(x))}_{s_n^{var}} \\ &= \frac{1}{n(h_H \mathbb{E}[K_1])^2} Var[\Gamma_1] + \frac{1}{(nh_H \mathbb{E}[K_1])^2} \sum_{i \neq j} Cov(\Gamma_i, \Gamma_j). \end{aligned}$$

Let us calculate the quantity  $Var[\Gamma_1(x)]$ . We have:

$$\begin{aligned} Var[\Gamma_1(x)] &= \mathbb{E}[K_1^2(x)H_1'^2(y)] - (\mathbb{E}[K_1(x)H_1'(y)])^2 \\ &= \mathbb{E}[K_1^2(x)] \frac{\mathbb{E}[K_1^2(x)H_1'^2(y)]}{\mathbb{E}[K_1^2(x)]} - (\mathbb{E}[K_1(x)])^2 \left( \frac{\mathbb{E}[K_1(x)H_1'(y)]}{\mathbb{E}[K_1(x)]} \right)^2. \end{aligned}$$

So, by using the same arguments as those used in pervious lemma we get

$$\begin{aligned} \frac{1}{\phi_x(h_K)} \mathbb{E}[K_1^2(x)] &= K^2(1) - \int_0^1 (K^2(s))' \beta_x(s) ds + o(1) \\ \frac{\mathbb{E}[K_1^2(x)H_1'^2(y)]}{\mathbb{E}[K_1^2(x)]} &= h_H f^x(y) \int H'^2(t) dt + o(h_H) \\ \frac{\mathbb{E}[K_1(x)H_1'(y)]}{\mathbb{E}[K_1(x)]} &= h_H f^x(y) + o(h_H) \end{aligned}$$

which implies that

$$Var[\Gamma_i(x)] = h_H \phi_x(h_K) f^x(y) \int H'^2(t) dt \left( K^2(1) - \int_0^1 (K^2(s))' \beta_x(s) ds \right) + o(h_H \phi_x(h_K)). \quad (11)$$

Now, let us focus on the covariance term. To do that, we need to calculate the asymptotic behavior of quantity defined as

$$\sum_{i \neq j} |Cov(\Gamma_i(x), \Gamma_j(x))| = \sum_{1 \leq |i-j| \leq c_n} |Cov(\Gamma_i(x), \Gamma_j(x))| = J_{1,n} + J_{2,n}.$$



with  $c_n \rightarrow \infty$ , as  $n \rightarrow \infty$ .

For all  $(i, j)$  we write

$$Cov(\Gamma_i(x), \Gamma_j(x)) = \mathbb{E} [K_i(x)K_j(x)H'_i(y)H'_j(y)] - (\mathbb{E} [K_i(x)H'_i(y)])^2$$

and we use the fact that  $\mathbb{E} [H'_i(y)H'_j(y)|(X_i, X_j)] = \mathcal{O}(h_H^2)$ ;  $\forall i \neq j$ ,  $\mathbb{E} [H'_i(y)|X_i] = \mathcal{O}(h_H)$ ;  $\forall i$ .

For  $J_{1,n}$ : by means of the integral realized above and under (H2) and (H5), we get

$$\mathbb{E} [K_i K_j H'_i H'_j] \leq Ch_H^2 \mathbb{P} [(X_i, X_j) \in B(x, h_K) \times B(x, h_K)]$$

and

$$\mathbb{E} [K_i(x)H'_i(y)] \leq Ch_H \mathbb{P} (X_i \in B(x, h_K)).$$

It follows that, the hypothesis (H0), (H2) and (H5), imply that

$$Cov(\Gamma_i(x), \Gamma_j(x)) \leq Ch_H^2 \phi_x(h_K) \left( \phi_x(h_K) + \left( \frac{\phi_x(h_K)}{n} \right)^{1/a} \right)$$

So

$$J_{1,n} \leq C \left( nc_n h_H^2 \left( \frac{\phi_x(h_K)}{n} \right)^{1/a} \phi_x(h_K) \right).$$

Hence

$$J_{1,n} = \mathcal{O} \left( nc_n h_H^2 \left( \frac{\phi_x(h_K)}{n} \right)^{1/a} \phi_x(h_K) \right).$$

On the other hand, these covariances can be controled by mean of the usual Davydov-Rios's covariance inequality for mixing processes (see Rio 2000, formula 1.12a). Together with (H1), this inequality leads to:

$$\forall i \neq j, |Cov(D_i(x), D_j(x))| \leq C |i - j|^{-a}.$$

By the fact,  $\sum_{k \geq c_n+1} k^{-a} \leq \int_{c_n}^{\infty} t^{-a} dt = \frac{c_n^{-a+1}}{a-1}$ , we get by applying (H1),

$$J_{2,n} \leq \sum_{|i-j| \geq c_n+1} |i-j|^{-a} \leq \frac{nc_n^{-a+1}}{a-1}$$

Thus, by using the following classical technique (see Bosq, 1998 [5]), we can write

$$s_n^{cov} = \sum_{0 < |i-j| \leq u_n} |Cov(\Gamma_i(x), \Gamma_j(x))| + \sum_{|i-j| > u_n} |Cov(\Gamma_i(x), \Gamma_j(x))|.$$

Thus

$$s_n^{cov} \leq Cn \left( c_n h_H^2 \left( \frac{\phi_x(h_K)}{n} \right)^{1/a} \phi_x(h_K) + \frac{c_n^{-a+1}}{a-1} \right)$$

Choosing  $c_n = h_H^{-2} \left( \frac{\phi_x(h_K)}{n} \right)^{-1/a}$ , and owing to the right inequality in (H7b), we can deduce

$$s_n^{cov} = o(nh_H \phi_x(h_K)). \quad (12)$$

Finally,

$$\begin{aligned} s_n^2 &= o(nh_H \phi_x(h_K)) + \mathcal{O}(nh_H \phi_x(h_K)) \\ &= \mathcal{O}(nh_H \phi_x(h_K)) \end{aligned}$$

In conclusion, we have

$$Var[\widehat{f}_N^x(y)] = \frac{f^x(y)}{nh_H \phi_x(h_K)} \left( \int H^2(t) dt \right) \left( \frac{\left( K^2(1) - \int_0^1 (K^2(s))' \beta_x(s) ds \right)}{\left( K(1) - \int_0^1 K'(s) \beta_x(s) ds \right)^2} \right) + o\left( \frac{1}{nh_H \phi_x(h_K)} \right) \quad (13)$$

Now, for  $\widehat{F}_N^x(y)$ , (resp.  $\widehat{F}_D^x$ ) we replace  $H_i'(y)$  by  $H_i(y)$  (resp. by 1) and we follow the same ideas, under the fact that  $H \leq 1$

$$Var[\widehat{F}_N^x(y)] = \frac{F^x(y)}{n\phi_x(h_K)} \left( \int H^2(t) dt \right) \left( \frac{\left( K^2(1) - \int_0^1 (K^2(s))' \beta_x(s) ds \right)}{\left( K(1) - \int_0^1 K'(s) \beta_x(s) ds \right)^2} \right) + o\left( \frac{1}{n\phi_x(h_K)} \right).$$

and

$$Var[\widehat{F}_D^x] = \frac{1}{n\phi_x(h_K)} \left( \frac{\left( K^2(1) - \int_0^1 (K^2(s))' \beta_x(s) ds \right)}{\left( K(1) - \int_0^1 K'(s) \beta_x(s) ds \right)^2} \right) + o\left( \frac{1}{n\phi_x(h_K)} \right).$$

This yields the proof.

*Proof of Lemma 4.* The proof of this lemma follows the same steps as the previous Lemma. For this, we keep the same notation and we write

$$\begin{aligned} Cov(\widehat{f}_N^x(y), \widehat{F}_N^x(y)) &= \frac{1}{nh_H(\mathbb{E}[K_1(x)])^2} Cov(\Gamma_1(x), \Delta_1(x)) \\ &\quad + \frac{1}{n^2 h_H(\mathbb{E}[K_1(x)])^2} \sum_{i \neq j} Cov(\Gamma_i(x), \Delta_j(x)) \end{aligned}$$

where

$$\Delta_i(x) = Ki(x)H_i(y) - \mathbb{E}[Ki(x)H_i(y)].$$

For the first term, we have under (H4)

$$\begin{aligned} Cov(\Gamma_1(x), \Delta_1(x)) &= \mathbb{E}[K_1^2(x)H_1(y)H_1'(y)] - \mathbb{E}[K_1(x)H_1(y)]\mathbb{E}[K_1(x)H_1'(y)] \\ &= \mathcal{O}(h_H\phi_x(h_K)) + \mathcal{O}(h_H\phi_x^2(h_K)) \\ &= \mathcal{O}(h_H\phi_x(h_K)) \end{aligned}$$

Therefore,

$$\begin{aligned} \frac{1}{nh_H(\mathbb{E}[K_1(x)])^2}Cov(\Gamma_1(x), \Delta_1(x)) &= \mathcal{O}\left(\frac{1}{n\phi_x(h_K)}\right) \\ &= o\left(\frac{1}{nh_H\phi_x(h_K)}\right) \end{aligned} \quad (14)$$

So, by using similar arguments as those invoked in the proof of Lemma 3, and we use once again the boundedness of  $K$  and  $H$ , and the fact that (H1) and (H6) imply that

$$\mathbb{E}(H_i'(y)|X_i) = \mathcal{O}(h_H).$$

Moreover, the right part of (H7b) implies that

$$Cov(\Gamma_i(x), \Delta_j(x)) = \mathcal{O}\left(h_H\phi_x(h_K)\left(\frac{\phi_x(h_K)}{n}\right)^{1/a} + \phi_x(h_K)\right),$$

Meanwhile, using the Davydov-Rio's inequality in Rio (2000) for mixing processes leads to

$$|Cov(\Gamma_i(x), \Delta_j(x))| \leq C\alpha(|i-j|) \leq C|i-j|^{-a},$$

we deduce easily that for any  $c_n > 0$  :

$$\begin{aligned} \sum_{i \neq j} Cov(\Gamma_i(x), \Delta_j(x)) &= \mathcal{O}\left(nc_n h_H\phi_x(h_K)\left(\frac{\phi_x(h_K)}{n}\right)^{1/a} + \phi_x(h_K)\right) \\ &+ \mathcal{O}(nh_H c_n^{-a}). \end{aligned}$$

It suffices now to take  $c_n = h_H^{-1}\left(\frac{\phi_x(h_K)}{n}\right)^{-1/a}$  to get the following expression for the sum of the covariances:

$$\sum_{i \neq j} Cov(\Gamma_i(x), \Delta_j(x)) = o(n\phi_x(h_K)). \quad (15)$$

From (14) and (15) we deduce that

$$\text{Cov}(\widehat{f}_N^x(y), \widehat{F}_N^x(y)) = o\left(\frac{1}{nh_H\phi_x(h_K)}\right)$$

The same arguments can be used to show that

$$\text{Cov}(\widehat{f}_N^x(y), \widehat{F}_D^x) = o\left(\frac{1}{nh_H\phi_x(h_K)}\right) \quad \text{and} \quad \text{Cov}(\widehat{F}_N^x(y), \widehat{F}_D^x) = o\left(\frac{1}{nh_H\phi_x(h_K)}\right).$$

*Proof of Lemma 7.* Let

$$S_n = \sum_{i=1}^n \Lambda_i(x)$$

where

$$\Lambda_i(x) := \frac{\sqrt{h_H\phi_x(h_K)}}{h_H\mathbb{E}[K_1(x)]} \Gamma_i(x). \quad (16)$$

Obviously, we have

$$\sqrt{nh_H\phi_x(h_K)} [\sigma_f(x, y)]^{-1} \left( \widehat{f}_N^x(y) - E\widehat{f}_N^x(y) \right) = (n(\sigma_f(x, y))^2)^{-1/2} S_n.$$

Thus, the asymptotic normality of  $(n(\sigma_f(x, y))^2)^{-1/2} S_n$ , is sufficient to show the proof of this Lemma. This last is shown by the blocking method, where the random variables  $\Lambda_i$  are grouped into blocks of different sizes defined.

We consider the classical big- and small-block decomposition. We split the set  $\{1, 2, \dots, n\}$  into  $2k_n + 1$  subsets with large blocks of size  $u_n$  and small blocks of size  $v_n$  and put

$$k_n := \left\lfloor \frac{n}{u_n + v_n} \right\rfloor.$$

Assumption (H7)(ii) allows us to define the large block size by

$$u_n =: \left\lfloor \left( \frac{nh_H\phi_x(h_K)}{q_n} \right)^{1/2} \right\rfloor.$$

Using Assumption (H7) and simple algebra allows us to prove that

$$\frac{v_n}{u_n} \rightarrow 0, \quad \frac{u_n}{n} \rightarrow 0, \quad \frac{u_n}{\sqrt{nh_H\phi_x(h_K)}} \rightarrow 0, \quad \text{and} \quad \frac{n}{u_n} \alpha(v_n) \rightarrow 0 \quad (17)$$

Now, let  $\Upsilon_j$ ,  $\Upsilon'_j$  and  $\Upsilon''_j$  be defined as follows:

$$\begin{aligned}\Upsilon_j &= \sum_{i=j(u+v)+1}^{j(u+v)+u} \Lambda_i(x), \quad 0 \leq j \leq k+1 \\ \Upsilon'_j &= \sum_{i=j(u+v)+u+1}^{(j+1)(u+v)+u} \Lambda_i(x), \quad 0 \leq j \leq k+1 \\ \Upsilon''_j &= \sum_{i=k(u+v)+1}^n \Lambda_i(x), \quad 0 \leq j \leq k+1\end{aligned}$$

Clearly, we can write

$$S_n = \sum_{j=0}^{k-1} \Upsilon_j + \sum_{j=0}^{k-1} \Upsilon'_j + \Upsilon''_k =: S'_n + S''_n + S'''_n.$$

We prove that

$$(i) \frac{1}{n} \mathbb{E}(S''_n)^2 \longrightarrow 0, \quad (ii) \frac{1}{n} \mathbb{E}(S'''_n)^2 \longrightarrow 0, \quad (18)$$

$$\left| \mathbb{E} \left\{ \exp \left( itn^{-1/2} S'_n \right) \right\} - \prod_{j=0}^{k-1} \mathbb{E} \left\{ \exp \left( itn^{-1/2} \Upsilon_j \right) \right\} \right| \longrightarrow 0, \quad (19)$$

$$\frac{1}{n} \sum_{j=0}^{k-1} \mathbb{E} \left( \Upsilon_j^2 \right) \longrightarrow \sigma_f^2(x, y), \quad (20)$$

$$\frac{1}{n} \sum_{j=0}^{k-1} \mathbb{E} \left( \Upsilon_j^2 \mathbf{1}_{\{|\Upsilon_j| > \varepsilon \sqrt{n\sigma_f^2(x, y)}\}} \right) \longrightarrow 0 \quad (21)$$

for every  $\varepsilon > 0$ .

Expression (18) show that the terms  $S''_n$  and  $S'''_n$  are negligible, while Equations (19) and (20) show that the  $\Upsilon_j$  are asymptotically independent, verifying that the sum of their variances tends to  $\sigma_f^2(x, y)$ . Expression (21) is the Lindeberg-Feller's condition for a sum of independent terms. Asymptotic normality of  $S_n$  is a consequence of Equations (18)-(21).

- **Proof of (18)** Because  $\mathbb{E}(\Lambda_j) = 0, \forall j$ , we have that

$$\mathbb{E}(S_n'')^2 = Var \left( \sum_{j=0}^{k-1} \Upsilon_j' \right) = \sum_{j=0}^{k-1} Var(\Upsilon_j') + \sum_{0 \leq i < j \leq k-1} Cov(\Upsilon_i', \Upsilon_j') := \Pi_1 + \Pi_2.$$

By the second-order stationarity we get

$$\begin{aligned} Var(\Upsilon_j') &= Var \left( \sum_{i=j(u_n+v_n)+u_n+1}^{(j+1)(u_n+v_n)} \Lambda_i(x) \right) \\ &= v_n Var(\Lambda_1(x)) + \sum_{i \neq j}^{v_n} Cov(\Lambda_i(x), \Lambda_j(x)). \end{aligned}$$

Then

$$\begin{aligned} \frac{\Pi_1}{n} &= \frac{kv_n}{n} Var(\Lambda_1(x)) + \frac{1}{n} \sum_{j=0}^{k-1} \sum_{i \neq j}^{v_n} Cov(\Lambda_i(x), \Lambda_j(x)) \\ &\leq \frac{kv_n}{n} \left\{ \frac{\phi_x(h_K)}{h_H \mathbb{E}^2 K_1(x)} Var(\Gamma_1(x)) \right\} + \frac{1}{n} \sum_{i \neq j}^n |Cov(\Lambda_i(x), \Lambda_j(x))| \\ &\leq \frac{kv_n}{n} \left\{ \frac{1}{h_H \phi_x(h_K)} Var(\Lambda_1(x)) \right\} + \frac{1}{n} \sum_{i \neq j}^n |Cov(\Lambda_i(x), \Lambda_j(x))| \end{aligned}$$

Simple algebra gives us

$$\frac{kv_n}{n} \cong \left( \frac{n}{u_n + v_n} \right) \frac{v_n}{n} \cong \frac{v_n}{u_n + v_n} \cong \frac{v_n}{u_n} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Using Equation (12) we have

$$\lim_{n \rightarrow \infty} \frac{\Pi_1}{n} = 0 \tag{22}$$

Now, let us turn to  $\Pi_2/n$ . We have

$$\begin{aligned} \frac{\Pi_2}{n} &= \frac{1}{n} \sum_{i=0, i \neq j}^{k-1} \sum_{j=0}^{k-1} Cov(\Upsilon_i(x), \Upsilon_j(x)) \\ &= \frac{1}{n} \sum_{i=0, i \neq j}^{k-1} \sum_{j=0}^{k-1} \sum_{l_1=1}^{v_n} \sum_{l_2}^{v_n} Cov(\Lambda_{m_j+l_1}, \Lambda_{m_j+l_2}) \end{aligned}$$

with  $m_i = i(u_n + v_n) + v_n$ . As  $i \neq j$ , we have  $|m_i - m_j + l_1 - l_2| \geq u_n$ . It follows that

$$\frac{\Pi_2}{n} \leq \frac{1}{n} \sum_{i=1}^n \sum_{\substack{j=1 \\ |i-j| \geq u_n}}^n Cov(\Lambda_i(x), \Lambda_j(x)),$$

then

$$\lim_{n \rightarrow \infty} \frac{\Pi_2}{n} = 0. \quad (23)$$

By Equations (22) and (23) we get Part(i) of the Equation(18).

We turn to (ii), we have

$$\begin{aligned} \frac{1}{n} \mathbb{E} \left( S_n''' \right)^2 &= \frac{1}{n} Var \left( \Upsilon_k'' \right) \\ &= \frac{\vartheta_n}{n} Var \left( \Lambda_1(x) \right) + \frac{1}{n} \sum_{i=1}^{\vartheta_n} \sum_{\substack{j=1 \\ i \neq j}}^{\vartheta_n} Cov \left( \Lambda_i(x), \Lambda_j(x) \right) \end{aligned}$$

where  $\vartheta_n = n - k_n(u_n + v_n)$ ; by the definition of  $k_n$ , we have  $\vartheta_n \leq u_n + v_n$ .

Then

$$\frac{1}{n} \mathbb{E} \left( S_n''' \right)^2 \leq \frac{u_n + v_n}{n} Var \left( \Lambda_1(x) \right) + \frac{1}{n} \sum_{i=1}^{\vartheta_n} \sum_{\substack{j=1 \\ i \neq j}}^{\vartheta_n} Cov \left( \Lambda_i(x), \Lambda_j(x) \right)$$

and by the definition of  $u_n$  and  $v_n$  we achieve the proof of (ii) of Equation (18).

- **Proof of (19)** We make use of Volkonskii and Rozanov's lemma (see the appendix in Masry, 2005) and the fact that the process  $(X_i, X_j)$  is strong mixing. Note that  $\Upsilon_a$  is  $\mathcal{F}_{i_a}^{j_a}$ -mesurable with  $i_a = a(u_n + v_n) + 1$  and  $j_a = a(u_n + v_n) + u_n$ ; hence, with  $V_j = \exp(itn^{-1/2}\Upsilon_j)$  we have

$$\left| \mathbb{E} \left\{ \exp \left( itn^{-1/2} S_n' \right) \right\} - \prod_{j=0}^{k-1} \mathbb{E} \left\{ \exp \left( itn^{-1/2} \Upsilon_j \right) \right\} \right| \leq 16k_n \alpha(v_n + 1) \cong \frac{n}{v_n} \alpha(v_n + 1)$$

which goes to zero by the last part of Equation (17). Now we establish Equation (20).

- **Proof of (20)** Note that  $Var(S_n') \rightarrow \sigma_f^2(x, y)$  by Equation (18)-(i) and (18)-(ii) and since  $Var(S_n) \rightarrow \sigma_f^2(x, y)$  (by the definition of the  $\Lambda_i$  and Equation

(13)). Then because

$$\mathbb{E} (S'_n)^2 = Var (S'_n) = \sum_{j=0}^{k-1} Var (\Upsilon_j) + \sum_{i=0}^{k-1} \sum_{i \neq j}^{k-1} Cov (\Upsilon_i, \Upsilon_j),$$

all we have to prove is that the double sum of covariances in the last equation tends to zero. Using the same arguments as those previously used for  $\Pi_2$  in the proof of first term of Equation (18) we obtain by replacing  $v_n$  by  $u_n$  we get

$$\frac{1}{n} \sum_{j=0}^{k-1} \mathbb{E} (\Upsilon_j^2) = \frac{ku_n}{n} Var (\Lambda_1) + o(1).$$

As  $Var (\Lambda_1) \rightarrow \sigma_f^2(x, y)$  and  $ku_n/n \rightarrow 1$ , we get the result.

Finally, we prove Equation (21).

- **Proof of (21)** Recall that

$$\Upsilon_j = \sum_{i=j(u_n+v_n)+1}^{j(u_n+v_n)+u_n} \Lambda_i.$$

Making use Assumptions (H5) and (H6), we have

$$|\Lambda_i| \leq C (h_H \phi_x(h_K))^{-1/2}$$

thus

$$|\Upsilon_j| \leq Cu_n (h_H \phi_x(h_K))^{-1/2},$$

which goes to zero as  $n$  goes to infinity by Equation (17). Then for  $n$  large enough, the set  $\{|\Upsilon_j| > \varepsilon (n\sigma_f^2(x, y))^{-1/2}\}$  becomes empty, this completes the proof and therefore that of the asymptotic normality of  $(n(\sigma_f(x, y))^2)^{-1/2} S_n$ ,

*Proof of Lemma 8.* It is clear that, the result of Lemma (2) and Lemma (3) permits us

$$\mathbb{E} \left( \widehat{F}_D^x - \widehat{F}_N^x - 1 + F^x(y) \right) \rightarrow 0$$

and

$$Var \left( \widehat{F}_D^x - \widehat{F}_N^x - 1 + F^x(y) \right) \rightarrow 0$$



then

$$\widehat{F}_D^x - \widehat{F}_N^x - 1 + F^x(y) \xrightarrow{\mathbb{P}} 0$$

Moreover, the asymptotic variance of  $\widehat{F}_D^x - \widehat{F}_N^x$  given in remark (5) allows to obtain

$$\frac{nh_H \phi_x(h_K)}{\sigma_h(x, y)^2} \text{Var} \left( \widehat{F}_D^x - \widehat{F}_N^x - 1 + \mathbb{E} \left( \widehat{F}_N^x(y) \right) \right) \longrightarrow 0.$$

By combining result with the fact that

$$\mathbb{E} \left( \widehat{F}_D^x - \widehat{F}_N^x - 1 + \mathbb{E} \left( \widehat{F}_N^x(y) \right) \right) = 0$$

we obtain the claimed result.

*Proof of lemma 11.* For  $i = 1, \dots, n$ , we consider the quantities  $K_i = K(h_K^{-1}d(z, Z_i))$ ,  $H_i''(x) = H''(h_H^{-1}(x - X_i))$  and let  $\widehat{f}'_N^Z(x)$  (resp.  $\widehat{F}_D^Z$ ) be defined as

$$\widehat{f}'_N^Z(x) = \frac{h_H^{-2}}{n \mathbb{E}K_1} \sum_{i=1}^n K_i H_i''(x) \quad (\text{resp. } \widehat{F}_D^Z = \frac{1}{n \mathbb{E}K_1} \sum_{i=1}^n K_i).$$

This proof is based on the following decomposition

$$\begin{aligned} \widehat{f}'^X(y) - f'^X(y) &= \frac{1}{\widehat{F}_D^X} \left\{ \left( \widehat{f}'_N^X(y) - \mathbb{E}\widehat{f}'_N^X(y) \right) - \left( f'^X(y) - \mathbb{E}\widehat{f}'_N^X(y) \right) \right\} + \\ &\quad \frac{f'^X(y)}{\widehat{F}_D^X} \left\{ \mathbb{E}\widehat{F}_D^X - \widehat{F}_D^X \right\} \end{aligned} \quad (24)$$

and on the following intermediate results.

$$\sqrt{nh_H^3 \phi_x(h_K)} \left( \widehat{f}'_N^X(y) - \mathbb{E}\widehat{f}'_N^X(y) \right) \xrightarrow{\mathcal{D}} N \left( 0, \sigma_{f'^X}^2(y) \right) \quad (25)$$

where  $\sigma_{f'^X}^2(y)$  is defined as in Lemma 11.

$$\lim_{n \rightarrow \infty} \sqrt{nh_H^3 \phi_x(h_K)} \left( \mathbb{E}\widehat{f}'_N^X(y) - f'^X(y) \right) = 0 \quad (26)$$

$$\sqrt{nh_H^3 \phi_x(h_K)} \left( \widehat{F}_D^X(y) - 1 \right) \xrightarrow{\mathbb{P}} 0, \quad \text{as } n \rightarrow \infty. \quad (27)$$

- Concerning (25). By the definition of  $\widehat{f}'_N^X(y)$ , it follows that

$$\sqrt{nh_H^3 \phi_x(h_K)} \left( \widehat{f}'_N^X(y) - \mathbb{E} \widehat{f}'_N^X(y) \right) = \sum_{i=1}^n \frac{\sqrt{\phi_x(h_K)}}{\sqrt{nh_H \mathbb{E} K_1}} (K_i H_i'' - \mathbb{E} K_i H_i'') = \sum_{i=1}^n \Delta_i,$$

which leads

$$\sum_{i=1}^n \mathbb{E} \Delta_i^2 = \frac{\phi_x(h_K)}{h_H \mathbb{E}^2 K_1} \mathbb{E} K_1^2 (H_1'')^2 - \frac{\phi_x(h_K)}{h_H \mathbb{E}^2 K_1} (\mathbb{E} K_1 H_1'')^2 = \Pi_{1n} - \Pi_{2n}. \quad (28)$$

As for  $\Pi_{1n}$ , by the property of conditional expectation, we get

$$\Pi_{1n} = \frac{\phi_x(h_K)}{\mathbb{E}^2 K_1} \mathbb{E} \left\{ K_1^2 \int H''^2(t) (f'^X(y - th_H) - f'^X(y) + f'^X(y)) dt \right\}.$$

Meanwhile, by (H0), (H3), (H4) and (H5), it follows that:

$$\frac{\phi_x(h_K) \mathbb{E} K_1^2}{\mathbb{E}^2 K_1} \xrightarrow{n \rightarrow \infty} \frac{a_2^z}{(a_1^z)^2},$$

which leads

$$\Pi_{1n} \xrightarrow{n \rightarrow \infty} \frac{a_2^z f^X(y)}{(a_1^z)^2} \int (H''(t))^2 dt, \quad (29)$$

Regarding  $\Pi_{2n}$ , by (H0), (H3) and (H6), we obtain

$$\Pi_{2n} \xrightarrow{n \rightarrow \infty} 0. \quad (30)$$

This result, combined with (28) and (29), allows us to get

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n \mathbb{E} \Delta_i^2 = \sigma_{f'^X}^2(y) \quad (31)$$

Secondly, by the boundedness of  $H''$ , we have

$$\begin{aligned} \mathbb{E} (|\Delta_i \Delta_j|) &\leq \frac{C \phi_x(h_K)}{n \mathbb{E}^2 K_1} (K_i K_j + \mathbb{E} K_i K_j) \\ &\leq \frac{C}{nh_H} \left\{ \left( \frac{\phi_x(h_K)}{n} \right)^{1/a} + \phi_x(h_K) \right\}, \quad \forall i \neq j. \end{aligned}$$

Then, taking

$$\delta_n = \max_{1 \leq i \neq j \leq n} \{\mathbb{E}(|\Delta_i \Delta_j|)\} = \frac{C}{nh_H} \left( \left( \frac{\phi_x(h_K)}{n} \right)^{1/a} + \phi_x(h_K) \right).$$

leads

$$nm_n \delta_n = \frac{Cm_n}{h_H} \left( \left( \frac{\phi_x(h_K)}{n} \right)^{1/a} + \phi_x(h_K) \right). \quad (32)$$

Similarly, the boundedness of  $H''$  and  $K$  allows us to take  $C_i = \mathcal{O}\left(\frac{1}{\sqrt{nh_H^3 \phi_x(h_K)}}\right)$ , which implies that

$$\left( \sum_{j=m_n+1}^{\infty} \alpha(j) \right) \sum_{i=1}^n C_i^2 \leq \frac{C}{h_H \phi_x(h_K)} \int_{t \geq m_n} t^{-a} dt = \frac{C}{h_H \phi_x(h_K)} \frac{m_n^{-a+1}}{a-1}. \quad (33)$$

Then, the sum of the right side of (32) and (33) is of type  $Am_n + Bm_n^{-a+1}$ , by talking  $m_n = (A/B)^{-1/a} = \{(a-1)\phi_x(h_K) \left( \left( \frac{\phi_x(h_K)}{n} \right)^{1/a} + \phi_x(h_K) \right)\}^{-1/a} \rightarrow \infty$ , it is clear that, under conditions (H9a) and (H9b), combining (32) and (33) allows us to get

$$nm_n \delta_n = o(1), \quad (34)$$

and

$$\left( \sum_{j=m_n+1}^{\infty} \alpha(j) \right) \sum_{i=1}^n C_i^2 = o(1), \quad (35)$$

respectively. Finally, by choosing  $\varrho_n = \sqrt{\frac{nh_H^3 \phi_x(h_K)}{\log n}}$ , under (H9a) again and  $a > 3$ , we have

$$\frac{\varrho_n}{\sqrt{n}} = o(1) \quad (36)$$

and

$$\begin{aligned} \frac{n}{\varrho_n} \alpha(\varepsilon \varrho_n) &\leq C \frac{(\log n)^{(a+1)/2}}{n^{(a-1)/2} (h_H^3 \phi_x(h_K))^{(a+1)/2}} \\ &\leq C \frac{(\log n)^{(a+1)/2}}{n^{(a-3)/2}} \rightarrow 0 \quad \text{as } n \rightarrow \infty. \end{aligned}$$

Therefore, combining (30)-(36) with Corollary 2.2 in Liebscher [17], (25) is valid.

- Concerning (26). The proof is completed along the same steps as that of  $\Pi_{1n}$ . We omit it here.
- Concerning (27). The idea is similar to that given by Ferraty *et al.* [29].

By definition of  $\widehat{F}_D^X(y)$ , we have

$$\sqrt{nh_H^3 \phi_x(h_K)} (\widehat{F}_D^X(y) - 1) = \Omega_n - \mathbb{E}\Omega_n,$$

where  $\Omega_n = \frac{\sqrt{nh_H^3 \phi_x(h_K)} \sum_{i=1}^n K_i}{n\mathbb{E}^2 K_1}$ . In order to prove (27), similar to Ferraty *et al.* [29], we only need to prove  $Var \Omega_n \rightarrow 0$ , as  $n \rightarrow \infty$ . In fact, since

$$\begin{aligned} Var \Omega_n &= \frac{nh_H^3 \phi_x(h_K)}{n\mathbb{E}^2 K_1} \left( nVar K_1 + \sum_{1 \leq i < j \leq n} \sum cov(K_i, K_j) \right) \\ &\leq \frac{nh_H^3 \phi_x(h_K)}{\mathbb{E}^2 K_1} \mathbb{E} K_1^2 + \frac{nh_H^3 \phi_x(h_K)}{n\mathbb{E}^2 K_1} \sum_{0 \leq |i-j| \leq v_n} \sum cov(K_i, K_j) \\ &\quad + \frac{nh_H^3 \phi_x(h_K)}{n\mathbb{E}^2 K_1} \sum_{0 \leq |i-j| \geq v_n} \sum cov(K_i, K_j) \\ &= \Psi_1 + \Psi_2 + \Psi_3, \end{aligned}$$

then, using the boundedness of function  $K$  allows us to get that:

$$\Psi_1 \leq Ch_H^3 \phi_x(h_K) \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

Meanwhile, by (H0) and (H1), it follows that

$$\Psi_2 \leq v_n h_H^3 \left\{ \left( \frac{\phi_x(h_K)}{n} \right)^{1/a} + \phi_x(h_K) \right\}. \quad (37)$$

Finally, using the Davydov-Rio's inequality in Rio [25] for mixing processes leads to

$$|cov(K_i, K_j)| \leq C\alpha(|i-j|),$$

for all  $i \neq j$ . Then, we have

$$\begin{aligned} \Psi_3 &\leq \frac{h_H^3 \phi_x(h_K)}{n\mathbb{E}^2 K_1} n^2 C\alpha(|i-j|) \leq C \frac{h_H^3 \phi_x(h_K)}{n\mathbb{E}^2 K_1} n^2 v_n^{-a+1} \\ &\leq Ch_H^3 n v_n^{-a+1}. \end{aligned} \quad (38)$$

Since the right side of (37) and (38) is also of type  $Av_n + Bv_n^{-a+1}$ , by choosing  $v_n = [n^{-1}((\frac{\phi_x(h_K)}{n})^{1/a} + \phi_x(h_K))]^{-1/a} \rightarrow \infty$  and simple calculations, we get that  $\Psi_2 \rightarrow 0$  and  $\Psi_3 \rightarrow 0$  as  $n \rightarrow \infty$ , respectively.

Therefore, the proof of this result is completed.

Therefore, the proof of this lemma is completed.

#### REFERENCES

- [1] I. A. Ahmad, *Uniform strong convergence of the generalized failure rate estimate*, Bull. Math. Stat. 17 (1976), 77-84.
- [2] K. Benhenni, F. Ferraty, M. Rachdi, P. Vieu, *Local smoothing regression with functional data*, Comput. Stat. 22 (2007), 353-369.
- [3] P. Besse, H. Cardot, D. Stephenson, *Autoregressive forecasting of some functional climatic variations*, Scand. J. Stat. 27 (2000), 673-687.
- [4] P. Besse, J.O. Ramsay, *Principal component analysis of sampled curves*, Psychometrika. 51 (1986), 285-311.
- [5] D. Bosq, *Nonparametric Statistics for Stochastic Processes. Estimation and Prediction*, (Second edition). Lecture Notes in Statistics, 10, Springer-Verlag, (1998).
- [6] M. Bouraine, A. Aït Saidi, F. Ferraty and P. Vieu, *Choix optimal de l'indice multi-fonctionnel: méthode de validation croisée*, R. Roumaine Math. Pures et Appl. 50, 5 (2010), 355-367.
- [7] H. Cardot, F. Ferraty, P. Sarda, *Functional linear model*, Stat. Proba. Lett. 45 (1999), 11-22.
- [8] J. Damon, S. Guillas, *The inclusion of exogenous variables in functional autoregressive ozone forecasting*, Environmetrics. 13 (2002), 759-774.
- [9] M. Ezzahrioui, E. Ould-Saïd, *Asymptotic results of a nonparametric conditional quantile estimator for functional time series*, Comm. Stat. Theory Methods. 37, 16-17 (2008), 2735-2759.
- [10] F. Ferraty, A. Mas, P. Vieu, *Advances in nonparametric regression for functional variables*, Australian and New Zealand Journal of Statistics. 49 (2007), 1-20.
- [11] F. Ferraty, A. Rabhi, P. Vieu, *Conditional quantiles for functional dependent data with application to the climatic El Nino phenomenon*, Sankhyā : The Indian Journal of Statistics, Special Issue on Quantile Regression and Related Methods. 67, 2 (2005), 378-399.

- [12] F. Ferraty, A. Rabhi, P. Vieu, *Estimation non paramétrique de la fonction de hasard avec variable explicative fonctionnelle*, Rom. J. Pure & Applied Math. 52 (2008), 1-18.
- [13] F. Ferraty, P. Vieu, *Non-parametric Functional Data Analysis*, Springer-Verlag, NewYork, (2006).
- [14] T. Gasser, P. Hall, B. Presnell, *Nonparametric estimation of the mode of a distribution of random curves*, Journal of the Royal Statistical Society, Ser. B. 60 (1998), 681-691.
- [15] P. Hall, R. C. L. Wolef, Q. Yao, *Methods for estimating a conditional distribution function*, Ann. Stat. Assoc. 94 (1999), 154-163.
- [16] R. J. Hyndman, D. M. Bashtannyk, G. K. Grunwald, *Estimating and visualizing conditional densities*, J. Comput. Graph. Stat. 5 (1996), 315-336.
- [17] E. Liebscher, *Central limit theorem for  $\alpha$ -mixing triangular arrays with applications to nonparametric statistics*, Mathematical Methods of Statistics. 10, 2 (2001), 194-214.
- [18] A. Mahiddine, A. A. Bouchentouf, A. Rabhi, *Nonparametric estimation of some characteristics of the conditional distribution in single functional index model*, Malaya Journal of Matematik (MJM). 2, 4 (2014), 392-410.
- [19] E. Masry, *Non-parametric regression estimation for dependent functional data: Asymptotic normality*, Stoch. Process. Appl. 115 (2005), 155-177.
- [20] A. Quintela, *Plug-in bandwidth selection in kernel hazard estimation from dependent data*, Comput. Stat. Data Anal. 51 (2007), 5800-5812.
- [21] A. Rabhi, S. Benaissa, E. H. Hamel, B. Mechab, *Mean square error of the estimator of the conditional hazard function*, Appl. Math. (Warsaw), 40, 4 (2013), 405-420.
- [22] M. Rachdi and P. Vieu, *Non-parametric regression for functional data: Automatic smoothing parameter selection*, J. Stat. Plan. Inference. 137 (2007), 2784-2801.
- [23] J.O. Ramsay, B.W. Silverman, *Functional Data Analysis*, 2nd ed., Springer-Verlag, NewYork, (2005).
- [24] J. Rice and B.W. Silverman, *Estimating the mean and covariance structure non-parametrically when the data are curves*, J. R. Stat. Soc., Ser. B. 53 (1991), 233-243.
- [25] E. Rio, *Théorie asymptotique des processus aléatoires dépendants*, (in french). Mathématiques et Applications. 31, Springer, (2000).
- [26] G. G. Roussas, *Hazard rate estimation under dependence conditions*, J. Stat. Plann. Inference. 22 (1989), 81-93.

- [27] N. D. Singpurwalla, M. Y. Wong, *Estimation of the failure rate-A survey of non-parametric methods. Part I: Non-Bayesian methods*, Commun. Stat. Theory & Meth. 12 (1983), 559-588.
- [28] L. Spierdijk, *Non-parametric conditional hazard rate estimation: A local linear approach*, Comput. Stat. Data Anal. 52 (2008), 2419-2434.
- [29] L. T. Tran, S. Yakowitz, *Nearest neighbor estimators for rom fields*, J. Multivariate. Anal. 44 (1993), 23-46.
- [30] G. S. Watson, M. R. Leadbetter, *Hazard analysis*, Sankhyā. 26 (1964), 101-116.

Naouel Belkhir  
Laboratory of Mathematics Abu Bakr Belkaid University of Tlemcen, Algeria.  
email: *n.belkhir@yahoo.com*

Abbes Rabhi  
Laboratory of Mathematics,  
Sidi Bel Abbes University,  
P. O. Box. 89 Sidi Bel Abbes 22000 Algeria.  
email: *rabhi\_abbes@yahoo.fr*

Sarah Soltani  
Stochastic Models Statistics and Applications Laboratory,  
Moulay Tahar University of Saida,  
P.O.Box 138 En-Nasr Saida 20 000 Algeria.  
email: *s.assala@hotmail.com*