

Editorial

MESSAGE FROM THE EDITORS

Saber Elaydi, *Mathematics and Mathematical Sciences Editor*

Department of Mathematics, Trinity University, San Antonio, Texas 78212, E-mail:

selaydi@trinity.edu

Abraham F. Jalbout, *Physical and Chemical Sciences Editor*

Department of Chemistry, The University of New Orleans, New Orleans, LA 70148

Department of Physics, Dillard University, 2 601 Gentilly Blvd., New Orleans, LA 70112

E-mail: Ajalbout@ejmaps.org

Bassel E. Sawaya, *Biological Sciences Editor*

Center for Neuro-Virology and Cancer Biology, College of Sciences and Technology, Temple

University, Philadelphia, PA. 19122, USA, E-mail: Sawaya@temple.edu

Received: 22 July 2002/ Accepted: 22 July 2002/ Published: 22 August 2002

Electronic Journal of
Mathematical
and Physical Sciences

EJMAPS

ISSN: 1538-263X

www.ejmaps.org

Editorial

MESSAGE FROM THE HONORARY EDITOR

Vangipuram Lakshmikantham

President and Founder - International Federation of Nonlinear Analysts
Department of Mathematical Sciences, Florida Institute of Technology, Melbourne, FL 32901
E-mail: dkermani@winnie.fit.edu, akshmik@winnie.fit.edu

Received: 22 July 2002/ Accepted: 22 July 2002/ Published: 22 August 2002

Editorial

**TEN YEARS OF THE ELECTRONIC JOURNAL OF
DIFFERENTIAL EQUATIONS**

Alfonso Castro

Department of Mathematics, University of Texas at San Antonio, TX, USA

E-mail: acastro@utsa.edu

Julio G. Dix

Department of Mathematics, Southwest Texas State University, San Marcos Texas, USA

E-mail: Julio@swt.edu

Received: 22 February 2002/ Accepted: 24 February 2002/ Published: 22 August 2002

Electronic Journal of
Mathematical
and Physical Sciences

EJMAPS

ISSN: 1538-263X

www.ejmaps.org

Editorial

LAUNCHING A NEW JOURNAL

Avshalom C. Elitzur

Unit for Interdisciplinary Studies, Bar-Ilan University, 52900 Ramat-Gan, Israel.

E-mail: avshalom.elitzur@weizmann.ac.il

Received: 5 May 2002/ Accepted: 6 May 2002/ Published: 22 August 2002

Editorial

WELCOMING MESSAGE FROM AN EDITORIAL BOARD MEMBER

Srinivasa Buddhudu

Department of Physics, S.V.University, Tirupati-517502, India

Received: 15 January 2002/ Accepted: 10 May 2002/ Published: 22 August 2002

STRANG-TYPE PRECONDITIONERS FOR SOLVING SYSTEM OF ODES BY BOUNDARY VALUE METHODS

Raymond H. Chan^{1*}, **Xiao-Qing Jin**², **Yue-Hung Tam**³

¹ Department of Mathematics, The Chinese University of Hong Kong, Shatin, Hong Kong.

² Faculty of Science and Technology, The University of Macau, Macau.

³ Department of Mathematics, The Chinese University of Hong Kong, Shatin, Hong Kong

*Corresponding Author. E-mail: rchan@math.cuhk.edu.hk

Received: 2 April 2002/ Accepted: 2 May 2002/ Published: 22 August 20002

Abstract: In this paper, we survey some of the latest developments in using boundary value methods for solving systems of ordinary differential equations with initial values. These methods require the solutions of one or more nonsymmetric, large and sparse linear systems. The GMRES method with the Strang-type preconditioner is proposed for solving these linear systems. One of the main results is that if an $A_{i, 2}$ stable boundary value method is used for an m -by- m system of ODEs, then the preconditioner is invertible and the preconditioned matrix can be decomposed as $I + L$ where I is the identity matrix and the rank of L is at most $2m(i_1 + i_2)$. It follows that when the GMRES method is applied to solving the preconditioned systems, the method will converge in at most $2m(i_1 + i_2) + 1$ iterations. Applications to differential algebraic equations and delay differential equations are also given.

Keywords: Boundary value method, GMRES, ordinary differential equation, Strang-type preconditioner

AMS Mathematical Subject Classification: 65L05, 65L06, 65N10, 65N22

ON THE ANALYSIS OF A VISCOPLASTIC CONTACT PROBLEM WITH TIME DEPENDENT TRESCA'S FRICTION LAW

Amina Amassad ^{*} and Caroline Fabre [#]
de Nice-Sophia Antipolis, Laboratoire J.-A. Dieudonné
UMR-CNRS 6621, Parc Valrose, F-06108 Nice, France
E-mail: ^{*} amassad@math.unice.fr and [#] cfabre@math.unice.fr

Received: 26 May 2002/ Accepted: 27 July 2002/ Published: 22 August 2002

Abstract: This paper deals with the study of a nonlinear problem of frictional contact between an elastic-viscoplastic body and a rigid obstacle. We model the frictional contact by a version of Tresca's friction law where the friction bound depends on time. Firstly, we obtain an existence and uniqueness result in a weak sense for a model including the bilateral contact. To this end we use a time discretization method and the Banach fixed-point theorem. Secondly, we show an existence result for a mechanical problem with the unilateral contact conditions (Signorini's contact) using an iterative method

Keywords: Quasistatic frictional contact, bilateral contact, unilateral contact, Tresca's friction law, fixed point, discretization

AMS Mathematical Subject Classification: 74D10, 74A55, 49B40

EXISTENCE RESULTS FOR IMPULSIVE FUNCTIONAL AND NEUTRAL FUNCTIONAL DIFFERENTIAL INCLUSIONS WITH LOWER SEMICONTINUOUS RIGHT HAND SIDE

Mouffak Benchohra¹, **Johnny Henderson**^{2*}, **Sotiris Ntouyas**³ and **Ahmed Ouahabi**¹

¹ Laboratoire de Mathématiques, Université de Sidi Bel Abbés, BP 89, 22000 Sidi Bel Abbés, Algérie

² Department of Mathematics, Baylor University, Waco, TX 76798-7328, USA

³ Department of Mathematics, University of Ioannina, 45110 Ioannina, Greece

*Corresponding Author. E-mail: Johnny.Henderson@baylor.edu

Received: 16 April 2002/ Accepted: 13 July 2002/ Published: 22 August 2002

Abstract: In this paper, Schaefer's fixed point theorem combined with a selection theorem due to Bressan and Colombo is used to investigate the existence of solutions for first and second order impulsive functional and neutral differential inclusions with lower semicontinuous and nonconvex-valued right-hand side.

Keywords: Impulsive differential inclusions, selection, existence, fixed point

AMS Mathematical Subject Classification: 34A37, 34A60

A FAST ALGORITHM FOR BLIND CHANNEL IDENTIFICATION

Michael K. Ng

Department of Mathematics, The University of Hong Kong, Pokfulam Road, Hong Kong.
E-mail: mng@maths.hku.hk.

Received: 26 June 2002/ Accepted: 15 July 2002/ Published: 22 August 20002

Abstract: In this paper, we address the problem of restoring a signal from its noisy convolutions with two unknown channels. When the transfer functions of these two channels have no common factors, the blind channel identification problem can be solved by finding the minimum eigenvalue of the Toeplitz-block matrix and its corresponding eigenvector. We present a fast algorithm to solve the numerical solution of the eigenvalue problem for Toeplitz-block matrices. Experimental results are given to illustrate the effectiveness of the method.

Keywords: Blind channel identification, Toeplitz-block matrix, eigenvalues

AMS Mathematical Subject Classification: 65F10, 65Y20

NIR EMISSION SPECTRA OF (Er³⁺: Yb³⁺) DOPED SiO₂-TiO₂-Al₂O₃ SOL-GEL GLASSES

Qing Xiang¹, Chan Hin Kam¹, Kantisara Pita¹ and Srinivasa Buddhudu^{2*}

¹Photonics Research Group, School of Electrical and Electronic Engineering
Nanyang Technological University, Nanyang Avenue, Singapore 639798

²Department of Physics, S.V.University, Tirupati-517502, India

* Corresponding Author. E-mail: profsb_svu@hotmail.com

Received: 20 February 2002/ Accepted: 12 May 2002/ Published: 22 August 2002

Abstract: The present paper reports the near infrared (NIR) emission spectra of Er³⁺: Yb³⁺ co-doped SiO₂-Al₂O₃-TiO₂ sol-gel glasses. Upon optical pumping either with 980nm(Diode Laser) or 488nm (Ar⁺ ion laser), the NIR emission transition (⁴I_{13/2}→⁴I_{15/2}) at 1550nm of (Er³⁺:Yb³⁺) glasses have revealed an improved intensity by many times compared with the Er³⁺ singly doped sol-gel glass. The dependence of the emission intensity and lifetimes on the Yb³⁺/Er³⁺ values and also the Er³⁺ concentration change has been investigated.

Keywords: (Er³⁺:Yb³⁺) sol-gel glasses-emission spectra

PACS: 81.20F; 32.30; 78.55

STRUCTURAL AND TEM ANALYSIS OF $Zn_{1-x}Mn_xTe$ CRYSTALS

K. Veera Brahmam

Department of Physics, Lenora College of Engineering, Rampachodavaram – 533 288, INDIA

E-mail Address: kveerabrahmam@rediffmail.com

Received: 22 June 2002/ Accepted: 12 August 2002/ Published: 22 August 20002

Abstract: Single crystals of $Zn_{1-x}Mn_xTe$ have been grown using vertical Bridgman growth technique. XRD analysis supports the zincblende structure of $Zn_{1-x}Mn_xTe$. As the concentration of Mn increases ($x > 2.5$), the single crystalline nature deteriorates (i.e grain size decreases) and polycrystalline nature with zincblende and hexagonal phases have been observed from x-ray diffractograms. This is supported by Transmission Electron Microscopic study of the samples. Surface morphology is studied from optical microscopic studies.

Keywords: $Zn_{1-x}Mn_xTe$ (ZMT)Crystals, Transmission electron microscopy (TEM), Bridgman growth technique, Optical microscopy.

PACS: 81.20F; 32.30; 78.55

CONDENSATION ON IONS

Borislav V. Toshev

Physical Chemistry Department, University of Sofia, 1, James Bourchier Blvd., 1126 Sofia, Bulgaria

E-mail: toshev@chem.uni-sofia.bg

Received: 26 February 2002/ Accepted: 15 May 2002/ Published: 22 August 2002

Abstract: The transition of a homogeneous system into a heterogeneous one may occur spontaneously in a labile region of supersaturations; in the metastable region of supersaturations it is realized as an energy barrier-determined process. The limit that separates the labile region from the metastable one is the *boundary supersaturation* $s_m \gg s_{cr}$, where s_{cr} is the *critical supersaturation*. The Ω -potential thermodynamic formalism, $\Omega = F - G$, where F and G are the Helmholtz and the Gibbs free energies, respectively, allows a straightforward derivation of the formula of Tohmfor and Volmer (1938) for a nucleus formation on an ion. For the case of charged liquid drops, expressions for the capillary pressure, the vapor pressure, the work of formation of equilibrium drops from vapors, and the drop surface tension, are obtained as well. The stability conditions for the charged equilibrium drops are also examined.

Keywords: Nucleation, charged drops, critical supersaturation, boundary supersaturation, thermodynamics, Ω - potential

PACS: 82.60.N Thermodynamics of nucleation
