



On the spectrum of a fourth order nonlinear eigenvalue problem with variable exponent and indefinite potential

Qing-Mei Zhou¹ and Ke-Qi Wang^{✉ 2}

¹Library, Northeast Forestry University, Harbin, 150040, P.R. China

²College of Mechanical and Electrical Engineering, Northeast Forestry University
Harbin, 150040, P.R. China

Received 14 January 2016, appeared 2 August 2016

Communicated by Dimitri Mugnai

Abstract. The present paper deals with the spectrum of a fourth order nonlinear eigenvalue problem involving variable exponent conditions and a sign-changing potential. The main result of this paper establishes the existence of two positive constants λ_0 and λ_1 with $\lambda_0 \leq \lambda_1$ such that every $\lambda \in [\lambda_1, +\infty)$ is an eigenvalue, while $\lambda \in (-\infty, \lambda_0)$ cannot be an eigenvalue of the above problem.

Keywords: fourth order elliptic equation, eigenvalue problem, variable exponent, indefinite potential.

2010 Mathematics Subject Classification: 35J35, 35P30, 46E35.

1 Introduction

The study of problems of elliptic equations and variational problems with $p(x)$ -growth condition has attracted more and more attention in recent years. It possesses a solid background in physics and originates from the study on electrorheological fluids (see Růžička [12]) and elastic mechanics (see Zhikov [14]). It also has wide applications in different research fields, such as image processing models (see e.g. [5, 9]), stationary thermorheological viscous flows (see [1]) and the mathematical description of the filtration processes of an ideal barotropic gas through a porous medium (see [2]).

In this paper, we are concerned with the study of the nonhomogeneous eigenvalue problem

$$\begin{cases} \Delta(|\Delta u|^{p_1(x)-2}\Delta u) + \Delta(|\Delta u|^{p_2(x)-2}\Delta u) + V(x)|u|^{\alpha(x)-2}u \\ \quad = \lambda(|u|^{q_1(x)-2}u + |u|^{q_2(x)-2}u), & \text{in } \Omega, \\ u = \Delta u = 0, & \text{on } \partial\Omega, \end{cases} \quad (P)$$

where Ω is a bounded domain of \mathbb{R}^N with smooth boundary, λ is a real number, V is an indefinite weight function, and $p_1, p_2, q_1, q_2, \alpha$ are continuous functions on $\bar{\Omega}$.

[✉] Corresponding author. Email: zdhwkq@gmail.com

The study of fourth order nonlinear eigenvalue problems involving variable exponents growth conditions has captured a special attention in the last few years; see e.g. [3, 4, 8, 11]. We give in what follows a concise but complete image of the actual stage of research on this topic.

In the case that $q_1(x) = q_2(x) = q(x)$ for any $x \in \overline{\Omega}$ and $V \equiv 0$ in Ω , Ge and Zhou [8] established the existence of two positive constants λ_0 and λ_1 with $\lambda_0 \leq \lambda_1$ such that any $\lambda \in [\lambda_1, +\infty)$ is an eigenvalue, while $\lambda \in (0, \lambda_0)$ is not an eigenvalue of the above problem.

The same problem, for $V(x) = 0$, $p_1(x) = p_2(x)$ and $q_1(x) = q_2(x)$ is studied by Ayoujil and Amrouss in [3]. The authors established the existence of infinitely many eigenvalues for problem (P) by using an argument based on the Ljusternik–Schnirelmann critical point theory. Denoting by Λ the set of all nonnegative eigenvalues, they showed that $\sup \Lambda = +\infty$ and they pointed out that only under special conditions, which are somehow connected with a kind of monotony of the function $p(x)$, we have $\inf \Lambda > 0$ (this is in contrast with the case when $p(x)$ is a constant; then, we always have $\inf \Lambda > 0$).

This paper is a natural outgrowth of the results in [8]. We consider the eigenvalues for a fourth order nonlinear eigenvalue problem (P) with the potential $V \neq 0$ and $q_1 \neq q_2$ on $\overline{\Omega}$ in the right-hand side.

In this paper we study problem (P) under the following assumptions:

$H(p_1, p_2, q_1, q_2, \alpha)$:

$$1 < p_2(x) < q_2^- \leq q_2^+ \leq \alpha(x) \leq q_1^- \leq q_1^+ < p_1(x) < \frac{N}{2}, \quad \forall x \in \overline{\Omega},$$

$$q_1^+ < \frac{Np_2(x)}{N - 2p_2(x)}, \quad \forall x \in \overline{\Omega},$$

where $q_i^- = \min_{x \in \overline{\Omega}} q_i(x)$ and $q_i^+ = \max_{x \in \overline{\Omega}} q_i(x)$ ($i = 1, 2$);

$H(V)$: $V \in L^{r(x)}(\Omega)$, with $r \in C(\overline{\Omega})$ and $r(x) > \frac{N}{\min_{\overline{\Omega}} \alpha}$, $\forall x \in \overline{\Omega}$.

Inspired by the above-mentioned papers, we study problem (P) from a more extensive viewpoint. More precisely, we will show the existence of two constants λ_0 and λ_1 with $\lambda_0 \leq \lambda_1$ such that any $\lambda \geq \lambda_1$ is an eigenvalue of problem (P) while any $\lambda < \lambda_0$ is not an eigenvalue of problem (P).

This paper is composed of three sections. In Section 2, we recall the definition of variable exponent Lebesgue spaces, $L^{p(x)}(\Omega)$, as well as Sobolev spaces, $W^{1,p(x)}(\Omega)$. Moreover, some properties of these spaces will be also exhibited to be used later. In Section 3, we give the main results and their proofs.

2 Preliminaries

In this section we first recall some facts on variable exponent spaces $L^{p(x)}(\Omega)$ and $W^{1,p(x)}(\Omega)$. For details, we refer to [6, 7, 10].

Set

$$C_+(\overline{\Omega}) = \{h \in C(\overline{\Omega}) : h(x) > 1 \text{ for any } x \in \overline{\Omega}\}.$$

Define

$$h^- = \min_{x \in \overline{\Omega}} h(x), \quad h^+ = \max_{x \in \overline{\Omega}} h(x) \quad \text{for any } h \in C_+(\overline{\Omega}).$$

For any $p \in C_+(\overline{\Omega})$, we define the variable exponent Lebesgue space:

$$L^{p(x)}(\Omega) = \left\{ u : u \text{ is a measurable real value function } \int_{\Omega} |u(x)|^{p(x)} dx < +\infty \right\},$$

with the norm $|u|_{L^{p(x)}(\Omega)} = |u|_{p(x)} = \inf \{ \lambda > 0 : \int_{\Omega} |\frac{u(x)}{\lambda}|^{p(x)} dx \leq 1 \}$, and define the variable exponent Sobolev space

$$W^{k,p(x)}(\Omega) = \{ u \in L^{p(x)}(\Omega) : D^{\alpha} u \in L^{p(x)}(\Omega), |\alpha| \leq k \},$$

with the norm $\|u\|_{W^{k,p(x)}(\Omega)} = \|u\|_{k,p(x)} = \sum_{|\alpha| \leq k} |D^{\alpha} u|_{p(x)}$.

We remember that spaces $L^{p(x)}(\Omega)$ and $W^{k,p(x)}(\Omega)$ are separable and reflexive Banach spaces. Denoting by $W_0^{k,p(x)}(\Omega)$ the closure of $C_0^{\infty}(\Omega)$ in $W^{k,p(x)}(\Omega)$.

For $p(x) \in C_+(\overline{\Omega})$, by $L^{q(x)}(\Omega)$ we denote the conjugate space of $L^{p(x)}(\Omega)$ with $\frac{1}{p(x)} + \frac{1}{q(x)} = 1$, then the Hölder's type inequality

$$\int_{\Omega} |uv| dx \leq \left(\frac{1}{p^-} + \frac{1}{q^-} \right) |u|_{L^{p(x)}(\Omega)} |v|_{L^{q(x)}(\Omega)}, \quad u \in L^{p(x)}(\Omega), v \in L^{q(x)}(\Omega) \quad (2.1)$$

holds. Furthermore, define mapping $\rho : L^{p(x)} \rightarrow \mathbb{R}$ by

$$\rho(u) = \int_{\Omega} |u|^{p(x)} dx,$$

then the following relations hold

$$|u|_{p(x)} < 1 (= 1, > 1) \Leftrightarrow \rho(u) < 1 (= 1, > 1), \quad (2.2)$$

$$|u|_{p(x)} > 1 \Rightarrow |u|_{p(x)}^{p^-} \leq \rho(u) \leq |u|_{p(x)}^{p^+}, \quad (2.3)$$

$$|u|_{p(x)} < 1 \Rightarrow |u|_{p(x)}^{p^+} \leq \rho(u) \leq |u|_{p(x)}^{p^-}. \quad (2.4)$$

$$|u_n - u|_{p(x)} \rightarrow 0 \Leftrightarrow \rho(u_n - u) \rightarrow 0. \quad (2.5)$$

Definition 2.1. Assume that spaces E, F are Banach spaces, we define the norm on the space $X := E \cap F$ as $\|u\|_X = \|u\|_E + \|u\|_F$.

In order to discuss problem (P), we need some theories on space

$$X_i := W_0^{1,p_i(x)}(\Omega) \cap W^{2,p_i(x)}(\Omega)$$

($i = 1, 2$). Since $p_1(x) > p_2(x)$ for any $x \in \overline{\Omega}$, so the space $W_0^{1,p_1(x)}(\Omega)$ is continuously embedded in $W_0^{1,p_2(x)}(\Omega)$, $W^{2,p_1(x)}(\Omega)$ is continuously embedded in $W^{2,p_2(x)}(\Omega)$, so X_1 is continuously embedded in X_2 .

From the Definition 2.1, it follows that for any $u \in X_1$, $\|u\|_1 = \|u\|_{1,p(x)} + \|u\|_{2,p(x)}$, thus $\|u\|_1 = |u|_{p(x)} + |\nabla u|_{p(x)} + \sum_{|\alpha|=2} |D^{\alpha} u|_{p(x)}$.

In Zang and Fu [13], the equivalence of the norms was proved, and it was even proved that the norm $|\Delta u|_{p(x)}$ is equivalent to the norm $\|u\|_1$ (see [11, Theorem 4.4]).

Let us choose on X_1 the norm defined by $\|u\|_1 = |\Delta u|_{p(x)}$. Note that, $(X_1, \|\cdot\|_1)$ is also a separable and reflexive Banach space. Similar to (2.2), (2.3), (2.4) and (2.5), we have the following, define mapping $\rho_1 : X_1 \rightarrow \mathbb{R}$ by

$$\rho_1(u) = \int_{\Omega} |\Delta u|^{p(x)} dx,$$

then the following relations hold

$$\|u\|_1 < 1 (= 1, > 1) \Leftrightarrow \rho_1(u) < 1 (= 1, > 1), \quad (2.6)$$

$$\|u\|_1 > 1 \Rightarrow \|u\|_1^{p^-} \leq \rho_1(u) \leq \|u\|_1^{p^+}, \quad (2.7)$$

$$\|u\|_1 < 1 \Rightarrow \|u\|_1^{p^+} \leq \rho_1(u) \leq \|u\|_1^{p^-}. \quad (2.8)$$

$$\|u_n - u\|_1 \rightarrow 0 \Leftrightarrow \rho_1(u_n - u) \rightarrow 0. \quad (2.9)$$

Hereafter, let

$$p_1^*(x) = \begin{cases} \frac{Np_1(x)}{N-2p_1(x)}, & p_1(x) < \frac{N}{2}, \\ +\infty, & p_1(x) \geq \frac{N}{2}, \end{cases}$$

and

$$p_2^*(x) = \begin{cases} \frac{Np_2(x)}{N-2p_2(x)}, & p_2(x) < \frac{N}{2}, \\ +\infty, & p_2(x) \geq \frac{N}{2}. \end{cases}$$

Remark 2.2. If $h \in C_+(\overline{\Omega})$ and $h(x) < p_i^*(x)$ ($i = 1$ or 2) for any $x \in \overline{\Omega}$, by Theorem 3.2 in [3], we deduce that X_i is continuously and compactly embedded in $L^{h(x)}(\Omega)$.

Remark 2.3. Since $p_2(x) < p_1(x)$ for any $x \in \Omega$ it follows that $p_2^*(x) < p_1^*(x)$, using condition $H(p_1, p_2, q_1, q_2, \alpha)$ we have a compact embedding $X_i \hookrightarrow L^{q_1(x)}(\Omega)$ and $X_i \hookrightarrow L^{q_2(x)}(\Omega)$ with ($i = 1, 2$).

3 Main results and proofs

Since $p_2(x) < p_1(x)$ for any $x \in \Omega$ it follows that $W_0^{1,p_1(x)}(\Omega)$ and $W^{2,p_1(x)}(\Omega)$ are continuously embedded in $W_0^{1,p_2(x)}(\Omega)$ and $W^{2,p_2(x)}(\Omega)$ respectively. Thus, a solution for a problem of type (P) will be sought in the variable exponent space $X_1 = W_0^{1,p_1(x)}(\Omega) \cap W^{2,p_1(x)}(\Omega)$.

We say that $\lambda \in \mathbb{R}$ is an eigenvalue of problem (P) if there exists $u \in X_1 \setminus \{0\}$ such that

$$\int_{\Omega} \left(|\Delta u|^{p_1(x)-2} + |\Delta u|^{p_2(x)-2} \right) \Delta u \Delta v dx + \int_{\Omega} V(x) |u|^{\alpha(x)-2} u v dx - \lambda \int_{\Omega} \left(|u|^{q_1(x)-2} + |u|^{q_2(x)-2} \right) u v dx = 0,$$

for all $v \in X_1$. We point out that if λ is an eigenvalue of problem (P), then the corresponding eigenfunction $u \in X_1 \setminus \{0\}$ is a weak solution of problem (P).

Define

$$\lambda_1 := \inf_{u \in X_1 \setminus \{0\}} \frac{\int_{\Omega} \frac{1}{p_1(x)} |\Delta u|^{p_1(x)} dx + \int_{\Omega} \frac{1}{p_2(x)} |\Delta u|^{p_2(x)} dx + \int_{\Omega} \frac{V(x)}{\alpha(x)} |u|^{\alpha(x)} dx}{\int_{\Omega} \frac{1}{q_1(x)} |u|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u|^{q_2(x)} dx}.$$

Our main result is given by the following theorem.

Theorem 3.1. *Assume that $\mathbf{H}(p_1, p_2, q_1, q_2, \alpha)$ holds. Then each $\lambda \in [\lambda_1, +\infty)$ is an eigenvalue of problem (P). Furthermore, there exists a constant λ_0 such that $\lambda_0 \leq \lambda_1$ and any $\lambda \in (-\infty, \lambda_0)$ is not an eigenvalue of problem (P).*

Proof. The proof is divided into the following five steps.

Step 1. For each $\varepsilon > 0$, there exists $c_\varepsilon > 0$ such that

$$\left| \int_{\Omega} \frac{V(x)}{\alpha(x)} |u|^{\alpha(x)} dx \right| \leq \varepsilon \left[\int_{\Omega} \frac{1}{p_1(x)} |\Delta u|^{p_1(x)} dx + \int_{\Omega} \frac{1}{p_2(x)} |\Delta u|^{p_2(x)} dx \right] + c_\varepsilon |V|_{r(x)} \int_{\Omega} (|u|^{\alpha^-} + |u|^{\alpha^+}) dx \quad (3.1)$$

for any $u \in X_1$.

Since $r(x) \geq r^-$ on $\bar{\Omega}$, it follows that $L^{r(x)}(\Omega) \subseteq L^{r^-(\Omega)}$. Note that $r(x) > \frac{N}{\alpha^-}$ for each $x \in \bar{\Omega}$, which implies that $V \in L^{r^-}(\Omega)$ and $r^- > \frac{N}{\alpha^-}$.

Applying the Hölder's inequality, we get

$$\int_{\Omega} |V(x)| |u|^{\alpha^-} dx \leq |V|_{r^-} \left| |u|^{\alpha^-} \right|_{(r^-)'} = |V|_{r^-} |u|_{\alpha^-(r^-)'}^{\alpha^-}, \quad (3.2)$$

where $\frac{1}{r^-} + \frac{1}{(r^-)'} = 1$.

Now for every $\varepsilon > 0$, we can show that for any $s \in (1, \frac{N\alpha^-}{N-2\alpha^-})$, there exists $C'_\varepsilon > 0$ such that

$$|u|_s \leq \varepsilon |\Delta u|_{\alpha^-} + C'_\varepsilon |u|_{\alpha^-}, \quad \forall u \in W_0^{1,\alpha^-}(\Omega) \cap W^{2,\alpha^-}(\Omega). \quad (3.3)$$

Indeed, assume it is not true for each $\varepsilon > 0$. Then there exists $\varepsilon_0 > 0$ and a sequence $\{v_n\} \subseteq W_0^{1,\alpha^-}(\Omega) \cap W^{2,\alpha^-}(\Omega)$ such that $|v_n|_s = 1$ and

$$\varepsilon_0 |\Delta v_n|_{\alpha^-} + n |v_n|_{\alpha^-} < 1, \quad \forall n \in N.$$

So, sequence $\{v_n\}$ is bounded in $W_0^{1,\alpha^-}(\Omega) \cap W^{2,\alpha^-}(\Omega)$ and $|v_n|_{\alpha^-} \rightarrow 0$, as $n \rightarrow +\infty$. Thus, up to a subsequence, still denote by $\{v_n\}$, we may assume that there exists $v \in W_0^{1,\alpha^-}(\Omega) \cap W^{2,\alpha^-}(\Omega)$, such that $v_n \rightharpoonup v$ in $W_0^{1,\alpha^-}(\Omega) \cap W^{2,\alpha^-}(\Omega)$ and actually $v = 0$. Since $s \in (1, \frac{N\alpha^-}{N-2\alpha^-})$, it follows by Remark 2.2 that $W_0^{1,\alpha^-}(\Omega) \cap W^{2,\alpha^-}(\Omega)$ is compactly embedded in $L^s(\Omega)$, which implies that $v_n \rightarrow 0$ in $L^s(\Omega)$. On the other hand, since $|v_n|_s = 1$ for each $n \in N$, we deduce that $|v|_s = 1$, which is a contradiction. Hence (3.3) is true.

Note that $r^- > \frac{N}{\alpha^-}$ and $(r^-)', \alpha^- < \frac{N\alpha^-}{N-2\alpha^-}$. From (3.2) and (3.3), we have

$$\begin{aligned} \int_{\Omega} |V(x)| |u|^{\alpha^-} dx &\leq |V|_{r^-} |u|_{\alpha^-(r^-)'}^{\alpha^-} \\ &\leq |V|_{r^-} (\varepsilon |\Delta u|_{\alpha^-} + C'_\varepsilon |u|_{\alpha^-})^{\alpha^-} \\ &\leq 2^{\alpha^- - 1} \varepsilon^{\alpha^-} |V|_{r^-} |\Delta u|_{\alpha^-}^{\alpha^-} + 2^{\alpha^- - 1} (C'_\varepsilon)^{\alpha^-} |V|_{r^-} |u|_{\alpha^-}^{\alpha^-}, \end{aligned} \quad (3.4)$$

for all $u \in W_0^{1,\alpha^-}(\Omega) \cap W^{2,\alpha^-}(\Omega)$.

Combining the similar arguments as those used in the proof of relation (3.4) with $r^- > \frac{N}{\alpha^+}$. We know that there exists $C_\varepsilon'' > 0$ such that

$$\int_{\Omega} |V(x)| |u|^{\alpha^+} dx \leq 2^{\alpha^+-1} \varepsilon^{\alpha^+} |V|_{r^-} |\Delta u|_{\alpha^+}^{\alpha^+} + 2^{\alpha^+-1} (C_\varepsilon'')^{\alpha^+} |V|_{r^-} |u|_{\alpha^+}^{\alpha^+}, \quad (3.5)$$

for all $u \in W_0^{1,\alpha^+}(\Omega) \cap W^{2,\alpha^+}(\Omega)$.

By virtue of hypothesis $\mathbf{H}(\mathbf{p}_1, \mathbf{p}_2, \mathbf{q}_1, \mathbf{q}_2, \alpha)$, we deduce that $\alpha^- \leq \alpha^+ < p_1(x)$ for any $x \in \bar{\Omega}$ and thus $X_1 \subseteq W_0^{1,\alpha^-}(\Omega) \cap W^{2,\alpha^-}(\Omega)$ and $X_1 \subseteq W_0^{1,\alpha^+}(\Omega) \cap W^{2,\alpha^+}(\Omega)$. Hence, relations (3.4) and (3.5) are true for any $u \in X_1$.

On the other hand, by $p_2(x) < \alpha^- \leq \alpha(x) \leq \alpha^+ < p_1(x)$ for each $x \in \bar{\Omega}$, we deduce that

$$\left| \int_{\Omega} V(x) |u|^{\alpha^+} dx \right| \leq \frac{1}{\alpha^-} \int_{\Omega} |V(x)| (|u|^{\alpha^-} + |u|^{\alpha^+}) dx, \quad \forall u \in X_1 \quad (3.6)$$

and

$$\int_{\Omega} (|\Delta u|^{\alpha^-} + |\Delta u|^{\alpha^+}) dx \leq 2p_1^+ \int_{\Omega} \left(\frac{1}{p_1(x)} |\Delta u|^{p_1(x)} + \frac{1}{p_2(x)} |\Delta u|^{p_2(x)} \right) dx. \quad (3.7)$$

From (3.4), (3.5), (3.6) and (3.7), we have

$$\begin{aligned} \left| \int_{\Omega} V(x) |u|^{\alpha^+} dx \right| &\leq \frac{1}{\alpha^-} \int_{\Omega} |V(x)| (|u|^{\alpha^-} + |u|^{\alpha^+}) dx \\ &\leq \frac{1}{\alpha^-} \left[2^{\alpha^+-1} \varepsilon^{\alpha^+} |V|_{r^-} |\Delta u|_{\alpha^-}^{\alpha^-} + 2^{\alpha^+-1} (C_\varepsilon'')^{\alpha^+} |V|_{r^-} |u|_{\alpha^-}^{\alpha^-} \right. \\ &\quad \left. + 2^{\alpha^+-1} \varepsilon^{\alpha^+} |V|_{r^-} |\Delta u|_{\alpha^+}^{\alpha^+} + 2^{\alpha^+-1} (C_\varepsilon'')^{\alpha^+} |V|_{r^-} |u|_{\alpha^+}^{\alpha^+} \right] \\ &\leq \frac{2^{\alpha^+-1} |V|_{r^-}}{\alpha^-} \left[\varepsilon^{\alpha^-} (|\Delta u|_{\alpha^-}^{\alpha^-} + |\Delta u|_{\alpha^+}^{\alpha^+}) + (C_\varepsilon'')^{\alpha^+} (|u|_{\alpha^-}^{\alpha^-} + |u|_{\alpha^+}^{\alpha^+}) \right] \\ &\leq \frac{2^{\alpha^+-1} |V|_{r^-}}{\alpha^-} \left[2p_1^+ \varepsilon^{\alpha^-} \int_{\Omega} \left(\frac{1}{p_1(x)} |\Delta u|^{p_1(x)} + \frac{1}{p_2(x)} |\Delta u|^{p_2(x)} \right) dx \right. \\ &\quad \left. + (C_\varepsilon'')^{\alpha^+} \int_{\Omega} (|u(x)|^{\alpha^-} + |u(x)|^{\alpha^+}) dx \right]. \end{aligned} \quad (3.8)$$

Combining the last inequality with arbitrariness of ε , we infer that relation (3.1) is true.

Step 2. We show that

$$\lim_{\|u\|_1 \rightarrow +\infty} \frac{\int_{\Omega} \frac{1}{p_1(x)} |\Delta u|^{p_1(x)} dx + \int_{\Omega} \frac{1}{p_2(x)} |\Delta u|^{p_2(x)} dx + \int_{\Omega} \frac{V(x)}{\alpha(x)} |u|^{\alpha(x)} dx}{\int_{\Omega} \frac{1}{q_1(x)} |u|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u|^{q_2(x)} dx} = +\infty \quad (3.9)$$

and

$$\lim_{\|u\|_1 \rightarrow 0} \frac{\int_{\Omega} \frac{1}{p_1(x)} |\Delta u|^{p_1(x)} dx + \int_{\Omega} \frac{1}{p_2(x)} |\Delta u|^{p_2(x)} dx + \int_{\Omega} \frac{V(x)}{\alpha(x)} |u|^{\alpha(x)} dx}{\int_{\Omega} \frac{1}{q_1(x)} |u|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u|^{q_2(x)} dx} = +\infty. \quad (3.10)$$

Since $q_1(x) < \alpha^-, \alpha^+ < q_2(x), \forall x \in \bar{\Omega}$, we deduce that for every $x \in \bar{\Omega}$ and $u \in X_1$,

$$|u(x)|^{\alpha^-} + |u(x)|^{\alpha^+} \leq 2(|u(x)|^{q_1(x)} + |u(x)|^{q_2(x)}),$$

which implies that

$$\frac{\int_{\Omega} (|u(x)|^{\alpha^-} + |u(x)|^{\alpha^+}) dx}{\int_{\Omega} (|u(x)|^{q_1(x)} + |u(x)|^{q_2(x)}) dx} \leq 2. \quad (3.11)$$

From Step 1, we find that for $\varepsilon \in (0, 1)$ there exists $c_\varepsilon > 0$ such that

$$\begin{aligned} & \frac{\int_{\Omega} \frac{1}{p_1(x)} |\Delta u|^{p_1(x)} dx + \int_{\Omega} \frac{1}{p_2(x)} |\Delta u|^{p_2(x)} dx + \int_{\Omega} \frac{V(x)}{\alpha(x)} |u|^{\alpha(x)} dx}{\int_{\Omega} \frac{1}{q_1(x)} |u|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u|^{q_2(x)} dx} \\ & \geq \frac{\frac{1-\varepsilon}{p_1^+} \int_{\Omega} (|\Delta u|^{p_1(x)} dx + |\Delta u|^{p_2(x)} dx) - c_\varepsilon |V|_{r(x)} \int_{\Omega} (|u|^{\alpha^-} + |u|^{\alpha^+}) dx}{\frac{1}{q_2} \int_{\Omega} (|u|^{q_1(x)} + |u|^{q_2(x)}) dx} \end{aligned} \quad (3.12)$$

for any $u \in X_1$.

By (3.11) and (3.12), there exist some positive constants c_1 and c_2 , such that

$$\begin{aligned} & \frac{\int_{\Omega} \frac{1}{p_1(x)} |\Delta u|^{p_1(x)} dx + \int_{\Omega} \frac{1}{p_2(x)} |\Delta u|^{p_2(x)} dx + \int_{\Omega} \frac{V(x)}{\alpha(x)} |u|^{\alpha(x)} dx}{\int_{\Omega} \frac{1}{q_1(x)} |u|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u|^{q_2(x)} dx} \\ & \geq c_1 \frac{\int_{\Omega} (|\Delta u|^{p_1(x)} dx + |\Delta u|^{p_2(x)} dx)}{\int_{\Omega} (|u|^{q_1(x)} + |u|^{q_2(x)}) dx} - c_2 |V|_{r(x)} \end{aligned} \quad (3.13)$$

for any $u \in X_1$.

Note that $q_1^+, q_2^+ < \frac{Np_2(x)}{N-2p_2(x)} < \frac{Np_1(x)}{N-2p_1(x)}$, $\forall x \in \bar{\Omega}$. So the embedding $X_1 \hookrightarrow L^{q_i^\pm}(\Omega)$ ($i = 1, 2$) is continuous and compact. Thus, for any $u \in X_1$, there positive constant c_3 , such that

$$\|u\|_{q_i^\pm} \leq c_3 \|u\|_1. \quad (3.14)$$

Thus, by (3.14), we have

$$\begin{aligned} & \int_{\Omega} \frac{1}{q_1(x)} |u|^{q_2(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u|^{q_2(x)} dx \\ & \leq \int_{\Omega} (|u|^{q_1(x)} + |u|^{q_2(x)}) dx \\ & \leq \int_{\Omega} (|u|^{q_1^-} + |u|^{q_1^+} + |u|^{q_2^-} + |u|^{q_2^+}) dx \\ & \leq c_3^{q_1^-} \|u\|_1^{q_1^-} + c_3^{q_1^+} \|u\|_1^{q_1^+} + c_3^{q_2^-} \|u\|_1^{q_2^-} + c_3^{q_2^+} \|u\|_1^{q_2^+}. \end{aligned} \quad (3.15)$$

Hence, from (3.13) and (3.15), for any $u \in X_1$ with $\|u\|_1 > 1$, there exists positive constant c_4 , such that

$$\begin{aligned} & \frac{\int_{\Omega} \frac{1}{p_1(x)} |\Delta u|^{p_1(x)} dx + \int_{\Omega} \frac{1}{p_2(x)} |\Delta u|^{p_2(x)} dx + \int_{\Omega} \frac{V(x)}{\alpha(x)} |u|^{\alpha(x)} dx}{\int_{\Omega} \frac{1}{q_1(x)} |u|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u|^{q_2(x)} dx} \\ & \geq c_1 \frac{\int_{\Omega} |\Delta u|^{p_1(x)} dx}{\int_{\Omega} \frac{1}{q_1(x)} |u|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u|^{q_2(x)} dx} - c_2 |V|_{r(x)} \\ & \geq c_4 \frac{\int_{\Omega} |\Delta u|^{p_1(x)} dx}{\|u\|_1^{q_1^-} + \|u\|_1^{q_1^+} + \|u\|_1^{q_2^-} + \|u\|_1^{q_2^+}} - c_2 |V|_{r(x)} \\ & \geq c_4 \frac{\|u\|_1^{p_1^-}}{\|u\|_1^{q_1^-} + \|u\|_1^{q_1^+} + \|u\|_1^{q_2^-} + \|u\|_1^{q_2^+}} - c_2 |V|_{r(x)} \\ & \rightarrow +\infty, \quad \text{as } \|u\|_1 \rightarrow +\infty, \end{aligned} \quad (3.16)$$

because $p_1^- > q_1^+ \geq q_1^- \geq q_2^+ \geq q_2^-$. So, the relation (3.9) holds.

On the other hand, using (3.13) and (3.14), for any $u \in X_1$ with $\|u\|_1 < 1$ small enough, there exists positive constant c_5 , such that

$$\begin{aligned}
& \frac{\int_{\Omega} \frac{1}{p_1(x)} |\Delta u|^{p_1(x)} dx + \int_{\Omega} \frac{1}{p_2(x)} |\Delta u|^{p_2(x)} dx + \int_{\Omega} \frac{V(x)}{\alpha(x)} |u|^{\alpha(x)} dx}{\int_{\Omega} \frac{1}{q_1(x)} |u|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u|^{q_2(x)} dx} \\
& \geq c_1 \frac{\int_{\Omega} |\Delta u|^{p_2(x)} dx}{\int_{\Omega} \frac{1}{q_1(x)} |u|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u|^{q_2(x)} dx} - c_2 |V|_{r(x)} \\
& \geq c_5 \frac{\int_{\Omega} |\Delta u|^{p_2(x)} dx}{\|u\|_1^{q_1^+} + \|u\|_1^{q_1^-} + \|u\|_1^{q_2^+} + \|u\|_1^{q_2^-}} - c_2 |V|_{r(x)} \\
& \geq c_4 \frac{\|u\|_1^{p_2^+}}{\|u\|_1^{q_1^-} + \|u\|_1^{q_1^+} + \|u\|_1^{q_2^-} + \|u\|_1^{q_2^+}} - c_2 |V|_{r(x)}.
\end{aligned} \tag{3.17}$$

Since $p_2^+ < q_2^- \leq q_2^+ \leq q_1^- \leq q_1^+$, passing to the limit as $\|u\|_1 \rightarrow 0$ in the above inequality we deduce that relation (3.10) holds true.

Step 3. There exists $u_0 \in X_1 \setminus \{0\}$ such that

$$\frac{\int_{\Omega} \frac{1}{p_1(x)} |\Delta u_0|^{p_1(x)} dx + \int_{\Omega} \frac{1}{p_2(x)} |\Delta u_0|^{p_2(x)} dx + \int_{\Omega} \frac{V(x)}{\alpha(x)} |u_0|^{\alpha(x)} dx}{\int_{\Omega} \frac{1}{q_1(x)} |u_0|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u_0|^{q_2(x)} dx} = \lambda_1.$$

In fact, let $\{u_n\} \subseteq X_1 \setminus \{0\}$ be a minimizing sequence for λ_1 , that is,

$$\lim_{n \rightarrow \infty} \frac{\int_{\Omega} \frac{1}{p_1(x)} |\Delta u_n|^{p_1(x)} dx + \int_{\Omega} \frac{1}{p_2(x)} |\Delta u_n|^{p_2(x)} dx + \int_{\Omega} \frac{V(x)}{\alpha(x)} |u_n|^{\alpha(x)} dx}{\int_{\Omega} \frac{1}{q_1(x)} |u_n|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u_n|^{q_2(x)} dx} = \lambda_1. \tag{3.18}$$

By Step 2, we have $\{u_n\}$ is bounded in X_1 . Since X_1 is reflexive it follows that there exists $u_0 \in X_1$ such that $u_n \rightharpoonup u_0$ in X_1 . On the other hand, the function $\int_{\Omega} \frac{1}{p_1(x)} |\Delta u|^{p_1(x)} dx + \int_{\Omega} \frac{1}{p_2(x)} |\Delta u|^{p_2(x)} dx : X_1 \rightarrow \mathbb{R}$ is a convex function, hence it is weakly lower semi-continuous, that is,

$$\begin{aligned}
& \liminf_{n \rightarrow \infty} \left(\int_{\Omega} \frac{1}{p_1(x)} |\Delta u_n|^{p_1(x)} dx + \int_{\Omega} \frac{1}{p_2(x)} |\Delta u_n|^{p_2(x)} dx \right) \\
& \geq \left(\int_{\Omega} \frac{1}{p_1(x)} |\Delta u_0|^{p_1(x)} dx + \int_{\Omega} \frac{1}{p_2(x)} |\Delta u_0|^{p_2(x)} dx \right).
\end{aligned} \tag{3.19}$$

Note that X_1 is compactly embedded in $L^{q_i(x)}(\Omega)$ ($i = 1, 2$), thus,

$$u_n \rightarrow u_0 \quad \text{in } L^{q_i(x)}(\Omega) \quad (i = 1, 2). \tag{3.20}$$

By (2.9) and (3.20), it is easy to see that

$$\begin{aligned}
& \lim_{n \rightarrow \infty} \int_{\Omega} \left[\frac{1}{q_1(x)} |u_n|^{q_1(x)} dx + \frac{1}{q_2(x)} |u_n|^{q_2(x)} dx \right] \\
& = \int_{\Omega} \frac{1}{q_1(x)} |u_0|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u_0|^{q_2(x)} dx.
\end{aligned} \tag{3.21}$$

On the other hand, by $X_1 \hookrightarrow L^{\alpha(x)r'(x)}(\Omega)$, we know that

$$\begin{aligned} \left| \int_{\Omega} V(x) |u_n - u_0|^{\alpha(x)} dx \right| &\leq 2 |V|_{r(x)} \left| |u_n - u_0|^{\alpha(x)} \right|_{r'(x)} \\ &\leq 2 |V|_{r(x)} |u_n - u_0|^{\alpha_{\alpha(x)r'(x)}} \\ &\rightarrow 0, \quad \text{as } n \rightarrow \infty, \end{aligned}$$

where

$$\tau = \begin{cases} +, & |u_n - u_0|_{\alpha(x)r'(x)} \geq 1, \\ -, & |u_n - u_0|_{\alpha(x)r'(x)} < 1, \end{cases}$$

which implies that

$$\lim_{n \rightarrow \infty} \int_{\Omega} V(x) |u_n|^{\alpha(x)} dx = \int_{\Omega} V(x) |u_0|^{\alpha(x)} dx. \quad (3.22)$$

In view of (3.19), (3.21) and (3.22), we obtain

$$\frac{\int_{\Omega} \frac{1}{p_1(x)} |\Delta u_0|^{p_1(x)} dx + \int_{\Omega} \frac{1}{p_2(x)} |\Delta u_0|^{p_2(x)} dx + \int_{\Omega} \frac{V(x)}{\alpha(x)} |u_0|^{\alpha(x)} dx}{\int_{\Omega} \frac{1}{q_1(x)} |u_0|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u_0|^{q_2(x)} dx} = \lambda_1,$$

if $u_0 \neq 0$.

It remains to show that u_0 is nontrivial. Assume the contrary. Then $u_n \rightarrow 0$ in X_1 and $u_n \rightarrow 0$ in $L^{\theta(x)}(\Omega)$ for any $\theta \in C(\bar{\Omega})$ with $1 < \theta(x) < p_1^*(x)$ on $\bar{\Omega}$. Thus, we have

$$\lim_{n \rightarrow \infty} \left[\int_{\Omega} \frac{1}{q_1(x)} |u_n|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u_n|^{q_2(x)} dx \right] = 0 \quad (3.23)$$

and

$$\lim_{n \rightarrow \infty} \int_{\Omega} \frac{V(x)}{\alpha(x)} |u_n|^{\alpha(x)} dx = 0. \quad (3.24)$$

Let $\varepsilon \in (0, \lambda_1)$ be fixed. By (3.18) we deduce that for n large enough we have

$$\begin{aligned} &\left| \int_{\Omega} \frac{1}{p_1(x)} |\Delta u_n|^{p_1(x)} dx + \int_{\Omega} \frac{1}{p_2(x)} |\Delta u_n|^{p_2(x)} dx + \int_{\Omega} \frac{V(x)}{\alpha(x)} |u_n|^{\alpha(x)} dx \right. \\ &\quad \left. - \lambda_1 \left(\int_{\Omega} \frac{1}{q_1(x)} |u_n|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u_n|^{q_2(x)} dx \right) \right| \\ &< \varepsilon \left(\int_{\Omega} \frac{1}{q_1(x)} |u_n|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u_n|^{q_2(x)} dx \right), \end{aligned}$$

which yields that

$$\begin{aligned} &(\lambda_1 - \varepsilon) \left(\int_{\Omega} \frac{1}{q_1(x)} |u_n|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u_n|^{q_2(x)} dx \right) \\ &< \int_{\Omega} \frac{1}{p_1(x)} |\Delta u_n|^{p_1(x)} dx + \int_{\Omega} \frac{1}{p_2(x)} |\Delta u_n|^{p_2(x)} dx + \int_{\Omega} \frac{V(x)}{\alpha(x)} |u_n|^{\alpha(x)} dx \\ &< (\lambda_1 + \varepsilon) \left(\int_{\Omega} \frac{1}{q_1(x)} |u_n|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u_n|^{q_2(x)} dx \right). \end{aligned} \quad (3.25)$$

Combining (3.23), (3.24) and (3.25), we have

$$\lim_{n \rightarrow \infty} \int_{\Omega} \frac{1}{p_1(x)} |\Delta u_n|^{p_1(x)} dx + \int_{\Omega} \frac{1}{p_2(x)} |\Delta u_n|^{p_2(x)} dx = 0. \quad (3.26)$$

By (2.9) and (3.26), we have

$$u_n \rightarrow 0 \quad \text{in } X_1, \quad \text{that is } \|u_n\|_1 \rightarrow 0.$$

From this information and relation (3.10), we get

$$\lim_{\|u_n\|_1 \rightarrow 0} \frac{\int_{\Omega} \frac{1}{p_1(x)} |\Delta u_n|^{p_1(x)} dx + \int_{\Omega} \frac{1}{p_2(x)} |\Delta u_n|^{p_2(x)} dx + \int_{\Omega} \frac{V(x)}{\alpha(x)} |u_n|^{\alpha(x)} dx}{\int_{\Omega} \frac{1}{q_1(x)} |u_n|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u_n|^{q_2(x)} dx} = +\infty$$

and this is a contradiction. Thus $u_0 \neq 0$.

By Step 3, we conclude that there exists $u_0 \in X_1 \setminus \{0\}$ such that

$$\begin{aligned} \lambda_1 &= \frac{\int_{\Omega} \frac{1}{p_1(x)} |\Delta u_0|^{p_1(x)} dx + \int_{\Omega} \frac{1}{p_2(x)} |\Delta u_0|^{p_2(x)} dx + \int_{\Omega} \frac{V(x)}{\alpha(x)} |u_0|^{\alpha(x)} dx}{\int_{\Omega} \frac{1}{q_1(x)} |u_0|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u_0|^{q_2(x)} dx} \\ &= \inf_{u \in X_1 \setminus \{0\}} \frac{\int_{\Omega} \frac{1}{p_1(x)} |\Delta u|^{p_1(x)} dx + \int_{\Omega} \frac{1}{p_2(x)} |\Delta u|^{p_2(x)} dx + \int_{\Omega} \frac{V(x)}{\alpha(x)} |u|^{\alpha(x)} dx}{\int_{\Omega} \frac{1}{q_1(x)} |u|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u|^{q_2(x)} dx}. \end{aligned}$$

Then for any $v \in X_1$ we have

$$\left. \frac{d}{dt} \left(\frac{\int_{\Omega} \frac{1}{p_1(x)} |\Delta(u_0 + tv)|^{p_1(x)} dx + \int_{\Omega} \frac{1}{p_2(x)} |\Delta(u_0 + tv)|^{p_2(x)} dx + \int_{\Omega} \frac{V(x)}{\alpha(x)} |(u_0 + tv)|^{\alpha(x)} dx}{\int_{\Omega} \frac{1}{q_1(x)} |u_0 + tv|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u_0 + tv|^{q_2(x)} dx} \right) \right|_{t=0} = 0.$$

A simple computation yields

$$\begin{aligned} & \left(\int_{\Omega} (|\Delta u_0|^{p_1(x)-2} + |\Delta u_0|^{p_2(x)-2}) \Delta u_0 \Delta v dx + \int_{\Omega} V(x) |u_0|^{\alpha(x)-2} u_0 v dx \right) \\ & \quad \times \left(\int_{\Omega} \frac{1}{q_1(x)} |u_0|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u_0|^{q_2(x)} dx \right) \\ &= \left(\int_{\Omega} \frac{1}{p_1(x)} |\Delta u_0|^{p_1(x)} dx + \int_{\Omega} \frac{1}{p_2(x)} |\Delta u_0|^{p_2(x)} dx + \int_{\Omega} \frac{1}{\alpha(x)} |u_0|^{\alpha(x)} dx \right) \\ & \quad \times \left(\int_{\Omega} |u_0|^{q_1(x)-2} u_0 v dx + \int_{\Omega} |u_0|^{q_2(x)-2} u_0 v dx \right) \end{aligned} \tag{3.27}$$

for any $v \in X_1$.

Returning to (3.27) and using

$$\frac{\int_{\Omega} \frac{1}{p_1(x)} |\Delta u_0|^{p_1(x)} dx + \int_{\Omega} \frac{1}{p_2(x)} |\Delta u_0|^{p_2(x)} dx + \int_{\Omega} \frac{V(x)}{\alpha(x)} |u_0|^{\alpha(x)} dx}{\int_{\Omega} \frac{1}{q_1(x)} |u_0|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u_0|^{q_2(x)} dx} = \lambda_1$$

and $\int_{\Omega} \frac{1}{q_1(x)} |u_0|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u_0|^{q_2(x)} dx \neq 0$, we obtain λ_1 is an eigenvalue of problem (P).

Step 4. We will show that any $\lambda \in (\lambda_1, +\infty)$ is an eigenvalue of problem (P).

Let $\lambda \in (\lambda_1, +\infty)$ be arbitrary but fixed. Define $D_{\lambda} : X_1 \rightarrow \mathbb{R}$ by

$$\begin{aligned} D_{\lambda}(u) &= \int_{\Omega} \frac{1}{p_1(x)} |\Delta u|^{p_1(x)} dx + \int_{\Omega} \frac{1}{p_2(x)} |\Delta u|^{p_2(x)} dx + \int_{\Omega} \frac{V(x)}{\alpha(x)} |u|^{\alpha(x)} dx \\ & \quad - \lambda \left(\int_{\Omega} \frac{1}{q_1(x)} |u|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u|^{q_2(x)} dx \right). \end{aligned}$$

Clearly, $D_\lambda \in C^1(X_1, \mathbb{R})$ with

$$\begin{aligned} \langle D'_\lambda(u), v \rangle &= \int_\Omega (|\Delta u|^{p_1(x)-2} + |\Delta u|^{p_2(x)-2}) \Delta u \Delta v dx + \int_\Omega V(x) |u|^{\alpha(x)-2} u v dx \\ &\quad - \lambda \int_\Omega (|u|^{q_1(x)-2} + |u|^{q_2(x)-2}) u v dx, \quad \forall u, v \in X_1. \end{aligned}$$

Thus, λ is an eigenvalue of problem (P) if and only if there exists $u_\lambda \in X_1 \setminus \{0\}$ a critical point of D_λ .

It follows from Step 1 and (3.15) that

$$\begin{aligned} D_\lambda(u) &= \int_\Omega \frac{1}{p_1(x)} |\Delta u|^{p_1(x)} dx + \int_\Omega \frac{1}{p_2(x)} |\Delta u|^{p_2(x)} dx + \int_\Omega \frac{V(x)}{\alpha(x)} |u|^{\alpha(x)} dx \\ &\quad - \lambda \left(\int_\Omega \frac{1}{q_1(x)} |u|^{q_1(x)} dx + \int_\Omega \frac{1}{q_2(x)} |u|^{q_2(x)} dx \right) \\ &\geq \frac{1}{p_1^+} \int_\Omega (|\Delta u|^{p_1(x)} + |\Delta u|^{p_2(x)}) dx + \int_\Omega \frac{V(x)}{\alpha(x)} |u|^{\alpha(x)} dx \\ &\quad - \lambda \left(\int_\Omega \frac{1}{q_1(x)} |u|^{q_1(x)} dx + \int_\Omega \frac{1}{q_2(x)} |u|^{q_2(x)} dx \right) \\ &\geq \frac{1-\varepsilon}{p_1^+} \int_\Omega (|\Delta u|^{p_1(x)} + |\Delta u|^{p_2(x)}) dx - c_\varepsilon |V|_{r(x)} \int_\Omega (|u|^{\alpha^-} + |u|^{\alpha^+}) dx \\ &\quad - \lambda c_3^{q_1^-} \|u\|_1^{q_1^-} + c_3^{q_1^+} \|u\|_1^{q_1^+} + c_3^{q_2^-} \|u\|_1^{q_2^-} + c_3^{q_2^+} \|u\|_1^{q_2^+} \\ &\geq \frac{1-\varepsilon}{p_1^+} \int_\Omega |\Delta u|^{p_1(x)} dx - c_\varepsilon |V|_{r(x)} (|u|^{\alpha^-} + |u|^{\alpha^+}) \\ &\quad - \lambda c_3^{q_1^-} \|u\|_1^{q_1^-} + c_3^{q_1^+} \|u\|_1^{q_1^+} + c_3^{q_2^-} \|u\|_1^{q_2^-} + c_3^{q_2^+} \|u\|_1^{q_2^+} \\ &\geq \frac{1-\varepsilon}{p_1^+} \|u\|_1^{p_1^-} - c_\varepsilon |V|_{r(x)} (c \|u\|_1^{\alpha^-} + c \|u\|_1^{\alpha^+}) - \lambda c_3^{q_1^-} \|u\|_1^{q_1^-} + c_3^{q_1^+} \|u\|_1^{q_1^+} \\ &\quad + c_3^{q_2^-} \|u\|_1^{q_2^-} + c_3^{q_2^+} \|u\|_1^{q_2^+} \\ &\rightarrow \infty, \quad \text{as } \|u\|_1 \rightarrow +\infty, \end{aligned}$$

since $1 < q_2^- \leq q_2^+ \leq \alpha^- \leq \alpha^+ \leq q_1^- \leq q_1^+ < p_1^-$, i.e. $\lim_{\|u\|_1 \rightarrow +\infty} D_\lambda(u) = \infty$.

On the other hand, with similar arguments as in the proof of Step 3, we can show that the functional D_λ is weakly lower semi-continuous. So by the Weierstrass theorem, there exists $u_\lambda \in X_1$ a global minimum point of D_λ and thus, a critical point of D_λ . Next we prove that u_λ is nontrivial.

Indeed, since

$$\lambda_1 = \inf_{u \in X_1 \setminus \{0\}} \frac{\int_\Omega \frac{1}{p_1(x)} |\Delta u|^{p_1(x)} dx + \int_\Omega \frac{1}{p_2(x)} |\Delta u|^{p_2(x)} dx + \int_\Omega \frac{V(x)}{\alpha(x)} |u|^{\alpha(x)} dx}{\int_\Omega \frac{1}{q_1(x)} |u|^{q_1(x)} dx + \int_\Omega \frac{1}{q_2(x)} |u|^{q_2(x)} dx}$$

and $\lambda > \lambda_1$ it follows that there exists $v_\lambda \in X_1$ such that $D_\lambda(v_\lambda) < 0$, that is

$$\begin{aligned} \int_\Omega \frac{1}{p_1(x)} |\Delta v_\lambda|^{p_1(x)} dx + \int_\Omega \frac{1}{p_2(x)} |\Delta v_\lambda|^{p_2(x)} dx + \int_\Omega \frac{V(x)}{\alpha(x)} |v_\lambda|^{\alpha(x)} dx \\ < \lambda \left(\int_\Omega \frac{1}{q_1(x)} |v_\lambda|^{q_1(x)} dx + \int_\Omega \frac{1}{q_2(x)} |v_\lambda|^{q_2(x)} dx \right). \end{aligned}$$

Thus, we conclude that

$$\inf_{u \in X_1} D_\lambda(u) < 0$$

and u_λ is a nontrivial critical point of D_λ , and then λ is an eigenvalue of problem (P).

Step 5. We will show that any $\lambda \in (0, \lambda_0)$ is not an eigenvalue of problem (P), where λ_0 is given by

$$\lambda_0 := \inf_{u \in X_1 \setminus \{0\}} \frac{\int_{\Omega} (|\Delta u|^{p_1(x)} dx + |\Delta u|^{p_2(x)} dx) \int_{\Omega} V(x) |u|^{\alpha(x)} dx}{\int_{\Omega} |u|^{q_1(x)} dx + \int_{\Omega} |u|^{q_2(x)} dx}.$$

Firstly, we verify that $\lambda_0 \leq \lambda_1$. Due to Step 2, λ_1 and u_0 is an eigenvalue and is an eigenfunction corresponding to λ_1 of (P), then for every $v \in X_1$ we have

$$\begin{aligned} \int_{\Omega} (|\Delta u_0|^{p_1(x)-2} + |\Delta u_0|^{p_2(x)-2}) \Delta u_0 \Delta v dx + \int_{\Omega} V(x) |u_0|^{\alpha(x)-2} u_0 v dx \\ = \lambda_1 \left(\int_{\Omega} |u_0|^{q_1(x)-2} u_0 v dx + \int_{\Omega} |u_0|^{q_2(x)-2} u_0 v dx \right), \end{aligned} \quad (3.28)$$

which implies that

$$\begin{aligned} \int_{\Omega} (|\Delta u_0|^{p_1(x)-2} + |\Delta u_0|^{p_2(x)-2}) \Delta u_0 \Delta u_0 dx + \int_{\Omega} V(x) |u_0|^{\alpha(x)-2} u_0 u_0 dx \\ = \lambda_1 \left(\int_{\Omega} |u_0|^{q_1(x)-2} u_0 u_0 dx + \int_{\Omega} |u_0|^{q_2(x)-2} u_0 u_0 dx \right), \end{aligned}$$

that is

$$\int_{\Omega} (|\Delta u_0|^{p_1(x)} + |\Delta u_0|^{p_2(x)}) dx + \int_{\Omega} V(x) |u_0|^{\alpha(x)} dx = \lambda_1 \left(\int_{\Omega} |u_0|^{q_1(x)} dx + \int_{\Omega} |u_0|^{q_2(x)} dx \right).$$

Then, it follows that $\lambda_0 \leq \lambda_1$.

Now we prove our assertion. Arguing by contradiction: assume that there exists $\lambda \in (0, \lambda_0)$ is an eigenvalue of problem (P). Thus, there exists $u_\lambda \in X_1 \setminus \{0\}$ such that

$$\begin{aligned} \int_{\Omega} (|\Delta u_\lambda|^{p_1(x)-2} + |\Delta u_\lambda|^{p_2(x)-2}) \Delta u_\lambda \Delta v dx + \int_{\Omega} V(x) |u_\lambda|^{\alpha(x)-2} u_\lambda v dx \\ = \lambda \left(\int_{\Omega} |u_\lambda|^{q_1(x)-2} u_\lambda v dx + \int_{\Omega} |u_\lambda|^{q_2(x)-2} u_\lambda v dx \right) \end{aligned} \quad (3.29)$$

for any $v \in X_1$. Thus, for $v = u_\lambda$ we have

$$\int_{\Omega} (|\Delta u_\lambda|^{p_1(x)} + |\Delta u_\lambda|^{p_2(x)}) dx + \int_{\Omega} V(x) |u_\lambda|^{\alpha(x)} dx = \lambda \left(\int_{\Omega} |u_\lambda|^{q_1(x)} dx + \int_{\Omega} |u_\lambda|^{q_2(x)} dx \right).$$

The fact that $u_\lambda \in X_1 \setminus \{0\}$ assures that $\int_{\Omega} |u_\lambda|^{q_1(x)} dx + \int_{\Omega} |u_\lambda|^{q_2(x)} dx > 0$. Since $\lambda < \lambda_0$, the above information yields

$$\begin{aligned} \int_{\Omega} (|\Delta u_\lambda|^{p_1(x)} + |\Delta u_\lambda|^{p_2(x)}) dx + \int_{\Omega} V(x) |u_\lambda|^{\alpha(x)} dx \\ \geq \lambda_0 \int_{\Omega} (|u_\lambda|^{q_1(x)} + |u_\lambda|^{q_2(x)}) dx \\ > \lambda \int_{\Omega} (|u_\lambda|^{q_1(x)} + |u_\lambda|^{q_2(x)}) dx \\ = \int_{\Omega} (|\Delta u_\lambda|^{p_1(x)} + |\Delta u_\lambda|^{p_2(x)}) dx + \int_{\Omega} V(x) |u_\lambda|^{\alpha(x)} dx. \end{aligned}$$

Obviously, this is contradiction. Therefore, we deduce that each $\lambda \in (-\infty, \lambda_0)$ is not an eigenvalue of problem (P). \square

Remark 3.2. In Theorem 3.1, we are not able to deduce whether $\lambda_0 = \lambda_1$ or $\lambda_0 < \lambda_1$. In the latter case an interesting question concerns the existence of eigenvalue of the problem (P) in the interval $[\lambda_0, \lambda_1)$.

Acknowledgements

This work is supported by the National Natural Science Foundation of China (No. 11201095), the Youth Scholar Backbone Supporting Plan Project of Harbin Engineering University (No. 307201411008), the Fundamental Research Funds for the Central Universities (No. 2016), Heilong Jiang Postdoctoral Funds for scientific research initiation (No. LBH-Q14044), the Science Research Funds for Overseas Returned Chinese Scholars of Heilongjiang Province (No. LC201502).

References

- [1] S. N. ANTONTSEV, J. F. RODRIGUES, On stationary thermo-rheological viscous flows, *Ann. Univ. Ferrara Sez. VII, Sci. Mat.* **52**(2006), 19–36. [MR2246902](#); [url](#)
- [2] S. N. ANTONTSEV, S. I. SHMAREV, A model porous medium equation with variable exponent of nonlinearity: existence uniqueness and localization properties of solutions, *Nonlinear Anal.* **60**(2005), 515–545. [MR2103951](#); [url](#)
- [3] A. AYOUIJIL, A. R. EL AMROUSS, On the spectrum of a fourth order elliptic equation with variable exponent, *Nonlinear Anal.* **71**(2009), 4916–4926. [MR2781059](#); [url](#)
- [4] A. AYOUIJIL, A. R. EL AMROUSS, Continuous spectrum of a fourth order nonhomogenous elliptic equation with variable exponent, *Electron. J. Differential Equations* **2011**, No. 24, 1–12. [MR2781059](#)
- [5] Y. CHEN, S. LEVINE, M. RAO, Variable exponent linear growth functionals in image restoration, *SIAM J. Appl. Math.* **66**(2006): 1383–1406. [MR2246061](#); [url](#)
- [6] X. L. FAN, D. ZHAO, On the space $L^{p(x)}(\Omega)$ and $W^{m,p(x)}(\Omega)$, *J. Math. Anal. Appl.* **263**(2001), 424–446. [MR1866056](#); [url](#)
- [7] X. L. FAN, Q. H. ZHANG, Existence of solutions for $p(x)$ -Laplacian Dirichlet problems, *Nonlinear Anal.* **52**(2003), 1843–1852. [MR1954585](#); [url](#)
- [8] B. GE, Q. M. ZHOU, Continuous spectrum of a fourth order nonhomogeneous differential operators with variable exponent, *Electron. J. Qual. Theory Differ.* **2013**, No. 18, 1–11. [MR3037220](#); [url](#)
- [9] P. HARJULEHTO, P. HÄSTÖ, V. LATVALA, Minimizers of the variable exponent, non-uniformly convex Dirichlet energy, *J. Math. Pures Appl. (9)* **89**(2008), 174–197. [MR2391646](#); [url](#)
- [10] O. KOVÁČIK, J. RÁKOSNÍK, On spaces $L^{p(x)}(\Omega)$ and $W^{m,p(x)}(\Omega)$, *Czechoslovak Math.* **41**(1991), 592–618. [MR1134951](#)

- [11] M. MIHĂILESCU, V. RĂDULESCU, Continuous spectrum for a class of nonhomogeneous differential operators, *Manuscripta Math.* **125**(2008), 157–167. [MR2373080](#); [url](#)
- [12] M. RŮŽIČKA, *Electrorheological fluids: modeling and mathematical theory*, Springer-Verlag, Berlin, 2000. [MR1810360](#); [url](#)
- [13] A. B. ZANG, Y. FU, Interpolation inequalities for derivatives in variable exponent Lebesgue–Sobolev spaces, *Nonlinear Anal.* **69**(2008), 3629–3636. [MR2450565](#); [url](#)
- [14] V. V. ZHIKOV, Averaging of functionals of the calculus of variations and elasticity theory, *Izv. Akad. Nauk SSSR Ser. Mat.* **50**(1986), 675–710. [MR0864171](#)