

## A certain class of quadratures with cubic B-spline as a weight function <sup>1</sup>

Zlatko Udovičić

### Abstract

We consider the quadrature rules of “practical type” (with five knots) for approximately computation of the integral

$$\int_{-2}^2 B(x)f(x)dx,$$

where  $B(\cdot)$  denotes centered cubic B-spline. We proved that maximal algebraic degree of exactness for this type of formulas is equal to five. At the end we gave numerical result.

**2010 Mathematics Subject Classification:** 65D30

**Key words and phrases:** Practical quadratures, Cubic B-spline

## 1 Introduction

Cardinal B-splines play very important role in the approximation theory (different methods for solving initial and boundary value problems, multiresolution approximation, spline interpolation, etc). Cardinal B-spline of order one is the characteristic function of the interval  $[0, 1)$ . Cardinal B-spline of order  $m, m \in \mathbb{N}, m \geq 2$ , is a function with the following properties:

- its support is interval  $[0, m]$ ;

---

<sup>1</sup>Received 16 December, 2009

Accepted for publication (in revised form) 11 January, 2010

- it belongs to the class  $C^{m-2}[0, m]$ ;
- at each interval  $[k, k + 1], k \in \{0, 1, \dots, m - 1\}$ , it is a polynomial of degree  $m - 1$ .

The most often used in practice is a cardinal B-spline of order four (cubic B-spline). Approximation obtained by using this spline is relatively good, and on the other hand, this spline is still simple for manipulation. Centered cubic B-spline is defined in the following way:

$$B(x) = \begin{cases} \frac{1}{6}(x+2)^3, & x \in [-2, -1), \\ \frac{1}{6}(-3x^3 - 6x^2 + 4), & x \in [-1, 0), \\ \frac{1}{6}(3x^3 - 6x^2 + 4), & x \in [0, 1), \\ \frac{1}{6}(-x+2)^3, & x \in [1, 2], \\ 0, & \text{otherwise.} \end{cases}$$

Thereat, the problem of calculation of the integral

$$\int_{-2}^2 B(x)f(x)dx$$

is unavoidable. In this paper we are investigating a certain class of quadratures (the so called quadratures of “practical type”) for approximate computation of the previous integral. The paper was motivated by results recently published in [1] and [2], where the same class of quadratures was considered, but without weight function. Similar problem (with hat function as a weight function) was considered in [4].

We say that quadrature formula

$$(1) \quad \int_{-2}^2 B(x)f(x)dx = \sum_{i=1}^5 A_i f(x_i) + R[f]$$

is of “practical type” if the following conditions hold:

1.  $A_1 = A_5$  and  $A_2 = A_4$ ;
2. nodes  $x_k, 1 \leq k \leq 5$  are symmetric and rational numbers from the interval  $[-2, 2]$ , i.e.  $x_1 = -r_1, x_2 = -r_2, x_3 = 0, x_4 = r_2$  and  $x_5 = r_1$ , for some  $r_1, r_2 \in (0, 2] \cap \mathbb{Q}, r_2 < r_1$  (as usuall,  $\mathbb{Q}$  denotes the set of rational numbers).

Hence, quadratures of “practical type” have the following form:

$$(2) \quad \int_{-2}^2 B(x)f(x)dx = A(f(-r_1) + f(r_1)) + B(f(-r_2) + f(r_2)) + Cf(0) + R[f],$$

for some  $r_1, r_2 \in (0, 2] \cap \mathbb{Q}, r_2 < r_1$ .

Quadrature rule (1) has algebraic degree of exactness equal to  $m, m \in \mathbb{N}$ , if and only if  $R[p] = 0$  whenever  $p(\cdot)$  is a polynomial of degree not greater than  $m$  and there exists the polynomial  $q(\cdot)$ , of degree  $m + 1$ , such that  $R[q] \neq 0$ . Our aim is construction of the quadrature rules of “practical type” with maximal algebraic degree of exactness.

We will finish this section with some well known facts from the theory of numerical integration.

**Lemma 1** *Quadrature rule (1) (i.e. (2)) has algebraic degree of exactness equal to  $m, m \in \mathbb{N}$  if and only if  $R[x^k] = 0$  for all  $k \in \{0, 1, \dots, m\}$ .*

**Lemma 2** *Quadrature rule (2) is exact for every odd function  $f(\cdot)$  (i.e.  $R[f] = 0$  for every odd function  $f(\cdot)$ ).*

From the previous lemmas follows that algebraic degree of exactness of the formula (2) has to be odd.

Finally, with the choice

$$x_{1245} = \pm \sqrt{\frac{14315 \pm \sqrt{70203301}}{11886}}, x_3 = 0$$

(expressions for the coefficients  $A_k, 1 \leq k \leq 5$ , are much more complicated, so we omit them here) formula (1) attains maximal algebraic degree of exactness (which is equal to nine), but this formula obviously is not of “practical type”. Hence, algebraic degree of exactness of the formula (2) can not be greater than seven.

## 2 Main result

Let us determine the coefficients  $A, B$  and  $C$  such that formula (2) has maximal algebraic degree of exactness.

From the condition that formula is exact for  $f(x) = 1$  (i.e. exact for any polynomial of zero degree) simply follows

$$C = 1 - 2(A + B).$$

Then formula (2) becomes

$$(3) \quad \int_{-2}^2 B(x)f(x)dx = A(f(-r_1) - 2f(0) + f(r_1)) + \\ B(f(-r_2) - 2f(0) + f(r_2)) + \\ f(0) + R[f]$$

and in accordance with the previous, this formula has algebraic degree of exactness equal to one. Furthermore, conditions that the last formula is exact for  $f(x) = x^2$  and  $f(x) = x^4$  give the following system of linear equations

$$\begin{aligned} 2r_1^2 A + 2r_2^2 B &= \frac{1}{3} \\ 2r_1^4 A + 2r_2^4 B &= \frac{3}{10}. \end{aligned}$$

which has the solution

$$(4) \quad A = \frac{9 - 10r_2^2}{60r_1^2(r_1^2 - r_2^2)} \text{ and } B = \frac{9 - 10r_1^2}{60r_2^2(r_2^2 - r_1^2)}.$$

Hence, with this choice of the coefficients  $A$  and  $B$  formula (3) has algebraic degree of exactness equal to five. Therein (in formula (3))

$$R[x^6] = \frac{17}{42} - \frac{9r_1^2 - 10r_1^2 r_2^2 + 9r_2^2}{30}.$$

It is natural to ask is it possible to choose rational nodes  $r_1$  and  $r_2$  such that formula (3) has algebraic degree of exactness equal to six, i.e. seven. Negative answer to this question gives the following lemma.

**Lemma 3** *There is no numbers  $r_1, r_2 \in (0, 2] \cap \mathbb{Q}$  such that*

$$(5) \quad \frac{9r_1^2 - 10r_1^2 r_2^2 + 9r_2^2}{30} = \frac{17}{42}.$$

**Proof.** Let us assume contrary, i.e. that  $r_1 = \frac{a}{b}$  and  $r_2 = \frac{c}{d}$ , for some  $a, b, c, d \in \mathbb{N}$  such that  $(a, b) = 1$  and  $(c, d) = 1$ . Putting this in equality (5), after simplification, gives

$$(6) \quad 7(9a^2d^2 - 10a^2c^2 + 9b^2c^2) = 85b^2d^2,$$

from which follows that  $b^2d^2 \equiv 0 \pmod{7}$ , i.e.  $bd \equiv 0 \pmod{7}$ .

Let  $b = 7k$  for some  $k \in \mathbb{N}$ . After simplification, equality (6) becomes

$$a^2(9d^2 - 10c^2) = 7k^2(85d^2 - 63c^2).$$

It can not be  $a^2 \equiv 0 \pmod{7}$  (because of  $(a, b) = 1$ ), so it has to be  $9d^2 - 10c^2 \equiv 0 \pmod{7}$ . Furthermore, because of  $9d^2 - 10c^2 = 7(d^2 - 2c^2) + 2(d^2 + 2c^2)$ , it also has to be  $d^2 + 2c^2 \equiv 0 \pmod{7}$ . Direct checking verify that this relation is impossible unless  $c \equiv 0 \pmod{7}$  and  $d \equiv 0 \pmod{7}$ , which together with an assumption  $(c, d) = 1$  gives the contradiction.

The case  $d = 7k$ , for some  $k \in \mathbb{N}$ , because of symmetry of the relation (6) can be proved analogue. This completes the proof.

Let us estimate the error of the formula (3), under assumption that the coefficients  $A$  and  $B$  are given by the equalities (4). Let  $H_5(\cdot)$  be Hermite's interpolating polynomial which interpolates the function  $f(\cdot)$  through the points  $\pm r_1, \pm r_2$  and 0, where the node 0 has multiplicity two. Then (see for example [3], p. 55),

$$f(x) - H_5(x) = \frac{f^{(vi)}(\xi(x))}{6!} x^2(x^2 - r_1^2)(x^2 - r_2^2),$$

and the error of the formula (3) is given by

$$\begin{aligned} R[f] &= \int_{-2}^2 B(x) \frac{f^{(vi)}(\xi(x))}{6!} x^2(x^2 - r_1^2)(x^2 - r_2^2) dx \\ &= \frac{1}{3 \cdot 6!} f^{(vi)}(\eta) (\eta^2 - r_1^2)(\eta^2 - r_2^2), \end{aligned}$$

for some  $\eta \in [-2, 2]$ , assuming  $f(\cdot) \in C^6[-2, 2]$ . Let

$$\Phi(\eta) = (\eta^2 - r_1^2)(\eta^2 - r_2^2).$$

It is easy to check that

$$\begin{aligned} \max_{\eta \in [-2, 2]} |\Phi(\eta)| &= \max \left\{ |\Phi(0)|, \left| \Phi\left(\sqrt{\frac{r_1^2 + r_2^2}{2}}\right) \right|, |\Phi(2)| \right\} \\ &= \max \left\{ r_1^2 r_2^2, \frac{(r_1^2 - r_2^2)^2}{4}, (4 - r_1^2)(4 - r_2^2) \right\}, \end{aligned}$$

so the error of the formula (3) can be estimated in the following way

$$(7) \quad |R[f]| \leq \frac{M_6}{3 \cdot 6!} \max \left\{ r_1^2 r_2^2, \frac{(r_1^2 - r_2^2)^2}{4}, (4 - r_1^2)(4 - r_2^2) \right\},$$

where  $M_6 = \max_{x \in [-2, 2]} |f^{(vi)}(x)|$ .

### 3 Numerical result

Estimation (7) naturally imposes the following problem

$$(8) \quad F(r_1, r_2) = \max \left\{ r_1^2 r_2^2, \frac{(r_1^2 - r_2^2)^2}{4}, (4 - r_1^2)(4 - r_2^2) \right\} \rightarrow \min .,$$

where  $r_1, r_2 \in (0, 2] \cap \mathbb{Q}$ ,  $r_2 < r_1$ . It is obvious that, for fixed  $r_1 \in (0, 2] \cap \mathbb{Q}$ , the function  $F(\cdot, \cdot)$  attains its minimum in one of the intersection points among three curves  $r_1^2 r_2^2$ ,  $\frac{(r_1^2 - r_2^2)^2}{4}$  and  $(4 - r_1^2)(4 - r_2^2)$ .

1. Curves  $r_1^2 r_2^2$  and  $\frac{(r_1^2 - r_2^2)^2}{4}$  ( $r_1$  is fixed) intersect each other at  $r_2 = \pm(1 \pm \sqrt{2})r_1$ , and since  $r_2 \notin \mathbb{Q}$  we will not consider this case.
2. Similarly, curves  $\frac{(r_1^2 - r_2^2)^2}{4}$  and  $(4 - r_1^2)(4 - r_2^2)$  ( $r_1$  is fixed) intersect each other at  $r_2 = \pm \sqrt{3r_1^2 - 8 \pm 2\sqrt{2}(r_1^2 - 1)}$ , and again because of  $r_2 \notin \mathbb{Q}$  we will not consider this case.
3. Finally, curves  $r_1^2 r_2^2$  and  $(4 - r_1^2)(4 - r_2^2)$  ( $r_1$  is still fixed) intersect each other at  $r_2 = \sqrt{4 - r_1^2}$ , and we will look for the nodes  $r_1$  and  $r_2$  among "rational points" from the centered circle with radius equal to two.

In the Table 1 we give some admissible values of the nodes  $r_1$  and  $r_2$  for which the function  $F(\cdot, \cdot)$  attains its local minimums. The corresponding rational numbers are round off to the five decimal places.

At the end, let us say that, by using any of the given choices for the nodes  $r_1$  and  $r_2$ , the error (7) can be estimated in the following way

$$|R[f]| \leq 0.2 \cdot 10^{-2} \cdot M_6.$$

Table 1: Some admissible values of the nodes

$r_1$	$r_2$	$F(r_1, r_2)$
$\frac{8}{5} = 1.60000$	$\frac{6}{5} = 1.20000$	3.68640
$\frac{42}{29} = 1.44828$	$\frac{40}{29} = 1.37931$	3.99049
$\frac{110}{73} = 1.50685$	$\frac{96}{73} = 1.31507$	3.92678
$\frac{144}{97} = 1.48454$	$\frac{130}{97} = 1.34021$	3.95845
$\frac{754}{505} = 1.49307$	$\frac{672}{505} = 1.33069$	3.94744
$\frac{1974}{1325} = 1.48981$	$\frac{1768}{1325} = 1.33434$	3.95180
$\frac{2584}{1733} = 1.49106$	$\frac{2310}{1733} = 1.33295$	3.95016
$\frac{13530}{9077} = 1.49058$	$\frac{12104}{9077} = 1.33348$	3.95079
$\frac{35422}{23761} = 1.49076$	$\frac{31680}{23761} = 1.33328$	3.95055
$\frac{46368}{31105} = 1.49069$	$\frac{41474}{31105} = 1.33335$	3.95064

## References

- [1] E. Constantinescu, *A certain class of quadratures*, Acta Univ. Apulensis Math. Inform., No. 7, 2004, 109-116.
- [2] E. Constantinescu, *A certain class of quadratures*, Gen. Math. 10, No. 1-2, 2002, 75-84.
- [3] G. Milovanović, *Numerička analiza II deo*, Naučna knjiga, Beograd, 1998 (in Serbian).
- [4] Z. Udovičić, *A certain class of quadratures with hat function as a weight function*, Acta Univ. Apulensis Math. Inform., No. 20, 2009, 251-258.

### Zlatko Udovičić

University of Sarajevo  
Faculty of Sciences, Department of Mathematics  
Zmaja od Bosne 33-35  
71 000 Sarajevo  
Bosnia and Herzegovina  
e-mail: zzlatko@pmf.unsa.ba