



*Gen. Math. Notes, Vol. 26, No. 2, February 2015, pp. 104-118*

*ISSN 2219-7184; Copyright © ICSRS Publication, 2015*

*www.i-csrs.org*

*Available free online at <http://www.geman.in>*

# **On $\alpha$ - $\eta$ - $\phi$ -Contraction in Fuzzy Metric Space and its Application**

Noori F. AL-Mayahi<sup>1</sup> and Sarim H. Hadi<sup>2</sup>

<sup>1,2</sup>Department of Mathematics, University of AL-Qadissiya  
College of Computer Science and Mathematics

<sup>1</sup>E-mail: [afm2005@yahoo.com](mailto:afm2005@yahoo.com)

<sup>2</sup>E-mail: [sarim.h2014@yahoo.com](mailto:sarim.h2014@yahoo.com)

(Received: 11-8-14 / Accepted: 3-11-14)

## **Abstract**

*The aim of this paper is to introduce new concepts of  $\alpha$ - $\eta$ -complete fuzzy metric space and  $\alpha$ - $\eta$ -continuous function and establish fixed point results for  $\alpha$ - $\eta$ - $\phi$ -contraction function in  $\alpha$ - $\eta$ -complete fuzzy metric space. As an application, we derive some Suzuki type fixed point theorems, fixed point in orbitally  $f$ -complete. Moreover, we introduce concept  $\alpha$ - $\psi$ - $\phi$ -contraction function and application on  $\alpha$ - $\eta$ - $\phi$ -contraction.*

**Keywords:**  *$\alpha$ - $\eta$ -complete,  $\alpha$ - $\eta$ -continuous,  $\alpha$ - $\eta$ - $\phi$ -contraction, orbitally  $f$ -complete, Suzuki Type Fixed Point Result,  $\alpha$ - $\psi$ - $\phi$ -Contraction Function, Application on  $\alpha$ - $\eta$ - $\phi$ -Contraction.*

## **1 Introduction**

The study of fixed points of functions in a fuzzy metric space satisfying certain contractive conditions has been at the center of vigorous research activity. In 1965, the concept of fuzzy sets was introduced by Zadeh [10]. With the concept of fuzzy sets, the fuzzy metric space was introduced by I. Kramosil and J. Michalek

[5] in 1975. Helpem [3] in 1981 first proved a fixed point theorem for fuzzy functions. Also M.Grabiec [2] in 1988 proved the contraction principle in the setting of the fuzzy metric spaces. Moreover, A. George and P. Veeramani [1] in 1994 modified the notion of fuzzy metric spaces with the help of t-norm.

This paper we introduce new concepts of  $\alpha$ - $\eta$ -complete fuzzy metric space and  $\alpha$ - $\eta$ -continuous function and establish fixed point results for  $\alpha$ - $\eta$ - $\phi$ -contraction function in  $\alpha$ - $\eta$ -complete fuzzy metric space. As an application, we derive some Suzuki type fixed point theorems, fixed point in orbitally  $f$ -complete. Moreover, we introduce concept  $\alpha$ - $\psi$ - $\phi$ -contraction function and application on  $\alpha$ - $\eta$ - $\phi$ -contraction.

## 2 Preliminaries

**Definition 2.1 [4]:** Let  $f: X \rightarrow X$  and  $\alpha: X \times X \rightarrow [0, \infty)$  be two function,  $f$  is said to be  $\alpha$ -admissible function if

$$\alpha(x, y) \geq 1 \implies \alpha(f(x), f(y)) \geq 1 \text{ for all } x, y \in X.$$

**Definition 2.2 [6]:** Let  $f: X \rightarrow X$  and  $\alpha, \beta: X \times X \rightarrow [0, \infty)$  be two function,  $f$  is said to be  $(\alpha, \beta)$ -admissible function if

$$\begin{cases} \alpha(x, y) \geq 1 \\ \beta(x, y) \geq 1 \end{cases} \implies \begin{cases} \alpha(f(x), f(y)) \geq 1 \\ \beta(f(x), f(y)) \geq 1 \end{cases} \text{ for all } x, y \in X.$$

**Definition 2.3 [4]:** Let  $f: X \rightarrow X$  and  $\alpha, \eta: X \times X \rightarrow [0, \infty)$  be two function,  $f$  is said to be  $\alpha$ -admissible function with respect  $\eta$  if

$$\alpha(x, y) \geq \eta(x, y) \implies \alpha(f(x), f(y)) \geq \eta(f(x), f(y)) , \text{ for all } x, y \in X$$

Not that if we take  $\eta(x, y) = 1$  , then this definition reduces to definition (2.1) .

**Definition 2.4:** Let  $(X, M, *)$  be a fuzzy metric space and

$\alpha, \eta: X \times X \rightarrow [0, \infty)$  ,  $X$  is said to be  $\alpha$ - $\eta$ -complete iff every Cauchy sequence  $\{x_n\}$  with  $\alpha(x_n, x_{n+1}) \geq \eta(x_n, x_{n+1})$  for all  $n \in \mathbb{N}$  converge in  $X$

Note:  $X$  is said to be  $\alpha$ -complete if  $\eta(x, y) = 1$  for all  $x, y \in X$ .

**Example 2.5:** Let  $X = [0,1]$  and  $M(x, y, t) = \frac{t}{t+|x-y|}$  , if  $t > 0$  with  $a * b = \min \{a, b\}$  for every  $a, b \in [0,1]$  , define  $\alpha, \eta: X \times X \rightarrow [0, \infty)$

By 
$$\alpha(x, y) = \begin{cases} (x + y)^2 & \text{if } x, y \in X \\ 0 & \text{if o.w} \end{cases}$$

$$\eta(x, y) = 2xy$$

Then  $(X, M, *)$  is  $\alpha$ - $\eta$ -complete fuzzy metric space.

**Definition 2.6:** Let  $(X, M, *)$  be a fuzzy metric space and let  $\alpha, \eta: X \times X \rightarrow [0, \infty)$  and  $f: X \rightarrow X$ ,  $f$  is said to be  $\alpha$ - $\eta$ -continuous function on  $X$  if for  $x \in X$  and  $\{x_n\}$  be a sequence in  $X$  with  $x_n \rightarrow x$  as  $n \rightarrow \infty$ ,  $\alpha(x_n, x_{n+1}) \geq \eta(x_n, x_{n+1})$  for all  $n \in \mathbb{N}$  implies  $f(x_n) \rightarrow f(x)$ .

**Definition 2.7 [9]:**

(1) Let  $f$  be a function of a fuzzy metric space  $(X, M, *)$  into itself.  $(X, M, *)$  is said to be  $f$ -orbitally complete if and only if every Cauchy sequence which is contained in  $\{x, f(x), f^2(x), f^3(x), \dots\}$  for some  $x \in X$  converges in  $X$ .

A  $f$ -orbitally complete fuzzy metric space may not be complete.

(2) A function  $f: X \rightarrow X$  is called orbitally continuous at  $x \in X$  if  $\lim_{n \rightarrow \infty} f^n(x) = x$  implies  $\lim_{n \rightarrow \infty} f^n(x) = f(x)$

The function  $f$  is orbitally continuous on  $X$  if  $f$  is orbitally continuous for all  $x \in X$ .

**Remark 2.8:**

(1) Let  $(X, M, *)$  be a fuzzy metric space and  $f: X \rightarrow X$  be a function and let  $X$  be an orbitally  $f$ -complete. Define  $\alpha, \eta: X \times X \rightarrow [0, \infty)$  by

$$\alpha(x, y) = \begin{cases} 3 & \text{if } x, y \in O(w) \\ 1 & \text{o.w} \end{cases}, \quad \eta(x, y) = 1$$

Where  $O(w)$  is an orbit of a point  $w \in X$ .  $(X, M, *)$  is an  $\alpha$ - $\eta$ -complete.

If  $\{x_n\}$  be a Cauchy sequence with  $\alpha(x_n, x_{n+1}) \geq \eta(x_n, x_{n+1})$  for all  $n \in \mathbb{N}$  then  $\{x_n\} \in O(w)$

Now, since  $X$  is an orbitally  $f$ -complete fuzzy metric space, then  $\{x_n\}$  converge in  $X$ . We can say that  $X$  is  $\alpha$ - $\eta$ -complete.

(2) Let  $(X, M, *)$  and  $\alpha, \eta: X \times X \rightarrow [0, \infty)$  is in (1), let  $f: X \rightarrow X$  be an orbitally continuous function on  $(X, M, *)$ . Then  $f$  is  $\alpha$ - $\eta$ -continuous function. Indeed if  $x_n \rightarrow x$  as  $n \rightarrow \infty$  and  $\alpha(x_n, x_{n+1}) \geq \eta(x_n, x_{n+1})$  for all  $n \in \mathbb{N}$ , so  $x_n \in O(w)$  for all  $n \in \mathbb{N}$ , then there exist sequence  $(k_i)_{i \in \mathbb{N}}$  of positive integer such that  $x_n \rightarrow f^{k_i}w \rightarrow x$  as  $i \rightarrow \infty$ . Now since  $f$  is an orbitally continuous on  $(X, M, *)$ , then  $f(x_n) = f(f^{k_i}w) \rightarrow f(x)$  as  $i \rightarrow \infty$ .

Denote with  $\phi$  the set of all the function  $\varphi: [0,1] \rightarrow [0,1]$  with the following properties:

- (1)  $\varphi$  is nondecreasing and continuous
- (2)  $\varphi(\tau) = 0$  iff  $\tau = 1$

**Definition 2.9:** Let  $(X, M, *)$  be a fuzzy metric space and  $f: X \rightarrow X$ , let  $\mathbb{M}(x, y) = \min \{M(x, y, t), M(x, f(x), t), M(y, f(y), t), M(x, f(y), t) * M(y, f(x), t)\}$

We say

- (1)  $f$  is an  $\alpha$ - $\eta$ - $\varphi$ -contraction function if  $\alpha(x, y) \geq \eta(x, y) \Rightarrow M(f(x), f(y), t) \geq \varphi(\mathbb{M}(x, y))$ .
- (2)  $f$  is  $\alpha$ - $\varphi$ -contraction function if  $\eta(x, y) = 1$  for all  $x, y \in X$ .

**Theorem 2.10:** Let  $(X, M, *)$  be a fuzzy metric space and let  $f: X \rightarrow X$ , suppose that  $\alpha, \eta: X \times X \rightarrow [0, \infty)$  are two function. Assume that the following assertions hold

- (i)  $(X, M, *)$  is an  $\alpha$ - $\eta$ -complete fuzzy metric space .
- (ii)  $f$  is an  $\alpha$ -admissible function with respect to  $\eta$  .
- (iii)  $f$  is  $\alpha$ - $\eta$ - $\varphi$ -contraction function on  $X$  .
- (iv)  $f$  is an  $\alpha$ - $\eta$ -continuous function .
- (v) There exist  $x_0 \in X$  such that  $\alpha(x_0, f(x_0)) \geq \eta(x_0, f(x_0))$ .

Then  $f$  has a fixed point in  $X$  .

**Proof:** Let  $x_0 \in X$  such that  $\alpha(x_0, f(x_0)) \geq \eta(x_0, f(x_0))$

Define a sequence  $\{x_n\}$  such that  $x_n = f(x_{n-1})$  for all  $n \in \mathbb{N}$

If  $x_n = x_{n+1}$  for some  $n$  , then  $x = x_n$  is a fixed point of  $f$ .

Suppose  $x_n \neq x_{n+1}$

Since  $f$  is  $\alpha$ -admissible function with respect  $\eta$  and

$$\alpha(x_0, f(x_0)) \geq \eta(x_0, f(x_0))$$

$$\text{Then } \alpha(x_1, x_2) = \alpha(f(x_0), f(x_1)) \geq \eta(f(x_0), f(x_1)) = \eta(x_1, x_2)$$

By continuing this process, we get

$$\alpha(x_n, f(x_n)) = \alpha(x_n, x_{n+1}) \geq \eta(x_n, x_{n+1}) = \eta(x_n, f(x_n))$$

By (1) in definition (2.9)

$$M(x_n, x_{n+1}, t) = M(f(x_{n-1}), f(x_n), t) \geq \alpha(\mathbb{M}(x_{n-1}, x_n))$$

Where

$$\mathbb{M}(x_{n-1}, x_n) = \min \{M(x_{n-1}, x_n, t), M(x_{n-1}, f(x_{n-1}), t), M(x_n, f(x_n), t), \\ M(x_{n-1}, f(x_n), t) * M(x_n, f(x_{n-1}), t)\}$$

$$= \min \{M(x_{n-1}, x_n, t), M(x_{n-1}, x_n, t), M(x_n, x_{n+1}, t), M(x_n, x_{n+1}, t), \\ M(x_{n-1}x_{n+1}, t) * M(x_n, x_n, t)\}$$

$$\geq \min \{M(x_{n-1}, x_n, t), M(x_n, x_{n+1}, t), M(x_{n-1}, x_n, t) * M(x_n, x_{n+1}, t)\}$$

$$= \min \{M(x_{n-1}, x_n, t), M(x_n, x_{n+1}, t)\}$$

Since  $\varphi$  is nondecreasing and continuous, we have

$$M(x_n, x_{n+1}, t) \geq \varphi(\min\{M(x_{n-1}, x_n, t), M(x_n, x_{n+1}, t)\})$$

$$\text{If } \min\{M(x_{n-1}, x_n, t), M(x_n, x_{n+1}, t)\} = M(x_n, x_{n+1}, t)$$

Then

$$M(x_n, x_{n+1}, t) \geq \varphi(\min\{M(x_{n-1}, x_n, t), M(x_n, x_{n+1}, t)\}) \geq \varphi(M(x_n, x_{n+1}, t)) \\ > M(x_n, x_{n+1}, t)$$

Which is contradiction.

$$\text{Therefore } \min\{M(x_{n-1}, x_n, t), M(x_n, x_{n+1}, t)\} = M(x_{n-1}, x_n, t)$$

Hence for all  $n \in \mathbb{N}$  we have

$$M(x_n, x_{n+1}, t) \geq \varphi(M(x_{n-1}, x_n, t)) \geq \varphi^2(M(x_{n-2}, x_{n-1}, t)) \\ \geq \dots \geq \varphi^n(M(x_0, x_1, t))$$

Let  $n, m \in \mathbb{N}$  with  $n > m$ , then

$$M(x_n, x_m, t) \geq \varphi(M(x_{n-1}, x_n, t)) \geq \dots \geq \varphi^n(M(x_0, x_1, t))$$

$$\text{Therefore } \lim_{n \rightarrow \infty} M(x_n, x_m, t) = 1$$

Hence  $\{x_n\}$  is a Cauchy sequence

Since  $X$  is an  $\alpha$ - $\eta$ -complete fuzzy metric space there is  $x \in X$  such that  $x_n \rightarrow x$  as  $n \rightarrow \infty$ .

Since  $f$  is an  $\alpha$ - $\eta$ -continuous function so  $x_{n+1} = f(x_n) \rightarrow f(x)$  as  $n \rightarrow \infty$

Hence  $f(x) = x$

Suppose  $y$  is a fixed point of  $f$  such that  $f(y) = y$

$$M(x, y, t) = M(f(x), f(y), t) \geq \varphi(M(x, y))$$

Hence  $x = y$

**Example 2.11:** Let  $X = [0,3]$  be equipped with the ordinary metric

$$d(x, y) = |x - y|, \varphi(\tau) = \sqrt{\tau} \text{ for all } \tau \in [0,1]. \text{ Define } M(x, y, t) = e^{-\frac{2|x-y|}{t}} \text{ for}$$

all  $x, y \in X$  and  $t > 0$ , and  $\alpha, \eta: X \times X \rightarrow [0, \infty)$

$$\text{By } \alpha(x, y) = \begin{cases} (x + y)^2 & \text{if } x, y \in X \\ 0 & \text{if o.w} \end{cases}$$

$$\eta(x, y) = 2xy,$$

Clearly,  $(X, M, *)$  is an  $\alpha$ - $\eta$ - complete fuzzy metric space with respect to  $t$  –norm  $a * b = ab$ . (by [8])

Let  $f: X \rightarrow X$  be defined as

$$f(x) = \begin{cases} 1 & x \in [0,1] \\ \frac{3-x}{2} & x \in (1,3] \end{cases}$$

$$d(f(x), f(y)) = |f(x) - f(y)| = \frac{1}{2}|x - y| \leq |x - y| = d(x, y)$$

$$\text{It follows that } M(f(x), f(y), t) = e^{-\frac{2|f(x)-f(y)|}{t}} \geq e^{-\frac{|x-y|}{t}} = \varphi(M(x, y))$$

Thus  $f$  is  $\alpha$ - $\eta$ - $\varphi$ -contraction function in fuzzy metric space  $(X, M, *)$ .

**Corollary 2.12:** Let  $(X, M, *)$  be a fuzzy metric space and let  $f: X \rightarrow X$ , suppose that  $\alpha, \eta: X \times X \rightarrow [0, \infty)$  are two function. Assume that the following assertions hold:

- (i)  $(X, M, *)$  is an  $\alpha$ -complete fuzzy metric space .
- (ii)  $f$  is an  $\alpha$ -admissible function .
- (iii)  $f$  is  $\alpha$ - $\varphi$ -contraction function on  $X$  .
- (iv)  $f$  is an  $\alpha$ -continuous function .
- (v) There exist  $x_0 \in X$  such that  $\alpha(x_0, f(x_0)) \geq 1$ .

Then  $f$  has a fixed point in  $X$  .

**Theorem 2.13:** Let  $(X, M, *)$  be a fuzzy metric space and let  $f: X \rightarrow X$ , suppose that  $\alpha, \eta: X \times X \rightarrow [0, \infty)$  are two function. Assume that the following assertions hold:

- (i)  $(X, M, *)$  is an  $\alpha$ -complete fuzzy metric space
- (ii)  $f$  is an  $\alpha$ -admissible function with respect to  $\eta$ .
- (iii)  $f$  is  $\alpha$ - $\eta$ - $\varphi$ -contraction function on  $X$ .
- (iv) There exist  $x_0 \in X$  such that  $\alpha(x_0, f(x_0)) \geq \eta(x_0, f(x_0))$ .
- (v) If  $\{x_n\}$  is a sequence in  $X$   $\alpha(x_n, x_{n+1}) \geq \eta(x_n, x_{n+1})$

With  $x_n \rightarrow x$  as  $n \rightarrow \infty$ , then

Either  $\alpha(f(x_n), x) \geq \eta(f(x_n), f^2(x_n))$

Or  $\alpha(f^2(x_n), x) \geq \eta(f^2(x_n), f^3(x_n))$  for all  $n \in \mathbb{N}$

Then  $f$  has a fixed point in  $X$ .

**Proof:** Let  $x_0 \in X$  such that  $\alpha(x_0, f(x_0)) \geq \eta(x_0, f(x_0))$

Define a sequence  $\{x_n\}$  in  $X$  by  $x_n = f^n(x_0) = f(x_{n-1})$  for all  $n \in \mathbb{N}$ .

Now is in the proof of theorem (2.10) we have  $\alpha(x_{n+1}, x_n) \geq \eta(x_{n+1}, x_n)$

There exist  $x \in X$  such that  $x_n \rightarrow x$  as  $n \rightarrow \infty$

Let  $M(x, f(x), t) \neq 1$ , from (v)

Either  $\alpha(f(x_{n-1}), x) \geq \eta(f(x_{n-1}), f^2(x_{n-1}))$

Or  $\alpha(f^2(x_{n-1}), x) \geq \eta(f^2(x_{n-1}), f^3(x_{n-1}))$

Then either  $\alpha(x_n, x) \geq \eta(x_n, x_{n+1})$

Or  $\alpha(x_{n+1}, x) \geq \eta(x_{n+1}, x_{n+2})$

Let  $\alpha(x_n, x) \geq \eta(x_n, x_{n+1})$  from definition (2.9) condition (1), we get

$$M(x_{n+1}, f(x), t) = M(f(x_n), f(x), t) \geq \varphi(\mathbb{M}(x_n, x))$$

Where

$$\mathbb{M}(x_n, x) = \min \{M(x_n, x, t), M(x_n, f(x_n), t), M(x, f(x), t), \\ M(x_n, f(x), t) * M(x, f(x_n), t)\}$$

$$= \varphi(\min \{M(x_n, x, t), M(x_n, x_{n+1}, t), M(x, f(x), t), \\ M(x_n, f(x), t) * M(x, x_{n+1}, t)\})$$

$$> \min\{M(x_n, x, t), M(x_n, x_{n+1}t), M(x, f(x), t), M(x_n, f(x), t) * M(x, x_{n+1}, t)\}$$

By taking  $n \rightarrow \infty$  in the above we get

$$M(x, f(x), t) = \varphi(\min \{1, M(x, f(x), t)\}) > M(x, f(x), t)$$

Which is contradiction.

Hence  $M(x, f(x), t) = 1 \implies f(x) = x$ .

**Corollary 2.14:** Let  $(X, M, *)$  be a fuzzy metric space and let  $f: X \rightarrow X$ , suppose that  $\alpha, \eta: X \times X \rightarrow [0, \infty)$  are two function. Assume that the following assertions hold:

- (i)  $(X, M, *)$  is an  $\alpha$ -complete fuzzy metric space.
- (ii)  $f$  is an  $\alpha$ -admissible function.
- (iii)  $f$  is  $\alpha$ - $\varphi$ -contraction function on  $X$ .
- (iv) There exist  $x_0 \in X$  such that  $\alpha(x_0, f(x_0)) \geq 1$ .
- (v) If  $\{x_n\}$  is a sequence in  $X$   $\alpha(x_n, x_{n+1}) \geq 1$

With  $x_n \rightarrow x$  as  $n \rightarrow \infty$ , then

Either  $\alpha(f(x_n), x) \geq 1$

Or  $\alpha(f^2(x_n), x) \geq 1$  for all  $n \in \mathbb{N}$

Then  $f$  has a fixed point in  $X$ .

**Corollary 2.15:** Let  $(X, M, *)$  be a complete fuzzy metric space and let  $f: X \rightarrow X$  be a continuous function such that  $f$  is contraction function that is

$$M(f(x), f(y), t) \geq \varphi(\mathbb{M}(x, y)) \text{ for all } x, y \in X$$

Then  $f$  has a fixed point in  $X$ .

## 2.2 Fixed Point in Orbitally $f$ - Complete Fuzzy Metric Space

**Theorem 2.2.16:** Let  $(X, M, *)$  be a fuzzy metric space and  $f: X \rightarrow X$  such that the following assertion hold:

- (i)  $(X, M, *)$  is an orbitally  $f$ -complete fuzzy metric space.
- (ii) there exist  $\varphi$  be a function such that  $M(f(x), f(y), t) \geq \varphi(\mathbb{M}(x, y))$  for all  $x, y \in O(w)$  for some  $w \in X$ .
- (iii) If  $\{x_n\}$  be a sequence .such that  $\{x_n\} \subseteq O(w)$  with  $x_n \rightarrow x$  as  $n \rightarrow \infty$ ,  $x \in O(w)$ .

Then  $f$  has fixed point in  $X$ .



**Proof:** Define  $\alpha: X \times X \rightarrow [0, \infty)$  from remark (2.8),  $(X, M, *)$  is an  $\alpha$ -complete fuzzy metric space and  $f$  is an  $\alpha$ -admissible function.

Let  $\alpha(x, y) \geq 1$  for all  $x, y \in O(w)$

Then from (ii)

$$M(f(x), f(y), t) \geq \varphi(M(x, y))$$

That is  $f$  is an  $\alpha$ - $\varphi$ -contraction function

Let  $\{x_n\}$  sequence such that  $\alpha(x_n, x_{n+1}) \geq 1$  with  $x_n \rightarrow x$

So  $\{x_n\} \subseteq O(w)$  from (iii) we have  $x \in O(w)$

That is  $\alpha(x_n, x) \geq 1$  by corollary (2.13)

Then  $f$  has unique fixed point in  $X$ .

**Corollary 2.2.17:** Let  $(X, M, *)$  be a fuzzy metric space and  $f: X \rightarrow X$  such that the following assertion hold:

- (i)  $(X, M, *)$  is an orbitally  $f$ -complete fuzzy metric space.
- (ii) there exist  $k \in (0, 1)$  such that  $M(f(x), f(y), t) \geq kM(x, y)$  for all  $x, y \in O(w)$  for some  $w \in X$ .
- (iii) If  $\{x_n\}$  be a sequence such that  $\{x_n\} \subseteq O(w)$  with  $x_n \rightarrow x$  as  $n \rightarrow \infty$ . Then  $x \in O(w)$ . Then  $f$  has fixed point in  $X$ .

### 2.3 Suzuki Type Fixed Point Result

From Theorem (2.10) we deduce the following Suzuki type fixed point result

**Theorem 2.3.18:** Let  $(X, M, *)$  be a complete fuzzy metric space and let  $f: X \rightarrow X$  continuous function. Assume that there exist  $k \in (0, 1)$  such that

$$M(x, f(x), t) \geq M(x, y, t) \implies M(f(x), f(y), t) \geq kM(x, y) \quad \text{---(1)}$$

For all  $x, y \in X$ , where

$$M(x, y) = \min \{M(x, y, t), M(x, f(x), t), M(y, f(y), t), M(x, f(y), t) * M(y, f(x), t)\}$$

Then  $f$  has a fixed point in  $X$ .

**Proof:** Define  $\alpha, \eta: X \times X \rightarrow [0, \infty)$  and  $\varphi: [0, 1] \rightarrow [0, 1]$  by

$$\alpha(x, y) = M(x, f(x), t) \quad , \quad \eta(x, y) = M(x, y, t)$$

Such that  $\alpha(x, y) \geq \eta(x, y)$  for all  $x, y \in X$

And let  $\varphi(\tau) = k\tau$  ,  $\tau \in [0,1]$

By condition (i) –(v) of theorem (2.10 )

Let  $\alpha(x, f(x)) \geq \eta(x, y)$

Then  $M(x, f(x), t) \geq M(x, y, t)$

From (1) we have  $M(f(x), f(y), t) \geq kM(x, y) = \varphi(M(x, y))$

Then  $f$  is  $\alpha$ - $\eta$ - $\varphi$ -contraction function by condition of theorem (2.10) and  $f$  has fixed point, then the unique fixed point follows from (1).

**Corollary 2.3.19:** *Let  $(X, M, *)$  be a complete fuzzy metric space and let  $f: X \rightarrow X$  continuous function. Assume that there exist  $k \in (0,1)$  such that*

$$M(x, f(x), t) \geq M(x, y, t) \implies M(f(x), f(y), t) \geq kM(x, y, t)$$

For all  $x, y \in X$ ,

Then  $f$  has a fixed point in  $X$  .

## 2.4 An $\alpha$ - $\psi$ - $\phi$ -Contraction Function

**Definition 2.4.20:** *Let  $(X, M, *)$  be a complete fuzzy metric space. Let  $f: X \rightarrow X$  be an  $\alpha$ -admissible which satisfies the following*

$$\psi(M(f(x), f(y), t)) \geq \psi(M(x, y)) - \phi(M(x, y))$$

Such that

- (i)  $\psi$  is continuous and decreasing with  $\psi(\tau) = 0$  iff  $\tau = 1$ .
- (ii)  $\phi$  is continuous with  $\phi(\tau) = 0$  iff  $\tau = 1$   
 $f$  is called  $\alpha$ -  $\psi$ -  $\phi$ - contraction function .

**Theorem 2.4.21:** *Let  $(X, M, *)$  be a complete fuzzy metric space . Let  $f: X \rightarrow X$  be an  $\alpha$ -admissible which satisfies the following*

$$\psi(M(f(x), f(y), t)) \geq \psi(M(x, y)) - \phi(M(x, y))$$

Such that

- (i)  $(X, M, *)$  is an  $\alpha$ -complete.
- (ii)  $f$  is  $\alpha$ -continuous function
- (iv) There exist  $x_0 \in X$  such that  $\alpha(x_0, f(x_0)) \geq 1$ .

**Proof:** Let  $x_0 \in X$  such that  $\alpha(x_0, f(x_0)) \geq 1$

Define a sequence  $\{x_n\}$  such that  $x_n = f(x_{n-1})$  for all  $n \in \mathbb{N}$

If  $x_n = x_{n+1}$  for some  $n$ , then  $x = x_n$  is a fixed point of  $f$ .

Suppose  $x_n \neq x_{n+1}$

Since  $f$  is  $\alpha$ -admissible function with respect  $\eta$  and

$$\alpha(x_0, f(x_0)) = \alpha(x_0, x_1) \geq 1$$

$$\text{Then } \alpha(x_1, x_2) = \alpha(f(x_0), f(x_1)) \geq \eta(f(x_0), f(x_1)) = \eta(x_1, x_2)$$

By continuing this process, we get

$$\alpha(x_n, f(x_n)) = \alpha(x_n, x_{n+1}) \geq 1 \text{ for all } n \in \mathbb{N}$$

$$\begin{aligned} \psi(M(x_n, x_{n+1}, t)) &= \psi(M(f(x_{n-1}), f(x_n), t)) \\ &\geq \psi(\mathbb{M}(x_{n-1}, x_n)) - \phi(\mathbb{M}(x_{n-1}, x_n)) \end{aligned}$$

Where

$$\begin{aligned} \mathbb{M}(x_{n-1}, x_n) &= \min \{M(x_{n-1}, x_n, t), M(x_{n-1}, f(x_{n-1}), t), M(x_n, f(x_n), t), \\ &\quad M(x_{n-1}, f(x_n), t) * M(x_n, f(x_{n-1}), t)\} \\ &= \min \{M(x_{n-1}, x_n, t), M(x_{n-1}, x_n, t), M(x_n, x_{n+1}, t), M(x_n, x_{n+1}, t), \\ &\quad M(x_{n-1}x_{n+1}, t) * M(x_n, x_n, t)\} \end{aligned}$$

$$\geq \min \{M(x_{n-1}, x_n, t), M(x_n, x_{n+1}, t), M(x_{n-1}, x_n, t) * M(x_n, x_{n+1}, t)\}$$

$$= \min \{M(x_{n-1}, x_n, t), M(x_n, x_{n+1}, t)\}$$

Since  $\psi$  decreasing, then  $M(x_{n-1}, x_n, t) < M(x_n, x_{n+1}, t)$  that is  $\{M(x_n, x_{n+1}, t)\}$  is an increasing sequence thus there exist  $l(t) \in (0, 1]$  such that  $\lim_{n \rightarrow \infty} M(x_n, x_{n+1}, t) = l(t) < 1$

Now taking  $n \rightarrow \infty$  we obtain for all  $t > 0$

$$\psi(l(t)) \geq \psi(l(t)) - \phi(l(t)) \text{ which is contradiction } l(t) = 1$$

Thus we conclude for all  $t > 0$

$$\lim_{n \rightarrow \infty} M(x_n, x_{n+1}, t) = 1$$

$\{x_n\}$  is Cauchy sequence .since  $X$  is  $\alpha$ -complete ,then  $x_n \rightarrow x$

$$\begin{aligned} \psi(M(x_{n+1}, f(x), t)) &= \psi(M(f(x_n), f(x), t)) \\ &\geq \psi(\mathbb{M}(x_n, x)) - \phi(\mathbb{M}(x_n, x)) \end{aligned}$$

Letting  $n \rightarrow \infty$  in the above inequality using properties  $\psi, \phi$

$$\psi(\lim_{n \rightarrow \infty} M(x_n, f(x), t)) \geq \psi(1) - \phi(1) = 0$$

Thus  $\lim_{n \rightarrow \infty} M(x_n, f(x), t) = 1$

Hence  $x_n \rightarrow f(x) \Rightarrow f(x) = x$  .

Now we assume  $y$  is a fixed point of  $f$  such that  $f(y) = y$

$$\begin{aligned} \psi(M(x, y, t)) &= \psi(M(f(x), f(y), t)) \geq \psi(\mathbb{M}(x, y)) - \phi(\mathbb{M}(x, y)) = 0 \\ \Rightarrow M(x, y, t) &= 1 \Rightarrow x = y . \end{aligned}$$

## 2.5 Application On $\alpha$ - $\eta$ - $\phi$ -Contraction

**Definition 2.5.22:** Let  $(X, M, *)$  be a fuzzy metric space and  $f: X \rightarrow X$  be an  $(\alpha, \beta)$ -admissible function,  $f$  is said to be

- (a)  $(\alpha, \beta)$ -contraction function of type ( I ) , if  $\alpha(x, y)\beta(x, y)M(f(x), f(y), t) \geq \phi(\mathbb{M}(x, y))$ .
- (b)  $(\alpha, \beta)$ -contraction function of type (II) , if there exist  $0 < \ell \leq 1$  such that  $(\alpha(x, y)\beta(x, y) + \ell)^{M(f(x), f(y), t)} \geq (1 + \ell)^{\phi(\mathbb{M}(x, y))}$

**Theorem 2.5.23:** Let  $(X, M, *)$  be a complete fuzzy metric space and let  $f: X \rightarrow X$  be an  $\alpha$ -continuous and  $(\alpha, \beta)$ -contraction function of type (I), (II), if there exist  $\alpha(x_0, f(x_0)) \geq 1$  and  $\beta(x_0, f(x_0)) \geq 1$  , then  $f$  has a unique fixed point in  $X$ .

**Proof:** Let  $x_0 \in X$  such that  $\alpha(x_0, f(x_0)) \geq 1$  and  $\beta(x_0, f(x_0)) \geq 1$

Define a sequence  $\{x_n\}$  such that  $x_n = f(x_{n-1})$  for all  $n \in \mathbb{N}$

Since  $f$  is  $(\alpha, \beta)$ -admissible function and  $\alpha(x_0, f(x_0)) \geq 1$

$$\text{Then } \alpha(x_1, x_2) = \alpha(f(x_0), f(x_1)) \geq 1$$

By continuing this process, we get

$$\alpha(x_n, x_{n+1}) = \alpha(x_n, f(x_n)) \geq 1$$

Similarly we have  $\beta(x_n, x_{n+1}) = \beta(x_n, f(x_n)) \geq 1$

If  $x_n = x_{n+1}$  for some  $n$ , then  $x = x_n$  is a fixed point of  $f$ .

Suppose  $x_n \neq x_{n+1}$

(a)

$$\begin{aligned} \alpha(x_n, x_{n+1})\beta(x_n, x_{n+1})M(x_n, x_{n+1}, t) \\ = \alpha(x_n, x_{n+1})\beta(x_n, x_{n+1})M(f(x_{n-1}), f(x_n), t) \\ \geq \varphi(\mathbb{M}(x_{n-1}, x_n)) \end{aligned}$$

Where

$$\mathbb{M}(x_{n-1}, x_n) = \min \{M(x_{n-1}, x_n, t), M(x_{n-1}, f(x_{n-1}), t), M(x_n, f(x_n), t), M(x_{n-1}, f(x_n), t) * M(x_n, f(x_{n-1}), t)\}$$

$$= \min \{M(x_{n-1}, x_n, t), M(x_{n-1}, x_n, t), M(x_n, x_{n+1}, t), M(x_n, x_{n+1}, t), M(x_{n-1}x_{n+1}, t) * M(x_n, x_n, t)\}$$

$$\geq \min \{M(x_{n-1}, x_n, t), M(x_n, x_{n+1}, t), M(x_{n-1}, x_n, t) * M(x_n, x_{n+1}, t)\}$$

$$= \min \{M(x_{n-1}, x_n, t), M(x_n, x_{n+1}, t)\}$$

$$\text{If } \min\{M(x_{n-1}, x_n, t), M(x_n, x_{n+1}, t)\} = M(x_n, x_{n+1}, t)$$

Then

$$\begin{aligned} M(x_n, x_{n+1}, t) &\geq \varphi(\min\{M(x_{n-1}, x_n, t), M(x_n, x_{n+1}, t)\}) \geq \varphi(M(x_n, x_{n+1}, t)) \\ &> M(x_n, x_{n+1}, t) \end{aligned}$$

Which is contradiction

$$\text{Therefore } \min\{M(x_{n-1}, x_n, t), M(x_n, x_{n+1}, t)\} = M(x_{n-1}, x_n, t)$$

Hence for all  $n \in \mathbb{N}$  we have

$$\begin{aligned} M(x_n, x_{n+1}, t) &\geq \varphi(M(x_{n-1}, x_n, t)) \geq \varphi^2(M(x_{n-2}, x_{n-1}, t)) \\ &\geq \dots \geq \varphi^n(M(x_0, x_1, t)) \end{aligned}$$

Let  $n, m \in \mathbb{N}$  with  $n > m$ , then

$$\begin{aligned} M(x_n, x_m, t) &= \alpha(x_n, x_{n+1})\beta(x_n, x_{n+1})M(f(x_{n-1}), f(x_n), t) \geq \varphi(\mathbb{M}(x_{n-1}, x_n)) \\ &\geq \dots \geq \varphi^n(M(x_0, x_1, t)) \end{aligned}$$

Therefore  $\lim_{n \rightarrow \infty} M(x_n, x_m, t) = 1$

Hence  $\{x_n\}$  is a Cauchy sequence

Since  $X$  is an  $\alpha$ - $\eta$ -complete fuzzy metric space there is  $x \in X$  such that  $x_n \rightarrow x$  as  $n \rightarrow \infty$ .

$$M(x_n, f(x), t) = \alpha(x_n, x)\beta(x_n, x)M(f(x_n), f(x), t) \geq \varphi(\mathbb{M}(x_n, x))$$

Hence  $f(x) = x$

Suppose  $y$  is a fixed point of  $f$  such that  $f(y) = y$

$$M(x, y, t) = \alpha(x, y)\beta(x, y)M(f(x), f(y), t) \geq \varphi(\mathbb{M}(x, y)).$$

Hence  $x = y$ .

(b)

$$\begin{aligned} & (\alpha(x_n, x_{n+1})\beta(x_n, x_{n+1}) + \ell)^{M(x_n, x_{n+1}, t)} \\ &= (\alpha(x_n, x_{n+1})\beta(x_n, x_{n+1}) + \ell)^{M(f(x_{n-1}), f(x_n), t)} \\ &\geq (1 + \ell)^{\varphi(\mathbb{M}(x_{n-1}, x_n))} \end{aligned}$$

Where

$$\mathbb{M}(x_{n-1}, x_n) = \min \{M(x_{n-1}, x_n, t), M(x_{n-1}, f(x_{n-1}), t), M(x_n, f(x_n), t), M(x_{n-1}, f(x_n), t) * M(x_n, f(x_{n-1}), t)\}$$

$$= \min \{M(x_{n-1}, x_n, t), M(x_{n-1}, x_n, t), M(x_n, x_{n+1}, t), M(x_n, x_{n+1}, t), M(x_{n-1}x_{n+1}, t) * M(x_n, x_n, t)\}$$

$$\geq \min \{M(x_{n-1}, x_n, t), M(x_n, x_{n+1}, t), M(x_{n-1}, x_n, t) * M(x_n, x_{n+1}, t)\}$$

$$= \min \{M(x_{n-1}, x_n, t), M(x_n, x_{n+1}, t)\}$$

$$\text{If } \min\{M(x_{n-1}, x_n, t), M(x_n, x_{n+1}, t)\} = M(x_n, x_{n+1}, t)$$

Then

$$M(x_n, x_{n+1}, t) \geq \varphi(\min\{M(x_{n-1}, x_n, t), M(x_n, x_{n+1}, t)\}) \geq \varphi(M(x_n, x_{n+1}, t)) > M(x_n, x_{n+1}, t)$$

Which is contradiction

$$\text{Therefore } \min\{M(x_{n-1}, x_n, t), M(x_n, x_{n+1}, t)\} = M(x_{n-1}, x_n, t)$$

Hence for all  $n \in \mathbb{N}$  we have

$$\begin{aligned} M(x_n, x_{n+1}, t) &\geq \varphi(M(x_{n-1}, x_n, t)) \geq \varphi^2(M(x_{n-2}, x_{n-1}, t)) \\ &\geq \dots \geq \varphi^n(M(x_0, x_1, t)) \end{aligned}$$

Let  $n, m \in \mathbb{N}$  with  $n > m$ , then

$$\begin{aligned} M(x_n, x_m, t) &= (\alpha(x_n, x_{n+1})\beta(x_n, x_{n+1}) + 1)^{M(f(x_{n-1}), f(x_n), t)} \\ &\geq (1 + \ell)^{\varphi(\mathbb{M}(x_{n-1}, x_n))} \geq \dots \geq (1 + \ell)^{\varphi^n(M(x_0, x_1, t))} \end{aligned}$$

Therefore  $\lim_{n \rightarrow \infty} M(x_n, x_m, t) = 1$

Hence  $\{x_n\}$  is a Cauchy sequence

Since  $X$  is an  $\alpha$ - $\eta$ -complete fuzzy metric space there is  $x \in X$  such that  $x_n \rightarrow x$  as  $n \rightarrow \infty$ .

$$\begin{aligned} (\alpha(x_n, x)\beta(x_n, x) + \ell)^{M(x_n, f(x), t)} &= (\alpha(x_n, x)\beta(x_n, x) + \ell)^{M(f(x_n), f(x), t)} \\ &\geq (1 + \ell)^{\varphi(\mathbb{M}(x_n, x))} \end{aligned}$$

Hence  $f(x) = x$

Suppose  $y$  is a fixed point of  $f$  such that  $f(y) = y$

$$(\alpha(x, y)\beta(x, y) + \ell)^{M(f(x), f(y), t)} \geq (1 + \ell)^{\varphi(\mathbb{M}(x, y))}.$$

Hence  $x = y$ .

## References

- [1] A. George and P. Veramani, On some results in fuzzy metric spaces, *Fuzzy Sets and Systems*, 64(1994), 395-399.
- [2] M. Grabiec, Fixed points in fuzzy metric space, *Fuzzy Sets and Systems*, 27(1988), 385-389.
- [3] S. Heipern, Fuzzy mappings and fixed point theorems, *J. Math. Anal. Appl.*, 83(1981), 566-569.
- [4] N. Hussain, M. Kutbi and P. Salimi, *Fixed Point Theory in  $\alpha$ -Complete Metric Space with Application*, Hindawi Publishing Corporation, (2014), 11.
- [5] I. Kramosil and J. Michalek, Fuzzy metric and statistical metric spaces, *Kybernetika*, 11(1975), 326-334.
- [6] P. Salimia, C. Vetro and P. Vetro, Fixed point theorems for twisted  $(\alpha, \beta)$ - $\psi$ -contractive type mappings and applications, *Faculty of Sciences and Mathematics*, University of Niš, Serbia, 24(4) (2013), 605-615.
- [7] P. Salimi, C. Vetro and P. Vetro, Some new fixed point results in non-Archimedean fuzzy metric space, *Modeling and Control*, 18(3) (2013), 344-358.
- [8] Y. Shen, D. Qiu and W. Chen, *On Convergence of Fixed Points in Fuzzy Metric Spaces*, Hindawi Publishing Corporation, (2013), 6.
- [9] B.C. Tripathy, S. Paul and N.R. Das, A fixed point theorem in a generalized fuzzy metric space, *Bol. Soc. Paran. Mat.*, 32(2) (2014), 221-227.
- [10] L.A. Zadeh, Fuzzy sets, *Inform. Control*, 8(1965), 338-353.