



Gen. Math. Notes, Vol. 17, No. 2, August, 2013, pp. 53-65
ISSN 2219-7184; Copyright © ICSRS Publication, 2013
www.i-csrs.org
Available free online at <http://www.geman.in>

Homomorphism and Anti Homomorphism on Bipolar Fuzzy Sub HX Groups

R. Muthuraj¹ and M. Sridharan²

¹ PG & Research Department of Mathematics
H.H. The Rajah's College
Pudukkottai – 622 001, Tamilnadu, India
E-mail: rmr1973@yahoo.co.in

² Department of Mathematics
PSNA College of Engineering and Technology
Dindigul – 624 622, Tamilnadu, India
E-mail: msridharan1972@gmail.com

(Received: 14-4-13 / Accepted: 29-5-13)

Abstract

In this paper, we introduce the concept of an image, pre image of a bipolar fuzzy subsets and discuss in detail a series of homomorphic and anti homomorphic properties of bipolar fuzzy sub groups.

Keywords: *Fuzzy set, HX group, Fuzzy HX group, Bipolar-valued fuzzy set, Bipolar fuzzy HX group, Homomorphism and Anti homomorphism of fuzzy HX group, Image and pre-image of bipolar fuzzy sets are discussed.*

1 Introduction

The concept of fuzzy sets was initiated by Zadeh [16]. Then it has become a vigorous area of research in engineering, medical science, social science, graph

theory etc. Rosenfeld [12] gave the idea of fuzzy subgroup. In fuzzy sets the membership degree of elements range over the interval $[0, 1]$. The membership degree expresses the degree of belongingness of elements to a fuzzy set. The membership degree 1 indicates that an element completely belongs to its corresponding fuzzy set and membership degree 0 indicates that an element does not belong to fuzzy set. The membership degrees on the interval $(0,1)$ indicate the partial membership to the fuzzy set. Sometimes, the membership degree means the satisfaction degree of elements to some property or constraint corresponding to a fuzzy set. Li Hongxing [3] introduced the concept of HX group and the authors Luo Chengzhong, Mi Honghai, Li Hongxing [4] introduced the concept of fuzzy HX group. The author W.R. Zhang [14], [15] commenced the concept of bipolar fuzzy sets as a generalization of fuzzy sets in 1994. In case of Bipolar-valued fuzzy sets membership degree range is enlarged from the interval $[0, 1]$ to $[-1, 1]$. In a bipolar-valued fuzzy set, the membership degree 0 means that the elements are irrelevant to the corresponding property, the membership degree $(0, 1]$ indicates that elements somewhat satisfy the property and the membership degree $[-1, 0)$ indicates that elements somewhat satisfy the implicit counter-property. M. Marudai, V. Rajendran [5] introduced the pre image of bipolar Q fuzzy subgroup. In this paper we define the image and pre image of a bipolar fuzzy subgroup and bipolar fuzzy sub HX group and discuss some of its properties.

2 Preliminaries

In this section, we site the fundamental definitions that will be used in the sequel. Throughout this paper, $G = (G, *)$ be a finite group, e is the identity element of G , and xy we mean $x * y$.

Definition 2.1[9]: Let X be any non-empty set. A fuzzy subset μ of X is a function $\mu : X \rightarrow [0,1]$.

Definition 2.2[3]: In $2^G - \{\emptyset\}$, a non-empty set $\vartheta \subset 2^G - \{\emptyset\}$ is called a HX group on G , if ϑ is a group with respect to the algebraic operation defined by $AB = \{ab / a \in A \text{ and } b \in B\}$, which its unit element is denoted by E .

Definition 2.3[10]: Let μ be a fuzzy subset defined on G . Let $\vartheta \subset 2^G - \{\emptyset\}$ be a HX group on G . A fuzzy set λ_μ defined on ϑ is said to be a fuzzy subgroup induced by μ on ϑ or a fuzzy HX subgroup on ϑ if for any $A, B \in \vartheta$,

- i. $\lambda_\mu(AB) \geq \min \{ \lambda_\mu(A), \lambda_\mu(B) \}$
- ii. $\lambda_\mu(A^{-1}) = \lambda_\mu(A)$.

where, $\lambda_\mu(A) = \max \{ \mu(x) / \text{for all } x \in A \subset G \}$.

Definition 2.4[10]: Let G be a non-empty set. A bipolar-valued fuzzy set or bipolar fuzzy set μ in G is an object having the form $\mu = \{\langle x, \mu^+(x), \mu^-(x) \rangle / \text{for all } x \in G\}$, where $\mu^+ : G \rightarrow [0,1]$ and $\mu^- : G \rightarrow [-1,0]$ are mappings. The positive membership degree $\mu^+(x)$ denotes the satisfaction degree of an element x to the property corresponding to a bipolar-valued fuzzy set $\mu = \{\langle x, \mu^+(x), \mu^-(x) \rangle / \text{for all } x \in G\}$ and the negative membership degree $\mu^-(x)$ denotes the satisfaction degree of an element x to some implicit counter property corresponding to a bipolar-valued fuzzy set $\mu = \{\langle x, \mu^+(x), \mu^-(x) \rangle / \text{for all } x \in G\}$. If $\mu^+(x) \neq 0$ and $\mu^-(x) = 0$, it is the situation that x is regarded as having only positive satisfaction for $\mu = \{\langle x, \mu^+(x), \mu^-(x) \rangle / \text{for all } x \in G\}$. If $\mu^+(x) = 0$ and $\mu^-(x) \neq 0$, it is the situation that x does not satisfy the property of $\mu = \{\langle x, \mu^+(x), \mu^-(x) \rangle / \text{for all } x \in G\}$, but somewhat satisfies the counter property of $\mu = \{\langle x, \mu^+(x), \mu^-(x) \rangle / \text{for all } x \in G\}$. It is possible for an element x to be such that $\mu^+(x) \neq 0$ and $\mu^-(x) \neq 0$ when the membership function of property overlaps that its counter property over some portion of G . For the sake of simplicity, we shall use the symbol $\mu = (\mu^+, \mu^-)$ for the bipolar-valued fuzzy set $\mu = \{\langle x, \mu^+(x), \mu^-(x) \rangle / \text{for all } x \in G\}$.

Definition 2.5[10]: A bipolar-valued fuzzy set or bipolar fuzzy set μ in G is a bipolar fuzzy subgroup of G if for all $x, y \in G$,

- i. $\mu^+(xy) \geq \min \{\mu^+(x), \mu^+(y)\}$,
- ii. $\mu^-(xy) \leq \max \{\mu^-(x), \mu^-(y)\}$,
- iii. $\mu^+(x^{-1}) = \mu^+(x)$, $\mu^-(x^{-1}) = \mu^-(x)$.

Definition 2.6[10]: Let ϑ be a non-empty set. A bipolar-valued fuzzy set or bipolar fuzzy set [BFS] λ_μ in ϑ is an object having the form $\lambda_\mu = \{\langle A, \lambda_\mu^+(A), \lambda_\mu^-(A) \rangle / \text{for all } x \in A \subset G\}$, where $\lambda_\mu^+ : \vartheta \rightarrow [0,1]$ and $\lambda_\mu^- : \vartheta \rightarrow [-1,0]$ are mappings. The positive membership degree $\lambda_\mu^+(A)$ denotes the satisfaction degree of an element A to the property corresponding to a bipolar-valued fuzzy set $\lambda_\mu = \{\langle A, \lambda_\mu^+(A), \lambda_\mu^-(A) \rangle / \text{for all } x \in A \subset G\}$ and the negative membership degree $\lambda_\mu^-(A)$ denotes the satisfaction degree of an element A to some implicit counter property corresponding to a bipolar-valued fuzzy set $\lambda_\mu = \{\langle A, \lambda_\mu^+(A), \lambda_\mu^-(A) \rangle / \text{for all } x \in A \subset G\}$. If $\lambda_\mu^+(A) \neq 0$ and $\lambda_\mu^-(A) = 0$, it is the situation that A is regarded as having only positive satisfaction for $\lambda_\mu = \{\langle A, \lambda_\mu^+(A), \lambda_\mu^-(A) \rangle / \text{for all } x \in A \subset G\}$. If $\lambda_\mu^+(A) = 0$ and $\lambda_\mu^-(A) \neq 0$, it is the situation that A does not satisfy the property of $\lambda_\mu = \{\langle A, \lambda_\mu^+(A), \lambda_\mu^-(A) \rangle / \text{for all } x \in A \subset G\}$, but somewhat satisfies the counter property of $\lambda_\mu = \{\langle A, \lambda_\mu^+(A), \lambda_\mu^-(A) \rangle / \text{for all } x \in A \subset G\}$. It is possible for an element A to be such that $\lambda_\mu^+(A) \neq 0$ and $\lambda_\mu^-(A) \neq 0$ when the membership function of property overlaps that its counter property over some portion of ϑ . For the sake of simplicity, we shall use the symbol $\lambda_\mu = (\lambda_\mu^+, \lambda_\mu^-)$. For the bipolar-valued fuzzy set $\lambda_\mu = \{\langle A, \lambda_\mu^+(A), \lambda_\mu^-(A) \rangle / \text{for all } x \in A \subset G\}$.

Definition 2.7[10]: Let μ be a bipolar fuzzy subset defined on G . Let $\vartheta \subset 2^G - \{\emptyset\}$ be a HX group on G . A bipolar fuzzy set λ_μ defined on ϑ is said to be a bipolar

fuzzy subgroup induced by μ on \mathcal{V} or a bipolar fuzzy HX subgroup on \mathcal{V} , if for any $A, B \in \mathcal{V}$,

- i. $\lambda_{\mu}^{+}(AB) \geq \min\{\lambda_{\mu}^{+}(A), \lambda_{\mu}^{+}(B)\}$,
- ii. $\lambda_{\mu}^{-}(AB) \leq \max\{\lambda_{\mu}^{-}(A), \lambda_{\mu}^{-}(B)\}$,
- iii. $\lambda_{\mu}^{+}(A^{-1}) = \lambda_{\mu}^{+}(A)$, $\lambda_{\mu}^{-}(A^{-1}) = \lambda_{\mu}^{-}(A)$.

Where, $\lambda_{\mu}^{+}(A) = \max\{\mu^{+}(x) / \text{for all } x \in A \subset G\}$ and
 $\lambda_{\mu}^{-}(A) = \min\{\mu^{-}(x) / \text{for all } x \in A \subset G\}$.

Definition 2.8[13]: A mapping f from a group G_1 to a group G_2 is said to be a homomorphism if $f(xy) = f(x)f(y)$ for all $x, y \in G_1$.

Definition 2.9[13]: A mapping f from a group G_1 to a group G_2 (G_1 and G_2 are not necessarily commutative) is said to be an anti homomorphism if $f(xy) = f(y)f(x)$ for all $x, y \in G_1$.

Definition 2.10[9]: A mapping f from a HX group \mathcal{V}_1 to a HX group \mathcal{V}_2 is said to be a homomorphism if $f(AB) = f(A)f(B)$ for all $A, B \in \mathcal{V}_1$.

Definition 2.11[9]: A mapping f from a HX group \mathcal{V}_1 to a HX group \mathcal{V}_2 (\mathcal{V}_1 and \mathcal{V}_2 are not necessarily commutative) is said to be an anti homomorphism if $f(AB) = f(B)f(A)$ for all $A, B \in \mathcal{V}_1$.

3 Image and Pre-Image of a Bipolar Fuzzy Sub Group of a Group under Homomorphism and Anti Homomorphism

In this section, we introduce the notion of image and pre-image of the bipolar fuzzy subgroup of a group, and discuss some of its properties. Throughout this section, We mean that G_1 and G_2 are finite groups and e_1 and e_2 are the identity elements of G_1 and G_2 respectively, and xy we mean $x * y$.

Definition 3.1: Let f be a mapping from a group G_1 to a group G_2 and let μ, φ be fuzzy subsets in G_1 and G_2 respectively. Then the image $f(\mu)$ of μ is the fuzzy subset of G_2 defined by for $u \in G_2$

$$(f(\mu))(u) = \begin{cases} \max\{\mu(x) : x \in f^{-1}(u)\}, & \text{if } f^{-1}(u) \neq \emptyset \\ 0 & \text{, Otherwise} \end{cases}$$

and the pre-image $f^{-1}(\varphi)$ of φ under f is the fuzzy subset of G_1 defined by for $x \in G_1$, $(f^{-1}(\varphi))(x) = \varphi(f(x))$.

Definition 3.2: Let f be a mapping a group G_1 to a group G_2 and let $\mu = (\mu^+, \mu^-)$ and $\varphi = (\varphi^+, \varphi^-)$ are bipolar fuzzy subsets in G_1 and G_2 respectively. Then the image $f(\mu)$ of μ is the bipolar fuzzy subset $f(\mu) = ((f(\mu))^+, (f(\mu))^-)$ of G_2 defined by for $u \in G_2$,

$$(f(\mu))^+(u) = \begin{cases} \max \{ \mu^+(x) : x \in f^{-1}(u) \}, & \text{if } f^{-1}(u) \neq \emptyset \\ 0 & , \text{Otherwise} \end{cases}$$

and

$$(f(\mu))^{-}(u) = \begin{cases} \max \{ \mu^{-}(x) : x \in f^{-1}(u) \}, & \text{if } f^{-1}(u) \neq \emptyset \\ 0 & , \text{Otherwise} \end{cases}$$

and the pre-image $f^{-1}(\varphi)$ of φ under f is the bipolar fuzzy subset of G_1 defined by for $x \in G_1$, $(f^{-1}(\varphi))^+(x) = \varphi^+(f(x))$, $(f^{-1}(\varphi))^{-}(x) = \varphi^{-}(f(x))$.

Theorem 3.1: Let f be a homomorphism from a group G_1 into a group G_2 . If $\mu = (\mu^+, \mu^-)$ be a bipolar fuzzy subgroup of G_1 then $f(\mu)$, the image of μ under f , is a bipolar fuzzy subgroup of G_2 .

Proof: Let $\mu = (\mu^+, \mu^-)$ be a bipolar fuzzy subgroup of G_1 , $\mu^+ : G_1 \rightarrow [0,1]$ and $\mu^- : G_1 \rightarrow [-1,0]$ are mappings.

Let $u, v \in G_2$, since f is homomorphism and so there exist $x, y \in G_1$ such that $f(x) = u$ and $f(y) = v$ it follows that $xy \in f^{-1}(uv)$.

$$\begin{aligned} \text{Now, } (f(\mu))^+(uv) &= \max \{ \mu^+(z) : z = xy \in f^{-1}(uv) \} \\ &\geq \max \{ \mu^+(xy) : x \in f^{-1}(u), y \in f^{-1}(v) \} \\ &\geq \max \{ \min \{ \mu^+(x), \mu^+(y) \} : x \in f^{-1}(u), y \in f^{-1}(v) \} \\ &= \min \{ \max \{ \mu^+(x) : x \in f^{-1}(u) \}, \max \{ \mu^+(y) : y \in f^{-1}(v) \} \} \\ &= \min \{ (f(\mu))^+(u), (f(\mu))^+(v) \} \end{aligned}$$

Therefore, $(f(\mu))^+(uv) \geq \min \{ (f(\mu))^+(u), (f(\mu))^+(v) \}$

$$\begin{aligned} \text{And } (f(\mu))^{-}(uv) &= \max \{ \mu^{-}(z) : z = xy \in f^{-1}(uv) \} \\ &\leq \max \{ \mu^{-}(xy) : x \in f^{-1}(u), y \in f^{-1}(v) \} \\ &\leq \max \{ \max \{ \mu^{-}(x), \mu^{-}(y) \} : x \in f^{-1}(u), y \in f^{-1}(v) \} \\ &= \max \{ \max \{ \mu^{-}(x) : x \in f^{-1}(u) \}, \max \{ \mu^{-}(y) : y \in f^{-1}(v) \} \} \\ &= \max \{ (f(\mu))^{-}(u), (f(\mu))^{-}(v) \} \end{aligned}$$

Therefore, $(f(\mu))^{-}(uv) \leq \max \{ (f(\mu))^{-}(u), (f(\mu))^{-}(v) \}$

$$\begin{aligned} \text{Now, } (f(\mu))^+(u^{-1}) &= \max \{ \mu^+(x) : x \in f^{-1}(u^{-1}) \} \\ &= \max \{ \mu^+(x^{-1}) : x^{-1} \in f^{-1}(u) \} \\ &= (f(\mu))^+(u) \end{aligned}$$

$$\text{And } (f(\mu))^{-}(u^{-1}) = \max \{ \mu^{-}(x) : x \in f^{-1}(u^{-1}) \}$$

$$\begin{aligned}
&= \max \{ \mu^- (x^{-1}) : x^{-1} \in f^{-1}(u) \} \\
&= (f(\mu))^- (u)
\end{aligned}$$

Therefore $f(\mu)$ is a bipolar fuzzy subgroup of G_2 .

Hence, if μ be a bipolar fuzzy subgroup of G_1 then $f(\mu)$ is a bipolar fuzzy subgroup of G_2 .

Theorem 3.2: *The homomorphic pre-image of a bipolar fuzzy subgroup $\varphi = (\varphi^+, \varphi^-)$ of a group of G_2 is a bipolar fuzzy subgroup of a group G_1 .*

Proof: Let $\varphi = (\varphi^+, \varphi^-)$ be a bipolar fuzzy subgroup of G_2 , $\varphi^+ : G_2 \rightarrow [0,1]$ and $\varphi^- : G_2 \rightarrow [-1,0]$ are mappings.

$$\begin{aligned}
\text{Now, } (f^{-1}(\varphi))^+ (xy) &= \varphi^+ (f(xy)) \\
&= \varphi^+ (f(x)f(y)) \\
&\geq \min \{ \varphi^+ (f(x)), \varphi^+ (f(y)) \} \\
&= \min \{ (f^{-1}(\varphi))^+ (x), (f^{-1}(\varphi))^+ (y) \}
\end{aligned}$$

Therefore, $(f^{-1}(\varphi))^+ (xy) \geq \min \{ (f^{-1}(\varphi))^+ (x), (f^{-1}(\varphi))^+ (y) \}$

$$\begin{aligned}
\text{And } (f^{-1}(\varphi))^- (xy) &= \varphi^- (f(xy)) \\
&= \varphi^- (f(x)f(y)) \\
&\leq \max \{ \varphi^- (f(x)), \varphi^- (f(y)) \} \\
&= \max \{ (f^{-1}(\varphi))^- (x), (f^{-1}(\varphi))^- (y) \}
\end{aligned}$$

Therefore, $(f^{-1}(\varphi))^- (xy) \leq \max \{ (f^{-1}(\varphi))^- (x), (f^{-1}(\varphi))^- (y) \}$

$$\begin{aligned}
\text{Now, } (f^{-1}(\varphi))^+ (x^{-1}) &= \varphi^+ (f(x^{-1})) \\
&= \varphi^+ (f(x)^{-1}) \\
&= \varphi^+ (f(x)) \\
&= (f^{-1}(\varphi))^+ (x)
\end{aligned}$$

$$\begin{aligned}
\text{And } (f^{-1}(\varphi))^- (x^{-1}) &= \varphi^- (f(x^{-1})) \\
&= \varphi^- (f(x)^{-1}) \\
&= \varphi^- (f(x)) \\
&= (f^{-1}(\varphi))^- (x)
\end{aligned}$$

Therefore, $f^{-1}(\varphi)$ is a bipolar fuzzy subgroup of G_1 .

Hence, if φ be a bipolar fuzzy subgroup of G_2 then $f^{-1}(\varphi)$ is a bipolar fuzzy subgroup of G_1 .

Theorem 3.3: *Let f be an anti homomorphism from a group G_1 into a group G_2 . If $\mu = (\mu^+, \mu^-)$ is a bipolar fuzzy subgroup of G_1 then $f(\mu)$, the image of μ under f , is a bipolar fuzzy subgroup of a group G_2 .*

Proof: Let $\mu = (\mu^+, \mu^-)$ be a bipolar fuzzy subgroup of G_1 , $\mu^+ : G_1 \rightarrow [0,1]$ and $\mu^- : G_1 \rightarrow [-1,0]$ are mappings.

Let $u, v \in G_2$, since f is an anti homomorphism and so there exist $x, y \in G_1$ such that $f(x) = u$ and $f(y) = v$ it follows that $xy \in f^{-1}(vu)$.

$$\begin{aligned} \text{Now, } (f(\mu))^+(uv) &= \max \{ \mu^+(z) : z = xy \in f^{-1}(vu) \} \\ &\geq \max \{ \mu^+(xy) : x \in f^{-1}(u), y \in f^{-1}(v) \} \\ &\geq \max \{ \min \{ \mu^+(x), \mu^+(y) \} : x \in f^{-1}(u), y \in f^{-1}(v) \} \\ &= \min \{ \max \{ \mu^+(x) : x \in f^{-1}(u) \}, \max \{ \mu^+(y) : y \in f^{-1}(v) \} \} \\ &= \min \{ (f(\mu))^+(u), (f(\mu))^+(v) \} \end{aligned}$$

$$\text{Therefore, } (f(\mu))^+(uv) \geq \min \{ (f(\mu))^+(u), (f(\mu))^+(v) \}$$

$$\begin{aligned} \text{And } (f(\mu))^-(uv) &= \max \{ \mu^-(z) : z = xy \in f^{-1}(vu) \} \\ &\leq \max \{ \mu^-(xy) : x \in f^{-1}(u), y \in f^{-1}(v) \} \\ &\leq \max \{ \max \{ \mu^-(x), \mu^-(y) \} : x \in f^{-1}(u), y \in f^{-1}(v) \} \\ &= \max \{ \max \{ \mu^-(x) : x \in f^{-1}(u) \}, \max \{ \mu^-(y) : y \in f^{-1}(v) \} \} \\ &= \max \{ (f(\mu))^-(u), (f(\mu))^-(v) \} \end{aligned}$$

$$\text{Therefore, } (f(\mu))^-(uv) \leq \max \{ (f(\mu))^-(u), (f(\mu))^-(v) \}$$

$$\begin{aligned} \text{Now, } (f(\mu))^+(u^{-1}) &= \max \{ \mu^+(x) : x \in f^{-1}(u^{-1}) \} \\ &= \max \{ \mu^+(x^{-1}) : x^{-1} \in f^{-1}(u) \} \\ &= (f(\mu))^+(u) \end{aligned}$$

$$\begin{aligned} \text{And } (f(\mu))^-(u^{-1}) &= \max \{ \mu^-(x) : x \in f^{-1}(u^{-1}) \} \\ &= \max \{ \mu^-(x^{-1}) : x^{-1} \in f^{-1}(u) \} \\ &= (f(\mu))^-(u) \end{aligned}$$

Therefore, $f(\mu)$ is a bipolar fuzzy subgroup of G_2 .

Hence, if μ be a bipolar fuzzy subgroup of G_1 then $f(\mu)$ is a bipolar fuzzy subgroup of G_2 .

Theorem 3.4: *The anti homomorphic pre-image of a bipolar fuzzy subgroup $\varphi = (\varphi^+, \varphi^-)$ of a group G_2 is a bipolar fuzzy subgroup of a group G_1 .*

Proof: Let $\varphi = (\varphi^+, \varphi^-)$ be a bipolar fuzzy subgroup of G_2 , $\varphi^+ : G_2 \rightarrow [0,1]$ and $\varphi^- : G_2 \rightarrow [-1,0]$ are mappings.

$$\begin{aligned} \text{Now, } (f^{-1}(\varphi))^+(xy) &= \varphi^+(f(xy)) \\ &= \varphi^+(f(y)f(x)) \\ &\geq \min \{ \varphi^+(f(y)), \varphi^+(f(x)) \} \\ &= \min \{ (f^{-1}(\varphi))^+(y), (f^{-1}(\varphi))^+(x) \} \\ &= \min \{ (f^{-1}(\varphi))^+(x), (f^{-1}(\varphi))^+(y) \} \end{aligned}$$

Therefore, $(f^{-1}(\varphi))^+(xy) \geq \min\{(f^{-1}(\varphi))^+(x), (f^{-1}(\varphi))^+(y)\}$

$$\begin{aligned} \text{And } (f^{-1}(\varphi))^{-}(xy) &= \varphi^{-}(f(xy)) \\ &= \varphi^{-}(f(y)f(x)) \\ &\leq \max\{\varphi^{-}(f(y)), \varphi^{-}(f(x))\} \\ &= \max\{(f^{-1}(\varphi))^{-}(y), (f^{-1}(\varphi))^{-}(x)\} \\ &= \max\{(f^{-1}(\varphi))^{-}(x), (f^{-1}(\varphi))^{-}(y)\} \end{aligned}$$

Therefore, $(f^{-1}(\varphi))^{-}(xy) \leq \max\{(f^{-1}(\varphi))^{-}(x), (f^{-1}(\varphi))^{-}(y)\}$

$$\begin{aligned} \text{Now, } (f^{-1}(\varphi))^+(x^{-1}) &= \varphi^{+}(f(x^{-1})) \\ &= \varphi^{+}(f(x)^{-1}) \\ &= \varphi^{+}(f(x)) \\ &= (f^{-1}(\varphi))^+(x) \end{aligned}$$

$$\begin{aligned} \text{And } (f^{-1}(\varphi))^{-}(x^{-1}) &= \varphi^{-}(f(x^{-1})) \\ &= \varphi^{-}(f(x)^{-1}) \\ &= \varphi^{-}(f(x)) \\ &= (f^{-1}(\varphi))^{-}(x) \end{aligned}$$

Therefore, $f^{-1}(\varphi)$ is a bipolar fuzzy subgroup of G_1 .

Hence, if φ be a bipolar fuzzy subgroup of G_2 then $f^{-1}(\varphi)$ is a bipolar fuzzy subgroup of G_1 .

4 Image and Pre-Image of a Bipolar Fuzzy HX Group of a HX Group under Homomorphism and Anti Homomorphism

In this section, we introduce the notion of image and pre-image of the bipolar fuzzy sub HX group of a HX group, and discuss some of its properties. Throughout this section, We mean that ϑ_1 and ϑ_2 are HX groups and E_1 and E_2 are the identity elements of ϑ_1 and ϑ_2 respectively, and XY we mean $X * Y$.

Definition 4.1: Let G_1 and G_2 be any two groups. Let $\vartheta_1 \subset 2^{G_1} - \{\emptyset\}$ and $\vartheta_2 \subset 2^{G_2} - \{\emptyset\}$ are HX groups defined on G_1 and G_2 respectively. Let $\mu = (\mu^+, \mu^-)$ and $\varphi = (\varphi^+, \varphi^-)$ are bipolar fuzzy subsets in G_1 and G_2 respectively, let $\lambda_\mu = (\lambda_\mu^+, \lambda_\mu^-)$, and $\lambda_\varphi = (\lambda_\varphi^+, \lambda_\varphi^-)$ are bipolar fuzzy subsets defined on ϑ_1 and ϑ_2 respectively induced by μ and φ . Let $f: \vartheta_1 \rightarrow \vartheta_2$ be a mapping then the image $f(\lambda_\mu)$ of λ_μ is the bipolar fuzzy subset $(f(\lambda_\mu)) = ((f(\lambda_\mu))^+, (f(\lambda_\mu))^-)$ of ϑ_2 defined by for $U \in \vartheta_2$,

$$(f(\lambda_\mu))^+(U) = \begin{cases} \max\{(\lambda_\mu)^+(X) : X \in f^{-1}(U)\}, & \text{if } f^{-1}(U) \neq \emptyset \\ 0 & , \text{ otherwise} \end{cases}$$

And

$$(f(\lambda_\mu))^- (U) = \begin{cases} \max \{ (\lambda_\mu)^- (X) : X \in f^{-1}(U) \}, & \text{if } f^{-1}(U) \neq \emptyset \\ 0 & , \text{ otherwise} \end{cases}$$

also the pre-image $f^{-1}(\lambda_\varphi)$ of λ_φ under f is bipolar fuzzy subset of ϑ_1 defined by $(f^{-1}(\lambda_\varphi))^+ (X) = \lambda_\varphi^+ (f(X))$, $(f^{-1}(\lambda_\varphi))^- (X) = \lambda_\varphi^- (f(X))$.

Theorem 4.1: Let f be a homomorphism from a HX group ϑ_1 into a HX group ϑ_2 . If $\lambda_\mu = (\lambda_\mu^+, \lambda_\mu^-)$ is a bipolar fuzzy sub HX group on ϑ_1 then $f(\lambda_\mu)$, the image of λ_μ under f , is a bipolar fuzzy sub HX group of ϑ_2 .

Proof: Let $\mu = (\mu^+, \mu^-)$ be a bipolar fuzzy subset of G_1 , $\mu^+ : G_1 \rightarrow [0,1]$ and $\mu^- : G_1 \rightarrow [-1,0]$ are mapping, and λ_μ is a bipolar fuzzy sub HX group on ϑ_1 .

Let $U, V \in \vartheta_2$, since f is homomorphism and so there exist $X, Y \in \vartheta_1$ such that $f(X) = U$ and $f(Y) = V$ it follows that $XY \in f^{-1}(UV)$.

Now,

$$\begin{aligned} (f(\lambda_\mu))^+ (UV) &= \max \{ \lambda_\mu^+ (Z) : Z = XY \in f^{-1}(UV) \} \\ &\geq \max \{ \lambda_\mu^+ (XY) : X \in f^{-1}(U), Y \in f^{-1}(V) \} \\ &\geq \max \{ \min \{ \lambda_\mu^+ (X), \lambda_\mu^+ (Y) \} : X \in f^{-1}(U), Y \in f^{-1}(V) \} \\ &= \min \{ \max \{ \lambda_\mu^+ (X) : X \in f^{-1}(U) \}, \max \{ \lambda_\mu^+ (Y) : Y \in f^{-1}(V) \} \} \\ &= \min \{ (f(\lambda_\mu))^+ (U), (f(\lambda_\mu))^+ (V) \} \end{aligned}$$

Therefore, $(f(\lambda_\mu))^+ (UV) \geq \min \{ (f(\lambda_\mu))^+ (U), (f(\lambda_\mu))^+ (V) \}$

And,

$$\begin{aligned} (f(\lambda_\mu))^- (UV) &= \max \{ \lambda_\mu^- (Z) : Z = XY \in f^{-1}(UV) \} \\ &\leq \max \{ \lambda_\mu^- (XY) : X \in f^{-1}(U), Y \in f^{-1}(V) \} \\ &\leq \max \{ \max \{ \lambda_\mu^- (X), \lambda_\mu^- (Y) \} : X \in f^{-1}(U), Y \in f^{-1}(V) \} \\ &= \max \{ \max \{ \lambda_\mu^- (X) : X \in f^{-1}(U) \}, \max \{ \lambda_\mu^- (Y) : Y \in f^{-1}(V) \} \} \\ &= \max \{ (f(\lambda_\mu))^- (U), (f(\lambda_\mu))^- (V) \} \end{aligned}$$

Therefore, $(f(\lambda_\mu))^- (UV) \leq \max \{ (f(\lambda_\mu))^- (U), (f(\lambda_\mu))^- (V) \}$

$$\begin{aligned} \text{Now, } (f(\lambda_\mu))^+ (U^{-1}) &= \max \{ \lambda_\mu^+ (X) : X \in f^{-1}(U^{-1}) \} \\ &= \max \{ \lambda_\mu^+ (X^{-1}) : X^{-1} \in f^{-1}(U) \} \\ &= (f(\lambda_\mu))^+ (U) \end{aligned}$$

$$\begin{aligned} \text{And } (f(\lambda_\mu))^- (U^{-1}) &= \max \{ \lambda_\mu^- (X) : X \in f^{-1}(U^{-1}) \} \\ &= \max \{ \lambda_\mu^- (X^{-1}) : X^{-1} \in f^{-1}(U) \} \\ &= (f(\lambda_\mu))^- (U) \end{aligned}$$

Therefore, $f(\lambda_\mu)$ is a bipolar fuzzy sub HX group of ϑ_2 .

Hence, if λ_μ be a bipolar fuzzy sub HX group on ϑ_1 then $f(\lambda_\mu)$ is a bipolar fuzzy sub HX group of ϑ_2 .

Theorem 4.2: *The homomorphic pre-image of a bipolar fuzzy sub HX group $\lambda_\varphi = (\lambda_\varphi^+, \lambda_\varphi^-)$ of a HX group ϑ_2 is a bipolar fuzzy sub HX group of a HX group ϑ_1 .*

Proof: Let $\varphi = (\varphi^+, \varphi^-)$ be a bipolar fuzzy subset of G_2 , $\varphi^+ : G_2 \rightarrow [0,1]$ and $\varphi^- : G_2 \rightarrow [-1,0]$ are mappings, and λ_φ be a bipolar fuzzy sub HX group on ϑ_2 .

$$\begin{aligned} \text{Now, } (f^{-1}(\lambda_\varphi))^+(XY) &= \lambda_\varphi^+(f(XY)) \\ &= \lambda_\varphi^+(f(X)f(Y)) \\ &\geq \min \{ \lambda_\varphi^+(f(X)), \lambda_\varphi^+(f(Y)) \} \\ &= \min \{ (f^{-1}(\lambda_\varphi))^+(X), (f^{-1}(\lambda_\varphi))^+(Y) \} \end{aligned}$$

$$\text{Therefore, } (f^{-1}(\lambda_\varphi))^+(XY) \geq \min \{ (f^{-1}(\lambda_\varphi))^+(X), (f^{-1}(\lambda_\varphi))^+(Y) \}$$

$$\begin{aligned} \text{And } (f^{-1}(\lambda_\varphi))^- (XY) &= \lambda_\varphi^-(f(XY)) \\ &= \lambda_\varphi^-(f(X)f(Y)) \\ &\leq \max \{ \lambda_\varphi^-(f(X)), \lambda_\varphi^-(f(Y)) \} \\ &= \max \{ (f^{-1}(\lambda_\varphi))^- (X), (f^{-1}(\lambda_\varphi))^- (Y) \} \end{aligned}$$

$$\text{Therefore, } (f^{-1}(\lambda_\varphi))^- (XY) \leq \max \{ (f^{-1}(\lambda_\varphi))^- (X), (f^{-1}(\lambda_\varphi))^- (Y) \}$$

$$\begin{aligned} \text{Now, } (f^{-1}(\lambda_\varphi))^+(X^{-1}) &= \lambda_\varphi^+(f(X^{-1})) \\ &= \lambda_\varphi^+(f(X)^{-1}) \\ &= \lambda_\varphi^+(f(X)) \\ &= (f^{-1}(\lambda_\varphi))^+(X) \end{aligned}$$

$$\begin{aligned} \text{And } (f^{-1}(\lambda_\varphi))^- (X^{-1}) &= \lambda_\varphi^-(f(X^{-1})) \\ &= \lambda_\varphi^-(f(X)^{-1}) \\ &= \lambda_\varphi^-(f(X)) \\ &= (f^{-1}(\lambda_\varphi))^- (X) \end{aligned}$$

Therefore, $f^{-1}(\lambda_\varphi)$ is a bipolar fuzzy sub HX group of ϑ_1 .

Hence, if λ_φ be a bipolar fuzzy sub HX group on ϑ_2 then $f^{-1}(\lambda_\varphi)$ is a bipolar fuzzy sub HX group of ϑ_1 .

Theorem 4.3: *Let f be an anti homomorphism from a HX group ϑ_1 into a HX group ϑ_2 . If $\lambda_\mu = (\lambda_\mu^+, \lambda_\mu^-)$ is a bipolar fuzzy sub HX group of ϑ_1 then $f(\lambda_\mu)$, the image of λ_μ under f , is a bipolar fuzzy sub HX group of ϑ_2 .*

Proof: Let $\mu = (\mu^+, \mu^-)$ be a bipolar fuzzy subgroup of G_1 , $\mu^+ : G_1 \rightarrow [0,1]$ and $\mu^- : G_1 \rightarrow [-1,0]$ are mappings, and λ_μ is a bipolar fuzzy sub HX group of ϑ_1 .

Let $U, V \in \vartheta_2$, since f is an anti homomorphism and so there exist $X, Y \in \vartheta_1$ such that $f(X) = U$ and $f(Y) = V$ it follows that $XY \in f^{-1}(VU)$.

Now,

$$\begin{aligned} (f(\lambda_\mu))^+(UV) &= \max \{ \lambda_\mu^+(Z) : Z = XY \in f^{-1}(VU) \} \\ &\geq \max \{ \lambda_\mu^+(XY) : X \in f^{-1}(U), Y \in f^{-1}(V) \} \\ &\geq \max \{ \min \{ \lambda_\mu^+(X), \lambda_\mu^+(Y) \} : X \in f^{-1}(U), Y \in f^{-1}(V) \} \\ &= \min \{ \max \{ \lambda_\mu^+(X) : X \in f^{-1}(U) \}, \max \{ \lambda_\mu^+(Y) : Y \in f^{-1}(V) \} \} \\ &= \min \{ (f(\lambda_\mu))^+(U), (f(\lambda_\mu))^+(V) \} \end{aligned}$$

Therefore, $(f(\lambda_\mu))^+(UV) \geq \min \{ (f(\lambda_\mu))^+(U), (f(\lambda_\mu))^+(V) \}$

And

$$\begin{aligned} (f(\lambda_\mu))^{-}(UV) &= \max \{ \lambda_\mu^{-}(Z) : Z = XY \in f^{-1}(VU) \} \\ &\leq \max \{ \lambda_\mu^{-}(XY) : X \in f^{-1}(U), Y \in f^{-1}(V) \} \\ &\leq \max \{ \max \{ \lambda_\mu^{-}(X), \lambda_\mu^{-}(Y) \} : X \in f^{-1}(U), Y \in f^{-1}(V) \} \\ &= \max \{ \max \{ \lambda_\mu^{-}(X) : X \in f^{-1}(U) \}, \max \{ \lambda_\mu^{-}(Y) : Y \in f^{-1}(V) \} \} \\ &= \max \{ (f(\lambda_\mu))^{-}(U), (f(\lambda_\mu))^{-}(V) \} \end{aligned}$$

Therefore, $(f(\lambda_\mu))^{-}(UV) \leq \max \{ (f(\lambda_\mu))^{-}(U), (f(\lambda_\mu))^{-}(V) \}$

$$\begin{aligned} \text{Now, } (f(\lambda_\mu))^+(U^{-1}) &= \max \{ \lambda_\mu^+(X) : X \in f^{-1}(U^{-1}) \} \\ &= \max \{ \lambda_\mu^+(X^{-1}) : X^{-1} \in f^{-1}(U) \} \\ &= (f(\lambda_\mu))^+(U) \end{aligned}$$

$$\begin{aligned} \text{And } (f(\lambda_\mu))^{-}(U^{-1}) &= \max \{ \lambda_\mu^{-}(X) : X \in f^{-1}(U^{-1}) \} \\ &= \max \{ \lambda_\mu^{-}(X^{-1}) : X^{-1} \in f^{-1}(U) \} \\ &= (f(\lambda_\mu))^{-}(U) \end{aligned}$$

Therefore, $f(\lambda_\mu)$ is a bipolar fuzzy sub HX group of ϑ_2 .

Hence, if λ_μ be a bipolar fuzzy sub HX group on ϑ_1 then $f(\lambda_\mu)$ is a bipolar fuzzy sub HX group of ϑ_2 .

Theorem 4.4: *The anti homomorphic pre-image of a bipolar fuzzy sub HX group $\lambda_\varphi = (\lambda_\varphi^+, \lambda_\varphi^-)$ of a HX group ϑ_2 is a bipolar fuzzy sub HX group of a HX group ϑ_1 .*

Proof: Let $\varphi = (\varphi^+, \varphi^-)$ be a bipolar fuzzy subgroup of G_2 , $\varphi^+ : G_2 \rightarrow [0,1]$ and $\varphi^- : G_2 \rightarrow [-1,0]$ are mappings and λ_φ be a bipolar fuzzy sub HX group on ϑ_2 .

$$\begin{aligned} \text{Now, } (f^{-1}(\lambda_\varphi))^+(XY) &= \lambda_\varphi^+(f(XY)) \\ &= \lambda_\varphi^+(f(Y)f(X)) \\ &\geq \min \{ \lambda_\varphi^+(f(Y)), \lambda_\varphi^+(f(X)) \} \\ &= \min \{ (f^{-1}(\lambda_\varphi))^+(Y), (f^{-1}(\lambda_\varphi))^+(X) \} \end{aligned}$$

$$= \min \{ (f^{-1}(\lambda_\varphi))^+(X), (f^{-1}(\lambda_\varphi))^+(Y) \}$$

Therefore, $(f^{-1}(\lambda_\varphi))^+(XY) \geq \min \{ (f^{-1}(\lambda_\varphi))^+(X), (f^{-1}(\lambda_\varphi))^+(Y) \}$

$$\begin{aligned} \text{And } (f^{-1}(\lambda_\varphi))^{-}(XY) &= \lambda_\varphi^{-}(f(XY)) \\ &= \lambda_\varphi^{-}(f(Y)f(X)) \\ &\leq \max \{ \lambda_\varphi^{-}(f(Y)), \lambda_\varphi^{-}(f(X)) \} \\ &= \max \{ (f^{-1}(\lambda_\varphi))^{-}(Y), (f^{-1}(\lambda_\varphi))^{-}(X) \} \\ &= \max \{ (f^{-1}(\lambda_\varphi))^{-}(X), (f^{-1}(\lambda_\varphi))^{-}(Y) \} \end{aligned}$$

Therefore, $(f^{-1}(\lambda_\varphi))^{-}(XY) \leq \max \{ (f^{-1}(\lambda_\varphi))^{-}(X), (f^{-1}(\lambda_\varphi))^{-}(Y) \}$

$$\begin{aligned} \text{Now, } (f^{-1}(\lambda_\varphi))^+(X^{-1}) &= \lambda_\varphi^{+}(f(X^{-1})) \\ &= \lambda_\varphi^{+}(f(X)^{-1}) \\ &= \lambda_\varphi^{+}(f(X)) \\ &= (f^{-1}(\lambda_\varphi))^+(X) \end{aligned}$$

$$\begin{aligned} \text{And } (f^{-1}(\lambda_\varphi))^{-}(X^{-1}) &= \lambda_\varphi^{-}(f(X^{-1})) \\ &= \lambda_\varphi^{-}(f(X)^{-1}) \\ &= \lambda_\varphi^{-}(f(X)) \\ &= (f^{-1}(\lambda_\varphi))^{-}(X) \end{aligned}$$

Therefore, $f^{-1}(\lambda_\varphi)$ is a bipolar fuzzy sub HX group of ϑ_1 .

Hence, if λ_φ be a bipolar fuzzy sub HX group on ϑ_2 then $f^{-1}(\lambda_\varphi)$ is a bipolar fuzzy sub HX group of ϑ_1 .

References

- [1] P.S. Das, Fuzzy groups and level subgroups, *J. Math. Anal. Appl.*, 84(1981), 264-269.
- [2] K.M. Lee, Bipolar-valued fuzzy sets and their operations, *Proc. Int. Conf. on Intelligent Technologies*, Bangkok, Thailand, (2000), 307-312.
- [3] L. Hongxing, HX group, *BUSEFAL*, 33(4) (October) (1987), 31-37.
- [4] L. Chengzhong, M. Honghai and L. Hongxing, Fuzzy HX group, *BUSEFAL*, 41(14) (Oct) (1989), 97-106.
- [5] M. Marudai and V. Rajendran, New constructions on bipolar anti Q-fuzzy groups and bipolar anti Q-fuzzy d-ideals under (t, s) norms, *Advances in Fuzzy Mathematics*, 6(1) (2011), 145-153.
- [6] M.S. Eroglu, The homomorphic image of a fuzzy subgroup is always a fuzzy subgroup, *Fuzzy Sets and System*, 33(1989), 255-256.
- [7] Md. Asaad, Groups and fuzzy subgroups, *Fuzzy Sets and Systems*, 39(1991), 323-328.
- [8] M. Akgul, Some properties of fuzzy groups, *J. Math. Anal. Appl.*, 133(1988), 93-100.

- [9] R. Muthuraj, M.S. Muthuraman and M. Sridharan, Bipolar fuzzy Sub-Bi HX group and its Bi Level Sub-Bi HX groups, *Antarctica J. Math.*, 8(5) (2011), 427-434.
- [10] R. Muthuraj and M. Sridharan, Bipolar anti fuzzy HX group and its lower level sub HX groups, *Journal of Physical Sciences*, 16(December) (2012), 157-169.
- [11] R. Kumar, Homomorphism and fuzzy (fuzzy normal) subgroups, *Fuzzy Set and System*, 44(1991), 165-168.
- [12] A. Rosenfeld, Fuzzy groups, *J. Math. Anal. Appl.*, 35(1971), 512-517.
- [13] K. Sunderrajan, R. Muthuraj, M.S. Muthuraman and M. Sridharan, Some characterization of anti Q-L-fuzzy l-group, *International Journal of Computer Application (0975-8887)*, 6(4) (September) (2010), 35-47.
- [14] W.R. Zhang, Bipolar fuzzy sets, *Proc. of FUZZ-IEEE*, (1998), 835-840.
- [15] W.R. Zhang, Bipolar fuzzy sets and relations: A computational frame work for cognitive modeling and multiple decision analysis, *Proc. of IEEE Conference*, (1994), 305-309.
- [16] L.A. Zadeh, Fuzzy sets, *Inform and Control*, 8(1965), 338-365.
- [17] H.J. Zimmermann, *Fuzzy Set Theory and its Applications*, Kluwer-Nijhoff Publishing, (1985).