

Research Article

On the Convergence of Multistep Iteration for Uniformly Continuous Φ -Hemicontractive Mappings

Zhiqun Xue,¹ Arif Rafiq,² and Haiyun Zhou³

¹ Department of Mathematics and Physics, Shijiazhuang Tiedao University, Shijiazhuang 050043, China

² Hajvery University, 43-52 Industrial Area, Gulberg III, Lahore 54660, Pakistan

³ Department of Mathematics, Shijiazhuang Mechanical Engineering College, Shijiazhuang 050003, China

Correspondence should be addressed to Zhiqun Xue, xuezhiqun@126.com

Received 18 July 2012; Revised 25 August 2012; Accepted 28 August 2012

Academic Editor: Xiaolong Qin

Copyright © 2012 Zhiqun Xue et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

It is shown that the convergence of the multistep iterative process with errors is obtained for uniformly continuous Φ -hemicontractive mappings in real Banach spaces. We also revise the problems of C. E. Chidume and C. O. Chidume (2005).

1. Introduction

Let E be a real Banach space with norm $\|\cdot\|$ and let E^* be its dual space. The normalized duality mapping $J : E \rightarrow 2^{E^*}$ is defined by

$$J(x) = \left\{ f \in E^* : \langle x, f \rangle = \|x\|^2 = \|f\|^2 \right\}, \quad \forall x \in E, \quad (1.1)$$

where $\langle \cdot, \cdot \rangle$ denotes the generalized duality pairing. The single-valued-normalized duality mapping is denoted by j .

A mapping T with domain $D(T)$ and range $R(T)$ in E is said to be strongly pseudocontractive if there is a constant $k \in (0, 1)$, and for all $x, y \in D(T)$, $\exists j(x - y) \in J(x - y)$ such that

$$\langle Tx - Ty, j(x - y) \rangle \leq k \|x - y\|^2. \quad (1.2)$$

The mapping T is called Φ -pseudocontractive if there exists a strictly increasing continuous function $\Phi : [0, +\infty) \rightarrow [0, +\infty)$ with $\Phi(0) = 0$ such that

$$\langle Tx - Ty, j(x - y) \rangle \leq \|x - y\|^2 - \Phi(\|x - y\|) \quad (1.3)$$

holds for all $x, y \in D(T)$. It is well known that the strongly pseudocontractive mapping must be the Φ -pseudocontractive mapping in the special case in which $\Phi(t) = (1 - k)t^2$, but the converse is not true in general. That is, the class of strongly pseudocontractive mappings is a proper subclass of the class of Φ -pseudocontractive mappings. Let $F(T) = \{x \in D(T) : Tx = x\} \neq \emptyset$. If the inequalities (1.2) and (1.3) hold for any $x \in D(T)$ and $y \in F(T)$, then the corresponding mapping T is called strongly hemiccontractive and Φ -hemiccontractive, respectively.

Let $N(T) = \{x \in E : Tx = 0\} \neq \emptyset$. An operator $T : D(T) \subseteq E \rightarrow E$ is called strongly quasiaccretive, Φ -quasiaccretive if and only if $I - T$ is strongly hemiccontractive, Φ -hemiccontractive, respectively, where I denotes the identity mapping on E . That is, if T is Φ -quasi-accretive, then $N(T) \neq \emptyset$ and there exists a strictly increasing continuous function $\Phi : [0, +\infty) \rightarrow [0, +\infty)$ with $\Phi(0) = 0$ such that

$$\langle Tx - Ty, j(x - y) \rangle \geq \Phi(\|x - y\|) \quad (1.4)$$

holds for all $x \in D(T)$ and $y \in N(T)$. Many authors have studied extensively the strongly convergence problems of the iterative algorithms for the class of operators.

In 2004, Rhoades and Soltuz [1] introduced the multistep iteration as follows.

Let D be a nonempty closed convex subset of real Banach space E and let $T : D \rightarrow D$ be a mapping. The multistep iteration $\{x_n\}$ is defined by

$$\begin{aligned} x_0 &\in D, \\ y_n^{p-1} &= (1 - b_n^{p-1})x_n + b_n^{p-1}Tx_n, \quad n \geq 0, \quad p \geq 2, \\ y_n^k &= (1 - b_n^k)x_n + b_n^kTy_n^{k+1}, \quad k = p-2, p-3, \dots, 2, 1, \\ x_{n+1} &= (1 - a_n)x_n + a_nTy_n^1, \quad n \geq 0, \end{aligned} \quad (1.5)$$

where $\{a_n\}, \{b_n^k\}$ ($k = 1, 2, \dots, p-1$) in $[0, 1]$ satisfy certain conditions. Obviously, the iteration defined above is generalization of Mann, Ishikawa, and Noor iterations.

Inspired and motivated by the work of Xu [2] and the iteration above, we discuss the following multistep iteration with errors:

$$\begin{aligned} x_0 &\in D, \\ y_n^{p-1} &= (1 - b_n^{p-1} - d_n^{p-1})x_n + b_n^{p-1}Tx_n + d_n^{p-1}w_n^{p-1}, \quad n \geq 0, \quad p \geq 2, \\ y_n^k &= (1 - b_n^k - d_n^k)x_n + b_n^kTy_n^{k+1} + d_n^kw_n^k, \quad k = p-2, p-3, \dots, 2, 1, \\ x_{n+1} &= (1 - a_n - c_n)x_n + a_nTy_n^1 + c_nu_n, \quad n \geq 0, \end{aligned} \quad (1.6)$$

where $\{a_n\}, \{c_n\}, \{b_n^k\}, \{d_n^k\}$ ($k = 1, 2, \dots, p-1$) in $[0, 1]$ with $a_n + c_n \leq 1$, $b_n^k + d_n^k \leq 1$, $\{u_n\}, \{w_n^k\}$ ($k = 1, 2, \dots, p-1$) are the bounded sequences of D .

In 2005, C. E. Chidume and C. O. Chidume [3] proved the convergence theorems for fixed points of uniformly continuous generalized Φ -hemicontractive mappings and published in [3]. However, there exists a gap in the proof course of their theorems.

The aim of this paper is to show the convergence of the multistep iteration with errors for fixed points of uniformly continuous Φ -hemicontractive mappings and revise the results of C. E. Chidume and C. O. Chidume [3]. For this, we need the following Lemmas.

Lemma 1.1 (see [4]). *Let E be a real Banach space and let $J : E \rightarrow 2^{E^*}$ be a normalized duality mapping. Then*

$$\|x + y\|^2 \leq \|x\|^2 + 2\langle y, j(x + y) \rangle, \tag{1.7}$$

for all $x, y \in E$ and for all $j(x + y) \in J(x + y)$.

Lemma 1.2 (see [5]). *Let $\{\delta_n\}_{n=0}^\infty$, $\{\lambda_n\}_{n=0}^\infty$ and $\{\gamma_n\}_{n=0}^\infty$ be three nonnegative real sequences and let $\Phi : [0, +\infty) \rightarrow [0, +\infty)$ be a strictly increasing and continuous function with $\Phi(0) = 0$ satisfying the following inequality:*

$$\delta_{n+1}^2 \leq \delta_n^2 - \lambda_n \Phi(\delta_{n+1}) + \gamma_n, \quad n \geq 0, \tag{1.8}$$

where $\lambda_n \in [0, 1]$ with $\sum_{n=0}^\infty \lambda_n = \infty$, $\gamma_n = o(\lambda_n)$. Then $\delta_n \rightarrow 0$ as $n \rightarrow \infty$.

2. Main Results

Theorem 2.1. *Let E be an arbitrary real Banach space, D a nonempty closed convex subset of E , and $T : D \rightarrow D$ a uniformly continuous Φ -hemicontractive mapping with $q \in F(T) \neq \emptyset$. Let $\{a_n\}$, $\{b_n^k\}$, $\{c_n\}$, $\{d_n^k\}$ be real sequences in $[0, 1]$ and satisfy the conditions:*

- (i) $a_n + c_n \leq 1$, $b_n^k + d_n^k \leq 1$, $k = 1, 2, \dots, p - 1$;
- (ii) $a_n, c_n, b_n^k, d_n^k \rightarrow 0$ as $n \rightarrow \infty$, $k = 1, 2, \dots, p - 1$;
- (iii) $c_n = o(a_n)$, $\sum_{n=0}^\infty a_n = \infty$.

For some $x_0 \in D$, let $\{u_n\}$, $\{w_n^1\}$, $\{w_n^2\}$, \dots , $\{w_n^{p-1}\}$ be any bounded sequences of D , and let $\{x_n\}$ be the multistep iterative sequence with errors defined by (1.6). Then (1.6) converges strongly to the fixed point q of T .

Proof. Since $T : D \rightarrow D$ is Φ -hemicontractive mapping, then there exists a strictly increasing continuous function $\Phi : [0, +\infty) \rightarrow [0, +\infty)$ with $\Phi(0) = 0$ such that

$$\langle Tx - Tq, j(x - q) \rangle \leq \|x - q\|^2 - \Phi(\|x - q\|), \tag{2.1}$$

for $x \in D$, $q \in F(T)$, that is

$$\langle Tx - x, j(x - q) \rangle \leq -\Phi(\|x - q\|). \tag{@}$$

Choose some $x_0 \in D$ and $x_0 \neq Tx_0$ such that $\|x_0 - Tx_0\| \cdot \|x_0 - q\| \in R(\Phi)$ and denote that $r_0 = \|x_0 - Tx_0\| \cdot \|x_0 - q\|$, $R(\Phi)$ is the range of Φ . Indeed, if $\Phi(r) \rightarrow +\infty$ as $r \rightarrow +\infty$, then

$r_0 \in R(\Phi)$; if $\sup\{\Phi(r) : r \in [0, +\infty)\} = r_1 < +\infty$ with $r_1 < r_0$ (here, we only give an example. If $r_0 = 2$, $\Phi(t) = t/(1+t)$, then $\sup\{\Phi(r) : r \in [0, +\infty)\} = r_1 = 1 < 2 = r_0$), then for $q \in D$, there exists a sequence $\{w_n\}$ in D such that $w_n \rightarrow q$ as $n \rightarrow \infty$ with $w_n \neq q$. Furthermore, we obtain that $Tw_n \rightarrow Tq$ as $n \rightarrow \infty$. So $\{w_n - Tw_n\}$ is the bounded sequence. Hence, there exists a natural number n_0 such that $\|w_n - Tw_n\| \cdot \|w_n - q\| < r_1/2$ for $n \geq n_0$, then we redefine $x_0 = w_{n_0}$ and $\|x_0 - Tx_0\| \cdot \|x_0 - q\| \in R(\Phi)$. This is to ensure that $\Phi^{-1}(r_0)$ is defined well.

Step 1. We show that $\{x_n\}$ is a bounded sequence.

Set $R = \Phi^{-1}(r_0)$, then from above formula (@), we obtain that $\|x_0 - q\| \leq R$. Denote

$$B_1 = \{x \in D : \|x - q\| \leq R\}, \quad B_2 = \{x \in D : \|x - q\| \leq 2R\}. \quad (2.2)$$

Since T is uniformly continuous, so T is a bounded mapping. We let

$$\begin{aligned} M = & \sup_{x \in B_2} \{\|Tx - q\| + 1\} \\ & + \max \left\{ \sup_n \{\|w_n^1 - q\|\}, \sup_n \{\|w_n^2 - q\|\}, \dots, \sup_n \{\|w_n^{p-1} - q\|\}, \sup_n \{\|u_n - q\|\} \right\}. \end{aligned} \quad (2.3)$$

Next, we want to prove that $x_n \in B_1$. If $n = 0$, then $x_0 \in B_1$. Now, assume that it holds for some n , that is, $x_n \in B_1$. We prove that $x_{n+1} \in B_1$. Suppose that it is not the case, then $\|x_{n+1} - q\| > R > R/2$. Since T is uniformly continuous, then for $\varepsilon_0 = \Phi(R/2)/8R$, there exists $\delta > 0$ such that $\|Tx - Ty\| < \varepsilon_0$ when $\|x - y\| < \delta$. Denote

$$\tau_0 = \min \left\{ 1, \frac{R}{M}, \frac{\Phi(R/2)}{8R(M+2R)}, \frac{\delta}{2M+4R} \right\}. \quad (2.4)$$

Since $a_n, b_n^k, c_n, d_n^k \rightarrow 0$ as $n \rightarrow \infty$ for $k = 1, 2, \dots, p-1$. Without loss of generality, we assume that $0 \leq a_n, b_n^k, c_n, d_n^k \leq \tau_0$ for any $n \geq 0$. Since $c_n = o(a_n)$, let $c_n < a_n \tau_0$. Now, estimate $\|y_n^k - q\|$ for $k = 1, 2, \dots, p-1$. By using (1.6), we have

$$\begin{aligned} \|y_n^{p-1} - q\| & \leq (1 - b_n^{p-1} - d_n^{p-1}) \|x_n - q\| + b_n^{p-1} \|Tx_n - q\| + d_n^{p-1} \|w_n^{p-1} - q\| \\ & \leq R + \tau_0 M \\ & \leq 2R, \end{aligned} \quad (2.5)$$

then $y_n^{p-1} \in B_2$. Similarly, we have

$$\begin{aligned} \|y_n^{p-2} - q\| & \leq (1 - b_n^{p-2} - d_n^{p-2}) \|x_n - q\| + b_n^{p-2} \|Ty_n^{p-1} - q\| + d_n^{p-2} \|w_n^{p-2} - q\| \\ & \leq R + \tau_0 M \\ & \leq 2R, \end{aligned} \quad (2.6)$$

then $y_n^{p-2} \in B_2, \dots$. We have

$$\begin{aligned} \|y_n^1 - q\| &\leq (1 - b_n^1 - d_n^1) \|x_n - q\| + b_n^1 \|Ty_n^2 - q\| + d_n^1 \|w_n^1 - q\| \\ &\leq R + \tau_0 M \\ &\leq 2R, \end{aligned} \quad (2.7)$$

then $y_n^1 \in B_2$. Therefore, we get

$$\begin{aligned} \|x_{n+1} - q\| &\leq (1 - a_n - c_n) \|x_n - q\| + a_n \|Ty_n^1 - q\| + c_n \|u_n - q\| \\ &\leq R + \tau_0 M \\ &\leq 2R. \end{aligned} \quad (2.8)$$

And we have

$$\begin{aligned} \|x_{n+1} - x_n\| &\leq a_n \|Ty_n^1 - x_n\| + c_n \|u_n - x_n\| \\ &\leq a_n (\|Ty_n^1 - q\| + \|x_n - q\|) + c_n (\|u_n - q\| + \|x_n - q\|) \\ &\leq \tau_0 [\|Ty_n^1 - q\| + \|u_n - q\| + 2\|x_n - q\|] \\ &\leq \tau_0 (M + 2R) \\ &\leq \frac{\Phi(R/2)}{8R}, \\ \|x_{n+1} - y_n^1\| &\leq a_n \|Ty_n^1 - x_n\| + c_n \|u_n - x_n\| + b_n^1 \|Ty_n^2 - x_n\| + d_n^1 \|w_n^1 - x_n\| \\ &\leq a_n (\|Ty_n^1 - q\| + \|x_n - q\|) + c_n (\|u_n - q\| + \|x_n - q\|) \\ &\quad + b_n^1 (\|Ty_n^2 - q\| + \|x_n - q\|) + d_n^1 (\|w_n^1 - q\| + \|x_n - q\|) \\ &\leq \tau_0 [\|Ty_n^1 - q\| + \|u_n - q\| + 2\|x_n - q\|] \\ &\quad + (\|Ty_n^2 - q\| + \|w_n^1 - q\| + 2\|x_n - q\|) \\ &\leq \tau_0 (2M + 4R) \\ &\leq \delta. \end{aligned} \quad (2.9)$$

Further, by using uniform continuity of T , we have

$$\|Tx_{n+1} - Ty_n^1\| < \frac{\Phi(R/2)}{8R}. \quad (2.10)$$

In view of Lemma 1.1 and the above formulas, we obtain

$$\begin{aligned}
& \|x_{n+1} - q\|^2 \\
&= \left\| (x_n - q) + a_n(Ty_n^1 - x_n) + c_n(u_n - x_n) \right\|^2 \\
&\leq \|x_n - q\|^2 + 2a_n \langle Ty_n^1 - x_n, j(x_{n+1} - q) \rangle + 2c_n \langle u_n - x_n, j(x_{n+1} - q) \rangle \\
&\leq \|x_n - q\|^2 + 2a_n \langle Tx_{n+1} - x_{n+1} + x_{n+1} - x_n - Tx_{n+1} + Ty_n^1, j(x_{n+1} - q) \rangle \\
&\quad + 2c_n \|u_n - x_n\| \cdot \|x_{n+1} - q\| \\
&\leq \|x_n - q\|^2 - 2a_n \Phi(\|x_{n+1} - q\|) + 2a_n \|x_{n+1} - x_n\| \cdot \|x_{n+1} - q\| \\
&\quad + 2a_n \|Tx_{n+1} - Ty_n^1\| \cdot \|x_{n+1} - q\| + 2c_n (\|u_n - q\| + \|x_n - q\|) \|x_{n+1} - q\| \\
&\leq \|x_n - q\|^2 - 2a_n \Phi\left(\frac{R}{2}\right) + 2a_n \frac{\Phi(R/2)}{8R} \cdot 2R + 2a_n \frac{\Phi(R/2)}{8R} \cdot 2R + 2a_n \tau_0 (R + M) 2R \\
&\leq \|x_n - q\|^2 - a_n \Phi\left(\frac{R}{2}\right) + 2a_n \frac{\Phi(R/2)}{8R(M + 2R)} (R + M) 2R \\
&\leq \|x_n - q\|^2 - \frac{a_n}{2} \Phi\left(\frac{R}{2}\right) \leq R^2,
\end{aligned} \tag{2.11}$$

which is a contradiction. Hence, $x_{n+1} \in B_1$, that is, $\{x_n\}$ is a bounded sequence; it leads to that $\{y_n^1\}, \{y_n^2\}, \dots, \{y_n^{p-1}\}$ are all bounded sequences as well.

Step 2. We want to prove $\|x_n - q\| \rightarrow 0$ as $n \rightarrow \infty$.

Since $a_n, b_n^k, c_n, d_n^k \rightarrow 0$ as $n \rightarrow \infty$ and $\{x_n\}, \{y_n^1\}$ are bounded. From (2.9), we obtain

$$\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0, \quad \lim_{n \rightarrow \infty} \|x_{n+1} - y_n^1\| = 0, \quad \lim_{n \rightarrow \infty} \|Tx_{n+1} - Ty_n^1\| = 0. \tag{2.12}$$

By (2.11), we have

$$\begin{aligned}
\|x_{n+1} - q\|^2 &= \left\| (x_n - q) + a_n(Ty_n^1 - x_n) + c_n(u_n - x_n) \right\|^2 \\
&\leq \|x_n - q\|^2 + 2a_n \langle Ty_n^1 - x_n, j(x_{n+1} - q) \rangle + 2c_n \langle u_n - x_n, j(x_{n+1} - q) \rangle \\
&\leq \|x_n - q\|^2 + 2a_n \langle Tx_{n+1} - x_{n+1} + x_{n+1} - x_n - Tx_{n+1} + Ty_n^1, j(x_{n+1} - q) \rangle \\
&\quad + 2c_n \|u_n - x_n\| \cdot \|x_{n+1} - q\|
\end{aligned}$$

$$\begin{aligned}
 &\leq \|x_n - q\|^2 - 2a_n\Phi(\|x_{n+1} - q\|) + 2a_n\|x_{n+1} - x_n\| \cdot \|x_{n+1} - q\| \\
 &\quad + 2a_n\|Tx_{n+1} - Ty_n^1\| \cdot \|x_{n+1} - q\| + 2c_n\|u_n - x_n\| \cdot \|x_{n+1} - q\| \\
 &= \|x_n - q\|^2 - 2a_n\Phi(\|x_{n+1} - q\|) + o(a_n),
 \end{aligned} \tag{2.13}$$

where $2a_n\|x_{n+1} - x_n\| \cdot \|x_{n+1} - q\| + 2a_n\|Tx_{n+1} - Ty_n^1\| \cdot \|x_{n+1} - q\| + 2c_n\|u_n - x_n\| \cdot \|x_{n+1} - q\| = o(a_n)$. By Lemma 1.2, we obtain that $\lim_{n \rightarrow \infty} \|x_n - q\| = 0$. \square

Theorem 2.2. *Let E be an arbitrary real Banach space and let $T : E \rightarrow E$ be a uniformly continuous Φ -quasi-accretive operator with $q \in N(T) \neq \emptyset$. Let $\{a_n\}, \{b_n^k\}, \{c_n\}, \{d_n^k\}$ be real sequences in $[0, 1]$ and satisfy the conditions:*

- (i) $a_n + c_n \leq 1, b_n^k + d_n^k \leq 1, k = 1, 2, \dots, p - 1$;
- (ii) $a_n, c_n, b_n^k, d_n^k \rightarrow 0$ as $n \rightarrow \infty, k = 1, 2, \dots, p - 1$;
- (iii) $c_n = o(a_n), \sum_{n=0}^{\infty} a_n = \infty$.

For some $x_0 \in E$, let $\{u_n\}, \{w_n^1\}, \{w_n^2\}, \dots, \{w_n^{p-1}\}$ be any bounded sequences of E , and let $\{x_n\}$ be the multistep iterative sequence with errors defined by

$$\begin{aligned}
 &x_0 \in D, \\
 &y_n^{p-1} = \left(1 - b_n^{p-1} - d_n^{p-1}\right)x_n + b_n^{p-1}Sx_n + d_n^{p-1}w_n^{p-1}, \quad n \geq 0, p \geq 2, \\
 &y_n^k = \left(1 - b_n^k - d_n^k\right)x_n + b_n^kSy_n^{k+1} + d_n^kw_n^k, \quad k = p - 2, p - 3, \dots, 2, 1, \\
 &x_{n+1} = (1 - a_n - c_n)x_n + a_nSy_n^1 + c_nu_n, \quad n \geq 0,
 \end{aligned} \tag{2.14}$$

where $S : E \rightarrow E$ is defined by $Sx = x - Tx$ for all $x \in E$. Then (2.14) converges strongly to the fixed point q of S .

Proof. We find easily that S is a uniformly continuous Φ -hemiccontractive. Then the conclusion of Theorem 2.2 is obtained directly by Theorem 2.1. \square

Remark 2.3. In Theorems 2.1 and 2.2, if $b_n^k = d_n^k = 0$ ($k = p - 1, p - 2, \dots, 2, 1$), then, the conclusions are as follows.

Corollary 2.4. *Let E be an arbitrary real Banach space, D a nonempty closed convex subset of E , and $T : D \rightarrow D$ a uniformly continuous Φ -hemiccontractive mapping with $q \in F(T) \neq \emptyset$. Let $\{a_n\}, \{c_n\}$ be real sequences in $[0, 1]$ and satisfy the conditions (i) $a_n + c_n \leq 1$; (ii) $a_n, c_n \rightarrow 0$ as $n \rightarrow \infty$; (iii) $c_n = o(a_n)$ and $\sum_{n=0}^{\infty} a_n = \infty$. For some $x_0 \in D$, let $\{u_n\}$ be any bounded sequence of D , and let $\{x_n\}$ be Mann iterative sequence with errors defined by $x_{n+1} = (1 - a_n - c_n)x_n + a_nTx_n + c_nu_n, n \geq 0$. Then $\{x_n\}$ converges strongly to the fixed point q of T .*

Corollary 2.5. *Let E be an arbitrary real Banach space and let $T : E \rightarrow E$ be a uniformly continuous Φ -quasi-accretive operator with $q \in N(T) \neq \emptyset$. Let $\{a_n\}, \{c_n\}$ be real sequences in $[0, 1]$ and satisfy the conditions (i) $a_n + c_n \leq 1$; (ii) $a_n, c_n \rightarrow 0$ as $n \rightarrow \infty$; (iii) $c_n = o(a_n)$ and $\sum_{n=0}^{\infty} a_n = \infty$. For*

some $x_0 \in E$, let $\{u_n\}$ be any bounded sequence of E , and let $\{x_n\}$ be Mann iterative sequence with errors defined by $x_{n+1} = (1 - a_n - c_n)x_n + a_n Sx_n + c_n u_n$, $n \geq 0$. where $S : E \rightarrow E$ is defined by $Sx = x - Tx$ for all $x \in E$. Then $\{x_n\}$ converges strongly to the fixed point q of S .

Remark 2.6. It is mentioned to notice that there exists a serious shortcoming in the proof process of Theorem 2.3 of [3]. That is, $M_1 c_n \leq (\Phi(\epsilon)/4)\alpha_n$ does not hold in line 15 of Claim 2 of page 552. The reason is that the conditions $\sum_{n=0}^{\infty} c_n < +\infty$ and $\sum_{n=0}^{\infty} b_n = +\infty$, $b_n \rightarrow 0$ as $n \rightarrow \infty$ can not obtain $c_n = o(b_n)$.

Counterexample, let the iteration parameters be $a_n = 1 - b_n - c_n$, b_n, c_n in the following:

$$\begin{aligned} \{b_n\} : b_0 = b_1 = 0, b_n = \frac{1}{n}, \quad n \geq 2, \\ \{c_n\} : 0, \frac{1}{1}, \frac{1}{2^2}, \frac{1}{3^2}, \frac{1}{4}, \frac{1}{5^2}, \frac{1}{6^2}, \frac{1}{7^2}, \frac{1}{8^2}, \frac{1}{9}, \frac{1}{10^2}, \frac{1}{11^2}, \frac{1}{12^2}, \frac{1}{13^2}, \frac{1}{14^2}, \frac{1}{15^2}, \frac{1}{16}, \\ \frac{1}{17^2}, \frac{1}{18^2}, \dots, \frac{1}{23^2}, \frac{1}{24^2}, \frac{1}{25}, \frac{1}{26^2}, \dots, \frac{1}{35^2}, \frac{1}{36}, \frac{1}{37^2}, \dots \end{aligned} \quad (2.15)$$

Then, $\sum_{n=0}^{\infty} b_n = +\infty$, $\sum_{n=0}^{\infty} c_n < 2 \sum_{n=1}^{\infty} (1/n^2) < +\infty$, but $c_n \neq o(b_n)$.

Application 1. Let $E = \mathbb{R}$ be a real number space with the usual norm and $D = [0, +\infty)$. Define $T : D \rightarrow D$ by

$$Tx = \frac{x^3}{1 + x^2} \quad (2.16)$$

for all $x \in D$. Then T is uniformly continuous with $F(T) = \{0\}$. Define $\Phi : [0, +\infty) \rightarrow [0, +\infty)$ by

$$\Phi(t) = \frac{t^2}{1 + t^2}, \quad (2.17)$$

then Φ is a strictly increasing function with $\Phi(0) = 0$. For all $x \in D$, $q \in F(T)$, we obtain that

$$\begin{aligned} \langle Tx - Tq, j(x - q) \rangle &= \left\langle \frac{x^3}{1 + x^2} - 0, j(x - 0) \right\rangle \\ &= \left\langle \frac{x^3}{1 + x^2}, x \right\rangle \\ &= \frac{x^4}{1 + x^2} \\ &= |x - q|^2 - \frac{|x - q|^2}{1 + |x - q|^2} \\ &= |x - q|^2 - \Phi(|x - q|). \end{aligned} \quad (2.18)$$

Therefore, T is a Φ -hemicontractive mapping. Set

$$a_n = \frac{1}{n+2}, \quad c_n = \frac{1}{(n+2)^2}, \quad b_n^k = d_n^k = \frac{1}{n+2}, \quad k = 1, 2, \dots, p-1 \quad (2.19)$$

for all $n \geq 0$.

Acknowledgment

This work is supported by Hebei Natural Science Foundation under Grant no. A2011210033.

References

- [1] B. E. Rhoades and S. M. Soltuz, "The equivalence between Mann-Ishikawa iterations and multistep iteration," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 58, no. 1-2, pp. 219–228, 2004.
- [2] Y. Xu, "Ishikawa and Mann iterative processes with errors for nonlinear strongly accretive operator equations," *Journal of Mathematical Analysis and Applications*, vol. 224, no. 1, pp. 91–101, 1998.
- [3] C. E. Chidume and C. O. Chidume, "Convergence theorems for fixed points of uniformly continuous generalized Φ -hemi-contractive mappings," *Journal of Mathematical Analysis and Applications*, vol. 303, no. 2, pp. 545–554, 2005.
- [4] K. Deimling, *Nonlinear Functional Analysis*, Springer, Berlin, Germany, 1985.
- [5] C. Moore and B. V. C. Nnoli, "Iterative solution of nonlinear equations involving set-valued uniformly accretive operators," *Computers & Mathematics with Applications*, vol. 42, no. 1-2, pp. 131–140, 2001.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

