

Research Article

The Atom-Bond Connectivity Index of Catacondensed Polyomino Graphs

Jinsong Chen,^{1,2} Jianping Liu,¹ and Qiaoliang Li²

¹ College of Mathematics and Computer Science, Fuzhou University, Fuzhou, Fujian 350108, China

² College of Mathematics and Computer Science, Hunan Normal University, Changsha, Hunan 410081, China

Correspondence should be addressed to Jinsong Chen; fzu_cjs@126.com

Received 3 January 2013; Accepted 4 February 2013

Academic Editor: Zhengkun Huan

Copyright © 2013 Jinsong Chen et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Let $G = (V, E)$ be a graph. The atom-bond connectivity (ABC) index is defined as the sum of weights $((d_u + d_v - 2)/d_u d_v)^{1/2}$ over all edges uv of G , where d_u denotes the degree of a vertex u of G . In this paper, we give the atom-bond connectivity index of the zigzag chain polyomino graphs. Meanwhile, we obtain the sharp upper bound on the atom-bond connectivity index of catacondensed polyomino graphs with h squares and determine the corresponding extremal graphs.

1. Introduction

One of the most active fields of research in contemporary chemical graph theory is the study of topological indices (graph topological invariants) that can be used for describing and predicting physicochemical and pharmacological properties of organic compounds. In chemistry and for chemical graphs, these invariant numbers are known as the topological indices. There are many publications on the topological indices, see [1–6].

Let $G = (V, E)$ be a simple graph of order n . A few years ago, Estrada et al. [7] introduced a further vertex-degree-based graph invariant, known as the atom-bond connectivity (ABC) index. It is defined as:

$$ABC(G) = \sum_{uv \in E(G)} \sqrt{\frac{d_u + d_v - 2}{d_u d_v}}. \quad (1)$$

The ABC index keeps the spirit of the Randić index, and it provides a good model for the stability of linear and branched alkanes as well as the strain energy of cycloalkanes [7]. Recently, the study of the ABC index attracts some research attention [6, 8–12].

Polyomino graphs [13], also called chessboards [14] or square-cell configurations [15] have attracted some mathematicians' considerable attention because many interesting combinatorial subjects are yielded from them such as domination problem and modeling problems of surface chemistry. A polyomino graph [16] is a connected geometric graph obtained by arranging congruent regular squares of side length 1 (called a cell) in a plane such that two squares are either disjoint or have a common edge. The polyomino graph has received considerable attentions.

Next, we introduce some graph definitions used in this paper.

Definition 1 (see [4]). Let G be a polyomino graph. If all vertices of G lie on its perimeter, then G is said to be catacondensed polyomino graph or tree-like polyomino graph. (see Figure 1).

Definition 2 (see [16]). Let G be a chain polyomino graph with h squares. If the subgraph obtained from G by deleting all the vertices of degree 2 and all the edges adjacent to the vertices is a path, then G is said to be the zigzag chain polyomino graph, denoted by Z_h (see Figure 1).

In this paper, we give the ABC indices of the zigzag chain polyomino graphs with h squares and obtain the sharp upper

bound on the ABC indices of catacondensed polyomino graphs with h squares and determine the corresponding extremal graphs.

2. The ABC Indices of Catacondensed Polyomino Graphs

Let

$$S_1 = \left\{ \sqrt{2} + \frac{2}{3}, \sqrt{2} + \frac{\sqrt{15}}{3} - \frac{2}{3}, 2 \right\},$$

$$S_2 = \left\{ 2, \frac{3\sqrt{2}}{2} + \frac{\sqrt{6}}{4} + \frac{\sqrt{15}}{3} - 2, \right.$$

$$\left. \frac{4}{3} + \frac{\sqrt{15}}{6}, \frac{2}{3} + \frac{\sqrt{15}}{3}, \sqrt{2} + \frac{\sqrt{15}}{3} - \frac{2}{3}, \right.$$

$$\left. \sqrt{2} + \frac{\sqrt{6}}{4} + \frac{\sqrt{15}}{6} - \frac{2}{3}, \sqrt{2} + \frac{\sqrt{6}}{4}, \right.$$

$$\left. \sqrt{2} + \frac{\sqrt{6}}{2} - \frac{\sqrt{15}}{6}, \sqrt{2} + \frac{\sqrt{6}}{4} + \frac{2}{3}, \right.$$

$$\left. \frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{4} + \frac{\sqrt{15}}{6}, \frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{2} + \frac{2}{3} - \frac{\sqrt{15}}{6}, \right.$$

$$\left. \frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{4} + \frac{\sqrt{15}}{3} - \frac{2}{3}, \frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{2}, \right.$$

$$\left. \frac{3\sqrt{2}}{2} + \frac{\sqrt{6}}{4} + \frac{\sqrt{15}}{2} - \frac{8}{3}, \frac{3\sqrt{2}}{2} + \frac{2\sqrt{6}}{4} + \frac{\sqrt{15}}{6} - 2, \right.$$

$$\left. \frac{3\sqrt{2}}{2} + \frac{\sqrt{6}}{4} + \frac{4\sqrt{15}}{6} - \frac{10}{3}, \right.$$

$$\left. \frac{3\sqrt{2}}{2} + \frac{\sqrt{6}}{2} + \frac{2\sqrt{15}}{6} - \frac{8}{3}, \frac{3\sqrt{2}}{2} + \frac{3\sqrt{6}}{4} - 2, \right.$$

$$\left. \frac{3\sqrt{2}}{2} + \frac{3\sqrt{6}}{4} - \frac{\sqrt{15}}{3} - \frac{2}{3}, \frac{3\sqrt{2}}{2} + \frac{3\sqrt{6}}{4} - \frac{\sqrt{15}}{6} - \frac{4}{3}, \right.$$

$$\left. \frac{3\sqrt{2}}{2} + \sqrt{6} - \frac{\sqrt{15}}{2} - \frac{2}{3}, \frac{3\sqrt{2}}{2} + \sqrt{6} - \frac{2\sqrt{15}}{6} - \frac{4}{3}, \right.$$

$$\left. \frac{3\sqrt{2}}{2} + \frac{5\sqrt{6}}{4} - \frac{2\sqrt{15}}{3} - \frac{2}{3}, \right.$$

$$\left. \frac{3\sqrt{2}}{2} + \frac{\sqrt{6}}{2} - \frac{4}{3}, \frac{3\sqrt{2}}{2} + \frac{3\sqrt{6}}{4} + \frac{\sqrt{15}}{6} - \frac{8}{3} \right\},$$

$$S = S_1 \cup S_2. \tag{2}$$

We call $\sqrt{(d_u + d_v - 2)/d_u d_v}$ the weight of the edge uv , denoted by W_{uv} .

Note that for any catacondensed polyomino graph H^* with h squares, it can be obtained by gluing a new square s to some catacondensed polyomino graph H with $h - 1$ squares. So, we have the following lemma.




FIGURE 1: The zigzag chain polyomino graph Z_h .




FIGURE 2




FIGURE 3




FIGURE 4

Lemma 3. Let H^* be a catacondensed polyomino graph with h squares which is obtained by gluing a new square s to some graph H , where H is a catacondensed polyomino graph with $h - 1$ squares. One has

- (i) If $2 \leq h \leq 3$, then $ABC(H^*) - ABC(H) \in S_1$,
- (ii) if $h \geq 4$, then $ABC(H^*) - ABC(H) \in S_2$.




FIGURE 5




FIGURE 6: A catacondensed polyomino graph with h ($h \leq 3$) squares.

Proof. Consider the following: (i) if $2 \leq h \leq 3$, by directly calculating, we have $ABC(H^*) - ABC(H) \in S_1$,

(ii) now, let $h \geq 4$. Without the loss of generality, let square s be adjacent to the edge AB in H (see Figure 2). In the following, if the weights of some edges of H have been changed when s is adjacent to the edge AB in H , then we marked these edges with thick lines in H^* . Let $D_i = ABC(H^*) - ABC(H)$ ($i = 1, 2, \dots, 35$). Note that except the edge AB of s , the summation of the weights of the remaining three edges is always $(3/2)\sqrt{2}$ in H^* . There are exactly three types of formations (see Figure 2).

Case 1. In Type I, $d_{A_1} = d_{B_1} = 2$ and $d_{A_2} = d_{B_2} = 3$ (see Figure 3).

By the definition of ABC index, we have $ABC(H^*) - ABC(H) = (3/2)\sqrt{2} + (W_{uA_2} - W_{uA_1}) + (W_{A_2B_2} - W_{A_1B_1}) + (W_{vB_2} - W_{vB_1}) = D_i$ ($i = 1, 2, 3$).

If $d_u = 3$ and $d_v = 3$, then $D_1 = 2$.

If $d_u = 3$ and $d_v = 4$ or $d_u = 4$ and $d_v = 3$, then $D_2 = 4/3 + \sqrt{15}/6$.

If $d_u = 4$ and $d_v = 4$, then $D_3 = 2/3 + \sqrt{15}/3$.

Case 2. In Type II, $d_{A_1} = 2$, $d_{B_1} = 3$, $d_{A_2} = 3$, and $d_{B_2} = 4$. (see Figure 4).

Let u adjacent to A and v, w adjacent to B (see Figure 4). Then $d_u \in \{2, 3, 4\}$, $d_v \in \{3, 4\}$, and $d_w \in \{2, 3, 4\}$. If $d_u = 2$, $d_v = 3$, and $d_w = 2$, which is in contradiction with $h \geq 4$; if $d_u = 3$ and $d_v = 3$, which is in contradiction with $d_{B_1} = 3$; if $d_u = 4$ and $d_v = 3$, which is in contradiction with $d_{A_1} = 2$ ($A \in V(H)$).

By the definition of ABC index, we have $ABC(H^*) - ABC(H) = (3/2)\sqrt{2} + (W_{uA_2} - W_{uA_1}) + (W_{A_2B_2} - W_{A_1B_1}) + (W_{vB_2} - W_{vB_1}) + (W_{wB_2} - W_{wB_1}) = D_i$ ($i = 4, 5, \dots, 14$).

If $d_u = 2$, $d_v = 3$, and $d_w = 3$, then $D_4 = \sqrt{2} + \sqrt{15}/2 - 4/3$.

If $d_u = 2$, $d_v = 3$, and $d_w = 4$, then $D_5 = \sqrt{2} + \sqrt{6}/4 + \sqrt{15}/6 - 2/3$.

If $d_u = 2$, $d_v = 4$, and $d_w = 2$, then $D_6 = \sqrt{2} + \sqrt{6}/4$.

If $d_u = 2$, $d_v = 4$, and $d_w = 3$, then $D_7 = \sqrt{2} + \sqrt{6}/4 + \sqrt{15}/6 - 2/3$.

If $d_u = 2$, $d_v = 4$, and $d_w = 4$, then $D_8 = \sqrt{2} + \sqrt{6}/2 - \sqrt{15}/6$.

If $d_u = 3$, $d_v = 4$, and $d_w = 2$, then $D_9 = \sqrt{2}/2 + \sqrt{6}/4 + 2/3$.

If $d_u = 3$, $d_v = 4$, and $d_w = 3$, then $D_{10} = \sqrt{2}/2 + \sqrt{6}/4 + \sqrt{5}/12$.

If $d_u = 3$, $d_v = 4$, and $d_w = 4$, then $D_{11} = \sqrt{2}/2 + \sqrt{6}/2 + 2/3 - \sqrt{5}/12$.

If $d_u = 4$, $d_v = 4$, and $d_w = 2$, then $D_{12} = \sqrt{2}/2 + \sqrt{6}/4 + \sqrt{5}/12$.

If $d_u = 4$, $d_v = 4$, and $d_w = 3$, then $D_{13} = \sqrt{2}/2 + \sqrt{6}/4 + 2\sqrt{5}/12 - 2/3$.

If $d_u = 4$, $d_v = 4$, and $d_w = 4$, then $D_{14} = \sqrt{2}/2 + \sqrt{6}/2$.

Case 3. In Type III, $d_{A_1} = d_{B_1} = 3$ and $d_{A_2} = d_{B_2} = 4$ (see Figure 5).

Let u, x adjacent to A and v, w adjacent to B (see Figure 5). Then, $d_u \in \{3, 4\}$, $d_v \in \{3, 4\}$, $d_w \in \{2, 3, 4\}$, and $d_x \in \{2, 3, 4\}$. Since the case $d_u = 3$, $d_v = 4$, $d_x = y_1$, and $d_w = y_2$ is the same as $d_u = 4$, $d_v = 3$, $d_x = y_2$, and $d_w = y_1$, where $y_1, y_2 \in \{2, 3, 4\}$. And note that if $d_u = d_v = 3$ or $d_u = d_v = 4$, the vertices x and w are symmetric.

By the definition of ABC index, we have $ABC(H^*) - ABC(H) = (3/2)\sqrt{2} + (W_{xA_2} - W_{xA_1}) + (W_{uA_2} - W_{uA_1}) + (W_{A_2B_2} - W_{A_1B_1}) + (W_{vB_2} - W_{vB_1}) + (W_{wB_2} - W_{wB_1}) = D_i$ ($i = 15, 16, \dots, 35$).

If $d_u = d_v = 3$, $d_x = 2$, and $d_w = 3$, then $D_{15} = 3\sqrt{2}/2 + \sqrt{6}/4 + \sqrt{15}/2 - 8/3$.

If $d_u = d_v = 3$, $d_x = 2$, and $d_w = 4$, then $D_{16} = 3\sqrt{2}/2 + \sqrt{6}/2 + \sqrt{15}/6 - 2$.

If $d_u = d_v = 3$, $d_x = 3$, and $d_w = 3$, then $D_{17} = 3\sqrt{2}/2 + \sqrt{6}/4 + 2\sqrt{15}/3 - 10/3$.

If $d_u = d_v = 3$, $d_x = 3$, and $d_w = 4$, then $D_{18} = 3\sqrt{2}/2 + \sqrt{6}/2 + \sqrt{15}/3 - 8/3$.

If $d_u = d_v = 3$, $d_x = 4$, and $d_w = 4$, then $D_{19} = 3\sqrt{2}/2 + 3\sqrt{6}/4 - 2$.

If $d_u = d_v = 4$, $d_x = 2$, and $d_w = 2$, then $D_{20} = 3\sqrt{2}/2 + 3\sqrt{6}/4 - \sqrt{15}/3 - 2/3$.

If $d_u = d_v = 4$, $d_x = 2$, and $d_w = 3$, then $D_{21} = 3\sqrt{2}/2 + 3\sqrt{6}/4 - \sqrt{15}/6 - 4/3$.

If $d_u = d_v = 4$, $d_x = 2$, and $d_w = 4$, then $D_{22} = 3\sqrt{2}/2 + \sqrt{6} - \sqrt{15}/2 - 2/3$.

If $d_u = d_v = 4$, $d_x = 3$, and $d_w = 3$, then $D_{23} = 3\sqrt{2}/2 + 3\sqrt{6}/4 - 2$.

If $d_u = d_v = 4$, $d_x = 3$, and $d_w = 4$, then $D_{24} = 3\sqrt{2}/2 + \sqrt{6} - \sqrt{15}/3 - 4/3$.

If $d_u = d_v = 4$, $d_x = 4$, and $d_w = 4$, then $D_{25} = 3\sqrt{2}/2 + 5\sqrt{6}/4 - 2\sqrt{15}/3 - 2/3$.

If $d_u = 3$, $d_v = 4$, $d_x = 2$, and $d_w = 2$, then $D_{26} = 3\sqrt{2}/2 + \sqrt{6}/2 - 4/3$.

If $d_u = 3$, $d_v = 4$, $d_x = 2$, and $d_w = 3$, then $D_{27} = 3\sqrt{2}/2 + \sqrt{6}/2 + \sqrt{15}/6 - 2$.

If $d_u = 3$, $d_v = 4$, $d_x = 2$, and $d_w = 4$, then $D_{28} = 3\sqrt{2}/2 + 3\sqrt{6}/4 - \sqrt{15}/6 - 4/3$.

If $d_u = 3$, $d_v = 4$, $d_x = 3$, and $d_w = 2$, then $D_{29} = 3\sqrt{2}/2 + \sqrt{6}/2 + \sqrt{15}/6 - 2$.

If $d_u = 3$, $d_v = 4$, $d_x = 3$, and $d_w = 3$, then $D_{30} = 3\sqrt{2}/2 + \sqrt{6}/2 + \sqrt{15}/3 - 8/3$.

If $d_u = 3$, $d_v = 4$, $d_x = 3$, and $d_w = 4$, then $D_{31} = 3\sqrt{2}/2 + 3\sqrt{6}/4 - 2$.

If $d_u = 3$, $d_v = 4$, $d_x = 4$, and $d_w = 2$, then $D_{32} = 3\sqrt{2}/2 + 3\sqrt{6}/4 - \sqrt{15}/6 - 4/3$.

If $d_u = 3$, $d_v = 4$, $d_x = 4$, and $d_w = 3$, then $D_{33} = 3\sqrt{2}/2 + 3\sqrt{6}/4 - 2$.

If $d_u = 3$, $d_v = 4$, $d_x = 4$, and $d_w = 4$, then $D_{34} = 3\sqrt{2}/2 + \sqrt{6} - \sqrt{15}/3 - 4/3$.

If $d_u = 3$, $d_v = 3$, $d_x = 2$, and $d_w = 2$, then $D_{35} = 3\sqrt{2}/2 + \sqrt{6}/4 + \sqrt{15}/3 - 2$.

By directly calculating, we have $D_5 = D_7$, $D_{10} = D_{12}$, $D_{16} = D_{27} = D_{29}$, $D_{18} = D_{30}$, $D_{19} = D_{23} = D_{31} = D_{33}$, $D_{21} = D_{28} = D_{32}$, $D_{24} = D_{34}$, and $D_6 = \max_{1 \leq i \leq 35} D_i$, $D_{14} = \min_{1 \leq i \leq 35} D_i$. So $ABC(H^*) - ABC(H) \in S_2$, where $h \geq 4$.

Therefore, $ABC(H^*) - ABC(H) \in S$. □

By Lemma 3, we have the following theorem.

Theorem 4. Let G be a catacondensed polyomino graph with h ($h \geq 2$) squares, then

$$\begin{aligned}
 ABC(G) &= 3\sqrt{2} + \frac{2}{3} + \left(\sqrt{2} + \frac{\sqrt{15}}{3} - \frac{2}{3} \right) a_1 \\
 &+ 2a_2 + \left(\frac{4}{3} + \frac{\sqrt{15}}{6} \right) a_3 \\
 &+ \left(\frac{2}{3} + \frac{\sqrt{15}}{3} \right) a_4 \\
 &+ \left(\sqrt{2} + \frac{\sqrt{15}}{2} - \frac{4}{3} \right) a_5 \\
 &+ \left(\sqrt{2} + \frac{\sqrt{6}}{4} + \frac{\sqrt{15}}{6} - \frac{2}{3} \right) a_6 \\
 &+ \left(\sqrt{2} + \frac{\sqrt{6}}{4} \right) a_7 \\
 &+ \left(\sqrt{2} + \frac{\sqrt{6}}{2} - \frac{\sqrt{15}}{6} \right) a_8 \\
 &+ \left(\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{4} + \frac{2}{3} \right) a_9 \\
 &+ \left(\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{4} + \frac{\sqrt{15}}{6} \right) a_{10} \\
 &+ \left(\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{2} + \frac{2}{3} - \frac{\sqrt{15}}{6} \right) a_{11} \\
 &+ \left(\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{4} + \frac{\sqrt{15}}{3} - \frac{2}{3} \right) a_{12} \\
 &+ \left(\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{2} \right) a_{13} \\
 &+ \left(\frac{3\sqrt{2}}{2} + \frac{\sqrt{6}}{4} + \frac{\sqrt{15}}{2} - \frac{8}{3} \right) a_{14} \\
 &+ \left(\frac{3\sqrt{2}}{2} + \frac{\sqrt{6}}{2} + \frac{\sqrt{15}}{6} - 2 \right) a_{15} \\
 &+ \left(\frac{3\sqrt{2}}{2} + \frac{\sqrt{6}}{4} + \frac{2\sqrt{15}}{3} - \frac{10}{3} \right) a_{16} \\
 &+ \left(\frac{3\sqrt{2}}{2} + \frac{\sqrt{6}}{2} + \frac{\sqrt{15}}{3} - \frac{8}{3} \right) a_{17} \\
 &+ \left(\frac{3\sqrt{2}}{2} + \frac{3\sqrt{6}}{4} - 2 \right) a_{18} \\
 &+ \left(\frac{3\sqrt{2}}{2} + \frac{3\sqrt{6}}{4} - \frac{\sqrt{15}}{3} - \frac{2}{3} \right) a_{19}
 \end{aligned}$$

$$\begin{aligned}
 & + \left(\frac{3\sqrt{2}}{2} + \frac{3\sqrt{6}}{4} - \frac{\sqrt{15}}{6} - \frac{4}{3} \right) a_{20} \\
 & + \left(\frac{3\sqrt{2}}{2} + \sqrt{6} - \frac{\sqrt{15}}{2} - \frac{2}{3} \right) a_{21} \\
 & + \left(\frac{3\sqrt{2}}{2} + \sqrt{6} - \frac{\sqrt{15}}{3} - \frac{4}{3} \right) a_{22} \\
 & + \left(\frac{3\sqrt{2}}{2} + \frac{5\sqrt{6}}{4} - \frac{2\sqrt{15}}{3} - \frac{2}{3} \right) a_{23} \\
 & + \left(\frac{3\sqrt{2}}{2} + \frac{\sqrt{6}}{2} - \frac{4}{3} \right) a_{24} \\
 & + \left(\frac{3\sqrt{2}}{2} + \frac{\sqrt{6}}{4} + \frac{\sqrt{15}}{3} - 2 \right) a_{25},
 \end{aligned} \tag{3}$$

where a_i is a nonnegative integer for $i = 1, 2, \dots, 25$ and $h = 2 + \sum_{i=1}^{25} a_i$.

Proof. We prove Theorem 4 by the induction on h . If $h = 2$, by directly calculating, we have $ABC(G) = 3\sqrt{2} + 2/3$, where $a_i = 0$ ($i = 1, 2, \dots, 25$). So, Theorem 4 holds for $h = 2$.

Assume that Theorem 4 holds for all catacondensed polyomino graphs with $h - 1$ ($h - 1 \geq 2$) squares, that is,

$$\begin{aligned}
 ABC(G) = & 3\sqrt{2} + \frac{2}{3} + \left(\sqrt{2} + \frac{\sqrt{15}}{3} - \frac{2}{3} \right) a_1 \\
 & + 2a_2 + \left(\frac{4}{3} + \frac{\sqrt{15}}{6} \right) a_3 \\
 & + \left(\frac{2}{3} + \frac{\sqrt{15}}{3} \right) a_4 \\
 & + \left(\sqrt{2} + \frac{\sqrt{15}}{2} - \frac{4}{3} \right) a_5 \\
 & + \left(\sqrt{2} + \frac{\sqrt{6}}{4} + \frac{\sqrt{15}}{6} - \frac{2}{3} \right) a_6 \\
 & + \left(\sqrt{2} + \frac{\sqrt{6}}{4} \right) a_7 \\
 & + \left(\sqrt{2} + \frac{\sqrt{6}}{2} - \frac{\sqrt{15}}{6} \right) a_8 \\
 & + \left(\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{4} + \frac{2}{3} \right) a_9 \\
 & + \left(\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{4} + \frac{\sqrt{15}}{6} \right) a_{10} \\
 & + \left(\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{2} + \frac{2}{3} - \frac{\sqrt{15}}{6} \right) a_{11} \\
 & + \left(\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{4} + \frac{\sqrt{15}}{3} - \frac{2}{3} \right) a_{12} \\
 & + \left(\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{2} \right) a_{13} \\
 & + \left(\frac{3\sqrt{2}}{2} + \frac{\sqrt{6}}{4} + \frac{\sqrt{15}}{2} - \frac{8}{3} \right) a_{14} \\
 & + \left(\frac{3\sqrt{2}}{2} + \frac{\sqrt{6}}{2} + \frac{\sqrt{15}}{6} - 2 \right) a_{15} \\
 & + \left(\frac{3\sqrt{2}}{2} + \frac{\sqrt{6}}{4} + \frac{2\sqrt{15}}{3} - \frac{10}{3} \right) a_{16} \\
 & + \left(\frac{3\sqrt{2}}{2} + \frac{\sqrt{6}}{2} + \frac{\sqrt{15}}{3} - \frac{8}{3} \right) a_{17} \\
 & + \left(\frac{3\sqrt{2}}{2} + \frac{3\sqrt{6}}{4} - 2 \right) a_{18} \\
 & + \left(\frac{3\sqrt{2}}{2} + \frac{3\sqrt{6}}{4} - \frac{\sqrt{15}}{3} - \frac{2}{3} \right) a_{19} \\
 & + \left(\frac{3\sqrt{2}}{2} + \frac{3\sqrt{6}}{4} - \frac{\sqrt{15}}{6} - \frac{4}{3} \right) a_{20} \\
 & + \left(\frac{3\sqrt{2}}{2} + \sqrt{6} - \frac{\sqrt{15}}{2} - \frac{2}{3} \right) a_{21} \\
 & + \left(\frac{3\sqrt{2}}{2} + \sqrt{6} - \frac{\sqrt{15}}{3} - \frac{4}{3} \right) a_{22} \\
 & + \left(\frac{3\sqrt{2}}{2} + \frac{5\sqrt{6}}{4} - \frac{2\sqrt{15}}{3} - \frac{2}{3} \right) a_{23} \\
 & + \left(\frac{3\sqrt{2}}{2} + \frac{\sqrt{6}}{2} - \frac{4}{3} \right) a_{24} \\
 & + \left(\frac{3\sqrt{2}}{2} + \frac{\sqrt{6}}{4} + \frac{\sqrt{15}}{3} - 2 \right) a_{25},
 \end{aligned} \tag{4}$$

where a_i is a nonnegative integer for $i = 1, 2, \dots, 25$ and $h-1 = 2 + \sum_{i=1}^{25} a_i$.

We will prove that Theorem 4 holds for h in the following. Let G^* be a catacondensed polyomino graph with h squares. Without the loss of generality, G^* can be obtained from some catacondensed polyomino graph G with $h - 1$ squares by gluing a new square s to G . By Lemma 3, we have $ABC(G^*) - ABC(G) \in S$. It means that $ABC(G^*) = ABC(G) + a$, where

$a \in S$. By the induction assumption and direct computation, we have

$$\begin{aligned}
ABC(G^*) &= 3\sqrt{2} + \frac{2}{3} + \left(\sqrt{2} + \frac{\sqrt{15}}{3} - \frac{2}{3}\right)a_1^* \\
&+ 2a_2^* + \left(\frac{4}{3} + \frac{\sqrt{15}}{6}\right)a_3^* \\
&+ \left(\frac{2}{3} + \frac{\sqrt{15}}{3}\right)a_4^* \\
&+ \left(\sqrt{2} + \frac{\sqrt{15}}{2} - \frac{4}{3}\right)a_5^* \\
&+ \left(\sqrt{2} + \frac{\sqrt{6}}{4} + \frac{\sqrt{15}}{6} - \frac{2}{3}\right)a_6^* \\
&+ \left(\sqrt{2} + \frac{\sqrt{6}}{4}\right)a_7^* \\
&+ \left(\sqrt{2} + \frac{\sqrt{6}}{2} - \frac{\sqrt{15}}{6}\right)a_8^* \\
&+ \left(\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{4} + \frac{2}{3}\right)a_9^* \\
&+ \left(\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{4} + \frac{\sqrt{15}}{6}\right)a_{10}^* \\
&+ \left(\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{2} + \frac{2}{3} - \frac{\sqrt{15}}{6}\right)a_{11}^* \\
&+ \left(\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{4} + \frac{\sqrt{15}}{3} - \frac{2}{3}\right)a_{12}^* \\
&+ \left(\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{2}\right)a_{13}^* \\
&+ \left(\frac{3\sqrt{2}}{2} + \frac{\sqrt{6}}{4} + \frac{\sqrt{15}}{2} - \frac{8}{3}\right)a_{14}^* \\
&+ \left(\frac{3\sqrt{2}}{2} + \frac{\sqrt{6}}{2} + \frac{\sqrt{15}}{6} - 2\right)a_{15}^* \\
&+ \left(\frac{3\sqrt{2}}{2} + \frac{\sqrt{6}}{4} + \frac{2\sqrt{15}}{3} - \frac{10}{3}\right)a_{16}^* \\
&+ \left(\frac{3\sqrt{2}}{2} + \frac{\sqrt{6}}{2} + \frac{\sqrt{15}}{3} - \frac{8}{3}\right)a_{17}^* \\
&+ \left(\frac{3\sqrt{2}}{2} + \frac{3\sqrt{6}}{4} - 2\right)a_{18}^* \\
&+ \left(\frac{3\sqrt{2}}{2} + \frac{3\sqrt{6}}{4} - \frac{\sqrt{15}}{3} - \frac{2}{3}\right)a_{19}^*
\end{aligned}$$

$$\begin{aligned}
&+ \left(\frac{3\sqrt{2}}{2} + \frac{3\sqrt{6}}{4} - \frac{\sqrt{15}}{6} - \frac{4}{3}\right)a_{20}^* \\
&+ \left(\frac{3\sqrt{2}}{2} + \sqrt{6} - \frac{\sqrt{15}}{2} - \frac{2}{3}\right)a_{21}^* \\
&+ \left(\frac{3\sqrt{2}}{2} + \sqrt{6} - \frac{\sqrt{15}}{3} - \frac{4}{3}\right)a_{22}^* \\
&+ \left(\frac{3\sqrt{2}}{2} + \frac{5\sqrt{6}}{4} - \frac{2\sqrt{15}}{3} - \frac{2}{3}\right)a_{23}^* \\
&+ \left(\frac{3\sqrt{2}}{2} + \frac{\sqrt{6}}{2} - \frac{4}{3}\right)a_{24}^* \\
&+ \left(\frac{3\sqrt{2}}{2} + \frac{\sqrt{6}}{4} + \frac{\sqrt{15}}{3} - 2\right)a_{25}^*.
\end{aligned} \tag{5}$$

There exists some $l \in \{1, 2, \dots, 25\}$ such that $a_l^* = a_l + 1$ and $a_j^* = a_j$ for $j \neq l$ ($j \in \{1, 2, \dots, 25\}$). Obviously, a_i^* is a nonnegative integer for $i = 1, 2, \dots, 25$ and $2 + \sum_{i=1}^{25} a_i^* = 2 + 1 + \sum_{i=1}^{25} a_i = h$. \square

Lemma 5. *Let H be a catacondensed polyomino graph with h squares. If $h \leq 3$, there are exactly four nonisomorphism catacondensed polyomino graphs (see Figure 6), where $ABC(H_1) = 2\sqrt{2}$, $ABC(H_2) = 3\sqrt{2} + 2/3$, $ABC(H_3) = 3\sqrt{2} + 8/3$, $ABC(H_4) = 4\sqrt{2} + \sqrt{15}/3$.*

Theorem 6. *Let Z_h be a zigzag chain polyomino graph with h squares, then*

$$ABC(Z_h) = \begin{cases} 2\sqrt{2}, & h = 1, \\ 3\sqrt{2} + \frac{2}{3}, & h = 2, \\ (h+1)\sqrt{2} + (h-3) \cdot \frac{\sqrt{6}}{4} + \frac{\sqrt{15}}{3}, & h \geq 3. \end{cases} \tag{6}$$

Proof. Obviously, Z_h can be obtained by gluing a new square s_h to Z_{h-1} . Let s_{h-1} be the square adjacent to s_h (see Figure 1). We will prove Theorem 6 by the induction on h .

If $h = 1, 2, 3$, then Theorem 6 holds (by Lemma 5). Assume that $ABC(Z_{h-1}) = (h-1+1)\sqrt{2} + (h-1-3) \cdot (\sqrt{6}/4) + (\sqrt{15}/3) = h\sqrt{2} + (h-4) \cdot (\sqrt{6}/4) + (\sqrt{15}/3)$ for $h-1 \geq 3$. By the induction assumption and the D_6 in Lemma 3, we have

$$\begin{aligned}
ABC(Z_h) &= ABC(Z_{h-1}) + D_6 \\
&= h\sqrt{2} + (h-4) \cdot \frac{\sqrt{6}}{4} + \frac{\sqrt{15}}{3} + \left(\sqrt{2} + \frac{\sqrt{6}}{4}\right) \\
&= (h+1)\sqrt{2} + (h-3) \cdot \frac{\sqrt{6}}{4} + \frac{\sqrt{15}}{3}.
\end{aligned} \tag{7}$$

So, Theorem 6 holds. \square

Note that $D_6 = \max_{1 \leq i \leq 35} D_i$ for $h \geq 4$ and by Lemma 5, we obtain the following Theorem 7.

Theorem 7. *Let G be a catacondensed polyomino graph with h squares, then $ABC(G) \leq (h+1)\sqrt{2} + (h-3) \cdot (\sqrt{6}/4) + (\sqrt{15}/3)$, with the equality if and only if $G \cong Z_h$.*

Acknowledgments

The project was supported by NSFC (no. 11071272; 11101086; 11101087) and the Foundation of Fuzhou University (no. 2012-XQ-30).

References

- [1] A. Behtoei, M. Jannesari, and B. Taeri, "Maximum Zagreb index, minimum hyper-Wiener index and graph connectivity," *Applied Mathematics Letters*, vol. 22, no. 10, pp. 1571–1576, 2009.
- [2] K. C. Das, I. Gutman, and B. Furtula, "Survey on geometric-arithmetic indices of graphs," *MATCH Communications in Mathematical and in Computer Chemistry*, vol. 65, no. 3, pp. 595–644, 2011.
- [3] B. Liu and I. Gutman, "On general Randić indices," *MATCH Communications in Mathematical and in Computer Chemistry*, vol. 58, no. 1, pp. 147–154, 2007.
- [4] B. Liu and Z. You, "A survey on comparing Zagreb indices," *MATCH Communications in Mathematical and in Computer Chemistry*, vol. 65, no. 3, pp. 581–593, 2011.
- [5] J. Liu and B. Liu, "E-L equienergetic graphs," *Communications in Mathematical and in Computer Chemistry*, vol. 66, no. 3, pp. 971–976, 2011.
- [6] B. Zhou and N. Trinajstić, "Further results on atom-bond connectivity index of trees," *Chemical Physics Letters*, vol. 455, pp. 120–123, 2008.
- [7] E. Estrada, L. Torres, L. Rodríguez, and I. Gutman, "An atom-bond connectivity index: modelling the enthalpy of formation of alkanes," *Indian Journal of Chemistry A*, vol. 37, pp. 849–855, 1998.
- [8] J. Chen and X. Guo, "Extreme atom-bond connectivity index of graphs," *MATCH Communications in Mathematical and in Computer Chemistry*, vol. 65, no. 3, pp. 713–722, 2011.
- [9] J. Chen and X. Guo, "The atom-bond connectivity index of chemical bicyclic graphs," *Applied Mathematics B*, vol. 27, no. 2, pp. 243–252, 2012.
- [10] J. Chen, J. Liu, and X. Guo, "Some upper bounds for the atom-bond connectivity index of graphs," *Applied Mathematics Letters*, vol. 25, no. 7, pp. 1077–1081, 2012.
- [11] B. Furtula, A. Graovac, and D. Vukičević, "Atom-bond connectivity index of trees," *Discrete Applied Mathematics*, vol. 157, no. 13, pp. 2828–2835, 2009.
- [12] R. Xing, B. Zhou, and Z. Du, "Further results on atom-bond connectivity index of trees," *Discrete Applied Mathematics*, vol. 158, no. 14, pp. 1536–1545, 2010.
- [13] C. Berge, C. C. Chen, V. Chvátal, and C. S. Seow, "Combinatorial properties of polyominoes," *Combinatorica*, vol. 1, no. 3, pp. 217–224, 1981.
- [14] E. J. Cockayne, "Chessboard domination problems," *Discrete Mathematics*, vol. 86, no. 1–3, pp. 13–20, 1990.
- [15] F. Harary and P. Mezey, "Cell-shedding transformations, equivalence relations, and similarity measures for square-cell configurations," *International Journal of Quantum Chemistry*, vol. 62, pp. 353–361, 1997.
- [16] R. Chen, "Perfect matchings of generalized polyomino graphs," *Graphs and Combinatorics*, vol. 21, no. 4, pp. 515–529, 2005.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

