

Research Article

On the Neutrix Composition of the Delta and Inverse Hyperbolic Sine Functions

Brian Fisher¹ and Adem Kılıçman²

¹ Department of Mathematics, University of Leicester, Leicester LE1 7RH, UK

² Department of Mathematics and Institute for Mathematical Research, University Putra Malaysia, 43400 UPM, Serdang, Selangor, Malaysia

Correspondence should be addressed to Adem Kılıçman, akilicman@putra.upm.edu.my

Received 24 December 2010; Accepted 8 April 2011

Academic Editor: A. A. Soliman

Copyright © 2011 B. Fisher and A. Kılıçman. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Let F be a distribution in \mathfrak{D}' and let f be a locally summable function. The composition $F(f(x))$ of F and f is said to exist and be equal to the distribution $h(x)$ if the limit of the sequence $\{F_n(f(x))\}$ is equal to $h(x)$, where $F_n(x) = F(x) * \delta_n(x)$ for $n = 1, 2, \dots$ and $\{\delta_n(x)\}$ is a certain regular sequence converging to the Dirac delta function. In the ordinary sense, the composition $\delta^{(s)}[(\sinh^{-1}x_+)^r]$ does not exist. In this study, it is proved that the neutrix composition $\delta^{(s)}[(\sinh^{-1}x_+)^r]$ exists and is given by $\delta^{(s)}[(\sinh^{-1}x_+)^r] = \sum_{k=0}^{sr+r-1} \sum_{i=0}^k \binom{k}{i} (-1)^k r c_{s,k,i} / 2^{k+1} k! \delta^{(k)}(x)$, for $s = 0, 1, 2, \dots$ and $r = 1, 2, \dots$, where $c_{s,k,i} = (-1)^s s! [(k-2i+1)^{rs-1} + (k-2i-1)^{rs+r-1}] / (2(rs+r-1)!)$. Further results are also proved.

1. Introduction

In the following, we let \mathfrak{D} be the space of infinitely differentiable functions with compact support, let $\mathfrak{D}[a, b]$ be the space of infinitely differentiable functions with support contained in the interval $[a, b]$, and let \mathfrak{D}' be the space of distributions defined on \mathfrak{D} .

Now, let $\rho(x)$ be a function in $\mathfrak{D}[-1, 1]$ having the following properties:

- (i) $\rho(x) \geq 0$,
- (ii) $\rho(x) = \rho(-x)$,
- (iii) $\int_{-1}^1 \rho(x) dx = 1$.

Putting $\delta_n(x) = n\rho(nx)$ for $n = 1, 2, \dots$, it follows that $\{\delta_n(x)\}$ is a regular sequence of infinitely differentiable functions converging to the Dirac delta-function $\delta(x)$. Further, if F is

an arbitrary distribution in \mathfrak{D}' and $F_n(x) = F(x) * \delta_n(x) = \langle F(x-t), \varphi(t) \rangle$, then $\{F_n(x)\}$ is a regular sequence converging to $F(x)$.

Since the theory of distributions is a linear theory, thus we can extend some of the operations which are valid for ordinary functions to the space of distributions and such operations are called regular operations such as: addition, multiplication by scalars; see [1]. Other operations can be defined only for a particular class of distributions or for certain restricted subclasses of distributions; these are called irregular operations such as: multiplication of distributions, convolution products, and composition of distributions; see [2-4]. Thus, there have been several attempts recently to define distributions of the form $F(f(x))$ in \mathfrak{D}' , where F and f are distributions in \mathfrak{D}' ; see for example [5-8]. In the following, we are going to consider an alternative approach. As a starting point, we look at the following definition which is a generalization of Gel'fand and Shilov's definition of the composition involving the delta function [9], and was given in [6].

Definition 1.1. Let F be a distribution in \mathfrak{D}' and let f be a locally summable function. We say that the neutrix composition $F(f(x))$ exists and is equal to h on the open interval (a, b) , with $-\infty < a < b < \infty$, if

$$N - \lim_{n \rightarrow \infty} \int_{-\infty}^{\infty} F_n(f(x)) \varphi(x) dx = \langle h(x), \varphi(x) \rangle, \quad (1.1)$$

for all φ in $\mathfrak{D}[a, b]$, where $F_n(x) = F(x) * \delta_n(x)$ for $n = 1, 2, \dots$ and N is the neutrix, see [10], having domain N' the positive and range N'' the real numbers, with negligible functions which are finite linear sums of the functions

$$n^\lambda \ln^{r-1} n, \quad \ln^r n : \lambda > 0, \quad r = 1, 2, \dots \quad (1.2)$$

and all functions which converge to zero in the usual sense as n tends to infinity.

In particular, we say that the composition $F(f(x))$ exists and is equal to h on the open interval (a, b) if

$$\lim_{n \rightarrow \infty} \int_{-\infty}^{\infty} F_n(f(x)) \varphi(x) dx = \langle h(x), \varphi(x) \rangle, \quad (1.3)$$

for all φ in $\mathfrak{D}[a, b]$.

Note that taking the neutrix limit of a function $f(n)$ is equivalent to taking the usual limit of Hadamard's finite part of $f(n)$. The definition of the neutrix composition of distributions was originally given in [10] but was then simply called the composition of distributions.

The following three theorems were proved in [11], [8], and [12], respectively.

Theorem 1.2. *The neutrix composition $\delta^{(s)}(\operatorname{sgn} x |x|^\lambda)$ exists and*

$$\delta^{(s)}(\operatorname{sgn} x |x|^\lambda) = 0, \quad (1.4)$$

for $s = 0, 1, 2, \dots$ and $(s + 1)\lambda = 1, 3, \dots$, and

$$\delta^{(s)}(\operatorname{sgn} x |x|^\lambda) = \frac{(-1)^{(s+1)(\lambda+1)} s!}{\lambda[(s+1)\lambda - 1]!} \delta^{((s+1)\lambda-1)}(x), \quad (1.5)$$

for $s = 0, 1, 2, \dots$, and $(s + 1)\lambda = 2, 4, \dots$

Theorem 1.3. The neutrix compositions $\delta^{(2s-1)}(\operatorname{sgn} x |x|^{1/s})$ and $\delta^{(s-1)}(|x|^{1/s})$ exist and

$$\begin{aligned} \delta^{(2s-1)}(\operatorname{sgn} x |x|^{1/s}) &= \frac{1}{2} (2s)! \delta'(x), \\ \delta^{(s-1)}(|x|^{1/s}) &= (-1)^{s-1} \delta(x), \end{aligned} \quad (1.6)$$

for $s = 1, 2, \dots$

Theorem 1.4. The neutrix composition $\delta^{(s)}(\sinh^{-1} x_+^{1/r})$ exists and

$$\delta^{(s)}\left[\left(\sinh^{-1} x_+\right)^{1/r}\right] = \sum_{k=0}^{(s+1)/r-1} \sum_{i=0}^k \binom{k}{i} \frac{(-1)^k r c_{s,k,i}}{2^{k+1} k!} \delta^{(k)}(x), \quad (1.7)$$

for $s = 0, 1, 2, \dots$ and $r = 1, 2, \dots$, where

$$c_{r,s,k,i} = \frac{(-1)^s s! \left[(k - 2i + 1)^{rs+r-1} + (k - 2i - 1)^{rs+r-1} \right]}{2(rs + r - 1)!}. \quad (1.8)$$

The next two theorems were proved in [13].

Theorem 1.5. The neutrix composition $\delta^{(s)}[\ln^r(1 + |x|)]$ exists and

$$\delta^{(s)}[\ln^r(1 + |x|)] = \sum_{k=0}^{sr+r-1} \sum_{i=0}^k \binom{k}{i} \frac{(-1)^{s-i} \left[1 + (-1)^k \right] s!(i+1)^{rs+r-1}}{2r(rs+r-1)! k!} \delta^{(k)}(x). \quad (1.9)$$

for $s = 0, 1, 2, \dots$, and $r = 1, 2, \dots$

In particular, the composition $\delta[\ln(1 + |x|)]$ exists and

$$\delta[\ln(1 + |x|)] = \delta(x). \quad (1.10)$$

Theorem 1.6. The neutrix composition $\delta^{(s)}[\ln(1 + |x|^{1/r})]$ exists and

$$\delta^{(s)}\left[\ln\left(1 + |x|^{1/r}\right)\right] = \sum_{k=0}^{m-1} \sum_{i=0}^{kr+r-1} \binom{kr+r-1}{i} \frac{(-1)^{r+s+i-1} \left[1 + (-1)^k \right] r(i+1)^s}{2k!} \delta^{(k)}(x), \quad (1.11)$$

for $s = 0, 1, 2, \dots$ and $r = 2, 3, \dots$, where m is the smallest non-negative integer greater than $(s-r+1)r^{-1}$.

In particular, the composition $\delta^{(s)}[\ln(1 + |x^{1/r}|)]$ exists and

$$\delta^{(s)}\left[\ln\left(1 + \left|x^{1/r}\right|\right)\right] = 0, \quad (1.12)$$

for $s = 0, 1, 2, \dots, r-2$ and $r = 2, 3, \dots$ and

$$\delta^{(r-1)}\left[\ln\left(1 + \left|x^{1/r}\right|\right)\right] = (-1)^{r-1} r! \delta(x), \quad (1.13)$$

for $r = 2, 3, \dots$

2. Main Results

We now prove the following theorem.

Theorem 2.1. The neutrix composition $\delta^{(s)}[(\sinh^{-1}x_+)^r]$ exists and

$$\delta^{(s)}\left[(\sinh^{-1}x_+)^r\right] = \sum_{k=0}^{sr+r-1} \sum_{i=0}^k \binom{k}{i} \frac{(-1)^k r c_{s,k,i}}{2^{k+1} k!} \delta^{(k)}(x), \quad (2.1)$$

for $s = 0, 1, 2, \dots$ and $r = 1, 2, \dots$, where

$$c_{r,s,k,i} = \frac{(-1)^s s! \left[(k-2i+1)^{rs+r-1} + (k-2i-1)^{rs+r-1} \right]}{2(rs+r-1)!}. \quad (2.2)$$

In particular, the neutrix composition $\delta(\sinh^{-1}x_+)$ exists and

$$\delta(\sinh^{-1}x_+) = \frac{1}{2} \delta(x). \quad (2.3)$$

Proof. To prove (2.1), we first of all evaluate

$$\int_{-1}^1 \delta_n^{(s)}\left[(\sinh^{-1}x_+)^r\right] x^k dx. \quad (2.4)$$

We have

$$\begin{aligned} \int_{-1}^1 \delta_n^{(s)}\left[(\sinh^{-1}x_+)^r\right] x^k dx &= n^{s+1} \int_{-1}^1 \rho^{(s)}\left[(n \sinh^{-1}x_+)^r\right] x^k dx \\ &= n^{s+1} \int_0^1 \rho^{(s)}\left[n(\sinh^{-1}x)\right] x^k dx \\ &\quad + n^{s+1} \int_{-1}^0 \rho^{(s)}(0) x^k dx \\ &= I_1 + I_2. \end{aligned} \quad (2.5)$$

It is obvious that

$$N - \lim_{n \rightarrow \infty} I_2 = N - \lim_{n \rightarrow \infty} \int_{-1}^0 \delta_n^{(s)} \left[\left(\sinh^{-1} x_+ \right)^r \right] x^k dx = 0, \quad (2.6)$$

for $k = 0, 1, 2, \dots$

Making the substitution $t = n(\sinh^{-1} x)^r$, we have for large enough n

$$\begin{aligned} I_1 &= \frac{n^{s-r+1}}{r} \int_0^1 t^{1/(r-1)} \sinh^k \left(\frac{t}{n} \right)^{1/r} \cosh \left(\frac{t}{n} \right)^{1/r} \rho^{(s)}(t) dt \\ &\times \int_0^1 t^{1/(r-1)} \left\{ \exp \left[(k-2i+1) \left(\frac{t}{n} \right)^{1/r} \right] + \exp \left[(k-2i-1) \left(\frac{t}{n} \right)^{1/r} \right] \right\} \rho^{(s)}(t) dt, \end{aligned} \quad (2.7)$$

where

$$\begin{aligned} &n^{(s-1)/(r+1)} \int_0^1 t^{1/(r-1)} \left\{ \exp \left[(k-2i+1) \left(\frac{t}{n} \right)^{1/r} \right] + \exp \left[(k-2i-1) \left(\frac{t}{n} \right)^{1/r} \right] \right\} \rho^{(s)}(t) dt \\ &= \sum_{p=0}^{\infty} \int_0^1 \frac{[(k-2i+1)^p + (k-2i-1)^p] t^{(p/r)+(1/r)-1}}{p! n^{(p/r)+(1/r)-s-1}} \rho^{(s)}(t) dt. \end{aligned} \quad (2.8)$$

It follows that

$$\begin{aligned} &N - \lim_{n \rightarrow \infty} n^{s-1/r+1} \int_0^1 t^{1/(r-1)} \left\{ \exp \left[(k-2i+1) \left(\frac{t}{n} \right)^{1/r} \right] + \exp \left[(k-2i-1) \left(\frac{t}{n} \right)^{1/r} \right] \right\} \rho^{(s)}(t) dt \\ &= \frac{(-1)^s s! [(k-2i+1)^{rs+r-1} + (k-2i-1)^{rs+r-1}]}{2(rs+r-1)!} \\ &= C_{r,s,k,i}, \end{aligned} \quad (2.9)$$

and by applying the neutrix limit we obtain

$$N - \lim_{n \rightarrow \infty} I_1 = N - \lim_{n \rightarrow \infty} \int_0^1 \delta_n^{(s)} \left[\left(\sinh^{-1} x_+ \right)^r \right] x^k dx = \frac{1}{2^{k+1} r} \sum_{i=0}^k \binom{k}{i} (-1)^i C_{r,s,k,i} \quad (2.10)$$

for $k = 0, 1, 2, \dots$

When $k = sr + r$, we have

$$\begin{aligned}
 |I_1| &= \int_0^1 \left| \delta_n^{(s)} \left[\left(\sinh^{-1} x_+ \right)^r \right] x^{sr+r} \right| dx \\
 &= n^{s+1} \int_0^1 \left| \rho_n^{(s)} \left[n \left(\sinh^{-1} x \right)^r \right] x^{sr+r} \right| dx \\
 &\leq \frac{n^{(s-1)/(r+1)}}{2^{sr+r}} \exp(sr+r+1) \int_0^1 \left| \left[1 - \exp \left[-2 \left(\frac{t}{n} \right)^{1/r} \right] \right]^{sr+r} \rho^{(s)}(t) \right| dt \\
 &= \frac{n^{(s-1)/(r+1)}}{2^{sr+r}} \exp(sr+r+1) \int_0^1 \left[2 \left(\frac{t}{n} \right)^{1/r} + O(n^{-2/r}) \right]^{sr+r} \left| \rho^{(s)}(t) \right| dt \\
 &\leq n^{-1/r} \exp(sr+r+1) \int_0^1 \left[1 + O(n^{-2/r}) \right] \left| \rho^{(s)}(t) \right| dt \\
 &= O(n^{-1/r}).
 \end{aligned} \tag{2.11}$$

Thus, if ψ is an arbitrary continuous function, then

$$\lim_{n \rightarrow \infty} \int_0^1 \delta_n^{(s)} \left[\left(\sinh^{-1} x_+ \right)^r \right] x^{rs+r} \psi(x) dx = 0. \tag{2.12}$$

We also have

$$\int_{-1}^0 \delta_n^{(s)} \left[\left(\sinh^{-1} x_+ \right)^r \right] \psi(x) dx = n^{s+1} \int_{-1}^0 \rho^{(s)}(0) \psi(x) dx, \tag{2.13}$$

and it follows that

$$N - \lim_{n \rightarrow \infty} \int_{-1}^0 \delta_n^{(s)} \left[\left(\sinh^{-1} x_+ \right)^r \right] \psi(x) dx = 0. \tag{2.14}$$

If now φ is an arbitrary function in $\mathfrak{D}[-1, 1]$, then by Taylor's Theorem, we have

$$\varphi(x) = \sum_{k=0}^{sr+r-1} \frac{\varphi^{(k)}(0)}{k!} x^k + \frac{x^{rs+r}}{(rs+r)!} \varphi^{(rs+r)}(\xi x), \tag{2.15}$$

where $0 < \xi < 1$, and so

$$\begin{aligned}
 & N - \lim_{n \rightarrow \infty} \left\langle \delta_n^{(s)} \left[\left(\sinh^{-1} x_+ \right)^{1/r} \right], \varphi(x) \right\rangle \\
 &= N - \lim_{n \rightarrow \infty} \sum_{k=0}^{sr+r-1} \frac{\varphi^{(k)}(0)}{k!} \int_0^1 \delta_n^{(s)} \left[\left(\sinh^{-1} x_+ \right)^r \right] x^k dx \\
 &+ N - \lim_{n \rightarrow \infty} \sum_{k=0}^{sr+r-1} \frac{\varphi^{(k)}(0)}{k!} \int_{-1}^0 \delta_n^{(s)} \left[\left(\sinh^{-1} x_+ \right)^r \right] x^k dx \\
 &+ \lim_{n \rightarrow \infty} \frac{1}{(sr+r)!} \int_0^1 \delta_n^{(s)} \left[\left(\sinh^{-1} x_+ \right)^r \right] x^{sr+r} \varphi^{(sr+r)}(\xi x) dx \tag{2.16} \\
 &+ \lim_{n \rightarrow \infty} \frac{1}{(sr+r)!} \int_{-1}^0 \delta_n^{(s)} \left[\left(\sinh^{-1} x_+ \right)^r \right] x^{sr+r} \varphi^{(sr+r)}(\xi x) dx \\
 &= \sum_{k=0}^{sr+r-1} \sum_{i=0}^k \binom{k}{i} \frac{r c_{r,s,k,i} \varphi^{(k)}(0)}{2^{k+1} k!} + 0 \\
 &= \sum_{k=0}^{sr+r-1} \sum_{i=0}^k \binom{k}{i} \frac{(-1)^k r c_{r,s,k,i}}{2^{k+1} k!} \langle \delta^{(k)}(x), \varphi(x) \rangle,
 \end{aligned}$$

on using (2.3) to (2.14). This proves (2.1) on the interval $(-1, 1)$.

It is clear that $\delta^{(s)}[(\sinh^{-1} x_+)^r] = 0$ for $x > 0$ and so (2.1) holds for $x > -1$.

Now, suppose that φ is an arbitrary function in $\mathfrak{D}[a, b]$, where $a < b < 0$. Then,

$$\int_a^b \delta_n^{(s)} \left[\left(\sinh^{-1} x_+ \right)^r \right] \varphi(x) dx = n^{s+1} \int_a^b \rho^{(s)}(0) \varphi(x) dx \tag{2.17}$$

and so

$$N - \lim_{n \rightarrow \infty} \int_a^b \delta_n^{(s)} \left[\left(\sinh^{-1} x_+ \right)^r \right] \varphi(x) dx = 0. \tag{2.18}$$

It follows that $\delta^{(s)}[(\sinh^{-1} x_+)^r] = 0$ on the interval (a, b) . Since a and b are arbitrary, we see that (2.1) holds on the real line. This completes the proof of the theorem. \square

Corollary 2.2. *The neutrix composition $\delta^{(s)}[(\sinh^{-1}|x|)^r]$ exists and*

$$\delta^{(s)} \left[\left(\sinh^{-1}|x| \right)^r \right] = \sum_{k=0}^{sr+r-1} \sum_{i=0}^k \binom{k}{i} \frac{[(-1)^k + 1] c_{r,s,k,i}}{2^{k+1} k!} \delta^{(k)}(x), \tag{2.19}$$

for $s = 0, 1, 2, \dots$ and $r = 1, 2, \dots$

In particular, the composition $\delta(\sinh^{-1}|x|)$ exists and

$$\delta(\sinh^{-1}|x|) = \frac{1}{2}\delta(x). \quad (2.20)$$

Proof. To prove (2.19), we note that

$$\begin{aligned} \int_{-1}^1 \delta_n^{(s)} \left[(\sinh^{-1}|x|)^r \right] x^k dx &= n^{s+1} \int_{-1}^1 \rho^{(s)} \left[(n \sinh^{-1}|x|)^r \right] x^k dx \\ &= n^{s+1} \left[1 + (-1)^k \right] \int_0^1 \rho^{(s)} \left[n(\sinh^{-1}x)^r \right] x^k dx, \end{aligned} \quad (2.21)$$

and (2.19) now follows as above.

Equation (2.20) follows on noting that in the particular case $s = 0$, the usual limit holds in (2.10). This completes the proof of the corollary. \square

Theorem 2.3. *The neutrix composition $\delta^{(2s-1)}[\sinh^{-1}(\operatorname{sgn} x \cdot x^2)]$ exists and*

$$\delta^{(2s-1)} \left[\sinh^{-1}(\operatorname{sgn} x \cdot x^2) \right] = \sum_{k=0}^{2s-1} \sum_{i=0}^{k+1} \binom{k}{i} \frac{(-1)^k b_{s,k,i}}{2^{k+1}(2k+1)!} \delta^{(k)}(x), \quad (2.22)$$

for $s = 1, 2, \dots$, where

$$b_{s,k,i} = (k - 2i + 1)^{2s-1} + (k - 2i - 1)^{2s-1}. \quad (2.23)$$

Proof. To prove (2.22), we now have to evaluate

$$\int_{-1}^1 \delta_n^{(2s-1)} \left[\sinh^{-1}(\operatorname{sgn} x \cdot x^2) \right] x^k dx. \quad (2.24)$$

We have

$$\begin{aligned} \int_{-1}^1 \delta_n^{(2s-1)} \left[\sinh^{-1}(\operatorname{sgn} x \cdot x^2) \right] x^k dx &= n^{2s} \int_{-1}^1 \rho^{(2s-1)} \left[n \sinh^{-1}(\operatorname{sgn} x \cdot x^2) \right] x^k dx \\ &= \begin{cases} 2n^{2s} \int_0^1 \rho^{(2s-1)} \left[n(\sinh^{-1}x^2) \right] x^k dx, & k \text{ odd,} \\ 0, & k \text{ even.} \end{cases} \end{aligned} \quad (2.25)$$

Making the substitution $t = n(\sinh^{-1} x^2)$, we have for large enough n

$$\begin{aligned} & \int_{-1}^1 \delta_n^{(2s-1)} \left[\sinh^{-1}(\operatorname{sgn} x \cdot x^2) \right] x^k dx \\ &= 2n^{2s} \int_0^1 \rho^{(2s-1)} \left[n(\sinh^{-1} x^2) \right] x^{2k+1} dx \\ &= \frac{n^{2s-1}}{2^{k+1}} \sum_{i=0}^k \binom{k}{i} (-1)^i \int_0^1 \left\{ \exp \left[\frac{(k-2i+1)t}{n} \right] + \exp \left[\frac{(k-2i-1)t}{n} \right] \right\} \rho^{(2s-1)}(t) dt, \end{aligned} \quad (2.26)$$

where

$$\begin{aligned} & n^{2s-1} \int_0^1 \left\{ \exp \left[\frac{(k-2i+1)t}{n} \right] + \exp \left[\frac{(k-2i-1)t}{n} \right] \right\} \rho^{(s)}(t) dt \\ &= \sum_{p=0}^{\infty} \int_0^1 \frac{[(k-2i+1)^p + (k-2i-1)^p] t^p}{p! n^{p-2s+1}} \rho^{(2s-1)}(t) dt. \end{aligned} \quad (2.27)$$

It follows that

$$\begin{aligned} & N \lim_{n \rightarrow \infty} n^{2s-1} \int_0^1 \left\{ \exp \left[\frac{(k-2i+1)t}{n} \right] + \exp \left[\frac{(k-2i-1)t}{n} \right] \right\} \rho^{(s)}(t) dt \\ &= N \lim_{n \rightarrow \infty} \sum_{p=0}^{\infty} \int_0^1 \frac{[(k-2i+1)^p + (k-2i-1)^p] t^p}{p! n^{p-2s+1}} \rho^{(2s-1)}(t) dt \\ &= \frac{-(k-2i+1)^{2s-1} + (k-2i-1)^{2s-1}}{2} \\ &= \frac{b_{s,k,i}}{2}, \end{aligned} \quad (2.28)$$

and so by using the neutrix limit, we have

$$N \lim_{n \rightarrow \infty} \int_{-1}^1 \delta_n^{(2s-1)} \left[\sinh^{-1}(\operatorname{sgn} x \cdot x^2) \right] x^{2k+1} dx = \sum_{i=0}^k \binom{k}{i} \frac{(-1)^{i+1} b_{s,k,i}}{2^{k+1}}, \quad (2.29)$$

for $k = 0, 1, 2, \dots$

When $k = 2s$, we have

$$\begin{aligned}
\int_{-1}^1 \left| \delta_n^{(2s-1)} \left[\sinh^{-1}(\operatorname{sgn} x \cdot x^2) \right] x^{4s+1} \right| dx &= n^{2s} \int_{-1}^1 \rho^{(2s-1)} \left[n \left(\sinh^{-1} x^2 \right) \right] x^{4s+1} dx \\
&\leq \frac{n^{2s-1}}{2^{s-1}} \exp(s+1) \int_{-1}^1 \left| \left[1 - \exp\left(-\frac{2t}{n}\right) \right]^{2s} \rho^{(2s-1)}(t) \right| dt \\
&= \frac{n^{2s-1}}{2^{s-1}} \exp(s+1) \int_{-1}^1 \left| \left[\frac{2t}{n} + O(n^{-2}) \right]^{2s} \rho^{(2s-1)}(t) \right| dt \\
&\leq 2^{2s+1} n^{-1} \exp(s+1) \int_{-1}^1 \left| 1 + O(n^{-2/r}) \right| \left| \rho^{(2s-1)}(t) \right| dt \\
&= O(n^{-1}).
\end{aligned} \tag{2.30}$$

Thus, if φ is an arbitrary continuous function, then

$$\lim_{n \rightarrow \infty} \int_{-1}^1 \delta_n^{(2s-1)} \left[\sinh^{-1}(\operatorname{sgn} x \cdot x^2) \right] x^{4s+1} \varphi(x) dx = 0. \tag{2.31}$$

If now φ is an arbitrary function in $\mathfrak{D}[-1, 1]$, then by Taylor's Theorem, we have

$$\varphi(x) = \sum_{k=0}^{4s} \frac{\varphi^{(k)}(0)}{k!} x^k + \frac{x^{4s+1}}{(4s+1)!} \varphi^{(4s+1)}(\xi x), \tag{2.32}$$

where $0 < \xi < 1$, and so

$$\begin{aligned}
&N - \lim_{n \rightarrow \infty} \left\langle \delta_n^{(2s-1)} \left[\sinh^{-1}(\operatorname{sgn} x \cdot x^2) \right], \varphi(x) \right\rangle \\
&= N - \lim_{n \rightarrow \infty} \sum_{k=0}^{2s-1} \frac{\varphi^{(2k+1)}(0)}{(2k+1)!} \int_{-1}^1 \delta_n^{(2s-1)} \left[\sinh^{-1}(\operatorname{sgn} x \cdot x^2) \right] x^{2k+1} dx \\
&\quad + \lim_{n \rightarrow \infty} \frac{1}{(4s+1)!} \int_{-1}^1 \delta_n^{(4s+1)} \left[\sinh^{-1}(\operatorname{sgn} x \cdot x^2) \right] x^{4s+1} \varphi^{(4s+1)}(\xi x) dx \\
&= \sum_{k=0}^{2s-1} \sum_{i=0}^k \binom{k}{i} \frac{(-1)^{i+1} b_{s,k,i} \varphi^{(k)}(0)}{2^{k+1} (2k+1)!} + 0 \\
&= \sum_{k=0}^{2s-1} \sum_{i=0}^k \binom{k}{i} \frac{(-1)^{i+k+1} b_{s,k,i}}{2^{k+1} (2k+1)!} \left\langle \delta^{(k)}(x), \varphi(x) \right\rangle,
\end{aligned} \tag{2.33}$$

on using (2.25) to (2.31), proving (2.22) on the interval $(-1, 1)$. However, it is clear that $\delta_n^{(2s-1)} \left[\sinh^{-1}(\operatorname{sgn} x \cdot x^2) \right] = 0$ for $|x| > 0$ and so (2.22) holds on the real line, completing the proof of the theorem. \square

Corollary 2.4. *The composition $\delta'[\sinh^{-1}\text{sgn } x \cdot x^2]$ exists and*

$$\delta'[\sinh^{-1}(\text{sgn } x \cdot x^2)] = \frac{\delta'(x)}{4 \cdot 3!} - 2\delta(x). \quad (2.34)$$

Proof. To prove (2.34) note that in the particular case $s = 1$, the usual limits hold and then (2.34) is a particular case of (2.22). This completes the proof of the corollary. \square

For further related results on the neutrix operation of distributions, see [12–22] and [2, 3, 23].

Acknowledgments

The authors would like to thank the referee(s) for the very constructive comments and suggestions that improved the paper. The paper was prepared when B. Fisher visited University Putra Malaysia and therefore the authors gratefully acknowledge that this research was partially supported by the University Putra Malaysia under the Research University Grant Scheme no. 05-01-09-0720RU.

References

- [1] H. Eltayeb, A. Kılıçman, and B. Fisher, "A new integral transform and associated distributions," *Integral Transforms and Special Functions*, vol. 21, no. 5-6, pp. 367–379, 2010.
- [2] A. Kılıçman, "On the commutative neutrix product of distributions," *Indian Journal of Pure and Applied Mathematics*, vol. 30, no. 8, pp. 753–762, 1999.
- [3] A. Kılıçman, "A comparison on the commutative neutrix convolution of distributions and the exchange formula," *Czechoslovak Mathematical Journal*, vol. 51(126), no. 3, pp. 463–471, 2001.
- [4] A. Kılıçman and H. Eltayeb, "A note on defining singular integral as distribution and partial differential equations with convolution term," *Mathematical and Computer Modelling*, vol. 49, no. 1-2, pp. 327–336, 2009.
- [5] P. Antosik, "Composition of distributions," Tech. Rep. 9, University of Wisconsin, 1988.
- [6] B. Fisher, "On defining the change of variable in distributions," *Rostocker Mathematisches Kolloquium*, no. 28, pp. 75–86, 1985.
- [7] B. Fisher, "On defining the distribution $(x_+^r)_-^s$," *Univerzitet u Novom Sadu. Zbornik Radova Prirodno-Matematičkog Fakulteta. Serija za Matemati*, vol. 15, no. 1, pp. 119–129, 1985.
- [8] B. Fisher, "The composition and neutrix composition of distributions," *Proceedings of Mathematical Methods of Engineering*, pp. 59–69, 2006.
- [9] I. M. Gelfand and G. E. Shilov, *Generalized Functions*, vol. 1, Academic Press, New York, NY, USA, 1st edition, 1964.
- [10] J. G. van der Corput, "Introduction to the neutrix calculus," *Journal d'Analyse Mathématique*, vol. 7, pp. 281–399, 1960.
- [11] B. Fisher, "The delta function and the composition of distributions," *Demonstratio Mathematica*, vol. 35, no. 1, pp. 117–123, 2002.
- [12] B. Fisher and A. Kılıçman, "On the composition and neutrix composition of the delta function and powers of the inverse hyperbolic sine function," *Integral Transforms and Special Functions*, vol. 21, no. 12, pp. 935–944, 2010.
- [13] B. Fisher, T. Kraiweeradechachai, and E. Özçağ, "Results on the neutrix composition of the delta function," *Hacetatepe Journal of Mathematics and Statistics*, vol. 36, no. 2, pp. 147–156, 2007.
- [14] B. Fisher, "On the composition of the distributions $x_+^{-s}\text{In}^m x_+$ and x_+^s ," *Applicable Analysis and Discrete Mathematics*, vol. 3, no. 2, pp. 212–223, 2009.
- [15] B. Fisher and B. Jolevska-Tuneska, "Two results on the composition of distributions," *Thai Journal of Mathematics*, vol. 3, no. 1, pp. 17–26, 2005.

- [16] B. Fisher, B. Jolevska-Tuneska, and E. Özçağ, "Further results on the compositions of distributions," *Integral Transforms and Special Functions*, vol. 13, no. 2, pp. 109–116, 2002.
- [17] B. Fisher and A. Kılıçman, "A commutative neutrix product of ultradistributions," *Integral Transforms and Special Functions*, vol. 4, no. 1-2, pp. 77–82, 1996.
- [18] B. Fisher and T. Kraiweeradechachai, "On the composition of the distributions $x_+^\lambda \text{In}^m x_+$ and $x_+^{-1/\lambda}$," *Sarajevo Journal of Mathematics*, vol. 4(17), no. 2, pp. 249–257, 2008.
- [19] B. Fisher, A. Kananthai, G. Sritanatana, and K. Nonlaopon, "The composition of the distributions $x_-^{ms} \text{In} x_-$ and $x_+^{r-p/m}$," *Integral Transforms and Special Functions*, vol. 16, no. 1, pp. 13–19, 2005.
- [20] B. Fisher and K. Taş, "On the composition of the distributions x_+^{-r} and x_+^μ ," *Indian Journal of Pure and Applied Mathematics*, vol. 36, no. 1, pp. 11–22, 2005.
- [21] B. Fisher and K. Taş, "On the composition of the distributions $x^{-1} \text{In}|x|$ and x_+^r ," *Integral Transforms and Special Functions*, vol. 16, no. 7, pp. 533–543, 2005.
- [22] B. Fisher and K. Taş, "Some results on the non-commutative neutrix product of distributions," *Integral Transforms and Special Functions*, vol. 20, no. 1-2, pp. 35–44, 2009.
- [23] B. Fisher, A. Kılıçman, and S. Pehlivan, "The neutrix convolution product of $x_+^\lambda \text{In}^r x_+$ and $x_-^\mu \text{In}^s x_-$," *Integral Transforms and Special Functions*, vol. 7, no. 3-4, pp. 237–246, 1998.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

