

## Research Article

# Asymptotic Behaviour of Eigenvalues and Eigenfunctions of a Sturm-Liouville Problem with Retarded Argument

Erdoğan Şen,<sup>1,2</sup> Jong Jin Seo,<sup>3</sup> and Serkan Araci<sup>4</sup>

<sup>1</sup> Department of Mathematics, Faculty of Arts and Science, Namik Kemal University, 59030 Tekirdağ, Turkey

<sup>2</sup> Department of Mathematics Engineering, Istanbul Technical University, Maslak, 34469 Istanbul, Turkey

<sup>3</sup> Department of Applied Mathematics, Pukyong National University, Busan 608-737, Republic of Korea

<sup>4</sup> Department of Mathematics, Faculty of Science and Arts, University of Gaziantep, 27310 Gaziantep, Turkey

Correspondence should be addressed to Jong Jin Seo; seo2011@pknu.ac.kr

Received 18 November 2012; Accepted 1 March 2013

Academic Editor: Suh-Yuh Yang

Copyright © 2013 Erdoğan Şen et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In the present paper, a discontinuous boundary-value problem with retarded argument at the two points of discontinuities is investigated. We obtained asymptotic formulas for the eigenvalues and eigenfunctions. This is the first work containing two discontinuities points in the theory of differential equations with retarded argument. In that special case the transmission coefficients  $\delta = \gamma = 1$  and retarded argument  $\Delta \equiv 0$  in the results obtained in this work coincide with corresponding results in the classical Sturm-Liouville operator.

## 1. Introduction

Delay differential equations arise in many areas of mathematical modelling, for example, population dynamics (taking into account the gestation times), infectious diseases (accounting for the incubation periods), physiological and pharmaceutical kinetics (modelling, for example, the body's reaction to CO<sub>2</sub>, and so forth, in circulating blood) and chemical kinetics (such as mixing reactants), the navigational control of ships and aircraft, and more general control problems. Also, differential equations and nonlinear differential equations have been studied by many mathematicians in several ways for a long time cf. [1–20].

Boundary value problems for differential equations of the second order with retarded argument were studied in [5–10, 13–16], and various physical applications of such problems can be found in [6].

In the papers [13–16], the asymptotic formulas for the eigenvalues and eigenfunctions of a discontinuous boundary value problem with retarded argument and a spectral parameter in the boundary conditions were derived. In spite of their being already a long years, these subjects are still today

enveloped in an aura of mystery within scientific community although they have penetrated numerous mathematical field.

The asymptotic formulas for the eigenvalues and eigenfunctions of the Sturm-Liouville problem with the spectral parameter in the boundary condition were obtained in [17–20].

In this paper we study the eigenvalues and eigenfunctions of a discontinuous boundary value problem with retarded argument. Namely, we consider the boundary value problem for the differential equation

$$y''(x) + q(x)y(x - \Delta(x)) + \lambda y(x) = 0, \quad (1)$$

on  $[0, h_1) \cup (h_1, h_2) \cup (h_2, \pi]$ , with boundary conditions

$$y(0) \cos \alpha + y'(0) \sin \alpha = 0, \quad (2)$$

$$y(\pi) \cos \beta + y'(\pi) \sin \beta = 0, \quad (3)$$

and transmission conditions

$$y(h_1 - 0) - \delta y(h_1 + 0) = 0, \tag{4}$$

$$y'(h_1 - 0) - \delta y'(h_1 + 0) = 0, \tag{5}$$

$$y(h_2 - 0) - \gamma y(h_2 + 0) = 0, \tag{6}$$

$$y'(h_2 - 0) - \gamma y'(h_2 + 0) = 0, \tag{7}$$

where the real-valued function  $q(x)$  is continuous in  $[0, h_1) \cup (h_1, h_2) \cup (h_2, \pi]$  and has finite limits

$$q(h_1 \pm 0) = \lim_{x \rightarrow h_1 \pm 0} q(x), \quad q(h_2 \pm 0) = \lim_{x \rightarrow h_2 \pm 0} q(x), \tag{8}$$

the real-valued function  $\Delta(x) \geq 0$  continuous in  $[0, h_1) \cup (h_1, h_2) \cup (h_2, \pi]$  and has finite limits

$$\Delta(h_1 \pm 0) = \lim_{x \rightarrow h_1 \pm 0} \Delta(x), \quad \Delta(h_2 \pm 0) = \lim_{x \rightarrow h_2 \pm 0} \Delta(x); \tag{9}$$

if  $x \in [0, h_1)$  then  $x - \Delta(x) \geq 0$ ; if  $x \in (h_1, h_2)$  then  $x - \Delta(x) \geq h_1$ ; if  $x \in (h_2, \pi)$  then  $x - \Delta(x) \geq h_2$ ;  $\lambda$  is a real spectral parameter;  $h_1, h_2, \alpha, \beta, \delta, \gamma \neq 0$  are arbitrary real numbers such that  $0 < h_1 < h_2 < \pi$  and  $\sin \alpha \sin \beta \neq 0$ .

It must be noted that some problems with transmission conditions which arise in mechanics (thermal condition problem for a thin laminated plate) were studied in [20].

Let  $w_1(x, \lambda)$  be a solution of (1) on  $[0, h_1]$ , satisfying the initial conditions

$$w_1(0, \lambda) = \sin \alpha, \quad w_1'(0, \lambda) = -\cos \alpha. \tag{10}$$

The conditions (10) define a unique solution of (1) on  $[0, h_1]$  ([5], page 12).

After defining the above solution, then we will define the solution  $w_2(x, \lambda)$  of (1) on  $[h_1, h_2]$  by means of the solution  $w_1(x, \lambda)$  using the initial conditions

$$w_2(h_1, \lambda) = \delta^{-1} w_1(h_1, \lambda), \quad w_2'(h_1, \lambda) = \delta^{-1} w_1'(h_1, \lambda). \tag{11}$$

The conditions (11) define a unique solution of (1) on  $[h_1, h_2]$ .

After describing the above solution, then we will give the solution  $w_3(x, \lambda)$  of (1) on  $[h_2, \pi]$  by means of the solution  $w_2(x, \lambda)$  using the initial conditions

$$w_3(h_2, \lambda) = \gamma^{-1} w_2(h_2, \lambda), \quad w_3'(h_2, \lambda) = \gamma^{-1} w_2'(h_2, \lambda). \tag{12}$$

The conditions (12) define a unique solution of (1) on  $[h_2, \pi]$ .

Consequently, the function  $w(x, \lambda)$  is defined on  $[0, h_1) \cup (h_1, h_2) \cup (h_2, \pi]$  by the equality

$$w(x, \lambda) = \begin{cases} w_1(x, \lambda), & x \in [0, h_1), \\ w_2(x, \lambda), & x \in (h_1, h_2), \\ w_3(x, \lambda), & x \in (h_2, \pi], \end{cases} \tag{13}$$

is a solution of (1) on  $[0, h_1) \cup (h_1, h_2) \cup (h_2, \pi]$ , which satisfies one of the boundary conditions and four transmission conditions.

**Lemma 1.** Let  $w(x, \lambda)$  be a solution of (1) and  $\lambda > 0$ . Then the following integral equations hold:

$$w_1(x, \lambda) = \sin \alpha \cos sx - \frac{\cos \alpha}{s} \sin sx - \frac{1}{s} \int_0^x q(\tau) \sin s(x - \tau) w_1(\tau - \Delta(\tau), \lambda) d\tau \tag{14}$$

$(s = \sqrt{\lambda}, \lambda > 0),$

$$w_2(x, \lambda) = \frac{1}{\delta} w_1(h_1, \lambda) \cos s(x - h_1) + \frac{w_1'(h_1, \lambda)}{s\delta} \sin s(x - h_1) - \frac{1}{s} \int_{h_1}^x q(\tau) \sin s(x - \tau) w_2(\tau - \Delta(\tau), \lambda) d\tau \tag{15}$$

$(s = \sqrt{\lambda}, \lambda > 0),$

$$w_3(x, \lambda) = \frac{1}{\gamma} w_2(h_2, \lambda) \cos s(x - h_2) + \frac{w_2'(h_2, \lambda)}{s\gamma} \sin s(x - h_2) - \frac{1}{s} \int_{h_2}^x q(\tau) \sin s(x - \tau) w_3(\tau - \Delta(\tau), \lambda) d\tau \tag{16}$$

$(s = \sqrt{\lambda}, \lambda > 0).$

*Proof.* To prove this lemma, it is enough to substitute  $-s^2 w_1(\tau, \lambda) - w_1''(\tau, \lambda)$ ,  $-s^2 w_2(\tau, \lambda) - w_2''(\tau, \lambda)$ , and  $-s^2 w_3(\tau, \lambda) - w_3''(\tau, \lambda)$  instead of  $-q(\tau) w_1(\tau - \Delta(\tau), \lambda)$ ,  $-q(\tau) w_2(\tau - \Delta(\tau), \lambda)$ , and  $-q(\tau) w_3(\tau - \Delta(\tau), \lambda)$  in the integrals in (14), (15), and (16), respectively, and integrate by parts twice.  $\square$

**Theorem 2.** The problem (1)–(7) can have only simple eigenvalues.

*Proof.* Let  $\tilde{\lambda}$  be an eigenvalue of the problem (1)–(7) and

$$\tilde{y}(x, \tilde{\lambda}) = \begin{cases} \tilde{y}_1(x, \tilde{\lambda}), & x \in [0, h_1), \\ \tilde{y}_2(x, \tilde{\lambda}), & x \in (h_1, h_2), \\ \tilde{y}_3(x, \tilde{\lambda}), & x \in (h_2, \pi], \end{cases} \tag{17}$$

be a corresponding eigenfunction. Then, from (2) and (10), it follows that the determinant

$$W[\tilde{y}_1(0, \tilde{\lambda}), w_1(0, \tilde{\lambda})] = \begin{vmatrix} \tilde{y}_1(0, \tilde{\lambda}) & \sin \alpha \\ \tilde{y}_1'(0, \tilde{\lambda}) & -\cos \alpha \end{vmatrix} = 0, \tag{18}$$

and, by Theorem 2.2.2, in [5] the functions  $\tilde{y}_1(x, \tilde{\lambda})$  and  $w_1(x, \tilde{\lambda})$  are linearly dependent on  $[0, h_1]$ . We can also prove that the functions  $\tilde{y}_2(x, \tilde{\lambda})$  and  $w_2(x, \tilde{\lambda})$  are linearly dependent on

$[h_1, h_2]$  and  $\tilde{y}_3(x, \tilde{\lambda})$  and  $w_3(x, \tilde{\lambda})$  are linearly dependent on  $[h_2, \pi]$ . Hence

$$\tilde{y}_i(x, \tilde{\lambda}) = K_i w_i(x, \tilde{\lambda}) \quad (i = 1, 2, 3), \quad (19)$$

for some  $K_1 \neq 0$ ,  $K_2 \neq 0$ , and  $K_3 \neq 0$ . We must show that  $K_1 = K_2$  and  $K_2 = K_3$ . Suppose that  $K_2 \neq K_3$ . From the equalities (6) and (19), we have

$$\begin{aligned} & \tilde{y}(h_2 - 0, \tilde{\lambda}) - \gamma \tilde{y}(h_2 + 0, \tilde{\lambda}) \\ &= \tilde{y}_2(h_2, \tilde{\lambda}) - \gamma \tilde{y}_3(h_2, \tilde{\lambda}) \\ &= K_2 w_2(h_2, \tilde{\lambda}) - \gamma K_3 w_3(h_2, \tilde{\lambda}) \\ &= K_2 \gamma w_3(h_2, \tilde{\lambda}) - K_3 \gamma w_3(h_2, \tilde{\lambda}) \\ &= \gamma (K_2 - K_3) w_3(h_2, \tilde{\lambda}) = 0. \end{aligned} \quad (20)$$

Since  $\gamma(K_2 - K_3) \neq 0$  it follows that

$$w_3(h_2, \tilde{\lambda}) = 0. \quad (21)$$

By the same procedure from equality (7) we can derive that

$$w_3'(h_2, \tilde{\lambda}) = 0. \quad (22)$$

From the fact that  $w_3(x, \tilde{\lambda})$  is a solution of the differential (1) on  $[h_2, \pi]$  and satisfies the initial conditions (21) and (22), it follows that  $w_3(x, \tilde{\lambda}) = 0$  identically on  $[h_2, \pi]$ .

By using this method, we may also find

$$\begin{aligned} w_2(h_2, \tilde{\lambda}) &= w_2'(h_2, \tilde{\lambda}) = 0, \\ w_1(h_1, \tilde{\lambda}) &= w_1'(h_1, \tilde{\lambda}) = 0. \end{aligned} \quad (23)$$

From the latter discussions of  $w_3(x, \tilde{\lambda})$ , it follows that  $w_2(x, \tilde{\lambda}) = 0$  and  $w_1(x, \tilde{\lambda}) = 0$  identically on  $(h_1, h_2)$  and  $[0, h_1)$ , but this contradicts (10), thus completing the proof.  $\square$

## 2. An Existence Theorem

The function  $w(x, \lambda)$  defined in Section 1 is a nontrivial solution of (1) satisfying conditions (2) and (4)–(7). Putting  $w(x, \lambda)$  into (3), we get the characteristic equation

$$F(\lambda) \equiv w(\pi, \lambda) \cos \beta + w'(\pi, \lambda) \sin \beta = 0. \quad (24)$$

By Theorem 2 the set of eigenvalues of boundary-value problem (1)–(7) coincides with the set of real roots of (24). Let

$$\begin{aligned} q_1 &= \int_0^{h_1} |q(\tau)| d\tau, & q_2 &= \int_{h_1}^{h_2} |q(\tau)| d\tau, \\ q_3 &= \int_{h_2}^{\pi} |q(\tau)| d\tau. \end{aligned} \quad (25)$$

**Lemma 3.** (1) Let  $\lambda \geq 4q_1^2$ . Then for the solution  $w_1(x, \lambda)$  of (14), the following inequality holds:

$$|w_1(x, \lambda)| \leq \frac{1}{q_1} \sqrt{4q_1^2 \sin^2 \alpha + \cos^2 \alpha}, \quad x \in [0, h_1]. \quad (26)$$

(2) Let  $\lambda \geq \max\{4q_1^2, 4q_2^2\}$ . Then for the solution  $w_2(x, \lambda)$  of (15), the following inequality holds:

$$|w_2(x, \lambda)| \leq \frac{4}{q_1 |\delta|} \sqrt{4q_1^2 \sin^2 \alpha + \cos^2 \alpha}, \quad x \in [h_1, h_2]. \quad (27)$$

(3) Let  $\lambda \geq \max\{4q_1^2, 4q_2^2, 4q_3^2\}$ . Then for the solution  $w_3(x, \lambda)$  of (16), the following inequality holds:

$$|w_3(x, \lambda)| \leq \frac{16}{q_1 |\delta \gamma|} \sqrt{4q_1^2 \sin^2 \alpha + \cos^2 \alpha}, \quad x \in [h_2, \pi]. \quad (28)$$

*Proof.* Let  $B_{1\lambda} = \max_{[0, h_1]} |w_1(x, \lambda)|$ . Then from (14), it follows that, for every  $\lambda > 0$ , the following inequality holds:

$$B_{1\lambda} \leq \sqrt{\sin^2 \alpha + \frac{\cos^2 \alpha}{s^2}} + \frac{1}{s} B_{1\lambda} q_1. \quad (29)$$

If  $s \geq 2q_1$  we get (26). Differentiating (14) with respect to  $x$ , we have

$$\begin{aligned} w_1'(x, \lambda) &= -s \sin \alpha \sin sx - \cos \alpha \cos sx \\ &\quad - \int_0^x q(\tau) \cos s(x - \tau) w_1(\tau - \Delta(\tau), \lambda) d\tau. \end{aligned} \quad (30)$$

From expressions of (30) and (26), it follows that, for  $s \geq 2q_1$ , the following inequality holds:

$$\frac{|w_1'(x, \lambda)|}{s} \leq \frac{1}{q_1} \sqrt{4q_1^2 \sin^2 \alpha + \cos^2 \alpha}. \quad (31)$$

Let  $B_{2\lambda} = \max_{[h_1, h_2]} |w_2(x, \lambda)|$ . Then from (11), (26), and (31) it follows that, for  $s \geq 2q_1$  and  $s \geq 2q_2$ , the following inequality holds:

$$B_{2\lambda} \leq \frac{2}{q_1 |\delta|} \sqrt{4q_1^2 \sin^2 \alpha + \cos^2 \alpha} + \frac{1}{2q_2} B_{2\lambda} q_2. \quad (32)$$

Hence, if  $\lambda \geq \max\{4q_1^2, 4q_2^2\}$ , it reduces to (27). Differentiating (15) with respect to, we get

$$\begin{aligned} w_2'(x, \lambda) &= -\frac{s}{\delta} w_1(h_1, \lambda) \sin s(x - h_1) \\ &\quad + \frac{w_1'(h_1, \lambda)}{\delta} \cos s(x - h_1) \\ &\quad - \int_{h_1}^x q(\tau) \cos s(x - \tau) w_2(\tau - \Delta(\tau), \lambda) d\tau. \end{aligned} \quad (s = \sqrt{\lambda}, \lambda > 0). \quad (33)$$

From (26) and (33), it follows that, for  $s \geq 2q_1$  and  $s \geq 2q_2$ , the following inequality holds:

$$\frac{|w'_2(x, \lambda)|}{s} \leq \frac{4}{|\delta|q_1} \sqrt{4q_1^2 \sin^2 \alpha + \cos^2 \alpha}. \quad (34)$$

Let  $B_{3\lambda} = \max_{[h_2, \pi]} |w_3(x, \lambda)|$ . Then from (16), (27), and (34) it follows that, for  $s \geq 2q_1$ ,  $s \geq 2q_2$  and  $s \geq 2q_3$ , the following inequality holds:

$$B_{3\lambda} \leq \frac{8}{q_1 |\delta \gamma|} \sqrt{4q_1^2 \sin^2 \alpha + \cos^2 \alpha} + \frac{1}{s} B_{3\lambda} q_3. \quad (35)$$

Hence, if  $\lambda \geq \max\{4q_1^2, 4q_2^2, 4q_3^2\}$  we procure (28).  $\square$

**Theorem 4.** *The problem (1)–(7) has an infinite set of positive eigenvalues.*

*Proof.* Differentiating (16) with respect to  $x$ , we readily see that

$$\begin{aligned} w'_3(x, \lambda) = & -\frac{s}{\gamma} w_2(h_2, \lambda) \sin s(x - h_2) \\ & + \frac{w'_2(h_2, \lambda)}{\gamma} \cos s(x - h_2) \\ & - \int_{h_2}^x q(\tau) \cos s(x - \tau) w_3(\tau - \Delta(\tau), \lambda) d\tau. \end{aligned} \quad (s = \sqrt{\lambda}, \lambda > 0). \quad (36)$$

With the helps of (14), (15), (16), (24), (30), and (36), we have the following:

$$\begin{aligned} & \left[ \frac{1}{\gamma} \left\{ \frac{1}{\delta} \left( \sin \alpha \cos sh_1 - \frac{\cos \alpha}{s} \sin sh_1 \right. \right. \right. \\ & \quad \left. \left. - \frac{1}{s} \int_0^{h_1} q(\tau) \sin s(h_1 - \tau) w_1(\tau - \Delta(\tau), \lambda) d\tau \right) \right. \\ & \quad \times \cos s(h_2 - h_1) \\ & \quad \left. - \frac{1}{s\delta} \left( s \sin \alpha \sin sh_1 + \cos \alpha \cos sh_1 \right. \right. \\ & \quad \left. \left. + \int_0^{h_1} q(\tau) \cos s(h_1 - \tau) w_1(\tau - \Delta(\tau), \lambda) d\tau \right) \right. \\ & \quad \times \sin s(h_2 - h_1) \\ & \quad \left. - \frac{1}{s} \int_{h_1}^{h_2} q(\tau) \sin s(h_2 - \tau) w_2(\tau - \Delta(\tau), \lambda) d\tau \right\} \end{aligned}$$

$$\begin{aligned} & \times \cos s(\pi - h_2) \\ & + \frac{1}{s\gamma} \left\{ -\frac{s}{\delta} \left( \sin \alpha \cos sh_1 - \frac{\cos \alpha}{s} \sin sh_1 \right. \right. \\ & \quad \left. \left. - \frac{1}{s} \int_0^{h_1} q(\tau) \sin s(h_1 - \tau) \right. \right. \\ & \quad \left. \left. \times w_1(\tau - \Delta(\tau), \lambda) d\tau \right) \right. \\ & \quad \times \sin s(h_2 - h_1) \\ & \quad \left. + \frac{1}{\delta} \left( -s \sin \alpha \sin sh_1 - \cos \alpha \cos sh_1 \right. \right. \\ & \quad \left. \left. - \int_0^{h_1} q(\tau) \cos s(h_1 - \tau) \right. \right. \\ & \quad \left. \left. \times w_1(\tau - \Delta(\tau), \lambda) d\tau \right) \right. \\ & \quad \times \cos s(h_2 - h_1) \\ & \quad \left. - \int_{h_1}^{h_2} q(\tau) \cos s(h_2 - \tau) w_2(\tau - \Delta(\tau), \lambda) d\tau \right\} \\ & \times \sin s(\pi - h_2) \\ & - \frac{1}{s} \int_{h_2}^{\pi} q(\tau) \sin s(\pi - \tau) w_3(\tau - \Delta(\tau), \lambda) d\tau \Big] \cos \beta \\ & + \left[ -\frac{s}{\gamma} \left\{ \frac{1}{\delta} \left( \sin \alpha \cos sh_1 - \frac{\cos \alpha}{s} \sin sh_1 \right. \right. \right. \\ & \quad \left. \left. - \frac{1}{s} \int_0^{h_1} q(\tau) \sin s(h_1 - \tau) \right. \right. \\ & \quad \left. \left. \times w_1(\tau - \Delta(\tau), \lambda) d\tau \right) \right. \\ & \quad \times \cos s(h_2 - h_1) \\ & \quad \left. - \frac{1}{s\delta} \left( s \sin \alpha \sin sh_1 + \cos \alpha \cos sh_1 \right. \right. \\ & \quad \left. \left. + \int_0^{h_1} q(\tau) \cos s(h_1 - \tau) \right. \right. \\ & \quad \left. \left. \times w_1(\tau - \Delta(\tau), \lambda) d\tau \right) \right. \\ & \quad \times \sin s(h_2 - h_1) \\ & \quad \left. - \frac{1}{s} \int_{h_1}^{h_2} q(\tau) \sin s(h_2 - \tau) w_2(\tau - \Delta(\tau), \lambda) d\tau \right\} \end{aligned}$$

$$\times \sin s(\pi - h_2)$$

$$\begin{aligned}
 & + \frac{1}{\gamma} \left\{ -\frac{s}{\delta} \times \left( \sin \alpha \cos sh_1 - \frac{\cos \alpha}{s} \sin sh_1 \right. \right. \\
 & \qquad \qquad \qquad \left. \left. - \frac{1}{s} \int_0^{h_1} q(\tau) \sin s(h_1 - \tau) \right. \right. \\
 & \qquad \qquad \qquad \left. \left. \times w_1(\tau - \Delta(\tau), \lambda) d\tau \right) \right. \\
 & \times \sin s(h_2 - h_1) \\
 & + \frac{1}{\delta} \left( -s \sin \alpha \sin sh_1 - \cos \alpha \cos sh_1 \right. \\
 & \qquad \qquad \qquad \left. - \int_0^{h_1} q(\tau) \cos s(h_1 - \tau) \right. \\
 & \qquad \qquad \qquad \left. \times w_1(\tau - \Delta(\tau), \lambda) d\tau \right) \\
 & \times \cos s(h_2 - h_1) \\
 & \left. - \int_{h_1}^{h_2} q(\tau) \cos s(h_2 - \tau) w_2(\tau - \Delta(\tau), \lambda) d\tau \right\} \\
 & \times \cos s(\pi - h_2) \\
 & - \int_{h_2}^{\pi} q(\tau) \cos s(\pi - \tau) w_3(\tau - \Delta(\tau), \lambda) d\tau \Big] \sin \beta = 0.
 \end{aligned} \tag{37}$$

Let  $\lambda$  be sufficiently big. Then, by (26), (27), and (28), (37) may be rewritten in the following form:

$$-\frac{s \sin \alpha \sin \beta}{\gamma \delta} \{ \cos sh_2 \sin s(\pi - h_2) \tag{38}$$

$$\begin{aligned}
 & + \sin sh_2 \cos s(\pi - h_2) \} + O(1) = 0, \\
 & s \sin s\pi + O(1) = 0.
 \end{aligned} \tag{39}$$

Obviously, for big  $s$  (39) has an infinite set of roots. Thus, the proof of the theorem is completed.  $\square$

### 3. Asymptotic Formulas for Eigenvalues and Eigenfunctions

Now we begin to study asymptotic properties of eigenvalues and eigenfunctions. In the following we will assume that is sufficiently big. From (14) and (26), we obtain

$$w_1(x, \lambda) = O(1) \quad \text{on } [0, h_1]. \tag{40}$$

By (15) and (27), this leads to

$$w_2(x, \lambda) = O(1) \quad \text{on } [h_1, h_2]. \tag{41}$$

By (16) and (28), this leads to

$$w_3(x, \lambda) = O(1) \quad \text{on } [h_2, \pi]. \tag{42}$$

The existence and continuity of the derivatives  $w'_{1s}(x, \lambda)$  for  $0 \leq x \leq h_1$ ,  $|\lambda| < \infty$ ,  $w'_{2s}(x, \lambda)$  for  $h_1 \leq x \leq h_2$ ,  $|\lambda| < \infty$  and  $w'_{3s}(x, \lambda)$  for  $h_2 \leq x \leq \pi$ ,  $|\lambda| < \infty$  follows from Theorem 1.4.1 in [5].

**Lemma 5.** *The following holds true:*

$$w'_{1s}(x, \lambda) = O(1), \quad x \in [0, h_1], \tag{43}$$

$$w'_{2s}(x, \lambda) = O(1), \quad x \in [h_1, h_2], \tag{44}$$

$$w'_{3s}(x, \lambda) = O(1), \quad x \in [h_2, \pi]. \tag{45}$$

*Proof.* By differentiating (16) with respect to  $s$ , we get, by (43) and (44) the following:

$$\begin{aligned}
 & w'_{3s}(x, \lambda) \\
 & = -\frac{1}{s} \int_{h_2}^x q(\tau) \sin s(x - \tau) w'_{3s}(\tau - \Delta(\tau), \lambda) + Z(x, \lambda), \\
 & \qquad \qquad \qquad (|Z(x, \lambda)| \leq Z_0).
 \end{aligned} \tag{46}$$

Let  $D_\lambda = \max_{[h_2, \pi]} |w'_{3s}(x, \lambda)|$ . Then the existence of  $D_\lambda$  follows from continuity of derivation for  $x \in [h_2, \pi]$ . From (46)

$$D_\lambda \leq \frac{1}{s} q_3 D_\lambda + Z_0. \tag{47}$$

Now let  $s \geq 2q_3$ . Then  $D_\lambda \leq 2Z_0$  and the validity of the asymptotic formula (45) follows. Formulas (43) and (44) may be proved analogically.  $\square$

**Theorem 6.** *Let  $n$  be a natural number. For each sufficiently big  $n$  there is exactly one eigenvalue of the problem (1)–(7) near  $n^2$ .*

*Proof.* We consider the expression which is denoted by  $O(1)$  in (39). If formulas (40)–(45) are taken into consideration, it can be shown by differentiation with respect to  $s$  that for big  $s$  this expression has bounded derivative. We will show that, for big  $n$ , only one root (39) lies near to each  $n$ . We consider the function  $\phi(s) = s \sin s\pi + O(1)$ . Its derivative, which has the form  $\phi'(s) = \sin s\pi + s\pi \cos s\pi + O(1)$ , does not vanish for  $s$  close to  $n$  for sufficiently big  $n$ . Thus our assertion follows by Rolle's Theorem.  $\square$

Let  $n$  be sufficiently big. In what follows we will denote by  $\lambda_n = s_n^2$  the eigenvalue of the problem (1)–(7) situated near  $n^2$ . We set  $s_n = n + \delta_n$ . Then from (39) it follows that  $\delta_n = O(1/n)$ . Consequently

$$s_n = n + O\left(\frac{1}{n}\right), \tag{48}$$

$$\lambda_n = n^2 + O(1). \tag{49}$$

Formula (48) makes it possible to obtain asymptotic expressions for eigenfunction of the problem (1)–(7). From (14), (30), and (40), we get

$$w_1(x, \lambda) = \sin \alpha \cos sx + O\left(\frac{1}{s}\right), \tag{50}$$

$$w'_1(x, \lambda) = -s \sin \alpha \sin sx + O(1). \tag{51}$$

From expressions of (15), (31), (39), and (41), we easily see that

$$\begin{aligned} w_2(x, \lambda) &= \frac{\sin \alpha}{\delta} \cos sx + O\left(\frac{1}{s}\right), \\ w_3(x, \lambda) &= \frac{\sin \alpha}{\delta\gamma} \cos sx + O\left(\frac{1}{s}\right). \end{aligned} \tag{52}$$

By substituting (48) into (50), (52), we find that

$$\begin{aligned} u_{1n} &= w_1(x, \lambda_n) = \sin \alpha \cos nx + O\left(\frac{1}{n}\right), \\ u_{2n} &= w_2(x, \lambda_n) = \frac{\sin \alpha}{\delta} \cos nx + O\left(\frac{1}{n}\right), \\ u_{3n} &= w_3(x, \lambda_n) = \frac{\sin \alpha}{\delta\gamma} \cos nx + O\left(\frac{1}{n}\right). \end{aligned} \tag{53}$$

Hence the eigenfunctions  $u_n(x)$  have the following asymptotic representation:

$$u_n(x) = \begin{cases} \sin \alpha \cos nx + O\left(\frac{1}{n}\right), & \text{for } x \in [0, h_1), \\ \frac{\sin \alpha}{\delta} \cos nx + O\left(\frac{1}{n}\right), & \text{for } x \in (h_1, h_2), \\ \frac{\sin \alpha}{\delta\gamma} \cos nx + O\left(\frac{1}{n}\right), & \text{for } x \in (h_2, \pi]. \end{cases} \tag{54}$$

Under some additional conditions the more exact asymptotic formulas which depend upon the retardation may be obtained. Let us assume that the following conditions are fulfilled.

(a) The derivatives  $q'(x)$  and  $\Delta''(x)$  exist and are bounded in  $[0, h_1) \cup (h_1, h_2) \cup (h_2, \pi]$  and have finite limits  $q'(h_1 \pm 0) = \lim_{x \rightarrow h_1 \pm 0} q'(x)$ ,  $q'(h_2 \pm 0) = \lim_{x \rightarrow h_2 \pm 0} q'(x)$  and  $\Delta''(h_1 \pm 0) = \lim_{x \rightarrow h_1 \pm 0} \Delta''(x)$ ,  $\Delta''(h_2 \pm 0) = \lim_{x \rightarrow h_2 \pm 0} \Delta''(x)$ , respectively.

(b)  $\Delta'(x) \leq 1$  in  $[0, h_1) \cup (h_1, h_2) \cup (h_2, \pi]$ ,  $\Delta(0) = 0$ ,  $\lim_{x \rightarrow h_1+0} \Delta(x) = 0$ , and  $\lim_{x \rightarrow h_2+0} \Delta(x) = 0$ .

It is easy to see that, using (b)

$$\begin{aligned} x - \Delta(x) &\geq 0, & x \in [0, h_1), \\ x - \Delta(x) &\geq h_1, & x \in (h_1, h_2), \\ x - \Delta(x) &\geq h_2, & x \in (h_2, \pi] \end{aligned} \tag{55}$$

are obtained.

By (50), (52), and (55), we have

$$w_1(\tau - \Delta(\tau), \lambda) = \sin \alpha \cos s(\tau - \Delta(\tau)) + O\left(\frac{1}{s}\right), \tag{56}$$

$$w_2(\tau - \Delta(\tau), \lambda) = \frac{\sin \alpha}{\delta} \cos s(\tau - \Delta(\tau)) + O\left(\frac{1}{s}\right), \tag{57}$$

$$w_3(\tau - \Delta(\tau), \lambda) = \frac{\sin \alpha}{\delta\gamma} \cos s(\tau - \Delta(\tau)) + O\left(\frac{1}{s}\right) \tag{58}$$

on  $[0, h_1)$ ,  $(h_1, h_2)$  and  $(h_2, \pi]$ , respectively.

Under the conditions (a) and (b) the following formulas:

$$\begin{aligned} \int_0^x q(\tau) \cos s(2\tau - \Delta(\tau)) d\tau &= O\left(\frac{1}{s}\right), \\ \int_0^x q(\tau) \sin s(2\tau - \Delta(\tau)) d\tau &= O\left(\frac{1}{s}\right) \end{aligned} \tag{59}$$

can be proved by the same technique in Lemma 3.3.3 in [5].

Putting the expressions (56), (57), and (58) into (37), and then using (59), after long operations we have

$$\begin{aligned} &\frac{\cos s\pi \sin(\alpha - \beta) - s \sin \alpha \sin \beta \sin s\pi}{\delta\gamma} \\ &- \frac{\sin \alpha \sin \beta}{2\delta\gamma} \int_0^\pi q(\tau) \cos s(\pi - \Delta(\tau)) d\tau \\ &+ O\left(\frac{1}{s}\right) = 0. \end{aligned} \tag{60}$$

Hence

$$\begin{aligned} \tan s\pi &= \frac{1}{s} \left( \frac{\sin(\alpha - \beta)}{\sin \alpha \sin \beta} \right. \\ &\quad \left. - \frac{1}{2} \int_0^\pi q(\tau) \cos s(\pi - \Delta(\tau)) d\tau \right) \\ &\quad + O\left(\frac{1}{s^2}\right). \end{aligned} \tag{61}$$

Again, if we take  $s_n = n + \delta_n$ , then from (48)

$$\begin{aligned} \tan((n + \delta_n)\pi) &= \tan \delta_n \pi \\ &= \frac{1}{n} \left( \frac{\sin(\alpha - \beta)}{\sin \alpha \sin \beta} \right. \\ &\quad \left. - \frac{1}{2} \int_0^\pi q(\tau) \cos s(\pi - \Delta(\tau)) d\tau \right) \\ &\quad + O\left(\frac{1}{n^2}\right). \end{aligned} \tag{62}$$

Hence for big  $n$ ,

$$\begin{aligned} \delta_n &= \frac{1}{n\pi} \left( \frac{\sin(\alpha - \beta)}{\sin \alpha \sin \beta} - \frac{1}{2} \int_0^\pi q(\tau) \cos s(\pi - \Delta(\tau)) d\tau \right) \\ &\quad + O\left(\frac{1}{n^2}\right) \end{aligned} \tag{63}$$

and finally

$$\begin{aligned} s_n &= n + \frac{1}{n\pi} \left( \frac{\sin(\alpha - \beta)}{\sin \alpha \sin \beta} - \frac{1}{2} \int_0^\pi q(\tau) \cos s(\pi - \Delta(\tau)) d\tau \right) \\ &\quad + O\left(\frac{1}{n^2}\right). \end{aligned} \tag{64}$$

Thus, we have proven the following theorem.

**Theorem 7.** *If conditions (a) and (b) are satisfied, then the eigenvalues  $\lambda_n = s_n^2$  of the problem (1)–(7) have the (64) asymptotic formula for  $n \rightarrow \infty$ .*

Now, we may obtain sharper asymptotic formulas for the eigenfunctions. From (14), (56), and (59), we have

$$w_1(x, \lambda) = \sin \alpha \cos sx \left[ 1 + \frac{1}{2s} \int_0^x q(\tau) \sin s\Delta(\tau) d\tau \right] - \frac{\sin sx}{s} \left[ \cos \alpha + \frac{\sin \alpha}{2} \int_0^x q(\tau) \cos s\Delta(\tau) d\tau \right] + O\left(\frac{1}{s^2}\right), \quad x \in [0, h_1]. \tag{65}$$

Now, replacing  $s$  by  $s_n$  and using (64), we have

$$u_{1n}(x) = w_1(x, \lambda_n) = \sin \alpha \left\{ \cos nx \left[ 1 + \frac{1}{2n} \int_0^x q(\tau) \sin(n\Delta(\tau)) d\tau \right] - \frac{\sin nx}{n\pi} \left[ \left( \frac{\sin(\alpha - \beta)}{\sin \alpha \sin \beta} - \frac{1}{2} \int_0^\pi q(\tau) \cos(n(\pi - \Delta(\tau))) d\tau \right) x + \left( \cot \alpha + \frac{1}{2} \int_0^x q(\tau) \cos(n\Delta(\tau)) d\tau \right) \pi \right] \right\} + O\left(\frac{1}{n^2}\right). \tag{66}$$

From (15), (57), (59), and (65) we have

$$w_2(x, \lambda) = \frac{\sin \alpha \cos sx}{\delta} \left[ 1 + \frac{1}{2s} \int_0^x q(\tau) \sin s\Delta(\tau) d\tau \right] - \frac{\sin sx}{s\delta} \left[ \cos \alpha + \frac{\sin \alpha}{2} \int_0^x q(\tau) \cos s\Delta(\tau) d\tau \right] + O\left(\frac{1}{s^2}\right), \quad x \in (h_1, h_2). \tag{67}$$

Now, replacing  $s$  by  $s_n$  and using (64), we have

$$u_{2n}(x) = w_2(x, \lambda_n) = \frac{\sin \alpha}{\delta} \left\{ \cos nx \left[ 1 + \frac{1}{2n} \int_0^x q(\tau) \sin(n\Delta(\tau)) d\tau \right] \right.$$

$$\left. - \frac{\sin nx}{n\pi} \left[ \left( \frac{\sin(\alpha - \beta)}{\sin \alpha \sin \beta} - \frac{1}{2} \int_0^\pi q(\tau) \cos(n(\pi - \Delta(\tau))) d\tau \right) x + \left( \cot \alpha + \frac{1}{2} \int_0^x q(\tau) \cos(n\Delta(\tau)) d\tau \right) \pi \right] \right\} + O\left(\frac{1}{n^2}\right). \tag{68}$$

From (16), (58), (59), (65), and (67) and after long operations, we have

$$w_3(x, \lambda) = \frac{\sin \alpha \cos sx}{\delta\gamma} \left[ 1 + \frac{1}{2s} \int_0^x q(\tau) \sin s\Delta(\tau) d\tau \right] - \frac{\sin sx}{s\delta\gamma} \left[ \cos \alpha + \frac{\sin \alpha}{2} \int_0^x q(\tau) \cos s\Delta(\tau) d\tau \right] + O\left(\frac{1}{s^2}\right), \quad x \in (h_2, \pi]. \tag{69}$$

Now replacing  $s$  by  $s_n$  and using (64), we have

$$u_{3n}(x) = w_3(x, \lambda_n) = \frac{\sin \alpha}{\delta\gamma} \left\{ \cos nx \left[ 1 + \frac{1}{2n} \int_0^x q(\tau) \sin(n\Delta(\tau)) d\tau \right] - \frac{\sin nx}{n\pi} \left[ \left( \frac{\sin(\alpha - \beta)}{\sin \alpha \sin \beta} - \frac{1}{2} \int_0^\pi q(\tau) \cos(n(\pi - \Delta(\tau))) d\tau \right) x + \left( \cot \alpha + \frac{1}{2} \int_0^x q(\tau) \cos(n\Delta(\tau)) d\tau \right) \pi \right] \right\} + O\left(\frac{1}{n^2}\right). \tag{70}$$

Thus, we have proven the following theorem.

**Theorem 8.** *If conditions (a) and (b) are satisfied, then the eigenfunctions  $u_n(x)$  of the problem (1)–(7) have the following asymptotic formula for  $n \rightarrow \infty$ :*

$$u_n(x) = \begin{cases} u_{1n}(x), & x \in [0, h_1], \\ u_{2n}(x), & x \in (h_1, h_2), \\ u_{3n}(x), & x \in (h_2, \pi], \end{cases} \tag{71}$$

where  $u_{1n}(x)$ ,  $u_{2n}(x)$ , and  $u_{3n}(x)$  are determined as in (49), (68), and (70), respectively.

## References

- [1] G. Teschl, *Ordinary Differential Equations and Dynamical Systems*, vol. 140, American Mathematical Society, Providence, RI, USA, 2012.
- [2] M. Shahriari, A. J. Akbarfam, and G. Teschl, “Uniqueness for inverse Sturm-Liouville problems with a finite number of transmission conditions,” *Journal of Mathematical Analysis and Applications*, vol. 395, no. 1, pp. 19–29, 2012.
- [3] T. Kim, “Identities involving Frobenius-Euler polynomials arising from non-linear differential equations,” *Journal of Number Theory*, vol. 132, no. 12, pp. 2854–2865, 2012.
- [4] S. B. Norkin, “On a boundary problem of Sturm-Liouville type for a second-order differential equation with a retarded argument,” *Izvestija Vysših Učebnyh Zavedenij Matematika*, vol. 6, no. 7, pp. 203–214, 1958 (Russian).
- [5] S. B. Norkin, *Differential Equations of the Second Order with Retarded Argument*, vol. 31 of *Translations of Mathematical Monographs*, American Mathematical Society, Providence, RI, USA, 1972.
- [6] R. Bellman and K. L. Cooke, *Differential-Difference Equations*, Academic Press, London, UK, 1963.
- [7] G. V. Demidenko and V. A. Likhoshvai, “On differential equations with retarded argument,” *Siberian Mathematical Journal*, vol. 46, no. 3, pp. 417–430, 2005.
- [8] G. V. Demidenko and I. I. Matveeva, “On numerical study of asymptotic stability of solutions to linear periodic differential equations with a parameter,” *Journal of Computational Mathematics and Optimization*, vol. 5, no. 3, pp. 163–173, 2009.
- [9] A. Bayramov, S. Öztürk Uslu, and S. Aliskan, “Computation of eigenvalues and eigenfunctions of a discontinuous boundary value problem with retarded argument,” *Applied Mathematics and Computation*, vol. 191, no. 2, pp. 592–600, 2007.
- [10] E. Şen, S. Araci, and M. Acikgoz, “Asymptotic behavior of eigenvalues and fundamental solutions of one discontinuous fourth-order boundary value problem,” *Research Journal of Mathematical and Statistical Sciences*, vol. 1, no. 1, pp. 23–32, 2013.
- [11] E. Şen and S. Araci, “Computation of Eigenvalues and fundamental solutions of a fourth-order boundary value,” *Proceedings of the Jangjeon Mathematical Society*, vol. 15, no. 4, pp. 455–464, 2012.
- [12] E. Şen and A. Bayramov, “On calculation of eigenvalues and eigenfunctions of a Sturm-Liouville type problem with retarded argument which contains a spectral parameter in the boundary condition,” *Journal of Inequalities and Applications*, vol. 2011, article 113, 2011.
- [13] E. Şen and A. Bayramov, “On a discontinuous Sturm-Liouville type problem with retarded argument,” *American Institute of Physics Conference Proceedings*, vol. 1389, pp. 1172–1175, 2011.
- [14] E. Şen and A. Bayramov, “Calculation of eigenvalues and eigenfunctions of a discontinuous boundary value problem with retarded argument which contains a spectral parameter in the boundary condition,” *Mathematical and Computer Modelling*, vol. 54, no. 11-12, pp. 3090–3097, 2011.
- [15] A. Bayramov and E. Şen, “On a Sturm-Liouville type problem with retarded argument,” *Mathematical Methods in the Applied Sciences*, vol. 36, no. 1, pp. 39–48, 2013.
- [16] C. T. Fulton, “Two-point boundary value problems with eigenvalue parameter contained in the boundary conditions,” *Proceedings of the Royal Society of Edinburgh A*, vol. 77, no. 3-4, pp. 293–308, 1977.
- [17] P. A. Binding, P. J. Browne, and B. A. Watson, “Sturm-Liouville problems with boundary conditions rationally dependent on the eigenparameter II,” *Journal of Computational and Applied Mathematics*, vol. 148, no. 1, pp. 147–168, 2002.
- [18] N. Altinisik, M. Kadakal, and O. Sh. Mukhtarov, “Eigenvalues and eigenfunctions of discontinuous Sturm-Liouville problems with eigenparameter-dependent boundary conditions,” *Acta Mathematica Hungarica*, vol. 102, no. 1-2, pp. 159–175, 2004.
- [19] N. B. Kerimov and Kh. R. Mamedov, “On a boundary value problem with a spectral parameter in the boundary conditions,” *Sibirskii Matematicheskii Zhurnal*, vol. 40, no. 2, pp. 325–335, 1999.
- [20] I. Titeux and Y. Yakubov, “Completeness of root functions for thermal conduction in a strip with piecewise continuous coefficients,” *Mathematical Models & Methods in Applied Sciences*, vol. 7, no. 7, pp. 1035–1050, 1997.





# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

