

Research Article

On a New Hilbert-Type Intergral Inequality with the Intergral in Whole Plane

Zheng Zeng¹ and Zitian Xie²

¹ Department of Mathematics, Shaoguan University, Shaoguan, Guangdong 512005, China

² Department of Mathematics, Zhaoqing University, Zhaoqing, Guangdong 526061, China

Correspondence should be addressed to Zitian Xie, gdzqxzt@163.com

Received 5 May 2010; Accepted 14 July 2010

Academic Editor: Andrea Laforgia

Copyright © 2010 Z. Zeng and Z. Xie. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

By introducing some parameters and estimating the weight functions, we build a new Hilbert's inequality with the homogeneous kernel of 0 order and the integral in whole plane. The equivalent inequality and the reverse forms are considered. The best constant factor is calculated using Complex Analysis.

1. Introduction

If $f(x), g(x) \geq 0$ and satisfy that $0 < \int_0^\infty f^2(x)dx < \infty$ and $0 < \int_0^\infty g^2(x)dx < \infty$, then we have [1]

$$\iint_0^\infty \frac{f(x)g(x)}{x+y} dx dy < \pi \left\{ \int_0^\infty f^2(x)dx \int_0^\infty g^2(x)dx \right\}^{1/2}, \quad (1.1)$$

where the constant factor π is the best possible. Inequality (1.1) is well known as Hilbert's integral inequality, which has been extended by Hardy-Riesz as [2].

If $p > 1$, $1/p + 1/q = 1$, $f(x), g(x) \geq 0$, such that $0 < \int_0^\infty f^p(x)dx < \infty$ and $0 < \int_0^\infty g^q(x)dx < \infty$, then we have the following Hardy-Hilbert's integral inequality:

$$\iint_0^\infty \frac{f(x)g(y)}{x+y} dx dy < \frac{\pi}{\sin(\pi/p)} \left\{ \int_0^\infty f^p(x)dx \right\}^{1/p} \left\{ \int_0^\infty g^q(x)dx \right\}^{1/q}, \quad (1.2)$$

where the constant factor $\pi / \sin(\pi/p)$ also is the best possible.

Both of them are important in Mathematical Analysis and its applications [3]. It attracts some attention in recent years. Actually, inequalities (1.1) and (1.2) have many generalizations and variations. Equation (1.1) has been strengthened by Yang and others (including double series inequalities) [4–21].

In 2008, Xie and Zeng gave a new Hilbert-type Inequality [4] as follows.

If $a > 0$, $b > 0$, $c > 0$, $p > 1$, $1/p + 1/q = 1$, $f(x), g(x) \geq 0$ such that $0 < \int_0^\infty x^{-1-p/2} f^p(x) dx < \infty$ and $0 < \int_0^\infty x^{-1-q/2} g^q(x) dx < \infty$, then

$$\begin{aligned} & \iint_0^\infty \frac{f(x)g(y)}{(x+a^2y)(x+b^2y)(x+c^2y)} dx dy \\ & < K \left\{ \int_0^\infty x^{-1-p/2} f^p(x) dx \right\}^{1/p} \left\{ \int_0^\infty x^{-1-q/2} g^q(x) dx \right\}^{1/q}, \end{aligned} \quad (1.3)$$

where the constant factor $K = \pi / (a+b)(a+c)(b+c)$ is the best possible.

The main purpose of this paper is to build a new Hilbert-type inequality with homogeneous kernel of degree 0, by estimating the weight function. The equivalent inequality is considered.

In the following, we always suppose that: $1/p + 1/q = 1$, $p > 1$, $r \in (-1, 0)$, $0 < \alpha < \beta < \pi$.

2. Some Lemmas

We start by introducing some lemmas.

Lemma 2.1. *If $k_1 := \int_0^\infty u^{-1+r} \ln((1+2u \cos \alpha + u^2)/(1+2u \cos \beta + u^2)) du$, $k_2 := \int_0^\infty u^{-1+r} \ln((1-2u \cos \beta + u^2)/(1-2u \cos \alpha + u^2)) du$, then*

$$\begin{aligned} k_1 &= \frac{4\pi \sin(r(\beta - \alpha)/2) \sin(r(\alpha + \beta)/2)}{r \sin r\pi}, \\ k_2 &= \frac{4\pi \sin(r(\beta - \alpha)/2) \sin(r\pi - r(\alpha + \beta)/2)}{r \sin r\pi}, \\ k &:= \int_{-\infty}^\infty |u|^{-1+r} \left| \ln \frac{1+2u \cos \alpha + u^2}{1+2u \cos \beta + u^2} \right| du \\ &= k_1 + k_2 = \frac{4\pi \sin(r(\beta - \alpha)/2) \cos((r/2)(\pi - \alpha - \beta))}{r \cos(r\pi/2)}. \end{aligned} \quad (2.1)$$

Proof. We have

$$\begin{aligned} A &:= \int_0^\infty x^{r-1} \ln(x^2 + 2x \cos \alpha + 1) dx = \frac{1}{r} x^r \ln(x^2 + 2x \cos \alpha + 1) \Big|_0^\infty \\ &\quad - \frac{2}{r} \int_0^\infty \frac{x^r (x + \cos \alpha)}{x^2 + 2x \cos \alpha + 1} dx \\ &:= -\frac{2}{r} B. \end{aligned} \quad (2.2)$$

Setting $f(z) = z^r(z + \cos \alpha)/(z^2 + 2z \cos \alpha + 1)$, $z_1 = -e^{i\alpha}$, $z_2 = -e^{-i\alpha}$, then

$$\begin{aligned} B &= \frac{2\pi i}{1 - e^{2\pi r i}} [\operatorname{Res}(f, z_1) + \operatorname{Res}(f, z_2)] \\ &= \frac{2\pi i}{1 - e^{2\pi r i}} \left[\frac{z_1^r(z_1 + \cos \alpha)}{z_1 - z_2} + \frac{z_2^r(z_2 + \cos \alpha)}{z_2 - z_1} \right] = -\frac{\pi \cos r\alpha}{\sin r\pi} \end{aligned} \tag{2.3}$$

we find that $A = -2B/r = 2\pi \cos r\alpha/r \sin r\pi$, then

$$\begin{aligned} k_1 &:= \int_0^\infty u^{-1+r} \ln \frac{1 + 2u \cos \alpha + u^2}{1 + 2u \cos \beta + u^2} du = \frac{4\pi \sin(r(\beta - \alpha)/2) \sin(r(\alpha + \beta)/2)}{r \sin r\pi}, \\ k_2 &:= \int_0^\infty u^{-1+r} \ln \frac{1 - 2u \cos \beta + u^2}{1 - 2u \cos \alpha + u^2} du = \int_0^\infty u^{-1+r} \ln \frac{1 + 2u \cos(\pi - \beta) + u^2}{1 + 2u \cos(\pi - \alpha) + u^2} du \\ &= \frac{4\pi \sin(r(\beta - \alpha)/2) \sin((r/2)(2\pi - \alpha - \beta))}{r \sin r\pi}, \\ k &= \int_{-\infty}^\infty |u|^{-1+r} \left| \ln \frac{1 + 2u \cos \alpha + u^2}{1 + 2u \cos \beta + u^2} \right| du \\ &= \int_0^\infty u^{-1+r} \ln \frac{1 + 2u \cos \alpha + u^2}{1 + 2u \cos \beta + u^2} du + \int_{-\infty}^0 (-u)^{-1+r} \ln \frac{1 + 2u \cos \beta + u^2}{1 + 2u \cos \alpha + u^2} du \\ &= k_1 + k_2 = \frac{4\pi \sin(r(\beta - \alpha)/2) \cos((r/2)(\pi - \alpha - \beta))}{r \cos(r\pi/2)}. \end{aligned} \tag{2.4}$$

The lemma is proved. □

Lemma 2.2. Define the weight functions as follow:

$$\begin{aligned} w(x) &:= \int_{-\infty}^\infty \frac{|x|^{-r}}{|y|^{1-r}} \left| \ln \frac{x^2 + 2xy \cos \alpha + y^2}{x^2 + 2xy \cos \beta + y^2} \right| dy, \\ \tilde{w}(y) &:= \int_{-\infty}^\infty \frac{|y|^r}{|x|^{1+r}} \left| \ln \frac{x^2 + 2xy \cos \alpha + y^2}{x^2 + 2xy \cos \beta + y^2} \right| dx, \end{aligned} \tag{2.5}$$

then $w(x) = \tilde{w}(y) = k = (4\pi \sin(r(\beta - \alpha)/2) \cos((r/2)(\pi - \alpha - \beta)))/[r \cos(r\pi/2)]$.

Proof. We only prove that $w(x) = k$ for $x \in (-\infty, 0)$.

Using Lemma 2.1, setting $y = ux$ and $y = -ux$,

$$\begin{aligned} w(x) &= \int_{-\infty}^0 \frac{(-x)^{-r}}{(-y)^{1-r}} \ln \frac{x^2 + 2xy \cos \alpha + y^2}{x^2 + 2xy \cos \beta + y^2} dy + \int_0^\infty \frac{(-x)^{-r}}{y^{1-r}} \ln \frac{x^2 + 2xy \cos \beta + y^2}{x^2 + 2xy \cos \alpha + y^2} dy \\ &= \int_0^\infty u^{-1+r} \ln \frac{1 + 2u \cos \alpha + u^2}{1 + 2u \cos \beta + u^2} du + \int_0^\infty u^{-1+r} \ln \frac{1 - 2u \cos \beta + u^2}{1 - 2u \cos \alpha + u^2} du = k_1 + k_2 = k. \end{aligned} \tag{2.6}$$

and the lemma is proved. □

Lemma 2.3. For $\varepsilon > 0$, and $(r - \max\{2\varepsilon/p, 2\varepsilon/q\}) \in (-1, 0)$, define both functions \tilde{f}, \tilde{g} as follows:

$$\tilde{f}(x) = \begin{cases} x^{-r-1-2\varepsilon/p}, & \text{if } x \in (1, \infty), \\ 0, & \text{if } x \in [-1, 1], \\ (-x)^{-r-1-2\varepsilon/p}, & \text{if } x \in (-\infty, -1), \end{cases} \quad \tilde{g}(x) = \begin{cases} x^{r-1-2\varepsilon/q}, & \text{if } x \in (1, \infty), \\ 0, & \text{if } x \in [-1, 1], \\ (-x)^{r-1-2\varepsilon/q}, & \text{if } x \in (-\infty, -1), \end{cases} \quad (2.7)$$

then

$$I(\varepsilon) := \varepsilon \left\{ \int_{-\infty}^{\infty} |x|^{p(r+1)-1} \tilde{f}^p(x) dx \right\}^{1/p} \left\{ \int_{-\infty}^{\infty} |x|^{q(r-1)-1} \tilde{g}^q(x) dx \right\}^{1/q} = 1, \quad (2.8)$$

$$\tilde{I}(\varepsilon) := \varepsilon \iint_{-\infty}^{\infty} \tilde{f}(x) \tilde{g}(y) \left| \ln \frac{x^2 + 2xy \cos \alpha + y^2}{x^2 + 2xy \cos \beta + y^2} \right| dx dy \rightarrow k \quad (\varepsilon \rightarrow 0^+).$$

Proof. Easily, we get the following:

$$I(\varepsilon) = \varepsilon \left\{ 2 \int_1^{\infty} x^{-1} x^{-2\varepsilon} dx \right\}^{1/p} \left\{ 2 \int_1^{\infty} x^{-1} x^{-2\varepsilon} dx \right\}^{1/q} = 1. \quad (2.9)$$

Let $y = -Y$, using $\tilde{f}(-x) = \tilde{f}(x)$, $\tilde{g}(-x) = \tilde{g}(x)$ and

$$\tilde{f}(-x) \int_{-\infty}^{\infty} \tilde{g}(y) \left| \ln \frac{x^2 - 2xy \cos \alpha + y^2}{x^2 - 2xy \cos \beta + y^2} \right| dy = \tilde{f}(x) \int_{-\infty}^{\infty} \tilde{g}(Y) \left| \ln \frac{x^2 + 2xY \cos \alpha + Y^2}{x^2 + 2xY \cos \beta + Y^2} \right| dY, \quad (2.10)$$

we have that $\tilde{f}(x) \int_{-\infty}^{\infty} \tilde{g}(y) |\ln((x^2 + 2xy \cos \alpha + y^2)/(x^2 + 2xy \cos \beta + y^2))| dy$ is an even function on x , then

$$\begin{aligned} \tilde{I}(\varepsilon) &= 2\varepsilon \int_0^{\infty} \tilde{f}(x) \left(\int_{-\infty}^{\infty} \tilde{g}(y) \left| \ln \frac{x^2 + 2xy \cos \alpha + y^2}{x^2 + 2xy \cos \beta + y^2} \right| dy \right) dx \\ &= 2\varepsilon \left[\int_1^{\infty} x^{-r-1-(2\varepsilon/p)} \left(\int_{-\infty}^{-1} (-y)^{r-1-(2\varepsilon/q)} \ln \frac{x^2 + 2xy \cos \beta + y^2}{x^2 + 2xy \cos \alpha + y^2} dy \right) dx \right. \\ &\quad \left. + \int_1^{\infty} x^{-r-1-(2\varepsilon/p)} \left(\int_1^{\infty} y^{r-1-(2\varepsilon/q)} \ln \frac{x^2 + 2xy \cos \alpha + y^2}{x^2 + 2xy \cos \beta + y^2} dy \right) dx \right] \\ &:= I_1 + I_2. \end{aligned} \quad (2.11)$$

Setting $y = tx$ then

$$\begin{aligned}
 I_1 &= 2\varepsilon \left[\int_1^\infty x^{-r-1-(2\varepsilon/p)} \left(\int_1^\infty y^{r-1-(2\varepsilon/q)} \ln \frac{x^2 - 2xy \cos \beta + y^2}{x^2 - 2xy \cos \alpha + y^2} dy \right) dx \right] \\
 &= 2\varepsilon \left[\int_1^\infty x^{-1-2\varepsilon} \left(\int_{1/x}^\infty t^{r-1-(2\varepsilon/q)} \ln \frac{1 - 2t \cos \beta + t^2}{1 - 2t \cos \alpha + t^2} dt \right) dx \right] \\
 &= 2\varepsilon \left[\int_1^\infty x^{-1-2\varepsilon} \left(\int_1^\infty t^{r-1-(2\varepsilon/q)} \ln \frac{1 - 2t \cos \beta + t^2}{1 - 2t \cos \alpha + t^2} dt \right) dx \right. \\
 &\quad \left. + \int_1^\infty x^{-1-2\varepsilon} \left(\int_{1/x}^1 t^{r-1-(2\varepsilon/q)} \ln \frac{1 - 2t \cos \beta + t^2}{1 - 2t \cos \alpha + t^2} dt \right) dx \right] \\
 &= \int_1^\infty t^{r-1-(2\varepsilon/q)} \ln \frac{1 - 2t \cos \beta + t^2}{1 - 2t \cos \alpha + t^2} dt \tag{2.12} \\
 &\quad + 2\varepsilon \int_0^1 t^{r-1-(2\varepsilon/q)} \ln \frac{1 - 2t \cos \beta + t^2}{1 - 2t \cos \alpha + t^2} \left(\int_{1/t}^\infty x^{-1-2\varepsilon} dx \right) dt \\
 &= \int_1^\infty t^{r-1-(2\varepsilon/q)} \ln \frac{1 - 2t \cos \beta + t^2}{1 - 2t \cos \alpha + t^2} dt + \int_0^1 t^{r-1+(2\varepsilon/p)} \ln \frac{1 - 2t \cos \beta + t^2}{1 - 2t \cos \alpha + t^2} dt \\
 &= \int_0^\infty t^{r-1-(2\varepsilon/q)} \ln \frac{1 - 2t \cos \beta + t^2}{1 - 2t \cos \alpha + t^2} dt + \int_0^1 \left(t^{(2\varepsilon/p)} - t^{-(2\varepsilon/q)} \right) t^{r-1} \ln \frac{1 - 2t \cos \beta + t^2}{1 - 2t \cos \alpha + t^2} dt \\
 &= \frac{4\pi \sin((r - (2\varepsilon/q))(\beta - \alpha)/2) \sin((r - (2\varepsilon/q))(2\pi - \alpha - \beta)/2)}{(r - (2\varepsilon/q)) \sin(r - (2\varepsilon/q))\pi} + \eta(\varepsilon),
 \end{aligned}$$

where $\lim_{\varepsilon \rightarrow 0^+} \eta(\varepsilon) = 0$, and we have $I_1 \rightarrow k_2$ ($\varepsilon \rightarrow 0^+$).

Similarly, $I_2 \rightarrow k_1$ ($\varepsilon \rightarrow 0^+$). The lemma is proved. □

Lemma 2.4. *If $f(x)$ is a nonnegative measurable function and $0 < \int_{-\infty}^\infty |x|^{p(1+r)-1} f^p(x) dx < \infty$, then*

$$J := \int_{-\infty}^\infty |y|^{pr-1} \left(\int_{-\infty}^\infty f(x) \left| \ln \frac{x^2 + 2xy \cos \alpha + y^2}{x^2 + 2xy \cos \beta + y^2} \right| dx \right)^p dy \leq k^p \int_{-\infty}^\infty |x|^{p(1+r)-1} f^p(x) dx. \tag{2.13}$$

Proof. By Lemma 2.2, we find that

$$\begin{aligned}
 &\left(\int_{-\infty}^\infty \left| \ln \frac{x^2 + 2xy \cos \alpha + y^2}{x^2 + 2xy \cos \beta + y^2} \right| dx \right)^p \\
 &= \left[\int_{-\infty}^\infty \left| \ln \frac{x^2 + 2xy \cos \alpha + y^2}{x^2 + 2xy \cos \beta + y^2} \right| \left(\frac{|x|^{(1+r)/q}}{|y|^{(1-r)/p}} f(x) \right) \left(\frac{|y|^{(1-r)/p}}{|x|^{(1+r)/q}} \right) dx \right]^p
 \end{aligned}$$

$$\begin{aligned}
&\leq \int_{-\infty}^{\infty} \left| \ln \frac{x^2 + 2xy \cos \alpha + y^2}{x^2 + 2xy \cos \beta + y^2} \right| \frac{|x|^{(1+r)(p-1)}}{|y|^{1-r}} f^p(x) dx \\
&\quad \times \left(\int_{-\infty}^{\infty} \left| \ln \frac{x^2 + 2xy \cos \alpha + y^2}{x^2 + 2xy \cos \beta + y^2} \right| \frac{|y|^{(1-r)(q-1)}}{|x|^{1+r}} dx \right)^{p-1} \\
&= k^{p-1} |y|^{-rp+1} \int_{-\infty}^{\infty} \left| \ln \frac{x^2 + 2xy \cos \alpha + y^2}{x^2 + 2xy \cos \beta + y^2} \right| \frac{|x|^{(1+r)(p-1)}}{|y|^{1-r}} f^p(x) dx, \\
J &\leq k^{p-1} \int_{-\infty}^{\infty} \left[\int_{-\infty}^{\infty} \left| \ln \frac{x^2 + 2xy \cos \alpha + y^2}{x^2 + 2xy \cos \beta + y^2} \right| \frac{|x|^{(1+r)(p-1)}}{|y|^{1-r}} f^p(x) dx \right] dy \\
&= k^{p-1} \int_{-\infty}^{\infty} \left[\int_{-\infty}^{\infty} \left| \ln \frac{x^2 + 2xy \cos \alpha + y^2}{x^2 + 2xy \cos \beta + y^2} \right| \frac{|x|^{(1+r)(p-1)}}{|y|^{1-r}} dy \right] f^p(x) dx \\
&= k^p \int_{-\infty}^{\infty} |x|^{p(1+r)-1} f^p(x) dx.
\end{aligned} \tag{2.14}$$

□

3. Main Results

Theorem 3.1. *If both functions, $f(x)$ and $g(x)$, are nonnegative measurable functions and satisfy $0 < \int_{-\infty}^{\infty} |x|^{p(1+r)-1} f^p(x) dx < \infty$ and $0 < \int_{-\infty}^{\infty} |x|^{q(1-r)-1} g^q(x) dx < \infty$, then*

$$\begin{aligned}
I^* &:= \iint_{-\infty}^{\infty} f(x) g(y) \left| \ln \frac{x^2 + 2xy \cos \alpha + y^2}{x^2 + 2xy \cos \beta + y^2} \right| dx dy \\
&< k \left(\int_{-\infty}^{\infty} |x|^{p(1+r)-1} f^p(x) dx \right)^{1/p} \left(\int_{-\infty}^{\infty} |x|^{q(1-r)-1} g^q(x) dx \right)^{1/q},
\end{aligned} \tag{3.1}$$

$$\begin{aligned}
J &= \int_{-\infty}^{\infty} |y|^{pr-1} \left(\int_{-\infty}^{\infty} f(x) \left| \ln \frac{x^2 + 2xy \cos \alpha + y^2}{x^2 + 2xy \cos \beta + y^2} \right| dx \right)^p dy \\
&< k^p \int_{-\infty}^{\infty} |x|^{p(1+r)-1} f^p(x) dx.
\end{aligned} \tag{3.2}$$

Inequalities (3.1) and (3.2) are equivalent, and where the constant factors k and k^p are the best possibles.

Proof. If (2.13) takes the form of equality for some $y \in (-\infty, 0) \cup (0, \infty)$, then there exists constants M and N , such that they are not all zero, and

$$M \frac{|x|^{(1+r)(p-1)}}{|y|^{1-r}} f^p(x) = N \frac{|y|^{(1-r)(q-1)}}{|x|^{1+r}} \quad \text{a.e. in } (-\infty, \infty) \times (-\infty, \infty). \tag{3.3}$$

Hence, there exists a constant C , such that

$$M|x|^{(1+r)p} f^p(x) = N|y|^{(1-r)q} = C \quad \text{a.e. in } (-\infty, \infty) \times (-\infty, \infty). \tag{3.4}$$

We claim that $M = 0$. In fact, if $M \neq 0$, then $|x|^{p(1+r)-1} f^p(x) = C/(M|x|^{-1})$ a.e. in $(-\infty, \infty)$ which contradicts the fact that $0 < \int_{-\infty}^{\infty} |x|^{p(1+r)-1} f^p(x) dx < \infty$. In the same way, we claim that $N = 0$. This is too a contradiction and hence by (2.13), we have (3.2).

By Hölder’s inequality with weight [22] and (3.2), we have the following:

$$\begin{aligned} I^* &= \int_{-\infty}^{\infty} \left[|y|^{-1+r+(1/q)} \int_{-\infty}^{\infty} f(x) \left| \ln \frac{x^2 + 2xy \cos \alpha + y^2}{x^2 + 2xy \cos \beta + y^2} \right| dx \right] \left[|y|^{1-r-(1/q)} g(y) \right] dy \\ &\leq (J)^{1/p} \left(\int_{-\infty}^{\infty} |y|^{q(1-r)-1} g^q(y) dy \right)^{1/q}. \end{aligned} \tag{3.5}$$

Using (3.2), we have (3.1).

Setting $g(y) = |y|^{r-1} (\int_{-\infty}^{\infty} f(x) |\ln((x^2 + 2xy \cos \alpha + y^2)/(x^2 + 2xy \cos \beta + y^2)) dx|)^{p-1}$, then $J = \int_{-\infty}^{\infty} |y|^{q(1-r)-1} g^q(y) dy$ by (2.13), we have $J < \infty$. If $J = 0$ then (3.2) is proved. If $0 < J < \infty$, by (3.1), we obtain

$$\begin{aligned} 0 &< \int_{-\infty}^{\infty} |y|^{q(1-r)-1} g^q(y) dy = J = I^* \\ &< k \left(\int_{-\infty}^{\infty} |x|^{p(1+r)-1} f^p(x) dx \right)^{1/p} \left(\int_{-\infty}^{\infty} |x|^{q(1-r)-1} g^q(x) dx \right)^{1/q}, \\ \left(\int_{-\infty}^{\infty} |x|^{q(1-r)-1} g^q(x) dx \right)^{1/p} &= J^{1/p} < k \left(\int_{-\infty}^{\infty} |x|^{p(1+r)-1} f^p(x) dx \right)^{1/p}. \end{aligned} \tag{3.6}$$

Inequalities (3.1) and (3.2) are equivalent.

If the constant factor k in (3.1) is not the best possible, then there exists a positive h (with $h < k$), such that

$$\begin{aligned} &\iint_{-\infty}^{\infty} f(x) g(y) \left| \ln \frac{x^2 + 2xy \cos \alpha + y^2}{x^2 + 2xy \cos \beta + y^2} \right| dx dy \\ &< h \left(\int_{-\infty}^{\infty} |x|^{p(1+r)-1} f^p(x) dx \right)^{1/p} \left(\int_{-\infty}^{\infty} |x|^{q(1-r)-1} g^q(x) dx \right)^{1/q}. \end{aligned} \tag{3.7}$$

For $\varepsilon > 0$, by (3.7), using Lemma 2.3, we have

$$k + o(1) < \varepsilon h \left(\int_{-\infty}^{\infty} |x|^{-1} \tilde{f}^p(x) dx \right)^{1/p} \left(\int_{-\infty}^{\infty} |x|^{-1} \tilde{g}^q(x) dx \right)^{1/q} = k. \tag{3.8}$$

Hence, we find $k + o(1) < h$. For $\varepsilon \rightarrow 0^+$, it follows that $k \leq h$, which contradicts the fact that $h < k$. Hence the constant k in (3.1) is the best possible.

Thus we complete the proof of the theorem. □

Remark 3.2. For $\alpha = \pi/4, \beta = \pi/3$ in (3.1), we have the following particular result:

$$\begin{aligned} & \iint_{-\infty}^{\infty} f(x)g(y) \left| \ln \frac{x^2 + \sqrt{2}xy + y^2}{x^2 + xy + y^2} \right| dx dy \\ & < \frac{4\pi \sin(\pi r/24) \sin(5\pi r/24)}{r \sin(\pi r/2)} \left(\int_{-\infty}^{\infty} |x|^{p(1+r)-1} f^p(x) dx \right)^{1/p} \left(\int_{-\infty}^{\infty} |x|^{q(1-r)-1} g^q(x) dx \right)^{1/q}. \end{aligned} \quad (3.9)$$

References

- [1] G. H. Hardy, J. E. Littlewood, and G. Pólya, *Inequalities*, Cambridge University Press, London, UK, 1952.
- [2] G. H. Hardy, "Note on a theorem of Hilbert concerning series of positive terms," *Proceedings of the London Mathematical Society*, vol. 23, no. 2, pp. 45–46, 1925.
- [3] D. S. Mitrinović, J. E. Pečarić, and A. M. Fink, *Inequalities Involving Functions and Their Integrals and Derivatives*, vol. 53, Kluwer Academic, Boston, Mass, USA, 1991.
- [4] Z. Xie and Z. Zeng, "A Hilbert-type integral inequality whose kernel is a homogeneous form of degree -3 ," *Journal of Mathematical Analysis and Applications*, vol. 339, no. 1, pp. 324–331, 2008.
- [5] Z. Xie and Z. Zeng, "A Hilbert-type integral inequality with a non-homogeneous form and a best constant factor," *Advances and Applications in Mathematical Science*, vol. 3, no. 1, pp. 61–71, 2010.
- [6] Z. Xie and Z. Zeng, "The Hilbert-type integral inequality with the system kernel of $-\lambda$ degree homogeneous form," *Kyungpook Mathematical Journal*, vol. 50, pp. 297–306, 2010.
- [7] B. Yang, "A new Hilbert-type integral inequality with some parameters," *Journal of Jilin University*, vol. 46, no. 6, pp. 1085–1090, 2008.
- [8] Z. Xie and B. Yang, "A new Hilbert-type integral inequality with some parameters and its reverse," *Kyungpook Mathematical Journal*, vol. 48, no. 1, pp. 93–100, 2008.
- [9] Z. Xie, "A new Hilbert-type inequality with the kernel of -3μ -homogeneous," *Journal of Jilin University*, vol. 45, no. 3, pp. 369–373, 2007.
- [10] Z. Xie and J. Murong, "A reverse Hilbert-type inequality with some parameters," *Journal of Jilin University*, vol. 46, no. 4, pp. 665–669, 2008.
- [11] Z. Xie, "A new reverse Hilbert-type inequality with a best constant factor," *Journal of Mathematical Analysis and Applications*, vol. 343, no. 2, pp. 1154–1160, 2008.
- [12] B. Yang, "A Hilbert-type inequality with a mixed kernel and extensions," *Journal of Sichuan Normal University*, vol. 31, no. 3, pp. 281–284, 2008.
- [13] Z. Xie and Z. Zeng, "A Hilbert-type inequality with parameters," *Natural Science Journal of Xiangtan University*, vol. 29, no. 3, pp. 24–28, 2007.
- [14] Z. Zeng and Z. Xie, "A Hilbert's inequality with a best constant factor," *Journal of Inequalities and Applications*, vol. 2009, Article ID 820176, 8 pages, 2009.
- [15] B. Yang, "A bilinear inequality with a -2 -order homogeneous kernel," *Journal of Xiamen University*, vol. 45, no. 6, pp. 752–755, 2006.
- [16] B. Yang, "On Hilbert's inequality with some parameters," *Acta Mathematica Sinica*, vol. 49, no. 5, pp. 1121–1126, 2006.
- [17] I. Brnetić and J. Pečarić, "Generalization of Hilbert's integral inequality," *Mathematical Inequalities and Application*, vol. 7, no. 2, pp. 199–205, 2004.
- [18] I. Brnetić, M. Krnić, and J. Pečarić, "Multiple Hilbert and Hardy-Hilbert inequalities with non-conjugate parameters," *Bulletin of the Australian Mathematical Society*, vol. 71, no. 3, pp. 447–457, 2005.
- [19] Z. Xie and F. M. Zhou, "A generalization of a Hilbert-type inequality with the best constant factor," *Journal of Sichuan Normal University*, vol. 32, no. 5, pp. 626–629, 2009.
- [20] Z. Xie and X. Liu, "A new Hilbert-type integral inequality and its reverse," *Journal of Henan University*, vol. 39, no. 1, pp. 10–13, 2009.
- [21] Z. Xie and B. L. Fu, "A new Hilbert-type integral inequality with a best constant factor," *Journal of Wuhan University*, vol. 55, no. 6, pp. 637–640, 2009.
- [22] J. Kang, *Applied Inequalities*, Shangdong Science and Technology Press, Jinan, China, 2004.