

# Degree bounds for type-A weight rings and Gelfand–Tsetlin semigroups

Benjamin J. Howard · Tyrrell B. McAllister

Received: 6 November 2009 / Accepted: 17 November 2010 / Published online: 4 December 2010  
© Springer Science+Business Media, LLC 2010

**Abstract** A weight ring in type A is the coordinate ring of the GIT quotient of the variety of flags in  $\mathbb{C}^n$  modulo a twisted action of the maximal torus in  $\mathrm{SL}(n, \mathbb{C})$ . We show that any weight ring in type A is generated by elements of degree strictly less than the Krull dimension, which is at worst  $O(n^2)$ . On the other hand, we show that the associated semigroup of Gelfand–Tsetlin patterns can have an essential generator of degree exponential in  $n$ .

**Keywords** Weight ring · Weight variety · Cohen–Macaulay ring · Toric degeneration · Gelfand–Tsetlin pattern

## 1 Introduction

Given a pair  $\lambda, \mu$  of weights for  $\mathrm{SL}_n(\mathbb{C})$  with  $\lambda$  dominant, let  $V_\lambda[\mu]$  denote the  $\mu$ -isotropic component of the irreducible representation  $V_\lambda$  with highest weight  $\lambda$ . The *weight ring*  $R(\lambda, \mu)$  is the graded ring  $\bigoplus_{N=0}^{\infty} V_{N\lambda}[N\mu]$ . We define the *weight variety*  $W(\lambda, \mu)$  as

$$W(\lambda, \mu) := \mathrm{Proj} R(\lambda, \mu).$$

---

Second author supported by the Netherlands Organization for Scientific Research (NWO) Mathematics Cluster DIAMANT.

B.J. Howard  
Mathematics Department, University of Michigan, Ann Arbor, MI 48109, USA  
e-mail: [howardbj@umich.edu](mailto:howardbj@umich.edu)

T.B. McAllister (✉)  
Department of Mathematics, University of Wyoming, Laramie, WY 82071-2000, USA  
e-mail: [tmcallis@uwyo.edu](mailto:tmcallis@uwyo.edu)

Alternatively, the weight ring is the projective coordinate ring of the GIT quotient of the flag variety modulo the  $\mu$ -twisted action of the maximal torus  $\mathbb{T}$  in  $\mathrm{SL}_n(\mathbb{C})$ .<sup>1</sup>

Our first theorem (Theorem 3.9 below) is that  $R(\lambda, \mu)$  is generated in degree strictly less than the Krull dimension of  $R(\lambda, \mu)$ , provided that the degree-one piece  $V_\lambda[\mu]$  is nonzero. The basic idea behind the proof is to show that the degree-one piece contains a system of parameters and that the  $a$ -invariant of  $R(\lambda, \mu)$  is negative.<sup>2</sup> (The  $a$ -invariant is the degree of the Hilbert series, which is a rational function.) The theorem then follows from the fact that  $R(\lambda, \mu)$  is Cohen–Macaulay.

It is well known (cf. [8, 9, 14]) that  $R(\lambda, \mu)$  has a flat degeneration to the semigroup algebra  $R'(\lambda, \mu)$  of the semigroup of Gelfand–Tsetlin patterns associated to semistandard tableaux of shape  $m\lambda$  and content  $m\mu$  for  $m \geq 0$ . In particular, the ring  $R'(\lambda, \mu)$  is the graded ring associated to a filtration of  $R(\lambda, \mu)$  by natural numbers. Generators for  $R'(\lambda, \mu)$  can be lifted to generators of  $R(\lambda, \mu)$ , so one might hope that  $R'(\lambda, \mu)$  is relatively simple. Unfortunately, we find pairs  $\lambda, \mu$  for which  $R'(\lambda, \mu)$  has essential generators of degree exponential in  $n$ .

Our second main result (Theorem 5.2 below) is that, in the case where  $n = 3k$  is a multiple of 3, the semigroup algebra  $R'(k\varpi_3, 0)$  has an essential generator of degree approximately  $(\sqrt{2})^n$ . This is in striking contrast to the upper bound of  $2n - 8$  for  $R(k\varpi_3, 0)$ , which follows from our first theorem (since the Krull dimension of  $R(k\varpi_3, 0)$  is  $2n - 7$ ). This case is particularly interesting because, via the Gelfand–MacPherson correspondence,  $R(k\varpi_3, 0)$  is the moduli space of  $n$ -tuples of points in the projective plane. This is a remarkable example of how a semigroup algebra produced by a promising toric degeneration can fail to serve as an effective proxy for the original ring.

Our motivation for studying the semigroup of Gelfand–Tsetlin patterns was to imitate the method of [12], which considered the case of  $n$  points on the projective line. Here one takes  $\lambda$  to be a multiple of the second fundamental weight  $\varpi_2$ . It was shown in [12] that the associated semigroup of Gelfand–Tsetlin patterns is generated in degree  $\leq 2$ . We had hoped to use the same method in the case of  $n$  points in the projective plane, but the second theorem indicates why this is not the right approach. However, there might still be another toric degeneration, perhaps among those discovered by Caldero [2], that yields a bound better than the one in Theorem 3.9

## 2 A description of the weight ring $R(\lambda, \mu)$

In this section, we give an explicit description of  $R(\lambda, \mu)$ . Let  $n \geq 2$ , and let  $B$  denote the Borel subgroup of  $\mathrm{SL}_n(\mathbb{C})$  consisting of the upper-triangular matrices in  $\mathrm{SL}_n(\mathbb{C})$ . Fix a nontrivial dominant weight  $\lambda$  of  $\mathrm{SL}_n(\mathbb{C})$ . We represent  $\lambda$  as a *partition*, i.e., as

<sup>1</sup>Weight varieties for arbitrary reductive Lie groups (not just those of type A) were studied by A. Knutson in his Ph.D. thesis [13]. Knutson also studied the symplectic geometry of these spaces.

<sup>2</sup>The  $a$ -invariant is negative in all types (see Remark 3.8 below); however, for types other than type A, the degree-one piece sometimes fails to contain a system of parameters. This condition is equivalent to the condition that all semistable flags lie in the supports of degree-one  $\mathbb{T}$ -invariants; see [11] for a counterexample in  $G = \mathrm{SO}_5(\mathbb{C})$ .

a weakly decreasing sequence  $(\lambda_1, \dots, \lambda_n)$  of nonnegative integers with  $\lambda_1 \geq 1$  and  $\lambda_n = 0$ . Let  $\mu$  be a weight of  $\mathrm{SL}_n(\mathbb{C})$  such that  $V_\lambda[\mu]$  is nonzero. Thus  $\mu$  may be expressed as a sequence  $(\mu_1, \dots, \mu_n)$  of nonnegative integers such that  $\sum_{i=1}^n \mu_i = \sum_{i=1}^n \lambda_i$ .

By the Borel–Weil construction, the irreducible representation with highest weight  $\lambda$  is the finite-dimensional vector space

$$V_\lambda = \{ \text{holomorphic } f : \mathrm{SL}_n(\mathbb{C}) \rightarrow \mathbb{C} \mid f(gb) = e^\lambda(b)f(g) \\ \text{for all } g \in \mathrm{SL}_n(\mathbb{C}) \text{ and } b \in B \},$$

where  $e^\lambda(b) := \prod_{i=1}^n b_{ii}^{\lambda_i}$  for  $b = (b_{ij})_{1 \leq i, j \leq n} \in B$ . The action of  $\mathrm{SL}_n(\mathbb{C})$  on  $V_\lambda$  is given by  $(g \cdot f)(h) = f(g^{-1}h)$  for  $g, h \in \mathrm{SL}_n(\mathbb{C})$ . We define

$$R(\lambda) := \bigoplus_{N=0}^{\infty} V_{N\lambda}.$$

Multiplication in  $R(\lambda)$  is the usual multiplication of functions  $\mathrm{SL}_n(\mathbb{C}) \rightarrow \mathbb{C}$ .

*Remark 2.1* As we will review in the next section, the dominant weight  $\lambda$  determines a line bundle  $L_\lambda \rightarrow \mathrm{SL}_n(\mathbb{C})/B$  such that the space  $\Gamma(\mathrm{SL}_n(\mathbb{C})/B, L_\lambda)$  of sections is isomorphic to  $V_\lambda$ . The multiplication in  $R(\lambda)$  coincides with multiplication of the corresponding sections. The ring  $R(\lambda)$  is the coordinate ring of the partial flag variety corresponding to the dominant weight  $\lambda$ .

We define

$$V_\lambda[\mu] := \{ f \in V_\lambda \mid f(tg) = e^\mu(t)f(g) \text{ for all } t \in \mathbb{T}, g \in \mathrm{SL}_n(\mathbb{C}) \}.$$

We now define a  $\mu$ -twisted action of  $\mathbb{T} \subset \mathrm{SL}_n(\mathbb{C})$  on  $R(\lambda)$ . For  $f \in V_{N\lambda} = R(\lambda)_N$  of degree  $N$ , the action of  $t \in \mathbb{T}$  on  $f$  is given by

$$(t \cdot f)(g) := e^{N\mu}(t)f(t^{-1}g).$$

Relative to this twisted action, the  $\mathbb{T}$ -invariant subring of  $R(\lambda)$  is exactly


$$R(\lambda, \mu) := \bigoplus_{N=0}^{\infty} V_{N\lambda}[N\mu].$$

*Remark 2.2* There is a unique  $\mathrm{SL}_n(\mathbb{C})$ -linearization of  $L_\lambda$ . This defines a canonical  $\mathbb{T}$ -linearization of  $L_\lambda$  by restriction to  $\mathbb{T} \hookrightarrow \mathrm{SL}_n(\mathbb{C})$ . The above action of  $\mathbb{T}$  coincides with the canonical  $\mathbb{T}$ -linearization twisted by  $\mu$ .

A fundamental fact from the representation theory of  $\mathrm{SL}_n(\mathbb{C})$  is that  $V_\lambda$  has a basis indexed by semistandard tableaux of shape  $\lambda$ . Furthermore, the elements in this basis indexed by semistandard tableaux with content  $\mu$  form a basis of  $V_\lambda[\mu]$ . We briefly review the construction.

A *Young diagram* of shape  $\lambda$  is a left-justified arrangement of  $\lambda_1 + \dots + \lambda_n$  boxes with  $\lambda_i$  boxes in the  $i$ th row. For example, if  $\lambda = (3, 3, 2, 1, 1, 0)$ , then the Young

diagram of shape  $\lambda$  is



A *semistandard tableaux* of shape  $\lambda$  is a filling of each box in a Young diagram of shape  $\lambda$  with a number from 1 through  $n$  such that the rows are weakly increasing and the columns are strictly increasing. For example, if  $\lambda = (3, 3, 2, 1, 0)$ , then

$$\tau = \begin{array}{|c|c|c|} \hline 1 & 1 & 5 \\ \hline 2 & 4 & 6 \\ \hline 3 & 5 & \\ \hline 5 & & \\ \hline 6 & & \\ \hline \end{array}$$

is a semistandard tableau of shape  $\lambda$ .

Such a tableau  $\tau$  determines a basis vector  $b_\tau \in V_\lambda$  as follows. Write  $\text{len}(I)$  for the length of a column  $I$  of  $\tau$ . We identify  $I$  with the  $\text{len}(I)$ -tuple of its entries, read from top to bottom. If  $I = (i_1, \dots, i_{\text{len}(I)})$ , let  $\det_I : \text{SL}_n(\mathbb{C}) \rightarrow \mathbb{C}$  be the function that returns the determinant of the  $\text{len}(I) \times \text{len}(I)$  submatrix of entries in rows  $i_1, \dots, i_{\text{len}(I)}$  and columns  $1, 2, \dots, \text{len}(I)$ . The basis vector  $b_\tau$  is then defined by

$$b_\tau := \prod_{\text{columns } I \text{ of } \tau} \det_I.$$

Hence, in the example above,  $b_\tau = \det_{1,2,3,5,6} \det_{1,4,5} \det_{5,6}$ .

We can also describe the  $\mathbb{T}$ -isotropic subspace  $V_\lambda[\mu]$  in terms of semistandard tableaux. The *content* of a tableau  $\tau$  is  $\mu = (\mu_1, \dots, \mu_n)$  if  $\mu_i$  is the number of boxes in  $\tau$  that contain the number  $i$  for  $1 \leq i \leq n$ . The subspace  $V_\lambda[\mu]$  is the span of the  $b_\tau$  such that  $\tau$  has shape  $\lambda$  and content  $\mu$ .

### 3 The first theorem: generators of $R(\lambda, \mu)$

We will derive an upper bound on the degree in which  $R(\lambda, \mu)$  is generated. We generally follow the method of [16] (also explained in [4]). The idea is to find a homogeneous system of parameters, together with an upper bound on the  $a$ -invariant of the ring; this yields an upper bound on the degree of a generating set.

We begin by referencing a result of [11] and showing why this implies the existence of a system of parameters in degree one. First we must introduce the notion of semistability. Recall that  $B$  is the Borel subgroup of  $\text{SL}_n(\mathbb{C})$ , and  $\text{SL}_n(\mathbb{C})/B$  is the flag variety. A function  $f \in V_\lambda$  defines a section of the line bundle  $L_\lambda := \text{SL}_n(\mathbb{C}) \times_B \mathbb{C} \rightarrow \text{SL}_n(\mathbb{C})/B$ , where  $\text{SL}_n(\mathbb{C}) \times_B \mathbb{C}$  denotes the quotient of  $\text{SL}_n(\mathbb{C}) \times \mathbb{C}$  by the equivalence relation  $(gb, e^\lambda(b)z) \sim (g, z)$ . The projection  $L_\lambda \rightarrow \text{SL}_n(\mathbb{C})/B$  is given by sending the equivalence class of  $(g, z)$  to  $gB$ . We define the  $\mu$ -twisted

linearization of  $\mathbb{T}$  on  $L_\lambda$  by  $t \cdot (g, z) := (t^{-1}g, e^\mu(t)z)$ . Given  $f \in V_\lambda$ , we define a global section  $s_f$  of  $L_\lambda$  by  $s_f(gB) = (g, f(g))$ . The map  $f \mapsto s_f$  is an isomorphism  $V_\lambda \cong \Gamma(\mathrm{SL}_n(\mathbb{C})/B, L_\lambda)$ . The  $\mu$ -twisted torus action on  $L_\lambda$  determines an action on global sections, which coincides with the  $\mu$ -twisted action on  $V_\lambda$  that we defined earlier.

A flag  $gB$  is *semistable* if, for some positive integer  $N$ , there is a global section  $s$  of  $L_{N\lambda}$  with  $s(gB) \neq 0$  that is invariant under the  $\mu$ -twisted action of  $\mathbb{T}$ . That is,  $gB$  is semistable if and only if there is an  $N > 0$  and an  $f \in V_{N\lambda}[N\mu]$  such that  $f(g) \neq 0$ . It was shown in [11] that we may take  $N = 1$ . That is,  $gB$  is semistable if and only if there exists an  $f \in V_\lambda[\mu]$  such that  $f(g) \neq 0$ . We will use this fact to show that there is a system of parameters within  $V_\lambda[\mu]$  for  $R(\lambda, \mu)$ .

We now recall some basic facts from commutative algebra. Our main references on Cohen–Macaulay rings and modules are [1, 15]. Let  $\mathbf{k}$  be an algebraically closed field. Suppose that  $A$  is a  $\mathbb{Z}_{\geq 0}$ -graded finitely-generated  $\mathbf{k}$ -algebra with  $A_0 = \mathbf{k}$ . Let  $\mathfrak{m}$  denote the graded ideal generated by the positive degree homogeneous elements of  $A$ . Then  $\mathfrak{m}$  is the unique graded ideal such that all other graded ideals are contained within it. A *homogeneous system of parameters* for  $A$  is a set of homogeneous elements  $x_1, \dots, x_s$  such that  $s$  is the Krull dimension of  $A$  and the ideal  $(x_1, \dots, x_s)$  is  $\mathfrak{m}$ -primary. By [1, Theorem 1.5.17] we have that  $x_1, \dots, x_s$  is a homogeneous system of parameters if and only if  $A$  is an integral extension of the subalgebra  $\mathbf{k}[x_1, \dots, x_s]$ , and that this is the case if and only if  $A$  is a finitely-generated  $\mathbf{k}[x_1, \dots, x_s]$ -module.

Let the *null cone*  $\mathcal{N}$  be the subvariety of points in  $\mathrm{Spec} R(\lambda)$  at which all positive-degree homogeneous elements of  $R(\lambda, \mu)$  vanish. The result of [11] translates into the following:

**Proposition 3.1** *The elements of  $V_\lambda[\mu]$  suffice to cut out the null cone set-theoretically. That is,  $\mathcal{N}$  is exactly the set of points at which all elements of  $V_\lambda[\mu]$  vanish.*

Now suppose that  $I \subset R(\lambda)$  is the ideal of elements vanishing on the null cone. By the above proposition,  $I$  is the radical closure in  $R(\lambda)$  of the ideal generated by  $V_\lambda[\mu] \subset R(\lambda)$ . Recall that  $R(\lambda, \mu)$  is the ring of polynomials in  $R(\lambda)$  that are invariant under the  $\mu$ -twisted action of  $\mathbb{T}$ . Since  $\mathbb{T}$  is linearly reductive, there is a canonical projection  $\pi : R(\lambda) \rightarrow R(\lambda, \mu)$ , called the *Reynolds operator*, which is  $R(\lambda, \mu)$ -linear. Following Hilbert (cf. [4, Proposition 3.1]), we have the following result.

**Proposition 3.2** *The invariant ring  $R(\lambda, \mu)$  is a finitely generated module over the subalgebra generated by  $V_\lambda[\mu] \subset R(\lambda, \mu)$ .*

*Proof* Let  $J$  and  $S$  be the ideal and subalgebra, respectively, generated by  $V_\lambda[\mu]$  in  $R(\lambda)$ . Then, since  $I = \mathrm{Rad}(J)$ , we have that  $I^m \subset J$  for some  $m > 0$ . Since  $\mathbb{T}$  is linearly reductive, the invariant ring  $R(\lambda, \mu)$  is finitely generated. Thus, there exist homogeneous  $y_1, \dots, y_t \in R(\lambda, \mu)$  such that  $y_1, \dots, y_t$  generate  $R(\lambda, \mu)$ . Suppose that  $h_1, \dots, h_\ell$  span  $V_\lambda[\mu]$ . We have that each  $y_i^m$  belongs to the ideal  $J$ , and so  $y_i^m = \sum_{j=1}^\ell f_j h_j$  for some homogeneous  $f_j \in R(\lambda)$ . Now we apply the Reynolds’s

operator  $\pi$  to obtain  $y_i^m = \sum_{j=1}^{\ell} \pi(f_j)h_j$ . Each coefficient  $\pi(f_j)$  is a homogeneous invariant of degree less than  $y_i^m$ . It follows that  $R(\lambda, \mu)$  is generated as an  $S$ -module by monomials  $\mathbf{m} = \prod_{i=1}^t y_i^{e_i}$ , where each  $e_i < m$ . There are only a finite number of such monomials, proving the claim.  $\square$

We now have the following (cf. [4, Proposition 3.2]):

**Proposition 3.3** *The degree-one piece  $V_\lambda[\mu]$  of  $R(\lambda, \mu)$  contains a homogeneous system of parameters.*

*Proof* We already know from the previous proposition that  $R(\lambda, \mu)$  is a finitely generated module over the subalgebra  $S$  generated by its degree-one piece  $V_\lambda[\mu]$ . Let  $s$  be the Krull dimension of  $R(\lambda, \mu)$ . Note that the Krull dimension of the subring  $S$  of  $R(\lambda, \mu)$  is also equal to  $s$ , since  $R(\lambda, \mu)$  is a finitely generated module over  $S$ .

By Noether normalization [7, Theorem 13.3], there exist  $f_1, \dots, f_s \in V_\lambda[\mu]$  that are algebraically independent and such that  $S$  is finitely generated over the polynomial subring  $\mathbb{C}[f_1, \dots, f_s] \subset S$ . (The  $f_i$  may be taken to be  $s$  generic linear combinations of a basis of  $V_\lambda[\mu]$ .)

Since furthermore  $R(\lambda, \mu)$  is finitely generated over  $S$ , we have that  $R(\lambda, \mu)$  is finitely generated over  $\mathbb{C}[f_1, \dots, f_s]$ . Finally, we may deduce from the Cayley–Hamilton theorem [7, Theorem 4.3 and Corollary 4.6] that  $R(\lambda, \mu)$  is an integral extension of  $\mathbb{C}[f_1, \dots, f_s]$ .  $\square$

**Proposition 3.4** *The weight ring  $R(\lambda, \mu)$  is Cohen–Macaulay.*

*Proof* The argument is the same as that given in [15, Corollary 14.25] to show that  $R(\lambda)$  is Cohen–Macaulay. We know that  $R(\lambda, \mu)$  has a Gröbner degeneration to a semigroup algebra of Gelfand–Tsetlin patterns (see Sect. 4). Such semigroup algebras are the invariant subrings of polynomial rings by the action of a torus (cf. [5]). By the theorem of Hochster [10], the subring of torus invariants in a polynomial ring is Cohen–Macaulay. A general principle regarding Gröbner degenerations is that any good property of the special fiber is shared by the general fiber. This is true in particular for the Cohen–Macaulay property [15, Corollary 8.31].  $\square$

**Proposition 3.5** *If  $f_1, \dots, f_s$  is a homogeneous system of parameters for  $R(\lambda, \mu)$ , then  $R(\lambda, \mu)$  is a free  $\mathbb{C}[f_1, \dots, f_s]$ -module.*

*Proof* Since  $f_1, \dots, f_s$  are algebraically independent, the ring  $\mathbb{C}[f_1, \dots, f_s]$  is regular, and so this proposition follows from [1, Proposition 2.2.11].  $\square$

For a graded module  $M$ , let  $H(M; t) := \sum_{d=0}^{\infty} \dim(M_d)t^d$  denote the *Hilbert series* of  $M$ . It is well known that, if  $M$  is finitely generated, then  $H(M; t)$  is a rational function in  $t$ . Let  $a(M)$  be the degree of  $H(M; t)$  as a rational function. The number  $a(M)$  is called the *a-invariant* of  $M$ .

Fix a homogeneous system of parameters  $f_1, \dots, f_s \in V_\lambda[\mu]$  for  $R(\lambda, \mu)$ . Let  $S = \mathbb{C}[f_1, \dots, f_s]$  be the subalgebra generated by the  $f_i$ . For brevity of notation, we

will write  $R := R(\lambda, \mu)$ . Let  $\mathbf{f}$  denote the  $s$ -tuple  $f_1, \dots, f_s$ . By Theorems 13.37(5) and 13.37(6) of [15],  $R$  is a free  $S$ -module, and

$$H(R/\mathbf{f}R; t) = H(R; t)(1 - t)^s.$$

But we can easily compute  $H(R/\mathbf{f}R; t)$ . Suppose that  $R = Sy_1 \oplus \dots \oplus Sy_m$ . Let  $k := \max_j(\deg y_j)$ . Now,  $H(R/\mathbf{f}R; t)$  is the polynomial  $p(t) = \sum_{i=0}^k h_d t^d$ , where  $h_d$  is the number of  $y_j$  such that  $\deg y_j = d$ . Therefore, we have proved the following.

**Proposition 3.6** *The ring  $R(\lambda, \mu)$  is generated in degree  $\leq k = \dim R(\lambda, \mu) + a(R(\lambda, \mu))$ .*

**Proposition 3.7** *The  $a$ -invariant of  $R(\lambda, \mu)$  is negative.*

*Proof* Let  $R := R(\lambda, \mu)$ . The dimension of the  $d$ th graded piece  $R_d$  of  $R$  is equal to the number of semistandard tableaux of shape  $d\lambda$  and content  $d\mu$ ; this coincides with the number of integer lattice points in the  $d$ th dilate of the rational polytope  $\text{GT}(\lambda, \mu)$  (see Definition 4.1 below). As a result of the theory of lattice point enumeration for rational polytopes (see, e.g., [17, Chap. 4]), we may conclude that the Hilbert series  $H(R; t) = \sum_{d=0}^{\infty} f(d)t^d$  is a rational function of negative degree. □

*Remark 3.8* In fact, in all types, given a pair of weights  $\lambda, \mu$  with  $\lambda$  dominant, the dimension of the  $d\mu$ -weight space in the irreducible representation  $V_{d\lambda}$  with highest weight  $d\lambda$  equals the number of integer lattice points in the  $d$ th dilate of a certain polytope (for example, the string polytope associated with the reduced word for the longest Weyl element). And so, in all types, the  $a$ -invariants of weight rings are negative.

The above propositions imply our first theorem:

**Theorem 3.9** *The algebra  $R(\lambda, \mu)$  is generated in degree strictly less than the Krull dimension of  $R(\lambda, \mu)$ .*

*Proof* This follows immediately from Propositions 3.6 and 3.7. □

Finally, we point out that the Krull dimension of  $R(\lambda, \mu)$  is one more than the dimension of the GIT quotient of the flag variety by  $\mathbb{T}$ . This is at most the dimension of the flag variety itself, which is  $n(n - 1)/2$ . In the case of  $n$  points in projective space  $\mathbb{P}^{m-1}$ , where  $\lambda$  is a multiple of the  $m$ th fundamental weight  $\varpi_m$  for  $\text{SL}_n(\mathbb{C})$ , the Krull dimension of  $R(\lambda, \mu)$  is at most  $n(m - 1) - (m^2 - 1) + 1$ .

#### 4 The toric degeneration to Gelfand–Tsetlin patterns

A *Gelfand–Tsetlin pattern*, or GT pattern, is a triangular array  $\mathbf{x} = (x_{ij})_{1 \leq i \leq j \leq n}$  of real numbers satisfying the *interlacing inequalities*  $x_{i,j+1} \geq x_{ij} \geq x_{i+1,j+1}$ . We ex-





the sum is over semistandard tableaux, then the term  $b_{\phi^{-1}(\mathbf{x}(\tau_1)+\mathbf{x}(\tau_2))}$  appears on the right-hand side with coefficient equal to 1. Furthermore, all other terms  $c^\tau b_\tau$  have strictly smaller filtration level. Thus, in  $R'(\lambda, \mu)$ , the multiplication rule becomes

$$b'_{\tau_1} b'_{\tau_2} = b'_{\phi^{-1}(\mathbf{x}(\tau_1)+\mathbf{x}(\tau_2))}.$$

Therefore  $R'(\lambda, \mu)$  is isomorphic to  $\mathbb{C}[S(\lambda, \mu)]$ , the semigroup algebra of GT patterns under addition of patterns.

Given an  $m$ -tuple of rational numbers  $q_1, \dots, q_m$ , define  $\text{den}(q_1, \dots, q_m)$  to be the least positive integer  $N$  such that  $Nq_i \in \mathbb{Z}$  for each  $i$ ,  $1 \leq i \leq m$ . We call this the *denominator* of the  $m$ -tuple. Now, if some vertex  $\mathbf{x}$  of the polytope  $\text{GT}(\lambda, \mu)$  has denominator  $N > 1$ , then the integer point  $N\mathbf{x}$  is an essential generator of the semigroup  $\mathbb{C}[S(\lambda, \mu)]$ , since  $\mathbf{x}$  cannot be written as a sum of other integral patterns in  $S(\lambda, \mu)$ . In the next section we show the existence of such a vertex with large denominator for the case where  $n$  is a multiple of 3,  $\lambda = \frac{n}{3}\varpi_3$ , and  $\mu = (1, 1, \dots, 1)$ .

### 5 The second theorem: $3k$ points on $\mathbb{P}^2$ and a nasty GT pattern

Suppose that  $n = 3k$ , where  $k \geq 2$  is an integer. Let  $\lambda = k\varpi_3$  be a multiple of the third fundamental weight for  $\text{SL}_n(\mathbb{C})$ . Thus, as a partition,  $\lambda = (k, k, k, 0, \dots, 0) \in \mathbb{R}^{3k}$ . Now let  $\mu$  be the “democratic” weight dominated by  $\lambda$ . That is, we represent  $\mu$  by the composition  $(1, \dots, 1) \in \mathbb{R}^{3k}$ . With this choice of  $\lambda$  and  $\mu$ , the projective variety  $\text{Proj } R(\lambda, \mu)$  is the moduli space of equally weighted  $3k$ -tuples of points in projective space  $\mathbb{P}^2$  (see [6] for more details).

We now construct a GT pattern that we claim will be a vertex of  $\text{GT}(\lambda, \mu)$ . Define the sequences  $\{T_j^{(1)}\}$  and  $\{T_j^{(2)}\}$  by the coupled recurrence relations

$$\begin{aligned} T_0^{(1)} &= k, & T_0^{(2)} &= k - 1/2, \\ T_j^{(1)} &= T_{j-1}^{(2)} - 1 \quad (j \geq 1), & T_j^{(2)} &= \frac{1}{2}(T_j^{(1)} + T_{j-1}^{(1)}) \quad (j \geq 1). \end{aligned} \tag{5.1}$$

Solving this system of recurrence relations yields the closed-form expressions

$$T_j^{(1)} = k - \frac{2}{3}j + \frac{5}{9}\left(\frac{-1}{2}\right)^j - \frac{5}{9}, \tag{5.2}$$

$$T_j^{(2)} = k - \frac{2}{3}j - \frac{5}{18}\left(\frac{-1}{2}\right)^j - \frac{2}{9}. \tag{5.3}$$

Let  $N = k + \lfloor k/2 \rfloor - 2$ . We will construct a triangular array  $\mathbf{x}$  by filling in the entries of  $\mathbf{x}$  in blocks from the upper left to the lower right using the values  $T_j^{(1)}$  and

$T_j^{(2)}$ . Begin by filling the entries in the upper left of the triangular array as follows:

$$\begin{array}{ccccccc}
 x_{1n} & & x_{2n} & & x_{3n} & & & k & & k & & k \\
 & x_{1,n-1} & & x_{2,n-1} & & x_{3,n-1} & & & k & & k & & k-1 \\
 & & x_{1,n-2} & & x_{2,n-2} & & & & & k & & k-\frac{1}{2} \\
 & & & x_{1,n-3} & & & & & & & k & & 
 \end{array} =$$

We then proceed from the upper left to the lower right of the triangular array by filling in blocks of entries as follows. For  $1 \leq j \leq N - 1$ , let

$$\begin{array}{ccccccc}
 & & & & & & & & & & T_j^{(1)} \\
 & & & & x_{3,n-2j} & & & & & & & & \\
 & & & x_{2,n-2j-1} & & x_{3,n-2j-1} & & & & T_j^{(1)} & & T_j^{(1)} \\
 & & & & & & & & & = & & & \\
 & & x_{1,n-2j-2} & & x_{2,n-2j-2} & & & & T_j^{(2)} & & T_j^{(1)} \\
 & & & x_{1,n-2j-3} & & & & & & & T_j^{(1)}
 \end{array}$$

If  $k$  is even, the final entries at the bottom of the array are filled in as follows:

$$\begin{array}{ccccccc}
 & & & & & & & & & & T_N^{(1)} \\
 & & & & x_{34} & & & & & & & & \\
 & & & x_{23} & & x_{33} & & & & T_N^{(1)} & & T_N^{(1)} \\
 & & & & & & & & & = & & & \\
 & & x_{12} & & x_{22} & & & & 2 - T_N^{(1)} & & T_N^{(1)} \\
 & & & x_{11} & & & & & & & 1
 \end{array}$$

On the other hand, if  $k$  is odd, then the final entries are filled in as follows:

$$\begin{array}{ccccccc}
 & & & & & & & & & & T_N^{(1)} \\
 & & & & x_{35} & & & & & & & & \\
 & & & & x_{24} & & x_{34} & & & T_N^{(1)} & & T_N^{(1)} \\
 & & & x_{13} & & x_{23} & & x_{33} & & = & T_N^{(1)} & T_N^{(1)} & 3 - 2T_N^{(1)} \\
 & & & & x_{12} & & x_{22} & & & & T_N^{(1)} & 2 - T_N^{(1)} \\
 & & & & & x_{11} & & & & & & 1
 \end{array}$$

All the remaining entries of the triangular array are assigned the value 0.

**Proposition 5.1** *The triangular array constructed above is a vertex of  $GT(\lambda, \mu)$  with denominator  $2^N$ .*

*Proof* To show that  $\mathbf{x} \in \text{GT}(\lambda, \mu)$ , we first check that  $\mathbf{x}$  is a GT-pattern. In this case, the interlacing inequalities to be verified are

$$\left. \begin{aligned} T_{j-1}^{(1)} > T_j^{(1)} > 0 \\ T_{j-2}^{(1)} > T_{j-1}^{(2)} > T_{j-1}^{(1)} \end{aligned} \right\} \text{ for } 2 \leq j \leq N,$$

$$\left. \begin{aligned} T_{N-1}^{(1)} > 2 - T_N^{(1)} > T_N^{(1)} \\ 2 - T_N^{(1)} > 1 > T_N^{(1)} \end{aligned} \right\} \text{ if } k \text{ is even,}$$

$$\left. \begin{aligned} T_N^{(1)} > 3 - 2T_N^{(1)} > 0 \\ T_N^{(1)} > 2 - T_N^{(1)} > 3 - 2T_N^{(1)} \\ T_N^{(1)} > 1 > 2 - T_N^{(1)} \end{aligned} \right\} \text{ if } k \text{ is odd.}$$

These are all straightforward consequences of the closed-form expressions (5.2) and (5.3) for  $T_j^{(1)}$  and  $T_j^{(2)}$ , respectively, so  $\mathbf{x}$  is a GT-pattern. Thus, to show that  $\mathbf{x} \in \text{GT}(\lambda, \mu)$ , we need only establish that the row-sums of  $\mathbf{x}$  are correct. This amounts to showing that


$$T_{j-1}^{(2)} + T_{j-1}^{(1)} + T_j^{(1)} = 3k - 2j,$$

$$T_{j-1}^{(1)} + 2T_j^{(1)} = 3k - 2j - 1 \tag{5.4}$$

for  $2 \leq j \leq N$ . These equalities may be shown using induction and the recursive definition (5.1) of  $T_j^{(1)}$  and  $T_j^{(2)}$ . It is clear from (5.4) that  $T_j^{(1)}$  has denominator  $2^j$  when written as a reduced fraction. Hence,  $\mathbf{x}$  has denominator  $2^N$ , as claimed.

It remains only to show that  $\mathbf{x}$  is a vertex of  $\text{GT}(\lambda, \mu)$ . We prove this by showing that, for any triangular array  $\varepsilon$ , if  $\mathbf{x} \pm \varepsilon \in \text{GT}(\lambda, \mu)$ , then  $\varepsilon = 0$ . This is most easily

**Fig. 1** Tiling of  $\mathbf{x}$  when  $k$  is even



seen by partitioning the entries of  $\mathbf{x}$  so that entries that are equal and adjacent are grouped together. Following [3], we call each group of entries in this partition a *tile*. See Fig. 1 for a depiction of the case when  $k$  is even. Each tile is labeled with the value shared by the entries that it contains.

Suppose that  $\mathbf{x} \pm \varepsilon \in \text{GT}(\lambda, \mu)$ . Note that, after the addition of  $\pm\varepsilon$ , the entries in each tile must still share a value, and the row-sums must be unchanged [3, Theorem 1.5]. We prove inductively that the entries in each tile cannot have changed, proceeding from the upper left to the lower right.

The entries in the tile labeled  $T_0^{(1)} = k$  cannot have changed because the top row is fixed. For the same reason, the 0 entries in  $\mathbf{x}$  are also fixed. Proceeding by induction, the entries in the tile labeled  $T_j^{(1)}$  cannot have changed because there is a row on which this is the only tile besides the tile labeled  $T_{j-1}^{(1)}$  and the tile of 0s, which have already been fixed. Hence, the entries in all the tiles labeled  $T_j^{(1)}$ ,  $0 \leq j \leq N$ , are fixed under the addition of  $\pm\varepsilon$ . Finally, for  $1 \leq j \leq N - 1$ , the tile labeled  $T_j^{(2)}$  lies on a row in which the other entries ( $T_j^{(1)}$ ,  $T_{j+1}^{(1)}$ , and 0) have been shown to be fixed, so the entry in this tile is also fixed under the addition of  $\pm\varepsilon$ . Therefore, we conclude that  $\varepsilon = 0$ , so that  $\mathbf{x}$  is a vertex, as claimed.  $\square$

The following theorem is an immediate consequence.

**Theorem 5.2** *The Gelfand–Tsetlin algebra  $R'(k\varpi_3, \mu)$  has essential generators of degree exceeding  $2^{(n/2)-3}$ .*

**Acknowledgements** We thank Harm Derksen, Ionuț Ciocan-Fontanine, and Mircea Mustață for their invaluable advice. We also thank the anonymous referees for several helpful suggestions.

## References

1. Bruns, W., Herzog, J.: Cohen–Macaulay Rings. Cambridge Studies in Advanced Mathematics, vol. 39. Cambridge University Press, Cambridge (1993)
2. Caldero, P.: Toric degenerations of Schubert varieties. *Transform. Groups* **7**(1), 51–60 (2002)
3. De Loera, J.A., McAllister, T.B.: Vertices of Gelfand–Tsetlin polytopes. *Discrete Comput. Geom.* **32**(4), 459–470 (2004)
4. Derksen, H., Kraft, H.: Constructive Invariant Theory, Algèbre non commutative, groupes quantiques et invariants, Reims, 1995. *Sémin. Congr.*, vol. 2, pp. 221–244. Soc. Math. France, Paris (1997)
5. Dolgachev, I.: Lectures on Invariant Theory. London Mathematical Society Lecture Note Series, vol. 296. Cambridge University Press, Cambridge (2003)
6. Dolgachev, I., Ortland, D.: Point sets in projective spaces and theta functions. *Astérisque* **165**, 210 (1988)
7. Eisenbud, D.: Commutative Algebra. With a View Toward Algebraic Geometry. Graduate Texts in Mathematics, vol. 150. Springer, New York (1995)
8. Foth, P., Hu, Y.: Toric Degenerations of Weight Varieties and Applications. *Travaux mathématiques. Fasc. XVI, Trav. Math.*, XVI, pp. 87–105. Univ. Luxemb., Luxembourg (2005)
9. Gonciulea, N., Lakshmibai, V.: Degenerations of flag and Schubert varieties to toric varieties. *Transform. Groups* **1**(3), 215–248 (1996)
10. Hochster, M.: Rings of invariants of tori. Cohen–Macaulay rings generated by monomials, and polytopes. *Ann. Math. (2)* **96**, 318–337 (1972)
11. Howard, B.J.: Matroids and geometric invariant theory of torus actions on flag spaces. *J. Algebra* **312**(1), 527–541 (2007)

12. Howard, B.J., Millson, J., Snowden, A., Vakil, R.: The equations for the moduli space of  $n$  points on the line. *Duke Math. J.* **146**(2), 175–226 (2009). doi:[10.1215/0012794-2008-063](https://doi.org/10.1215/0012794-2008-063)
13. Knutson, A.: Weight varieties. Ph.D. thesis, Massachusetts Institute of Technology, May 1996
14. Kogan, M., Miller, E.: Toric degeneration of Schubert varieties and Gelfand–Tsetlin polytopes. *Adv. Math.* **193**(1), 1–17 (2005)
15. Miller, E., Sturmfels, B.: *Combinatorial Commutative Algebra*. Graduate Texts in Mathematics, vol. 227. Springer, New York (2005)
16. Popov, V.L.: Constructive Invariant Theory, Young tableaux and Schur functors in algebra and geometry, Toruń, 1980. *Astérisque*, vol. 87, pp. 303–334. Soc. Math. France, Paris (1981)
17. Stanley, R.P.: *Enumerative Combinatorics*. Vol. 1. Cambridge Studies in Advanced Mathematics, vol. 49. Cambridge University Press, Cambridge (1997). With a foreword by Gian-Carlo Rota, Corrected reprint of the 1986 original