

Journal of Inequalities in Pure and Applied Mathematics

BEST GENERALIZATION OF HARDY-HILBERT'S INEQUALITY WITH MULTI-PARAMETERS

DONGMEI XIN

Department of Mathematics
Guangdong Education College
Guangzhou, Guangdong 510303
People's Republic Of China.

EMail: xdm77108@gdei.edu.cn

©2000 Victoria University
ISSN (electronic): 1443-5756
034-06



volume 7, issue 4, article 153,
2006.

*Received 07 February, 2006;
accepted 13 April, 2006.*

Communicated by: B. Yang

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

Abstract

By introducing some parameters and the β function and improving the weight function, we obtain a generalization of Hilbert's integral inequality with the best constant factor. As its applications, we build its equivalent form and some particular results.

2000 Mathematics Subject Classification: 26D15.

Key words: Hardy-Hilbert's inequality, Hölder's inequality, weight function, β function.

Contents

1	Introduction	3
2	Some Lemmas	7
3	Main Results and Applications	9
	References	



Best Generalization of Hardy-Hilbert's Inequality With Multi-Parameters

Dongmei Xin

Title Page

Contents



Go Back

Close

Quit

Page 2 of 18

1. Introduction

If $p > 1$, $\frac{1}{p} + \frac{1}{q} = 1$, f, g are non-negative functions such that $0 < \int_0^\infty f^p(t)dt < \infty$ and $0 < \int_0^\infty g^q(t)dt < \infty$, then we have


$$(1.1) \quad \int_0^\infty \int_0^\infty \frac{f(x)g(y)}{x+y} dx dy < \frac{\pi}{\sin(\frac{\pi}{p})} \left\{ \int_0^\infty f^p(t)dt \right\}^{\frac{1}{p}} \left\{ \int_0^\infty g^q(t)dt \right\}^{\frac{1}{q}};$$

$$(1.2) \quad \int_0^\infty \left[\int_0^\infty \frac{f(x)}{x+y} dx \right]^p dy < \left[\frac{\pi}{\sin(\frac{\pi}{p})} \right]^p \int_0^\infty f^p(t)dt,$$

where the constant factors $\frac{\pi}{\sin(\frac{\pi}{p})}$ and $\left[\frac{\pi}{\sin(\frac{\pi}{p})} \right]^p$ are the best possible (see [1]). Inequality (1.1) is well known as Hardy-Hilbert's integral inequality, which is important in analysis and applications (see [2]). Inequality (1.1) is equivalent to (1.2).

In 2002, Yang [3] gave some generalizations of (1.1) and (1.2) by introducing a parameter $\lambda > 0$ as:

$$(1.3) \quad \int_0^\infty \int_0^\infty \frac{f(x)g(y)}{x^\lambda + y^\lambda} dx dy < \frac{\pi}{\lambda \sin(\frac{\pi}{p})} \left\{ \int_0^\infty t^{(p-1)(1-\lambda)} f^p(t)dt \right\}^{\frac{1}{p}} \times \left\{ \int_0^\infty t^{(q-1)(1-\lambda)} g^q(t)dt \right\}^{\frac{1}{q}};$$



Best Generalization of
Hardy-Hilbert's Inequality With
Multi-Parameters

Dongmei Xin

Title Page

Contents



Go Back

Close

Quit

Page 3 of 18

$$(1.4) \quad \int_0^\infty y^{\lambda-1} \left[\int_0^\infty \frac{f(x)}{x^\lambda + y^\lambda} dx \right]^p dy < \left[\frac{\pi}{\lambda \sin(\frac{\pi}{p})} \right]^p \int_0^\infty t^{(p-1)(1-\lambda)} f^p(t) dt,$$


where the constant factors $\frac{\pi}{\lambda \sin(\frac{\pi}{p})}$ and $\left[\frac{\pi}{\lambda \sin(\frac{\pi}{p})} \right]^p$ are the best possible. Inequality (1.3) is equivalent to (1.4).

When $\lambda = 1$, both (1.3) and (1.4) change to (1.1) and (1.2). Yang [4] gave another generalization of (1.1) by introducing a parameter λ and a β function.

In 2004, by introducing some parameters and estimating the weight function, Yang [5] gave some extensions of (1.1) and (1.2) with the best constant factors as:

$$(1.5) \quad \int_0^\infty \int_0^\infty \frac{f(x)g(y)}{x^\lambda + y^\lambda} dx dy < \frac{\pi}{\lambda \sin(\frac{\pi}{r})} \left\{ \int_0^\infty x^{p(1-\frac{\lambda}{r})-1} f^p(x) dx \right\}^{\frac{1}{p}} \left\{ \int_0^\infty x^{q(1-\frac{\lambda}{s})-1} g^q(x) dx \right\}^{\frac{1}{q}};$$

$$(1.6) \quad \int_0^\infty y^{\frac{p\lambda}{s}-1} \left[\int_0^\infty \frac{f(x)}{x^\lambda + y^\lambda} dx \right]^p dy < \left[\frac{\pi}{\lambda \sin(\frac{\pi}{r})} \right]^p \int_0^\infty x^{p(1-\frac{\lambda}{r})-1} f^p(x) dx,$$



Best Generalization of Hardy-Hilbert's Inequality With Multi-Parameters

Dongmei Xin

Title Page	
Contents	
◀◀	▶▶
◀	▶
Go Back	
Close	
Quit	
Page 4 of 18	

where the constant factors $\frac{\pi}{\lambda \sin(\pi/r)}$ and $\left[\frac{\pi}{\lambda \sin(\pi/r)}\right]^p$ are the best possible. Inequality (1.5) is equivalent to (1.6). Recently, [6, 7, 8, 9] considered some multiple extensions of (1.1).

Under the same conditions with (1.1), we still have (see [1, Th. 342]):


$$(1.7) \quad \int_0^\infty \int_0^\infty \frac{\ln\left(\frac{x}{y}\right) f(x)g(y)}{x-y} dx dy < \left[\frac{\pi}{\sin(\frac{\pi}{p})}\right]^2 \left\{\int_0^\infty f^p(t) dt\right\}^{\frac{1}{p}} \left\{\int_0^\infty g^q(t) dt\right\}^{\frac{1}{q}};$$

$$(1.8) \quad \int_0^\infty \left[\frac{\ln\left(\frac{x}{y}\right) f(x)}{x-y} dx\right]^p dy < \left[\frac{\pi}{\sin(\frac{\pi}{p})}\right]^{2p} \int_0^\infty f^p(t) dt,$$

where the constant factors $\left[\frac{\pi}{\sin(\pi/p)}\right]^2$ and $\left[\frac{\pi}{\sin(\pi/p)}\right]^{2p}$ are the best possible. Inequality (1.7) is equivalent to (1.8). In recent years, by introducing a parameter λ , Kuang [10] gave a new extension of (1.7).

In 2003, by introducing a parameter $\lambda > 0$ and the weight function, Yang [11] gave another generalisation of (1.7) and the extended equivalent form as:

$$(1.9) \quad \int_0^\infty \int_0^\infty \frac{\ln\left(\frac{x}{y}\right) f(x)g(y)}{x^\lambda - y^\lambda} dx dy$$



Best Generalization of Hardy-Hilbert's Inequality With Multi-Parameters

Dongmei Xin

Title Page

Contents




Go Back

Close

Quit

Page 5 of 18



Best Generalization of Hardy-Hilbert's Inequality With Multi-Parameters

Dongmei Xin

Title Page	
Contents	
◀◀	▶▶
◀	▶
Go Back	
Close	
Quit	
Page 6 of 18	

$$< \left[\frac{\pi}{\lambda \sin(\frac{\pi}{p})} \right]^2 \left\{ \int_0^\infty t^{(p-1)(1-\lambda)} f^p(t) dt \right\}^{\frac{1}{p}} \left\{ \int_0^\infty t^{(q-1)(1-\lambda)} g^q(t) dt \right\}^{\frac{1}{q}} ;$$

$$(1.10) \quad \int_0^\infty y^{\lambda-1} \left[\frac{\ln\left(\frac{x}{y}\right) f(x)}{x^\lambda - y^\lambda} dx \right]^p dy < \left[\frac{\pi}{\lambda \sin(\frac{\pi}{p})} \right]^{2p} \int_0^\infty t^{(p-1)(1-\lambda)} f^p(t) dt,$$

where the constant factors $\left[\frac{\pi}{\lambda \sin(\pi/p)} \right]^2$ and $\left[\frac{\pi}{\lambda \sin(\pi/p)} \right]^{2p}$ are the best possible. Inequality (1.9) is equivalent to (1.10).

In this paper, by using the β function and obtaining the expression of the weight function, we give a new extension of (1.7) with some parameters as (1.5). As applications, we also consider the equivalent form and some other particular results.

2. Some Lemmas

Lemma 2.1. *If $p > 1$, $\frac{1}{p} + \frac{1}{q} = 1$, $r > 1$, $\frac{1}{s} + \frac{1}{r} = 1$, $\lambda > 0$, define the weight function $\omega_\lambda(s, p, x)$ as*

$$(2.1) \quad \omega_\lambda(s, p, x) = \int_0^\infty \frac{\ln\left(\frac{x}{y}\right)}{x^\lambda - y^\lambda} \cdot \frac{x^{(p-1)(1-\frac{\lambda}{r})}}{y^{1-\frac{\lambda}{s}}} dy, \quad x \in (0, \infty).$$


Then we have

$$(2.2) \quad \omega_\lambda(s, p, x) = x^{p(1-\frac{\lambda}{r})-1} \left[\frac{\pi}{\lambda \sin(\frac{\pi}{r})} \right]^2.$$

Proof. For fixed x , setting $u = (\frac{y}{x})^\lambda$ in the integral (2.1) and by [1] (see [1, Th. 342 Remark]), we have

$$(2.3) \quad \begin{aligned} \omega_\lambda(s, p, x) &= \frac{1}{\lambda^2} \int_0^\infty \frac{\ln u}{x^\lambda(u-1)} \cdot \frac{x^{(p-1)(1-\frac{\lambda}{r})}}{(u^{\frac{1}{\lambda}}x)^{1-\frac{\lambda}{s}}} x u^{\frac{1}{\lambda}-1} du \\ &= \frac{1}{\lambda^2} x^{p(1-\frac{\lambda}{r})-1} \int_0^\infty \frac{\ln u}{u-1} \cdot u^{-\frac{1}{r}} du \\ &= \left[\frac{\pi}{\lambda \sin(\frac{\pi}{r})} \right]^2 x^{p(1-\frac{\lambda}{r})-1}. \end{aligned}$$

Hence, (2.2) is valid and the lemma is proved. □



Best Generalization of
Hardy-Hilbert's Inequality With
Multi-Parameters

Dongmei Xin

Title Page

Contents



Go Back


Close

Quit

Page 7 of 18

Note. By (2.3), we still have

$$(2.4) \quad \begin{aligned} \omega_\lambda(r, q, y) &= \int_0^\infty \frac{\ln\left(\frac{x}{y}\right)}{x^\lambda - y^\lambda} \cdot \frac{x^{(q-1)(1-\frac{\lambda}{s})}}{y^{1-\frac{\lambda}{r}}} dx \\ &= y^{q(1-\frac{\lambda}{s})-1} \left[\frac{\pi}{\lambda \sin(\frac{\pi}{s})} \right]^2. \end{aligned}$$



**Best Generalization of
Hardy-Hilbert's Inequality With
Multi-Parameters**

Dongmei Xin

Title Page

Contents



Go Back

Close

Quit

Page 8 of 18

3. Main Results and Applications

Theorem 3.1. If $p > 1$, $\frac{1}{p} + \frac{1}{q} = 1$, $r > 1$, $\frac{1}{s} + \frac{1}{r} = 1$, $\lambda > 0$, $f, g \geq 0$ such that $0 < \int_0^\infty x^{p(1-\frac{\lambda}{r})-1} f^p(x) dx < \infty$, and $0 < \int_0^\infty x^{q(1-\frac{\lambda}{s})-1} g^q(x) dx < \infty$, then we have

$$(3.1) \quad \int_0^\infty \int_0^\infty \frac{\ln\left(\frac{x}{y}\right) f(x)g(y)}{x^\lambda - y^\lambda} dx dy$$

$$< \left[\frac{\pi}{\lambda \sin\left(\frac{\pi}{r}\right)} \right]^2 \left\{ \int_0^\infty x^{p(1-\frac{\lambda}{r})-1} f^p(x) dx \right\}^{\frac{1}{p}}$$

$$\times \left\{ \int_0^\infty x^{q(1-\frac{\lambda}{s})-1} g^q(x) dx \right\}^{\frac{1}{q}},$$


where the constant factor $\left[\frac{\pi}{\lambda \sin(\pi/r)} \right]^2$ is the best possible. In particular,

(a) for $r = s = 2$, we have

$$(3.2) \quad \int_0^\infty \int_0^\infty \frac{\ln\left(\frac{x}{y}\right) f(x)g(y)}{x^\lambda - y^\lambda} dx dy$$

$$< \left(\frac{\pi}{\lambda}\right)^2 \left\{ \int_0^\infty x^{p(1-\frac{\lambda}{2})-1} f^p(x) dx \right\}^{\frac{1}{p}}$$

$$\times \left\{ \int_0^\infty x^{q(1-\frac{\lambda}{2})-1} g^q(x) dx \right\}^{\frac{1}{q}},$$



Best Generalization of
Hardy-Hilbert's Inequality With
Multi-Parameters

Dongmei Xin

Title Page

Contents



Go Back

Close

Quit


Page 9 of 18

(b) for $\lambda = 1$, we have

$$(3.3) \quad \int_0^\infty \int_0^\infty \frac{\ln\left(\frac{x}{y}\right) f(x)g(y)}{x-y} dx dy < \left[\frac{\pi}{\sin\left(\frac{\pi}{r}\right)} \right]^2 \left\{ \int_0^\infty x^{\frac{p}{s}-1} f^p(x) dx \right\}^{\frac{1}{p}} \left\{ \int_0^\infty x^{\frac{q}{r}-1} g^q(x) dx \right\}^{\frac{1}{q}}.$$

Proof. By Hölder's inequality and Lemma 2.1, we have

$$(3.4) \quad \begin{aligned} & \int_0^\infty \int_0^\infty \frac{\ln\left(\frac{x}{y}\right) f(x)g(y)}{x^\lambda - y^\lambda} dx dy \\ &= \int_0^\infty \int_0^\infty \left\{ \left[\frac{\ln\left(\frac{x}{y}\right)}{x^\lambda - y^\lambda} \right]^{\frac{1}{p}} \cdot \frac{x^{(1-\frac{\lambda}{r})/q}}{y^{(1-\frac{\lambda}{s})/p}} f(x) \right\} \\ & \quad \times \left\{ \left[\frac{\ln\left(\frac{x}{y}\right)}{x^\lambda - y^\lambda} \right]^{\frac{1}{q}} \cdot \frac{y^{(1-\frac{\lambda}{s})/p}}{x^{(1-\frac{\lambda}{r})/q}} g(y) \right\} dx dy \\ &\leq \left\{ \int_0^\infty \left[\int_0^\infty \frac{\ln\left(\frac{x}{y}\right)}{x^\lambda - y^\lambda} \cdot \frac{x^{(p-1)(1-\frac{\lambda}{r})}}{y^{(1-\frac{\lambda}{s})}} dy \right] f^p(x) dx \right\}^{\frac{1}{p}} \\ & \quad \times \left\{ \int_0^\infty \left[\int_0^\infty \frac{\ln\left(\frac{x}{y}\right)}{x^\lambda - y^\lambda} \cdot \frac{y^{(q-1)(1-\frac{\lambda}{s})}}{x^{(1-\frac{\lambda}{r})}} dx \right] g^q(y) dy \right\}^{\frac{1}{q}} \end{aligned}$$



Best Generalization of Hardy-Hilbert's Inequality With Multi-Parameters

Dongmei Xin

Title Page

Contents



Go Back

Close

Quit

Page 10 of 18

$$= \left\{ \int_0^\infty \omega_\lambda(s, p, x) f^p(x) dx \right\}^{\frac{1}{p}} \left\{ \int_0^\infty \omega_\lambda(r, q, y) g^q(y) dy \right\}^{\frac{1}{q}}.$$

If (3.4) takes the form of equality, then there exist constants A and B , such that they are not all zero and (see [12])

$$A \frac{\ln\left(\frac{x}{y}\right)}{x^\lambda - y^\lambda} \cdot \frac{x^{(p-1)(1-\frac{\lambda}{r})}}{y^{1-\frac{\lambda}{s}}} f^p(x) = B \frac{\ln\left(\frac{x}{y}\right)}{x^\lambda - y^\lambda} \cdot \frac{y^{(q-1)(1-\frac{\lambda}{s})}}{x^{1-\frac{\lambda}{r}}} g^q(y),$$

a.e. in $(0, \infty) \times (0, \infty)$.


We find that $Ax \cdot x^{p(1-\frac{\lambda}{r})-1} f^p(x) = By \cdot y^{q(1-\frac{\lambda}{s})-1} g^q(y)$, a.e. in $(0, \infty) \times (0, \infty)$. Hence there exists a constant C , such that

$$Ax \cdot x^{p(1-\frac{\lambda}{r})-1} f^p(x) = C = By \cdot y^{q(1-\frac{\lambda}{s})-1} g^q(y), \quad \text{a.e. in } (0, \infty).$$

Without loss of generality, suppose $A \neq 0$, we may get $x^{p(1-\frac{\lambda}{r})-1} f^p(x) = C/(Ax)$, a.e. in $(0, \infty)$, which contradicts $0 < \int_0^\infty x^{p(1-\frac{\lambda}{r})-1} f^p(x) dx < \infty$. Hence (3.4) takes strict inequality as follows:

$$(3.5) \quad \int_0^\infty \int_0^\infty \frac{\ln\left(\frac{x}{y}\right)}{x^\lambda - y^\lambda} f(x)g(y) dx dy < \left\{ \int_0^\infty \omega_\lambda(s, p, x) f^p(x) dx \right\}^{\frac{1}{p}} \left\{ \int_0^\infty \omega_\lambda(r, q, y) g^q(y) dy \right\}^{\frac{1}{q}}.$$

In view of (2.2) and (2.4), we have (3.1).



Best Generalization of
Hardy-Hilbert's Inequality With
Multi-Parameters

Dongmei Xin

Title Page

Contents



Go Back

Close

Quit

Page 11 of 18

If the constant factor $\left[\frac{\pi}{\lambda \sin(\pi/r)} \right]^2$ in (3.1) is not the best possible, then there exists a positive constant K (with $K < \left[\frac{\pi}{\lambda \sin(\pi/r)} \right]^2$) and an $a > 0$. We have

$$(3.6) \quad \int_a^\infty \int_0^\infty \frac{\ln\left(\frac{x}{y}\right) f(x)g(y)}{x^\lambda - y^\lambda} dx dy < K \left\{ \int_a^\infty x^{p(1-\frac{\lambda}{r})-1} f^p(x) dx \right\}^{\frac{1}{p}} \left\{ \int_a^\infty x^{q(1-\frac{\lambda}{s})-1} g^q(x) dx \right\}^{\frac{1}{q}}.$$

For $\varepsilon > 0$ small enough ($\varepsilon < \frac{p\lambda}{r}$) and $0 < b < a$, setting f_ε and g_ε as:

$$f_\varepsilon(x) = g_\varepsilon(x) = 0, \quad x \in (0, b);$$

$$f_\varepsilon = x^{-1-\frac{\varepsilon}{p}+\frac{\lambda}{r}}, \quad g_\varepsilon = x^{-1-\frac{\varepsilon}{q}+\frac{\lambda}{s}}, \quad x \in [b, \infty),$$

then we find

$$(3.7) \quad \int_a^\infty \int_b^\infty \frac{\ln\left(\frac{x}{y}\right) f_\varepsilon(x) \cdot g_\varepsilon(y)}{x^\lambda - y^\lambda} dx dy = \int_a^\infty \int_b^\infty \frac{\ln\left(\frac{x}{y}\right) x^{-1-\frac{\varepsilon}{p}+\frac{\lambda}{r}} \cdot y^{-1-\frac{\varepsilon}{q}+\frac{\lambda}{s}}}{x^\lambda - y^\lambda} dx dy.$$

In (3.7), for $b \rightarrow 0^+$, by (3.6), we have

$$\frac{1}{\lambda^2 a^\varepsilon} \int_0^\infty \frac{\ln u}{u-1} u^{-1+\frac{1}{s}-\frac{\varepsilon}{q\lambda}} du = \varepsilon \int_a^\infty \int_0^\infty \frac{\ln\left(\frac{x}{y}\right) f_\varepsilon(x)g_\varepsilon(y)}{x^\lambda - y^\lambda} dx dy \leq \frac{K}{a^\varepsilon}.$$



**Best Generalization of
Hardy-Hilbert's Inequality With
Multi-Parameters**

Dongmei Xin

Title Page

Contents



Go Back

Close

Quit

Page 12 of 18

For $\varepsilon^+ \rightarrow 0$, by [1] (see [1, Th. 342 Remark]), it follows that $\left[\frac{\pi}{\lambda \sin(\pi/r)}\right]^2 \leq K$, which contradicts the fact that $K < \left[\frac{\pi}{\lambda \sin(\pi/r)}\right]^2$. Hence the constant factor $\left[\frac{\pi}{\lambda \sin(\pi/r)}\right]^2$ in (3.1) is the best possible. The theorem is proved. \square


Theorem 3.2. *If $p > 1$, $\frac{1}{p} + \frac{1}{q} = 1$, $r > 1$, $\frac{1}{s} + \frac{1}{r} = 1$, $\lambda > 0$, $f \geq 0$ such that $0 < \int_0^\infty x^{p(1-\frac{\lambda}{r})-1} f^p(x) dx < \infty$, then we have*

$$(3.8) \quad \int_0^\infty y^{\frac{p\lambda}{s}-1} \left[\int_0^\infty \frac{\ln\left(\frac{x}{y}\right) f(x)}{x^\lambda - y^\lambda} dx \right]^p dy < \left[\frac{\pi}{\lambda \sin\left(\frac{\pi}{r}\right)}\right]^{2p} \int_0^\infty x^{p(1-\frac{\lambda}{r})-1} f^p(x) dx,$$

where the constant $\left[\frac{\pi}{\lambda \sin(\pi/r)}\right]^{2p}$ is the best possible. Inequality (3.8) is equivalent to (3.1). In particular,

(a) for $r = s = 2$, we have

$$(3.9) \quad \int_0^\infty y^{\frac{p\lambda}{2}-1} \left[\int_0^\infty \frac{\ln\left(\frac{x}{y}\right) f(x)}{x^\lambda - y^\lambda} dx \right]^p dy < \left(\frac{\pi}{\lambda}\right)^{2p} \int_0^\infty x^{p(1-\frac{\lambda}{2})-1} f^p(x) dx,$$



Best Generalization of Hardy-Hilbert's Inequality With Multi-Parameters

Dongmei Xin

Title Page

Contents



Go Back

Close

Quit

Page 13 of 18

(b) for $\lambda = 1$, we have


$$(3.10) \quad \int_0^\infty y^{\frac{p}{s}-1} \left[\int_0^\infty \frac{\ln\left(\frac{x}{y}\right) f(x)}{x-y} dx \right]^p dy < \left[\frac{\pi}{\sin\left(\frac{\pi}{r}\right)} \right]^{2p} \int_0^\infty x^{\frac{p}{s}-1} f^p(x) dx.$$

Proof. Setting a real function $g(y)$ as

$$g(y) = y^{\frac{p\lambda}{s}-1} \left[\int_0^\infty \frac{\ln\left(\frac{x}{y}\right) f(x)}{x^\lambda - y^\lambda} dx \right]^{p-1}, \quad y \in (0, \infty),$$

then by (3.1), we find

$$(3.11) \quad \begin{aligned} & \left[\int_0^\infty y^{q(1-\frac{\lambda}{s})-1} g^q(y) dy \right]^p \\ &= \left\{ \int_0^\infty y^{\frac{p\lambda}{s}-1} \left[\int_0^\infty \frac{\ln\left(\frac{x}{y}\right) f(x)}{x^\lambda - y^\lambda} dx \right]^p dy \right\}^p \\ &= \left[\int_0^\infty \int_0^\infty \frac{\ln\left(\frac{x}{y}\right) f(x) g(y)}{x^\lambda - y^\lambda} dx dy \right]^p \\ &\leq \left[\frac{\pi}{\lambda \sin\left(\frac{\pi}{r}\right)} \right]^{2p} \left\{ \int_0^\infty x^{p(1-\frac{\lambda}{r})-1} f^p(x) dx \right\} \end{aligned}$$



**Best Generalization of
Hardy-Hilbert's Inequality With
Multi-Parameters**

Dongmei Xin

Title Page

Contents



Go Back

Close

Quit

Page 14 of 18

$$\times \left\{ \int_0^\infty x^{q(1-\frac{\lambda}{s})-1} g^q(x) dx \right\}^{p-1}.$$


Hence we obtain

$$(3.12) \quad 0 < \int_0^\infty y^{q(1-\frac{\lambda}{s})-1} g^q(y) dy \\ \leq \left[\frac{\pi}{\lambda \sin(\frac{\pi}{r})} \right]^{2p} \int_0^\infty x^{p(1-\frac{\lambda}{r})-1} f^p(x) dx < \infty.$$

By (3.1), both (3.11) and (3.12) take the form of strict inequality, and we have (3.8).

On the other hand, suppose that (3.8) is valid. By Hölder's inequality, we find

$$(3.13) \quad \int_0^\infty \int_0^\infty \frac{\ln\left(\frac{x}{y}\right) f(x)g(y)}{x^\lambda - y^\lambda} dx dy \\ = \int_0^\infty \left[y^{\frac{\lambda}{s}-\frac{1}{p}} \int_0^\infty \frac{\ln\left(\frac{x}{y}\right) f(x)}{x^\lambda - y^\lambda} dx \right] \left[y^{-\frac{\lambda}{s}+\frac{1}{p}} g(y) \right] dy \\ \leq \left\{ \int_0^\infty y^{\frac{p\lambda}{s}-1} \left[\int_0^\infty \frac{\ln\left(\frac{x}{y}\right) f(x)}{x^\lambda - y^\lambda} dx \right]^p dy \right\}^{\frac{1}{p}} \\ \times \left\{ \int_0^\infty y^{q(1-\frac{\lambda}{s})-1} g^q(y) dy \right\}^{\frac{1}{q}}.$$



Best Generalization of
Hardy-Hilbert's Inequality With
Multi-Parameters

Dongmei Xin

Title Page

Contents



Go Back

Close

Quit


Page 15 of 18

Then by (3.8), we have (3.1). Hence (3.1) and (3.8) are equivalent.

If the constant $\left[\frac{\pi}{\lambda \sin(\pi/r)}\right]^{2p}$ in (3.8) is not the best possible, by using (3.13), we may get a contradiction that the constant factor in (3.1) is not the best possible. Thus we complete the proof of the theorem. \square

Remark 1.

- (a) For $r = q, s = p$, Inequality (3.1) reduces to (1.9) and (3.8) reduces to (1.10).
- (b) Inequality (3.1) is an extension of (1.7) with parameters (λ, r, s) .
- (c) It is interesting that inequalities (1.9) and (3.2) are different, although they have the same parameters and possess a best constant factor.



Best Generalization of
Hardy-Hilbert's Inequality With
Multi-Parameters

Dongmei Xin

Title Page

Contents



Go Back


Close

Quit

Page 16 of 18

References

- [1] G.H. HARDY, J.E. LITTLEWOOD AND G. POLYA, *Inequalities*, Cambridge University Press, Cambridge, UK, 1952.
- [2] D.S. MITRINOVIĆ, J.E. PEČARIĆ AND A.M. FINK, *Inequalities Involving Functions and their Integrals and Derivatives*, Kluwer Academic Publishers, Boston, 1991.
- [3] BICHENG YANG, On an extension of Hardy-Hilbert's inequality, *J. Math. Anal. Appl.*, **23A**(2) (2002), 247–254.
- [4] BICHENG YANG, On Hardy-Hilbert's integral inequality, *J. Math. Anal. Appl.*, **261** (2001), 295–306.
- [5] BICHENG YANG, On the extended Hilbert's integral inequality and applications, *International Review of Pure and Applied Mathematics*, (2005), 10–11.
- [6] YONG HONG , All-sided generalization about Hardy-Hilbert's integral inequalities, *Acta Math. Sinica*, **44**(4) (2001), 619–626.
- [7] LEPING HE, JIANGMING YU AND MINZHE GAO, An extension of Hilbert's integral inequality, *J. Shaoguan Univ.*, (Natural Science), **23**(3) (2002), 25–30.
- [8] JICHANG KUANG, On new extensions of Hilbert's integral inequality, *J. Math. Anal. Appl.*, **235** (1999), 608–614.
- [9] M. KRNIĆ AND J. PEČARIĆ, General Hilbert's and Hardy's inequalities, *Math. Inequal. & Applics.*, **8**(1) (2005), 29–51.



Best Generalization of
Hardy-Hilbert's Inequality With
Multi-Parameters

Dongmei Xin

Title Page

Contents




Go Back

Close

Quit

Page 17 of 18

- [10] JICHANG KUANG AND L. DEBNATH, On new generalizations of Hilbert's inequality and their applications, *J. Math. Anal. Appl.*, **245** (2000), 248–265.
- [11] BICHENG YANG, On a generalization of a Hilbert's type integral inequality and its applications, *Mathematics Applications*, **16**(2) (2003), 82–86.
- [12] JICHANG KUANG, *Applied Inequalities*, Shandong Science and Technology Press, Jiinan, (2004).



**Best Generalization of
Hardy-Hilbert's Inequality With
Multi-Parameters**

Dongmei Xin

Title Page

Contents



Go Back

Close

Quit

Page 18 of 18