

NAK EUN CHO AND SHIGEYOSHI OWA

Department of Applied Mathematics  
Pukyong National University  
Pusan 608-737, KOREA.  
*EMail:* [necho@pknu.ac.kr](mailto:necho@pknu.ac.kr)

Department of Mathematics  
Kinki University  
Higashi-Osaka, Osaka 577-8502  
JAPAN.  
*EMail:* [owa@math.kindai.ac.jp](mailto:owa@math.kindai.ac.jp)



---

volume 5, issue 2, article 30,  
2004.

*Received 03 March, 2004;  
accepted 30 March, 2004.*

*Communicated by: H.M. Srivastava*

---

Abstract

Contents



Home Page

Go Back

Close

Quit

## Abstract

The purpose of the present paper is to establish some results concerning the partial sums of meromorphic starlike and meromorphic convex functions analogous to the results due to H. Silverman [*J. Math. Anal. Appl.* **209** (1997), 221-227]. Furthermore, we consider the partial sums of certain integral operator.


*2000 Mathematics Subject Classification:* Primary 30C45.

*Key words:* Partial sums, Meromorphic starlike functions, Meromorphic convex functions, Meromorphic close-to-convex functions, Integral operators.

This work was supported by the *Korea Research Foundation Grant* (KRF-2003-015-C00024).

## Contents

1	Introduction .....	3
2	Main Results .....	6
	References	



### Partial Sums Of Certain Meromorphic Functions

Nak Eun Cho and Shigeyoshi Owa

Title Page

Contents



Go Back

Close

Quit

Page 2 of 15

# 1. Introduction

Let  $\Sigma$  be the class consisting of functions of the form

$$(1.1) \quad f(z) = \frac{1}{z} + \sum_{k=1}^{\infty} a_k z^k$$

which are analytic in the punctured open unit disk  $\mathcal{D} = \{z : 0 < |z| < 1\}$ . Let  $\Sigma^*(\alpha)$  and  $\Sigma_c(\alpha)$  be the subclasses of  $\Sigma$  consisting of all functions which are, respectively, meromorphic starlike and meromorphic convex of order  $\alpha$  ( $0 \leq \alpha < 1$ ) in  $\mathcal{D}$ . We also denote by  $\Sigma_c(\alpha)$  the subclass of  $\Sigma$  which satisfies


$$-\operatorname{Re}\{z^2 f'(z)\} > \alpha \quad (0 \leq \alpha < 1; z \in \mathcal{U} = \mathcal{D} \cup \{0\}).$$

We note that every function belonging to the class  $\Sigma_c(\alpha)$  is meromorphic close-to-convex of order  $\alpha$  in  $\mathcal{D}$  (see [2]). If  $f(z) = \sum_{k=0}^{\infty} a_k z^k$  and  $g(z) = \sum_{k=0}^{\infty} b_k z^k$  are analytic in  $\mathcal{U}$ , then their Hadamard product (or convolution), denoted by  $f * g$ , is the function defined by the power series

$$(f * g)(z) = \sum_{k=0}^{\infty} a_k b_k z^k \quad (z \in \mathcal{U}).$$

A sufficient condition for a function  $f$  of the form (1.1) to be in  $\Sigma^*(\alpha)$  is that

$$(1.2) \quad \sum_{k=1}^{\infty} (k + \alpha) |a_k| \leq 1 - \alpha$$



---

## Partial Sums Of Certain Meromorphic Functions

Nak Eun Cho and Shigeyoshi Owa

---

Title Page

Contents



Go Back

Close

Quit


Page 3 of 15

and to be in  $\Sigma_k(\alpha)$  is that

$$(1.3) \quad \sum_{k=1}^{\infty} k(k + \alpha)|a_k| \leq 1 - \alpha.$$

Further, we note that these sufficient conditions are also necessary for functions of the form (1.1) with positive or negative coefficients ([6, 13], also see [7]). Recently, Silverman [10] determined sharp lower bounds on the real part of the quotients between the normalized starlike or convex functions and their sequences of partial sums. Also, Li and Owa [4] obtained the sharp radius which for the normalized univalent functions in  $\mathcal{U}$ , the partial sums of the well-known Libera integral operator [5] imply starlikeness. Further, for various other interesting developments concerning partial sums of analytic univalent functions, the reader may be (for examples) referred to the works of Brickman et al. [1], Sheil-Small [9], Silvia [11], Singh and Singh [12] and Yang and Owa [14].

Since to a certain extent the work in the meromorphic univalent case has paralleled that of the analytic univalent case, one is tempted to search results analogous to those of Silverman [10] for meromorphic univalent functions in  $\mathcal{D}$ . In the present paper, motivated essentially by the work of Silverman [10], we will investigate the ratio of a function of the form (1.1) to its sequence of partial sums  $f_n(z) = \frac{1}{z} + \sum_{k=1}^n a_k z^k$  when the coefficients are sufficiently small to satisfy either condition (1.2) or (1.3). More precisely, we will determine sharp lower bounds for  $\operatorname{Re}\{f(z)/f_n(z)\}$ ,  $\operatorname{Re}\{f_n(z)/f(z)\}$ ,  $\operatorname{Re}\{f'(z)/f'_n(z)\}$ , and  $\operatorname{Re}\{f'_n(z)/f'(z)\}$ . Further, we give a property for the partial sums of certain integral operators in connection with meromorphic close-to-convex functions. In the sequel, we will make use of the well-known result that  $\operatorname{Re}\{(1 + w(z))/(1 -$




---

**Partial Sums Of Certain  
Meromorphic Functions**

Nak Eun Cho and Shigeyoshi Owa

---

Title Page

Contents




Go Back

Close

Quit

Page 4 of 15

$w(z) \} > 0 (z \in \mathcal{U})$  if and only if  $w(z) = \sum_{k=1}^{\infty} c_k z^k$  satisfies the inequality  $|w(z)| < |z|$ . Unless otherwise stated, we will assume that  $f$  is of the form (1.1) and its sequence of partial sums is denoted by  $f_n(z) = \frac{1}{z} + \sum_{k=1}^n a_k z^k$ .




---

**Partial Sums Of Certain Meromorphic Functions**

Nak Eun Cho and Shigeyoshi Owa

---

Title Page

Contents



Go Back

Close

Quit

Page 5 of 15

## 2. Main Results

**Theorem 2.1.** *If  $f$  of the form (1.1) satisfies condition (1.2), then*

$$\operatorname{Re} \left\{ \frac{f(z)}{f_n(z)} \right\} \geq \frac{n+2\alpha}{n+1+\alpha} \quad (z \in \mathcal{U}).$$

*The result is sharp for every  $n$ , with extremal function*

$$(2.1) \quad f(z) = \frac{1}{z} + \frac{1-\alpha}{n+1+\alpha} z^{n+1} \quad (n \geq 0).$$

*Proof.* We may write


$$\begin{aligned} & \frac{n+1+\alpha}{1-\alpha} \left[ \frac{f(z)}{f_n(z)} - \frac{n+2\alpha}{n+1+\alpha} \right] \\ &= \frac{1 + \sum_{k=1}^n a_k z^{k+1} + \frac{n+1+\alpha}{1-\alpha} \sum_{k=n+1}^{\infty} a_k z^{k+1}}{1 + \sum_{k=1}^n a_k z^{k+1}} \\ &:= \frac{1 + A(z)}{1 + B(z)}. \end{aligned}$$

Set  $(1 + A(z))/(1 + B(z)) = (1 + w(z))/(1 - w(z))$ , so that  $w(z) = (A(z) - B(z))/(2 + A(z) + B(z))$ . Then

$$w(z) = \frac{\frac{n+1+\alpha}{1-\alpha} \sum_{k=n+1}^{\infty} a_k z^{k+1}}{2 + 2 \sum_{k=1}^n a_k z^{k+1} + \frac{n+1+\alpha}{1-\alpha} \sum_{k=n+1}^{\infty} a_k z^{k+1}}$$

and

$$|w(z)| \leq \frac{\frac{n+1+\alpha}{1-\alpha} \sum_{k=n+1}^{\infty} |a_k|}{2 - 2 \sum_{k=1}^n |a_k| - \frac{n+1+\alpha}{1-\alpha} \sum_{k=n+1}^{\infty} |a_k|}.$$



Partial Sums Of Certain  
Meromorphic Functions

Nak Eun Cho and Shigeyoshi Owa

Title Page

Contents



Go Back

Close

Quit

Page 6 of 15

Now  $|w(z)| \leq 1$  if and only if

$$2 \left( \frac{n+1+\alpha}{1-\alpha} \right) \sum_{k=n+1}^{\infty} |a_k| \leq 2 - 2 \sum_{k=1}^n |a_k|,$$

which is equivalent to

$$(2.2) \quad \sum_{k=1}^n |a_k| + \frac{n+1+\alpha}{1-\alpha} \sum_{k=n+1}^{\infty} |a_k| \leq 1.$$

It suffices to show that the left hand side of (2.2) is bounded above by  $\sum_{k=1}^{\infty} ((k+\alpha)/(1-\alpha))|a_k|$ , which is equivalent to

$$\sum_{k=1}^n \left( \frac{k+2\alpha-1}{1-\alpha} \right) |a_k| + \sum_{k=n+1}^{\infty} \left( \frac{k-n-1}{1-\alpha} \right) |a_k| \geq 0.$$


To see that the function  $f$  given by (2.1) gives the sharp result, we observe for  $z = re^{\pi i/(n+2)}$  that

$$\frac{f(z)}{f_n(z)} = 1 + \frac{1-\alpha}{n+1+\alpha} z^{n+2} \longrightarrow 1 - \frac{1-\alpha}{n+1+\alpha} = \frac{n+2\alpha}{n+1+\alpha} \text{ when } r \rightarrow 1^-.$$

Therefore we complete the proof of Theorem 2.1. □

**Theorem 2.2.** *If  $f$  of the form (1.1) satisfies condition (1.3), then*

$$\operatorname{Re} \left\{ \frac{f(z)}{f_n(z)} \right\} \geq \frac{(n+2)(n+\alpha)}{(n+1)(n+1+\alpha)} \quad (z \in \mathcal{U}).$$



### Partial Sums Of Certain Meromorphic Functions

Nak Eun Cho and Shigeyoshi Owa

Title Page

Contents



Go Back

Close

Quit

Page 7 of 15

The result is sharp for every  $n$ , with extremal function

$$(2.3) \quad f(z) = \frac{1}{z} + \frac{1 - \alpha}{(n + 1)(n + 1 + \alpha)} z^{n+1} \quad (n \geq 0).$$

*Proof.* We write

$$\begin{aligned} & \frac{(n + 1)(n + 1 + \alpha)}{1 - \alpha} \left[ \frac{f(z)}{f_n(z)} - \frac{(n + 2)(n + \alpha)}{(n + 1)(n + 1 + \alpha)} \right] \\ &= \frac{1 + \sum_{k=1}^n a_k z^{k+1} + \frac{(n+1)(n+1+\alpha)}{1-\alpha} \sum_{k=n+1}^{\infty} a_k z^{k+1}}{1 + \sum_{k=1}^n a_k z^{k+1}} \\ &:= \frac{1 + w(z)}{1 - w(z)}, \end{aligned}$$

where


$$w(z) = \frac{\frac{(n+1)(n+1+\alpha)}{1-\alpha} \sum_{k=n+1}^{\infty} a_k z^{k+1}}{2 + 2 \sum_{k=1}^n a_k z^{k+1} + \frac{(n+1)(n+1+\alpha)}{1-\alpha} \sum_{k=n+1}^{\infty} a_k z^{k+1}}.$$

Now

$$|w(z)| \leq \frac{\frac{(n+1)(n+1+\alpha)}{1-\alpha} \sum_{k=n+1}^{\infty} |a_k|}{2 - 2 \sum_{k=1}^n |a_k| - \frac{(n+1)(n+1+\alpha)}{1-\alpha} \sum_{k=n+1}^{\infty} |a_k|} \leq 1,$$

if

$$(2.4) \quad \sum_{k=1}^n |a_k| + \frac{(n + 1)(n + 1 + \alpha)}{1 - \alpha} \sum_{k=n+1}^{\infty} |a_k| \leq 1.$$



## Partial Sums Of Certain Meromorphic Functions

Nak Eun Cho and Shigeyoshi Owa

Title Page

Contents



Go Back

Close

Quit

Page 8 of 15



The left hand side of (2.4) is bounded above by  $\sum_{k=1}^{\infty} (k(k + \alpha)/(1 - \alpha))|a_k|$  if

$$\frac{1}{1 - \alpha} \left\{ \sum_{k=1}^n (k(k + \alpha) - (1 - \alpha)|a_k| + \sum_{k=n+1}^{\infty} (k(k + \alpha) - (n + 1)(n + 1 + \alpha))|a_k| \right\} \geq 0,$$

and the proof is completed. □

We next determine bounds for  $\operatorname{Re}\{f_n(z)/f(z)\}$ .

**Theorem 2.3.** (a) If  $f$  of the form (1.1) satisfies condition (1.2), then


$$\operatorname{Re} \left\{ \frac{f_n(z)}{f(z)} \right\} \geq \frac{n + 1 + \alpha}{n + 2} \quad (z \in \mathcal{U}).$$

(b) If  $f$  of the form (1.1) satisfies condition (1.3), then

$$\operatorname{Re} \left\{ \frac{f_n(z)}{f(z)} \right\} \geq \frac{(n + 1)(n + 1 + \alpha)}{(n + 1)(n + 2) - n(1 - \alpha)} \quad (z \in \mathcal{U}).$$

Equalities hold in (a) and (b) for the functions given by (2.1) and (2.3), respectively.

*Proof.* We prove (a). The proof of (b) is similar to (a) and will be omitted. We



**Partial Sums Of Certain Meromorphic Functions**

Nak Eun Cho and Shigeyoshi Owa

Title Page	
Contents	
◀◀	▶▶
◀	▶
Go Back	
Close	
Quit	
Page 9 of 15	

write

$$\begin{aligned} & \frac{n+2}{1-\alpha} \left[ \frac{f_n(z)}{f(z)} - \frac{n+1+\alpha}{n+2} \right] \\ &= \frac{1 + \sum_{k=1}^n a_k z^{k+1} + \frac{n+1+\alpha}{1-\alpha} \sum_{k=n+1}^{\infty} a_k z^{k+1}}{1 + \sum_{k=1}^n a_k z^{k+1}} \\ &:= \frac{1+w(z)}{1-w(z)}, \end{aligned}$$

where

$$|w(z)| \leq \frac{\frac{n+2}{1-\alpha} \sum_{k=n+1}^{\infty} |a_k|}{2 - 2 \sum_{k=1}^n |a_k| - \frac{n+2\alpha}{1-\alpha} \sum_{k=n+1}^{\infty} |a_k|} \leq 1.$$

This last inequality is equivalent to


$$(2.5) \quad \sum_{k=1}^n |a_k| + \frac{n+1+\alpha}{1-\alpha} \sum_{k=n+1}^{\infty} |a_k| \leq 1.$$

Since the left hand side of (2.5) is bounded above by  $\sum_{k=1}^{\infty} ((k+\alpha)/(1-\alpha))|a_k|$ , the proof is completed.  $\square$

We turn to ratios involving derivatives. The proof of Theorem 2.4 below follows the pattern of those in Theorem 2.1 and (a) of Theorem 2.3 and so the details may be omitted.

**Theorem 2.4.** *If  $f$  of the form (1.1) satisfies condition (1.2) with  $\alpha = 0$ , then*

$$(a) \quad \operatorname{Re} \left\{ \frac{f'(z)}{f_n'(z)} \right\} \geq 0 \quad (z \in \mathcal{U}),$$



### Partial Sums Of Certain Meromorphic Functions

Nak Eun Cho and Shigeyoshi Owa

Title Page

Contents



Go Back

Close

Quit

Page 10 of 15

$$(b) \quad \operatorname{Re} \left\{ \frac{f'_n(z)}{f'(z)} \right\} \geq \frac{1}{2} \quad (z \in \mathcal{U}).$$

In both cases, the extremal function is given by (2.1) with  $\alpha = 0$ .

**Theorem 2.5.** If  $f$  of the form (1.1) satisfies condition (1.3), then

$$(a) \quad \operatorname{Re} \left\{ \frac{f'(z)}{f'_n(z)} \right\} \geq \frac{n + 2\alpha}{n + 1 + \alpha} \quad (z \in \mathcal{U}),$$

$$(b) \quad \operatorname{Re} \left\{ \frac{f'_n(z)}{f'(z)} \right\} \geq \frac{n + 1 + \alpha}{n + 2} \quad (z \in \mathcal{U}).$$

In both cases, the extremal function is given by (2.3).

*Proof.* It is well known that  $f \in \Sigma_k(\alpha) \Leftrightarrow -zf' \in \Sigma^*(\alpha)$ . In particular,  $f$  satisfies condition (1.3) if and only if  $-zf'$  satisfies condition (1.2). Thus, (a) is an immediate consequence of Theorem 2.1 and (b) follows directly from (a) of Theorem 2.3.  $\square$


For a function  $f \in \Sigma$ , we define the integral operator  $F$  as follows:

$$F(z) = \frac{1}{z^2} \int_0^z t f(t) dt = \frac{1}{z} + \sum_{k=1}^{\infty} \frac{1}{k+2} a_k z^k \quad (z \in \mathcal{D}).$$

The  $n$ -th partial sum  $F_n$  of the integral operator  $F$  is given by

$$F_n(z) = \frac{1}{z} + \sum_{k=1}^n \frac{1}{k+2} a_k z^k \quad (z \in \mathcal{D}).$$

The following lemmas will be required for the proof of Theorem 2.8 below.



## Partial Sums Of Certain Meromorphic Functions

Nak Eun Cho and Shigeyoshi Owa

Title Page

Contents



Go Back

Close

Quit

Page 11 of 15

**Lemma 2.6.** For  $0 \leq \theta \leq \pi$ ,

$$\frac{1}{2} + \sum_{k=1}^m \frac{\cos(k\theta)}{k+1} \geq 0.$$

**Lemma 2.7.** Let  $P$  be analytic in  $\mathcal{U}$  with  $P(0) = 1$  and  $\operatorname{Re}\{P(z)\} > \frac{1}{2}$  in  $\mathcal{U}$ . For any function  $Q$  analytic in  $\mathcal{U}$ , the function  $P * Q$  takes values in the convex hull of the image on  $\mathcal{U}$  under  $Q$ .

Lemma 2.6 is due to Rogosinski and Szegő [8] and Lemma 2.7 is a well-known result (c.f. [3, 12]) that can be derived from the Herglotz' representation for  $P$ .

Finally, we derive


**Theorem 2.8.** If  $f \in \Sigma_c(\alpha)$ , then  $F_n \in \Sigma_c(\alpha)$

*Proof.* Let  $f$  be of the form (1.1) and belong to the class  $\Sigma_c(\alpha)$  for  $0 \leq \alpha < 1$ . Since  $-\operatorname{Re}\{z^2 f'(z)\} > \alpha$ , we have

$$(2.6) \quad \operatorname{Re} \left\{ 1 - \frac{1}{2(1-\alpha)} \sum_{k=1}^{\infty} k a_k z^{k+1} \right\} > \frac{1}{2} \quad (z \in \mathcal{U}).$$

Applying the convolution properties of power series to  $F'_n$ , we may write

$$(2.7) \quad \begin{aligned} -z^2 F'_n(z) &= 1 - \sum_{k=1}^n \frac{k}{k+2} a_k z^{k+1} \\ &= \left( 1 - \frac{1}{2(1-\alpha)} \sum_{k=1}^{\infty} k a_k z^{k+1} \right) * \left( 1 + 2(1-\alpha) \sum_{k=1}^{n+1} \frac{1}{k+1} z^k \right). \end{aligned}$$



## Partial Sums Of Certain Meromorphic Functions

Nak Eun Cho and Shigeyoshi Owa

Title Page

Contents



Go Back

Close

Quit

Page 12 of 15

Putting  $z = re^{i\theta}$  ( $0 \leq r < 1$ ,  $0 \leq |\theta| \leq \pi$ ), and making use of the minimum principle for harmonic functions along with Lemma 2.6, we obtain


$$(2.8) \quad \operatorname{Re} \left\{ 1 + 2(1 - \alpha) \sum_{k=1}^{n+1} \frac{1}{k+1} z^k \right\} = 1 + 2(1 - \alpha) \sum_{k=1}^{n+1} \frac{r^k \cos k\theta}{k+1}$$

$$> 1 + 2(1 - \alpha) \sum_{k=1}^{n+1} \frac{\cos k\theta}{k+1} \geq \alpha.$$

In view of (2.6), (2.7), (2.8) and Lemma 2.7, we deduce that

$$-\operatorname{Re}\{z^2 F'_n(z)\} > \alpha \quad (0 \leq \alpha < 1; z \in \mathcal{U}).$$

Therefore we complete the proof of Theorem 2.8. □




---

**Partial Sums Of Certain Meromorphic Functions**

Nak Eun Cho and Shigeyoshi Owa

---

Title Page

Contents



Go Back


Close

Quit

Page 13 of 15

## References

- [1] L. BRICKMAN, D.J. HALLENBECK, T.H. MacGREGOR AND D. WILKEN, Convex hulls and extreme points of families of starlike and convex mappings, *Trans. Amer. Math. Soc.*, **185** (1973), 413–428.
- [2] M.D. GANIGI AND B.A. URALEGADDI, Subclasses of meromorphic close-to-convex functions, *Bull. Math. Soc. Sci. Math. R. S. Roumanie* (N. S.), **33(81)** (1989), 105–109.
- [3] A.W. GOODMAN, *Univalent functions*, Vol. I, Mariner Publ. Co., Tampa, Fl., 1983.
- [4] J.L. LI AND S. OWA, On partial sums of the Libera integral operator, *J. Math. Anal. Appl.*, **213** (1997), 444–454.
- [5] R.J. LIBERA, Some classes of regular univalent functions, *Proc. Amer. Math. Soc.*, **16** (1965), 755–758.
- [6] M.L. MOGRA, T.R. REDDY AND O.P. JUNEJA, Meromorphic univalent functions, *Bull. Austral. Math. Soc.*, **32** (1985), 161–176.
- [7] M.L. MOGRA, Hadamard product of certain meromorphic univalent functions, *J. Math. Anal. Appl.*, **157** (1991), 10–16.
- [8] W. ROGOSINSKI AND G. SZEGÖ, Über die abschnitliche von potenzreihen die in einem kreise beschränkt bleiben, *Math. Z.*, **28** (1928), 73–94.
- [9] T. SHEIL-SMALL, A note on partial sums of convex schlicht functions, *Bull. London Math. Soc.*, **2** (1970), 165–168.



---

### Partial Sums Of Certain Meromorphic Functions

Nak Eun Cho and Shigeyoshi Owa

---

Title Page

Contents




Go Back

Close

Quit

Page 14 of 15

- [10] H. SILVERMAN, Partial sums of starlike and convex functions, *J. Math. Anal. Appl.*, **209** (1997), 221–227.
- [11] E.M. SILVIA, On partial sums of convex functions of order  $\alpha$ , *Houston J. Math.*, **11** (1985), 397–404.
- [12] R. SINGH AND S. SINGH, Convolution properties of a class of starlike functions, *Proc. Amer. Math. Soc.*, **106** (1989), 145–152.
- [13] B.A. URALEGADDI AND M.D. GANIGI, Meromorphic convex functions with negative coefficients, *J. Math. Res. & Exposition*, **7** (1987), 21–26.
- [14] D. YANG AND S. OWA, Subclasses of certain analytic functions, *Hokkaido Math. J.*, **32** (2003), 127–136.



---

**Partial Sums Of Certain  
Meromorphic Functions**

Nak Eun Cho and Shigeyoshi Owa

---

Title Page

Contents



Go Back

Close

Quit

Page 15 of 15