

Journal of Inequalities in Pure and Applied Mathematics

SOME NEW DISCRETE INEQUALITIES AND THEIR APPLICATIONS

Sh. SALEM AND K.R. RASLAN

Department of Mathematics,
Faculty of Science,
Al-Azhar University,
Nasr City, Cairo, Egypt.
EMail: kamal_raslan@yahoo.com



volume 5, issue 1, article 2,
2004.

*Received 27 May, 2002;
accepted 05 August, 2002.*

Communicated by: D. Bainov

Abstract

Contents



Home Page

Go Back

Close

Quit

Abstract


The aim of the present paper is to establish some new linear and nonlinear discrete inequalities in two independent variables. We give some examples in difference equations and we also give numerical test problems for our results.

2000 Mathematics Subject Classification: 26D15

Key words: Discrete inequalities, two independent variables, difference equations, nondecreasing.

Contents

1	Introduction	3
2	Linear Inequality in Two Independent Variables	4
3	Nonlinear Inequalities in Two Independent Variables	7
4	Some Applications	15
5	Conclusions	21
	References	



Some New Discrete Inequalities and Their Applications

Sh. Salem and K.R. Raslan

Title Page

Contents



Go Back


Close

Quit

Page 2 of 22

1. Introduction

The role played by linear and nonlinear discrete inequalities in one and more than one variable in the theory of difference equations and numerical analysis is well known. During the last few years there have been a number of papers written on the discrete inequalities of the Gronwall inequality and its nonlinear version to the Bhiari type, see [1, 2, 3, 4]. In this paper we present several new linear and nonlinear discrete inequalities in two independent variables. Finally, we give two examples to illustrate the importance of our results. Also, we give some numerical examples and compare our theoretical results with the numerical results.



Some New Discrete Inequalities and Their Applications

Sh. Saleem and K.R. Raslan

Title Page

Contents



Go Back

Close

Quit

Page 3 of 22

2. Linear Inequality in Two Independent Variables

Theorem 2.1. Let $u(m, n)$, $a(m, n)$, $b(m, n)$ be nonnegative functions and $a(m, n)$ nondecreasing for $m, n \in \mathbb{N}$. If

$$(2.1) \quad u(m, n) \leq a(m, n) + \sum_{s=0}^{m-1} \sum_{t=0}^{n-1} b(s, t)u(s, t)$$

for $m, n \in \mathbb{N}$, then

$$(2.2) \quad u(m, n) \leq a(m, n) \prod_{t=0}^{n-1} \left[1 + \sum_{s=0}^{m-1} b(s, t) \right].$$

Proof. Define a function $z(m, n)$ by


$$(2.3) \quad z(m, n) = a(m, n) + \sum_{s=0}^{m-1} \sum_{t=0}^{n-1} b(s, t)u(s, t).$$

From (2.1) and (2.3), we have

$$(2.4) \quad u(m, n) \leq z(m, n).$$

Since $a(m, n)$ is nonnegative for $m, n \in \mathbb{N}$, then from (2.3) and (2.4), we get

$$(2.5) \quad \frac{z(m, n)}{a(m, n)} \leq 1 + \sum_{s=0}^{m-1} \sum_{t=0}^{n-1} b(s, t) \frac{z(s, t)}{a(s, t)}.$$



Some New Discrete Inequalities
and Their Applications

Sh. Salem and K.R. Raslan

Title Page

Contents



Go Back

Close

Quit

Page 4 of 22

Define a function $v(m, n)$ by

$$(2.6) \quad v(m, n) = 1 + \sum_{s=0}^{m-1} \sum_{t=0}^{n-1} b(s, t) \frac{z(s, t)}{a(s, t)},$$

then, from (2.5) and (2.6), we get

$$(2.7) \quad z(m, n) \leq a(m, n)v(m, n).$$

From (2.6), we obtain


$$(2.8) \quad v(m+1, n+1) = 1 + b(m, n) \frac{z(m, n)}{a(m, n)} + \sum_{t=0}^{n-1} b(m, t) \frac{z(m, t)}{a(m, t)} \\ + \sum_{s=0}^{m-1} b(s, n) \frac{z(s, n)}{a(s, n)} + \sum_{s=0}^{m-1} \sum_{t=0}^{n-1} b(s, t) \frac{z(s, t)}{a(s, t)},$$

then from (2.6) and (2.8), we get

$$(2.9) \quad v(m+1, n+1) - v(m, n) \\ = b(m, n) \frac{z(m, n)}{a(m, n)} + \sum_{t=0}^{n-1} b(m, t) \frac{z(m, t)}{a(m, t)} + \sum_{s=0}^{m-1} b(s, n) \frac{z(s, n)}{a(s, n)}.$$

Also from (2.7), we have

$$(2.10) \quad v(m+1, n) - v(m, n) = \sum_{t=0}^{n-1} b(m, t) \frac{z(m, t)}{a(m, t)},$$



**Some New Discrete Inequalities
and Their Applications**

Sh. Salem and K.R. Raslan

Title Page

Contents



Go Back

Close

Quit

Page 5 of 22

and

$$(2.11) \quad v(m, n + 1) - v(m, n) = \sum_{s=0}^{m-1} b(s, n) \frac{z(s, n)}{b(s, n)}.$$

From (2.9), (2.10) and (2.11), we get

$$(2.12) \quad [v(m + 1, n + 1) - v(m, n + 1)] - [v(m + 1, n) - v(m, n)] \leq b(m, n)v(m, n).$$


Suppose n is fixed, then from (2.12), we get

$$v(m, n + 1) \leq \left[1 + \sum_{s=0}^{m-1} b(s, n) \right] v(m, n),$$

from which we have

$$(2.13) \quad v(m, n) \leq \prod_{t=0}^{n-1} \left(1 + \sum_{s=0}^{m-1} b(s, n) \right).$$

The required inequality (2.2) follows from (2.4), (2.7) and (2.13). □



**Some New Discrete Inequalities
and Their Applications**

Sh. Salem and K.R. Raslan

Title Page

Contents



Go Back

Close

Quit

Page 6 of 22

3. Nonlinear Inequalities in Two Independent Variables

Theorem 3.1. Let $u(m, n)$, $a(m, n)$, $b(m, n)$ be nonnegative functions and $a(m, n)$ nondecreasing for $m, n \in \mathbb{N}$. If


$$(3.1) \quad u^{m_1}(m, n) \leq a(m, n) + \sum_{s=0}^{m-1} \sum_{t=0}^{n-1} b(s, t) u^{m_2}(s, t).$$

Then

$$(3.2) \quad u(m, n) \leq a^{\frac{1}{m_1}}(m, n) \prod_{t=0}^{n-1} \left[1 + \sum_{s=0}^{m-1} b(s, t) \right]^{\frac{1}{m_1}}; \quad m_1 = m_2,$$

$$(3.3) \quad u(m, n) \leq a^{\frac{1}{m_1}}(m, n) \prod_{t=0}^{n-1} \left[1 + \sum_{s=0}^{m-1} b(s, t) a^{\frac{m_2-m_1}{m_1}}(s, t) \right]^{\frac{m_2(n-t-1)}{m_1^2}}; \quad m_1 < m_2;$$

$$(3.4) \quad u(m, n) \leq a^{\frac{1}{m_1}}(m, n) \prod_{t=0}^{n-1} \left[1 + \sum_{s=0}^{m-1} b(s, t) a^{\frac{m_2-m_1}{m_1}}(s, t) \right]^{\frac{1}{m_1}}; \quad m_1 > m_2.$$



Some New Discrete Inequalities and Their Applications

Sh. Salem and K.R. Raslan

Title Page

Contents



Go Back

Close

Quit

Page 7 of 22

Proof. Define a function $z(m, n)$ by

$$(3.5) \quad z^{m_1}(m, n) = a(m, n) + \sum_{s=0}^{m-1} \sum_{t=0}^{n-1} b(s, t) u^{m_2}(s, t).$$

From (3.1), (3.5), we have

$$(3.6) \quad u(m, n) \leq z(m, n).$$

Since $a(m, n)$ is nonnegative and nondecreasing for $m, n \in \mathbb{N}$; then we get

$$(3.7) \quad \frac{z^{m_1}(m, n)}{a(m, n)} \leq 1 + \sum_{s=0}^{m-1} \sum_{t=0}^{n-1} b(s, t) \frac{u^{m_2}(s, t)}{a(s, t)}.$$

Define function $v(m, n)$ by

$$(3.8) \quad v(m, n) = 1 + \sum_{s=0}^{m-1} \sum_{t=0}^{n-1} b(s, t) \frac{z^{m_2}(s, t)}{a(s, t)},$$


so, we obtain from (3.7) and (3.8) that

$$(3.9) \quad z^{m_1}(m, n) \leq a(m, n) v(m, n).$$

As in Theorem 2.1, from (3.8), we get

$$(3.10) \quad [v(m+1, n+1) - v(m, n+1)] - [v(m+1, n) - v(m, n)] \\ \leq b(m, n) a^{\frac{m_2-m_1}{m_1}}(m, n) v^{\frac{m_2}{m_1}}(m, n).$$

Now, we consider the following cases:



**Some New Discrete Inequalities
and Their Applications**

Sh. Salem and K.R. Raslan

Title Page

Contents




Go Back

Close

Quit

Page 8 of 22



Some New Discrete Inequalities
and Their Applications

Sh. Salem and K.R. Raslan

Title Page

Contents



Go Back

Close

Quit

Page 9 of 22

Case 1. If $m_1 = m_2$, then from (3.10), we have

$$(3.11) \quad v(m+1, n+1) - v(m+1, n) - v(m, n+1) \leq (-1 + b(m, n)) v(m, n),$$

keeping n fixed in (3.11), set $m = 0, 1, 2, \dots, m-1$, then we get

$$(3.12) \quad v(m, n+1) \leq \left[1 + \sum_{s=0}^{m-1} b(s, n) \right] v(m, n).$$

From (3.12), we have

$$(3.13) \quad v(m, n) \leq \prod_{t=0}^{n-1} \left[1 + \sum_{s=0}^{m-1} b(s, t) \right].$$

The required result (3.2) follows from (3.6), (3.9) and (3.13).

Case 2. If $m_2 > m_1$ then as in Case 1 from (3.10), we have

$$(3.14) \quad v(m+1, n+1) - v(m, n+1) - v(m+1, n) + v(m, n) \leq b(m, n) a^{\frac{m_2-m_1}{m_1}}(m, n) v^{\frac{m_2}{m_1}}(m, n),$$

when n is fixed and $m = 0, 1, 2, \dots, m-1$, we obtain from (3.14) that

$$(3.15) \quad v(m, n+1) \leq \left[1 + \sum_{s=0}^{m-1} b(s, n) a^{\frac{m_2-m_1}{m_1}}(s, n) \right] v^{\frac{m_2}{m_1}}(m, n).$$

Lemma 3.2. *If*

$$(3.16) \quad v(m, n + 1) \leq (1 + b(m, n)) v^p(m, n); \quad p > 1,$$

then

$$(3.17) \quad v(m, n) \leq \prod_{t=0}^{n-1} (1 + b(m, t))^{(n-t-1)p}.$$

Then from (3.15), (3.16), (3.17), we get

$$(3.18) \quad v(m, n) \leq \prod_{t=0}^{n-1} \left[1 + \sum_{s=0}^{m-1} b(s, t) a^{\frac{m_2-m_1}{m_1}}(s, t) \right]^{\frac{m_2(n-t-1)}{m_1}}.$$


The required result (3.3) follows from (3.6), (3.9) and (3.18).

Case 3. *If $m_2 < m_1$, then $v^{\frac{m_2}{m_1}}(m, n) \leq v(m, n)$, then, as in the last two cases, we get*

$$(3.19) \quad v(m, n + 1) \leq \left[1 + \sum_{s=0}^{m-1} b(s, n) a^{\frac{m_2-m_1}{m_1}}(s, n) \right] v(m, n).$$

Then from (3.19), we obtain

$$(3.20) \quad v(m, n) \leq \prod_{t=0}^{n-1} \left[1 + \sum_{s=0}^{m-1} b(s, t) a^{\frac{m_2-m_1}{m_1}}(s, t) \right].$$



**Some New Discrete Inequalities
and Their Applications**

Sh. Salem and K.R. Raslan

Title Page

Contents



Go Back

Close

Quit

Page 10 of 22

From (3.6), (3.9) and (3.20), we have

$$u(m, n) \leq a^{\frac{1}{m_1}}(m, n) \prod_{t=0}^{n-1} \left[1 + \sum_{s=0}^{m-1} b(s, t) a^{\frac{m_2-m_1}{m_1}}(s, t) \right]^{\frac{1}{m_1}},$$

which is the required result (3.4). □

Remark 3.1.

1. If $m_1 = m_2 = 1$, then from (3.1) and (3.2), we get the same result as that of Theorem 2.1.
2. If $m_1 = 1, m_2 > 1$, then from (3.1) and (3.2), we get
if


$$(3.21) \quad u(m, n) \leq a(m, n) + \sum_{s=0}^{m-1} \sum_{t=0}^{n-1} b(s, t) u^{m_2}(s, t),$$

then

$$(3.22) \quad u(m, n) \leq a(m, n) \prod_{t=0}^{n-1} \left[1 + \sum_{s=0}^{m-1} b(s, t) a^{m_2-1}(s, t) \right]^{m_2(n-t-1)}.$$

3. Let $m_2 = 1, m_1 > 1$, then from (3.1) and (3.4), we get
if

$$(3.23) \quad u^{m_1}(m, n) \leq a(m, n) + \sum_{s=0}^{m-1} \sum_{t=0}^{n-1} b(s, t) u(s, t),$$



Some New Discrete Inequalities and Their Applications

Sh. Salem and K.R. Raslan

Title Page

Contents




Go Back

Close

Quit

Page 11 of 22



Some New Discrete Inequalities and Their Applications

Sh. Salem and K.R. Raslan

Title Page

Contents



Go Back

Close

Quit

Page 12 of 22

then

$$(3.24) \quad u(m, n) \leq a^{\frac{1}{m_1}}(m, n) \prod_{t=0}^{n-1} \left[1 + \sum_{s=0}^{m-1} b(s, t) a^{\frac{1-m_1}{m_1}}(s, t) \right]^{\frac{1}{m_1}}.$$

Theorem 3.3. Let $u(m, n)$, $a(m, n)$, $b(m, n)$ and $c(m, n)$ be nonnegative and $a(m, n)$ is nondecreasing for $m, n \in \mathbb{N}$, if $m_1, m_2 \in \mathbb{R}^+$, and

$$(3.25) \quad u^{m_1}(m, n) \leq a(m, n) + \sum_{s=0}^{m-1} \sum_{t=0}^{n-1} b(s, t) u(s, t) + \sum_{s=0}^{m-1} \sum_{t=0}^{n-1} c(s, t) u^{m_2}(s, t),$$

then

$$(3.26) \quad u(m, n) \leq a(m, n) \prod_{t=0}^{n-1} \left[1 + \sum_{s=0}^{m-1} (b(s, t) + c(s, t)) \right], \quad m_1 = m_2 = 1,$$

$$(3.27) \quad u(m, n) \leq a^{\frac{1}{m_1}}(m, n) \prod_{t=0}^{n-1} \left[1 + \sum_{s=0}^{m-1} (c(s, t) + b(s, t)) a^{\frac{1-m_1}{m_1}}(s, t) \right]^{\frac{1}{m_1}},$$

$m_1 = m_2 > 1,$


$$(3.28) \quad u(m, n) \leq a^{\frac{1}{m_1}}(m, n) \prod_{t=0}^{n-1} \left[1 + \sum_{s=0}^{m-1} (c(s, t) + b(s, t)) \times a^{\frac{1-m_1}{m_1}}(s, t) \right]^{\frac{n-t-1}{m_1^2}}, \quad 0 < m_1 = m_2 < 1,$$

$$(3.29) \quad u(m, n) \leq a^{\frac{1}{m_1}}(m, n) \prod_{t=0}^{n-1} \left[1 + \sum_{s=0}^{m-1} \left(b(s, t) a^{\frac{1-m_1}{m_1}}(s, t) + c(s, t) a^{\frac{m_2-m_1}{m_1}}(s, t) \right) \right]^{\frac{m_2(n-t-1)}{m_1^2}}, \quad m_2 > m_1,$$

$$(3.30) \quad u(m, n) \leq a^{\frac{1}{m_1}}(m, n) \prod_{t=0}^{n-1} \left[1 + \sum_{s=0}^{m-1} \left(b(s, t) a^{\frac{1-m_1}{m_1}}(s, t) + c(s, t) a^{\frac{m_2-m_1}{m_1}}(s, t) \right) \right]^{\frac{1}{m_1}}, \quad 1 \leq m_2 < m_1,$$

and

$$(3.31) \quad u(m, n) \leq a^{\frac{1}{m_1}}(m, n) \prod_{t=0}^{n-1} \left[1 + \sum_{s=0}^{m-1} \left(b(s, t) a^{\frac{1-m_1}{m_1}}(s, t) + c(s, t) a^{\frac{m_2-m_1}{m_1}}(s, t) \right) \right]^{\frac{n-t-1}{m_1^2}}, \quad 0 < m_2 < m_1 < 1.$$



**Some New Discrete Inequalities
and Their Applications**

Sh. Salem and K.R. Raslan

Title Page

Contents



Go Back

Close


Quit

Page 13 of 22

Proof. The proof of this theorem is similar to the proof of Theorem 3.1. Here we leave the details to the reader. \square

Remark 3.2.

1. If $c(m, n) = 0$, $m_1 = m_2$, then we get Theorem 2.1.
2. If $b(m, n) = 0$, then we get Theorem 3.1.



**Some New Discrete Inequalities
and Their Applications**

Sh. Salem and K.R. Raslan

Title Page

Contents



Go Back

Close

Quit

Page 14 of 22

4. Some Applications

There are many possible applications of the inequality established in this paper, but those presented here are sufficient to convey the importance of our results.

Example 4.1. Consider the difference equation

$$(4.1) \quad u(m, n) = a(m, n) + \sum_{s=0}^{m-1} \sum_{t=0}^{n-1} k(s, t, u(s, t)).$$

Let

$$(4.2) \quad k(s, t, u(s, t)) \leq t u(s, t),$$

from (4.1), (4.2), we get

$$(4.3) \quad u(m, n) \leq a(m, n) + \sum_{s=0}^{m-1} \sum_{t=0}^{n-1} t u(s, t).$$


From (2.1), (2.2) and (4.1) we get

$$(4.4) \quad u(m, n) \leq a(m, n) \prod_{t=0}^{n-1} (1 + m t).$$

Remark 4.1.

1. If

$$(4.5) \quad k(s, t, u(s, t)) \leq 2 s t u(s, t),$$



Some New Discrete Inequalities
and Their Applications

Sh. Salem and K.R. Raslan

Title Page

Contents




Go Back

Close

Quit

Page 15 of 22



Some New Discrete Inequalities
and Their Applications

Sh. Salem and K.R. Raslan

Title Page

Contents



Go Back

Close

Quit

Page 16 of 22

then, we get

$$(4.6) \quad u(m, n) \leq a(m, n) \prod_{t=0}^{n-1} (1 + m(m-1)t).$$

2. If

$$(4.7) \quad k(s, t, u(s, t)) \leq u(s, t),$$

then, we get

$$(4.8) \quad u(m, n) \leq a(m, n) \prod_{t=0}^{n-1} (1 + m) = a(m, n)(1 + m)^n.$$

Example 4.2. Consider the difference equation


$$(4.9) \quad u^{m_1}(m, n) = a(m, n) + \sum_{s=0}^{m-1} \sum_{t=0}^{n-1} k(s, t, u(s, t)).$$

let

$$(4.10) \quad k(s, t, u(s, t)) \leq b(s, t) u(s, t),$$

if we consider $a(s, t) = b(s, t) = t$, from (3.23) and (3.24) we get

$$(4.11) \quad u(m, n) \leq n^{\frac{1}{m_1}} \prod_{t=0}^{n-1} \left[1 + mt^{\frac{1}{m_1}} \right]^{\frac{1}{m_1}}.$$



Title Page

Contents



Go Back

Close

Quit

Page 17 of 22

Example 4.3. Consider the difference equation

$$(4.12) \quad u^{m_1}(m, n) = a(m, n) + \sum_{s=0}^{m-1} \sum_{t=0}^{n-1} k(s, t, u(s, t)).$$

Let

$$(4.13) \quad k(s, t, u(s, t)) \leq b(s, t) u(s, t) + b(s, t) u^{m_2}(s, t),$$


if we take $m_1 = 3$, $m_2 = 2$, $a(s, t) = b(s, t) = c(s, t) = t^3$, then from (3.30) we have

$$(4.14) \quad u(m, n) \leq n \prod_{t=0}^{n-1} [1 + mt(t+1)]^{\frac{1}{3}},$$

As special cases of (4.14), let $m = 2$ and $n = 2$, then $u(2, 2) \leq 2\sqrt[3]{5}$, if we take $m = 2$ and $n = 3$, then $u(2, 3) \leq 3\sqrt[3]{45}$, also for $m = 3$ and $n = 2$ then $u(3, 2) \leq 2\sqrt[3]{7}$.

Example 4.4. Consider the difference inequality as in (2.1) with $a(s, t) = \alpha(st + 5)$, $b(s, t) = \alpha(2t + s^2 + 1)$, $\alpha = 10^{-6}$, and we compute the values of $u(m, n)$ from (2.1) and also we find the values of $u(m, n)$ by using the result (2.2). In our computations we use (2.1) and (2.2) as equations and as we see in the Table 1 the computation values as in (2.1) are less than the values of the result (2.2).

Example 4.5. Consider the difference as in (3.1) with $a(s, t) = \alpha(t + s^2 + st)$, $b(s, t) = \beta(t + s^2 + 6)$, $\beta = 10^{-6}$, $\alpha = 10^{-5}$, and we compute the values of $u(m, n)$ from (3.1) and also we find the values of $u(m, n)$ by using the results



Some New Discrete Inequalities
and Their Applications

Sh. Salem and K.R. Raslan

Title Page

Contents



Go Back

Close

Quit

Page 18 of 22


m, n	(2.1)	(2.2)	m, n	(2.1)	(2.2)
1,1	6.0e-6	6.0e-6	6,1	1.103019e-5	1.616208e-5
1,10	1.5e-5	1.513245e-5	6,5	3.503161e-5	5.506815e-5
2,1	7.000005e-5	7.082850e-5	6,10	6.504472e-5	1.124603e-4
2,5	1.500034e-5	1.537153e-5	7,1	1.204472e-5	2.119437e-5
2,10	2.500372e-5	2.623212e-6	7,5	4.004651e-5	7.685613e-5
3,1	8.003725e-6	8.445905e-6	7,10	7.506240e-5	1.616860e-4
3,5	2.000427e-5	2.166051e-5	8,5	4.506461e-5	1.099740e-4
3,10	3.500988e-5	3.945630e-5	8,10	8.508343e-5	2.380212e-4
4,1	9.009876e-6	1.024249e-6	9,1	1.408343e-5	4.035491e-5
4,5	2.501068e-5	2.958550e-5	9,5	5.008608e-5	1.621632e-4
4,10	4.501864e-5	5.640386e-5	9,10	9.510797e-5	3.608566e-4
5,1	1.001864e-5	1.269826e-5	10,1	1.510797e-5	5.891117e-5
5,5	3.001973e-5	4.018442e-5	10,5	5.511112e-5	2.474333e-4
5,10	5.503019e-5	7.947053e-5	10,10	1.051362e-4	5.659792e-4

Table 1:

(3.2) – (3.4) and tabled them in the following Table 2.

Example 4.6. Consider the difference as in (4.1) with $a(s, t) = \alpha(t^2 + s + st)$, $b(s, t) = \alpha(t^2 + s + 6)$, $c(s, t) = \alpha(s + t + 1)$, $\alpha = 10^{-6}$, and we compute the values of $u(m, n)$ from (3.25) and also we find the values of $u(m, n)$ by using the results (3.26) – (3.31) and tabled them in the following Table 3.

From Tables 1, 2 and 3, we can say that the numerical solution agrees with the analytical solution for some discrete linear and nonlinear inequalities. The programs for each case are written in the programming language Fortran.



**Some New Discrete Inequalities
and Their Applications**

Sh. Salem and K.R. Raslan

Case m, n	$m_1 = m_2 = 2$		$1 = m_1 < m_2 = 4$		$1 = m_1 > m_2 = 0.6$	
	(3.1)	(3.2)	(3.1)	(3.3)	(3.1)	(3.4)
1,1	5.477225e-3	5.477390e-3	3.0e-5	3.0e-5	3.0e-5	3.0e-5
1,10	1.449138e-2	1.452727e-2	2.1e-4	2.1e-4	2.1e-4	2.1e-4
2,1	8.366600e-3	8.392021e-3	6.7e-5	6.7e-5	6.999999e-5	7.000654e-5
2,10	1.843944e-2	1.863311e-2	3.4e-4	3.4e-4	3.404703e-4	3.401309e-4
3,1	1.140233e-2	1.153497e-2	1.3e-4	1.3e-4	1.304703e-4	1.300566e-4
3,10	2.213686e-2	2.269679e-2	4.9e-4	4.9e-4	4.912776e-4	4.905238e-4
4,1	1.449278e-2	1.488674e-2	2.1e-4	2.1e-4	2.112776e-4	2.102450e-4
4,10	2.569232e-2	2.695817e-2	6.6e-4	6.6e-4	6.626532e-4	6.615494e-4
5,1	1.760952e-2	1.852931e-2	3.1e-4	3.1e-4	3.126533e-4	3.107818e-4
5,10	2.915815e-2	3.167530e-2	8.5e-4	8.5e-4	8.549279e-4	8.538832e-4
6,1	2.074121e-2	2.262540e-2	4.3e-4	4.3e-4	4.349280e-4	4.320908e-4
6,10	3.256347e-2	3.719871e-2	1.06e-3	1.06e-3	1.068537e-3	1.068736e-3
7,1	2.388262e-2	2.744415e-2	5.7e-4	5.7e-4	5.785374e-4	5.749682e-4
7,10	3.592609e-2	4.405006e-2	1.29e-3	1.29e-3	1.304028e-3	1.308159e-3
8,1	2.703117e-2	3.341747e-2	7.3e-4	7.3e-4	7.440277e-4	7.408159e-4
8,10	3.925774e-2	5.304486e-2	1.54e-3	1.54e-3	1.562061e-3	1.575471e-3
9,1	3.018559e-2	4.124418e-2	9.1e-4	9.1e-4	9.320608e-4	9.319749e-4
9,10	4.256656e-2	6.551374e-2	1.81e-3	1.81e-3	1.843420e-3	1.875835e-3
10,1	3.334534e-2	5.208647e-2	1.11e-3	1.11e-3	1.143420e-3	1.152192e-3
10,10	4.585852e-2	8.372936e-2	2.1e-3	2.1e-3	2.149017e-3	2.217082e-3

Table 2:

Title Page

Contents




Go Back

Close

Quit

Page 19 of 22



**Some New Discrete Inequalities
and Their Applications**

Sh. Salem and K.R. Raslan

Case	$m_1 = m_2 = 1$		$m_1 = m_2 = 2 > 1$		$0 < m_1 = m_2 = 0.5 < 1$	
m, n	(3.25)	(3.26)	(3.25)	(3.27)	(3.25)	(3.28)
1,1	3.000000e-5	3.000210e-5	5.477225e-3	5.477390e-3	9.0e-10	9.01946e-10
1,10	2.100000e-4	2.115060e-4	1.449138e-2	1.452747e-2	4.41e-8	4.649055e-8
2,10	3.400212e-4	3.508332e-4	1.843980e-2	1.863446e-2	1.156178e-7	1.562039e-7
3,10	4.900642e-4	5.285408e-4	2.213770e-2	2.268906e-2	2.401761e-7	5.146701e-7
4,10	6.601446e-4	7.631531e-4	2.569380e-2	2.688507e-2	4.358260e-7	1.866012e-6
5,10	8.502871e-4	1.087983e-3	2.916045e-2	3.137589e-2	7.230648e-7	7.930772e-6
6,10	1.060527e-3	1.564760e-3	3.256680e-2	3.631432e-2	1.124866e-6	4.124445e-5
7,10	1.210846e-3	2.312758e-3	3.593069e-2	4.187295e-2	1.666718e-6	2.713262e-4
8,10	1.541507e-3	3.575617e-3	3.926384e-2	4.826101e-2	2.376683e-6	2.321973e-3
9,10	1.812392e-3	5.886483e-3	4.257445e-2	5.574329e-2	3.285458e-6	2.650311e-2
10,10	2.103667e-3	1.051122e-2	4.586849e-2	6.466483e-2	4.426470e-6	4.128818e-1
Case	$2 = m_2 > m_1 = 1$		$1 \leq m_2 = 1.5 < m_1 = 2$		$0 < m_2 = 0.2 < m_1 = 0.8$	
m, n	(3.25)	(3.29)	(3.25)	(3.27)	(3.25)	(3.31)
1,1	3.000000e-5	3.003242e-5	5.477225e-3	5.477225e-3	2.220248e-6	2.220248e-6
1,10	2.100000e-4	2.156167e-4	1.449138e-2	1.449402e-2	2.527983e-5	2.530985e-5
2,10	3.400000e-4	3.952264e-4	1.844572e-2	1.845583e-2	4.996473e-5	4.658744e-5
3,10	4.900000e-4	7.221420e-4	2.215105e-2	2.218821e-2	8.272198e-5	7.483793e-5
4,10	6.600000e-4	1.427778e-3	2.571710e-2	2.581235e-2	1.247796e-4	1.120071e-4
5,10	8.500001e-4	3.288740e-3	2.919728e-2	2.939673e-2	1.776342e-4	1.617969e-4
6,10	1.060000e-3	9.571671e-3	3.262191e-2	3.299124e-2	2.431235e-4	2.313847e-4
7,10	1.290001e-3	3.864970e-2	3.601008e-2	3.664057e-2	3.234317e-4	3.350727e-4
8,10	1.450001e-3	2.411131e-1	3.937489e-2	4.039110e-2	4.211098e-4	5.026789e-4
9,10	1.810002e-3	2.625509	4.272596e-2	4.429572e-2	5.391026e-4	8.007679e-4
10,10	2.100004e-3	57.185780	4.607081e-2	4.831796e-2	6.807797e-4	1.392293e-3

Table 3:

Title Page

Contents



Go Back


Close

Quit

Page 20 of 22

5. Conclusions

This study presents the design and implementation of new discrete linear and nonlinear inequalities in one and two independent variables. We give new theoretical studies for those inequalities as in Section 3. We give test problems to demonstrate our results with different cases as we have shown in Section 4. We believe that the present studies can be useful for other applications and be extended to more complicated problems in higher dimensions.



Some New Discrete Inequalities and Their Applications

Sh. Salem and K.R. Raslan

Title Page

Contents



Go Back


Close

Quit

Page 21 of 22

References

- [1] D. BAINOV AND P. SIMEONOV, *Integral Inequalities and Applications*, Kluwer Academic Publishers, Dordrecht, 1992.
- [2] E.H. YANG, On some new discrete inequalities of the Bellman-Bihari type, *Nonlinear Anal.*, **7** (1983), 1238–1246.
- [3] E.H. YANG, On some new discrete generalizations of Gronwall's inequality, *J. Math. Anal. Appl.*, **129** (1988), 505–516.
- [4] Sh. SALEM, On Some System of Two Discrete Inequalities of Gronwall Type, *J. Math. Anal. Appl.*, **208** (1997), 553–566.



Some New Discrete Inequalities and Their Applications

Sh. Salem and K.R. Raslan

Title Page

Contents



Go Back

Close

Quit

Page 22 of 22