

# INEQUALITIES INVOLVING THE INNER PRODUCT

**DORIAN POPA AND IOAN RAȘA**

Technical University of Cluj-Napoca

Department of Mathematics

Str. C. Daicoviciu 15

Cluj-Napoca, Romania

E-Mail: {[@Popa.Dorian](mailto:Popa.Dorian), [@Ioan.Rasa](mailto:Ioan.Rasa)}@math.utcluj.ro

*Received:* 20 February, 2007

*Accepted:* 16 July, 2007

*Communicated by:* **S.S. Dragomir**

*2000 AMS Sub. Class.:* 26D20.

*Key words:* Inner product, Schwarz inequality.

*Abstract:* The paper contains inequalities related to generalizations of Schwarz's inequality.





**Inequalities Involving  
the Inner Product**

Dorian Popa and Ioan Rașa

vol. 8, iss. 3, art. 86, 2007

[Title Page](#)

[Contents](#)



Page 1 of 8

[Go Back](#)

[Full Screen](#)


[Close](#)

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-5756

# Contents

1	Introduction	3
2	The Results	4



---

**Inequalities Involving  
the Inner Product**

Dorian Popa and Ioan Raşa

vol. 8, iss. 3, art. 86, 2007

---

Title Page

Contents



Page 2 of 8


Go Back

Full Screen

Close

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-5756



Title Page

Contents



Page 3 of 8

Go Back

Full Screen

Close

## 1. Introduction


Let  $(H, \langle \cdot, \cdot \rangle)$  be a Hilbert space over the field  $K = \mathbb{R}$  or  $K = \mathbb{C}$ . Then for all  $x, a, b \in H$  the following inequality holds:

$$(1.1) \quad \left| \langle a, x \rangle \langle x, b \rangle - \frac{1}{2} \langle a, b \rangle \|x\|^2 \right| \leq \frac{1}{2} \|a\| \|b\| \|x\|^2.$$

In particular, for  $a = b$  (1.1) reduces to Schwarz's inequality.

For historical remarks, proofs, extensions, generalizations and applications of (1.1), see [1] – [5] and the references given therein.

In this paper we consider a suitable quadratic form and derive inequalities related to (1.1). More precisely, we obtain estimates involving the real and the imaginary part of the expression whose absolute value is contained in the left hand of (1.1).



## 2. The Results

Let  $v_1, \dots, v_n$  ( $n \geq 2$ ) be linearly independent vectors in  $H$ , and  $v := v_1 + \dots + v_n$ .

Consider the matrix

$$A := \begin{pmatrix} \langle v_1, v - v_1 \rangle & \langle v_2, v - v_1 \rangle & \cdots & \langle v_n, v - v_1 \rangle \\ \langle v_1, v - v_2 \rangle & \langle v_2, v - v_2 \rangle & \cdots & \langle v_n, v - v_2 \rangle \\ \vdots & \vdots & \cdots & \vdots \\ \langle v_1, v - v_n \rangle & \langle v_2, v - v_n \rangle & \cdots & \langle v_n, v - v_n \rangle \end{pmatrix}$$

### Theorem 2.1.

(i) The matrix  $A$  has real eigenvalues  $\lambda_1 \leq \dots \leq \lambda_n$ ; moreover,  $\lambda_1 \leq 0$  and  $\lambda_n \geq 0$ .

(ii) The following inequalities hold:


$$(2.1) \quad \lambda_1 \|x\|^2 \leq \sum_{\substack{i,j=1 \\ i \neq j}}^n \langle v_i, x \rangle \langle x, v_j \rangle \leq \lambda_n \|x\|^2, \quad x \in H.$$

*Proof.* Let  $H_n$  denote the linear subspace of  $H$  generated by  $v_1, \dots, v_n$ . Consider the linear operator  $T : H_n \rightarrow H_n$  defined by

$$Tx = \langle x, v \rangle v - \sum_{i=1}^n \langle x, v_i \rangle v_i, \quad x \in H_n.$$

Then for all  $x, y \in H_n$  we have

$$\langle Tx, y \rangle = \langle x, v \rangle \langle v, y \rangle - \sum_{i=1}^n \langle x, v_i \rangle \langle v_i, y \rangle$$



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 5 of 8

Go Back

Full Screen

Close

$$\begin{aligned}
 &= \overline{\langle y, v \rangle} \langle x, v \rangle - \sum_{i=1}^n \overline{\langle y, v_i \rangle} \langle x, v_i \rangle \\
 &= \left\langle x, \langle y, v \rangle v - \sum_{i=1}^n \langle y, v_i \rangle v_i \right\rangle = \langle x, Ty \rangle.
 \end{aligned}$$

We conclude that  $T$  is self-adjoint, hence it has real eigenvalues  $\lambda_1 \leq \dots \leq \lambda_n$  and:

$$(2.2) \quad \lambda_1 \|x\|^2 \leq \langle Tx, x \rangle \leq \lambda_n \|x\|^2, \quad x \in H_n.$$

On the other hand,

$$\begin{aligned}
 Tv_j &= \langle v_j, v \rangle v - \sum_{i=1}^n \langle v_j, v_i \rangle v_i \\
 &= \sum_{i=1}^n (\langle v_j, v \rangle - \langle v_j, v_i \rangle) v_i = \sum_{i=1}^n \langle v_j, v - v_i \rangle v_i
 \end{aligned}$$

for all  $j = 1, 2, \dots, n$ .

This means that  $A$  is the matrix of  $T$  with respect to the basis  $\{v_1, v_2, \dots, v_n\}$  of  $H_n$ , and so  $\lambda_1 \leq \dots \leq \lambda_n$  are the eigenvalues of the matrix  $A$ .

Now remark that

$$\begin{aligned}
 \langle Tx, x \rangle &= \langle x, v \rangle \langle v, x \rangle - \sum_{i=1}^n \langle x, v_i \rangle \langle v_i, x \rangle \\
 &= \sum_{i=1}^n \langle v_i, x \rangle \sum_{i=1}^n \langle x, v_i \rangle - \sum_{i=1}^n \langle v_i, x \rangle \langle x, v_i \rangle
 \end{aligned}$$

$$= \sum_{\substack{i,j=1 \\ i \neq j}}^n \langle v_i, x \rangle \langle x, v_j \rangle, \quad x \in H_n.$$

Combined with (2.2), this gives

$$(2.3) \quad \lambda_1 \|x\|^2 \leq \sum_{\substack{i,j=1 \\ i \neq j}}^n \langle v_i, x \rangle \langle x, v_j \rangle \leq \lambda_n \|x\|^2, \quad x \in H_n.$$

Let  $x \in H_n$ ,  $x \neq 0$ ,  $\langle x, v_i \rangle = 0$ ,  $i = 1, 2, \dots, n-1$ .

From (2.3) we infer that

$$(2.4) \quad \lambda_1 \leq 0 \leq \lambda_n.$$

Let  $y \in H$ . Then  $y = x + z$ ,  $x \in H_n$ ,  $z \in H_n^\perp$  and  $\|y\|^2 = \|x\|^2 + \|z\|^2$ , so that  $\|y\|^2 \geq \|x\|^2$ . Moreover,

$$\langle v_i, y \rangle = \langle v_i, x + z \rangle = \langle v_i, x \rangle, \quad i = 1, \dots, n.$$


Using (2.3) and (2.4), we get

$$\lambda_1 \|y\|^2 \leq \lambda_1 \|x\|^2 \leq \sum_{\substack{i,j=1 \\ i \neq j}}^n \langle v_i, y \rangle \langle y, v_j \rangle \leq \lambda_n \|x\|^2 \leq \lambda_n \|y\|^2$$

and this concludes the proof. □

**Corollary 2.2.** *Let  $a, b, x \in H$ . Then*

$$(2.5) \quad \left| \operatorname{Re} \left( \langle a, x \rangle \langle x, b \rangle - \frac{1}{2} \|x\|^2 \langle a, b \rangle \right) \right| \leq \frac{1}{2} \|x\|^2 \sqrt{\|a\|^2 \|b\|^2 - (\operatorname{Im} \langle a, b \rangle)^2}.$$



Title Page

Contents

◀◀ ▶▶


◀ ▶

Page 6 of 8

Go Back



Full Screen

Close



Title Page

Contents



Page 7 of 8

Go Back

Full Screen

Close

*Proof.* If  $a$  and  $b$  are linearly dependent, (2.5) can be verified directly. Otherwise it is a consequence of Theorem 2.1.

Indeed for  $n = 2$ ,  $v_1 = a$ ,  $v_2 = b$ , the eigenvalues of the matrix  $A$  are

$$\lambda_{1,2} = \operatorname{Re}\langle a, b \rangle \pm \sqrt{\|a\|^2\|b\|^2 - (\operatorname{Im}\langle a, b \rangle)^2}$$

and

$$\langle Tx, x \rangle = 2 \operatorname{Re}\langle x, a \rangle \langle b, x \rangle, \quad x \in H.$$

□

*Remark 1.*

(i) When  $K = \mathbb{R}$ , (2.5) coincides with (1.1).

(ii) Let  $K = \mathbb{C}$ . Applying Corollary 2.2 to the vectors  $ia, b, x$  we get

$$(2.6) \quad \left| \operatorname{Im} \left( \langle a, x \rangle \langle x, b \rangle - \frac{1}{2} \|x\|^2 \langle a, b \rangle \right) \right| \leq \frac{1}{2} \|x\|^2 \sqrt{\|a\|^2\|b\|^2 - (\operatorname{Re}\langle a, b \rangle)^2}.$$

## References

- [1] M.L. BUZANO, Generalizzazione della disuguglianza di Cauchy-Schwarz (Italian), *Rend. Sem. Mat. Univ. e Politech. Torino*, **31**(1971/73), 405–409(1974).
- [2] S.S. DRAGOMIR, Refinements of Buzano's and Kurepa's inequalities in inner product spaces, *Facta Univ. (Niš), Ser. Math. Inform.*, **20** (2005), 65–73.
- [3] S.S. DRAGOMIR, A potpourri of Schwarz related inequalities in inner product spaces (I), *J. Inequal. Pure Appl. Math.*, **6**(3) (2005), Art. 59. [ONLINE: <http://jipam.vu.edu.au/article.php?sid=532>].
- [4] J.E. PEČARIĆ, On some classical inequalities in unitary spaces, *Mat. Bilten*, **16** (1992), 63–72.
- [5] T. PRECUPANU, On a generalization of Cauchy-Buniakowski-Schwarz inequality, *Anal. St. Univ. "Al. I. Cuza" Iași*, **22**(2) (1976), 173–175.



Inequalities Involving  
the Inner Product

Dorian Popa and Ioan Raşa

vol. 8, iss. 3, art. 86, 2007

Title Page

Contents



Page 8 of 8

Go Back

Full Screen

Close

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-5756