



## A NOTE ON THE MAGNITUDE OF WALSH FOURIER COEFFICIENTS

B. L. GHODADRA AND J. R. PATADIA

DEPARTMENT OF MATHEMATICS  
FACULTY OF SCIENCE  
THE MAHARAJA SAYAJIRAO UNIVERSITY OF BARODA  
VADODARA - 390 002 (GUJARAT), INDIA.  
bhikhu\_ghodadra@yahoo.com

jamanadaspat@gmail.com

*Received 11 March, 2008; accepted 07 May, 2008*

*Communicated by L. Leindler*

---

**ABSTRACT.** In this note, the order of magnitude of Walsh Fourier coefficients for functions of the classes  $BV^{(p)}$  ( $p \geq 1$ ),  $\phi BV$ ,  $\Lambda BV^{(p)}$  ( $p \geq 1$ ) and  $\phi \Lambda BV$  is studied. For the classes  $BV^{(p)}$  and  $\phi BV$ , Taibleson-like technique for Walsh Fourier coefficients is developed.

However, for the classes  $\Lambda BV^{(p)}$  and  $\phi \Lambda BV$  this technique seems to be not working and hence classical technique is applied. In the case of  $\Lambda BV$ , it is also shown that the result is best possible in a certain sense.

---

*Key words and phrases:* Functions of  $p$ -bounded variation,  $\phi$ -bounded variation,  $p - \Lambda$ -bounded variation and of  $\phi - \Lambda$ -bounded variation, Walsh Fourier coefficients, Integral modulus continuity of order  $p$ .

2000 *Mathematics Subject Classification.* 42C10, 26D15.

### 1. INTRODUCTION

It appears that while the study of the order of magnitude of the trigonometric Fourier coefficients for the functions of various classes of generalized variations such as  $BV^{(p)}$  ( $p \geq 1$ ) [9],  $\phi BV$  [2],  $\Lambda BV$  [8],  $\Lambda BV^{(p)}$  ( $p \geq 1$ ) [5],  $\phi \Lambda BV$  [4], etc. has been carried out, such a study for the Walsh Fourier coefficients has not yet been done. The only result available is due to N.J. Fine [1], who proves, using the second mean value theorem that, if  $f \in BV[0, 1]$  then its Walsh Fourier coefficients  $\hat{f}(n) = O(\frac{1}{n})$ . In this note we carry out this study. Interestingly, here, no use of the second mean value theorem is made. We also prove that for the class  $\Lambda BV$ , our result is best possible in a certain sense.

**Definition 1.1.** Let  $I = [a, b]$ ,  $p \geq 1$  be a real number,  $\{\lambda_k\}$ ,  $k \in \mathbb{N}$ , be a sequence of non-decreasing positive real numbers such that  $\sum_{k=1}^{\infty} \frac{1}{\lambda_k}$  diverges and  $\phi : [0, \infty) \rightarrow \mathbb{R}$ , be a strictly increasing function. We say that:

(1)  $f \in BV^{(p)}(I)$  (that is,  $f$  is of  $p$ -bounded variation over  $I$ ) if

$$V(f, p, I) = \sup_{\{I_k\}} \left\{ \left( \sum_k |f(b_k) - f(a_k)|^p \right)^{\frac{1}{p}} \right\} < \infty,$$

(2)  $f \in \phi BV(I)$  (that is,  $f$  is of  $\phi$ -bounded variation over  $I$ ) if

$$V(f, \phi, I) = \sup_{\{I_k\}} \left\{ \sum_k \phi(|f(b_k) - f(a_k)|) \right\} < \infty,$$

(3)  $f \in \Lambda BV^{(p)}(I)$  (that is,  $f$  is of  $p - \Lambda$ -bounded variation over  $I$ ) if

$$V_{\Lambda}(f, p, I) = \sup_{\{I_k\}} \left\{ \left( \sum_k \frac{|f(b_k) - f(a_k)|^p}{\lambda_k} \right)^{\frac{1}{p}} \right\} < \infty,$$

(4)  $f \in \phi \Lambda BV(I)$  (that is,  $f$  is of  $\phi - \Lambda$ -bounded variation over  $I$ ) if

$$V_{\Lambda}(f, \phi, I) = \sup_{\{I_k\}} \left\{ \sum_k \frac{\phi(|f(b_k) - f(a_k)|)}{\lambda_k} \right\} < \infty,$$

in which  $\{I_k = [a_k, b_k]\}$  is a sequence of non-overlapping subintervals of  $I$ .

In (2) and (4), it is customary to consider  $\phi$  a convex function such that

$$\phi(0) = 0, \quad \frac{\phi(x)}{x} \rightarrow 0 \quad (x \rightarrow 0_+), \quad \frac{\phi(x)}{x} \rightarrow \infty \quad (x \rightarrow \infty);$$

such a function is necessarily continuous and strictly increasing on  $[0, \infty)$ .

Let  $\{\varphi_n\}$  ( $n = 0, 1, 2, 3, \dots$ ) denote the complete orthonormal Walsh system [7], where the subscript denotes the number of zeros (that is, sign-changes) in the interior of the interval  $[0, 1]$ . For a 1-periodic  $f$  in  $L[0, 1]$  its Walsh Fourier series is given by

$$f(x) \sim \sum_{n=0}^{\infty} \hat{f}(n) \varphi_n(x),$$

where the  $n^{\text{th}}$  Walsh Fourier coefficient  $\hat{f}(n)$  is given by

$$\hat{f}(n) = \int_0^1 f(x) \varphi_n(x) dx \quad (n = 0, 1, 2, 3, \dots).$$

The Walsh system can be realized [1] as the full set of characters of the dyadic group  $G = Z_2^{\infty}$ , in which  $Z_2 = \{0, 1\}$  is the group under addition modulo 2. We denote the operation of  $G$  by  $\dot{+}$ .  $(G, \dot{+})$  is identified with  $([0, 1], +)$  under the usual convention for the binary expansion of elements of  $[0, 1]$  [1].

## 2. RESULTS

We prove the following theorems. In Theorem 2.5 it is shown that Theorem 2.3 with  $p = 1$  is best possible in a certain sense.

**Theorem 2.1.** *If  $f \in BV^{(p)}[0, 1]$  then  $\hat{f}(n) = O\left(1 / \left(n^{\frac{1}{p}}\right)\right)$ .*

**Note.** Theorem 2.1 with  $p = 1$  gives the result of Fine [1, Theorem VI].

**Theorem 2.2.** *If  $f \in \phi BV[0, 1]$  then  $\hat{f}(n) = O(\phi^{-1}(1/n))$ .*

**Theorem 2.3.** *If 1-periodic  $f \in \Lambda BV^{(p)}[0, 1]$  ( $p \geq 1$ ) then*

$$\hat{f}(n) = O \left( 1 / \left( \sum_{j=1}^n \frac{1}{\lambda_j} \right)^{\frac{1}{p}} \right).$$

**Theorem 2.4.** *If 1-periodic  $f \in \phi \Lambda BV[0, 1]$  then*

$$\hat{f}(n) = O \left( \phi^{-1} \left( 1 / \left( \sum_{j=1}^n \frac{1}{\lambda_j} \right) \right) \right).$$

**Theorem 2.5.** *If  $\Gamma BV[0, 1] \supseteq \Lambda BV[0, 1]$  properly then*

$$\exists f \in \Gamma BV[0, 1] \ni \hat{f}(n) \neq O \left( 1 / \left( \sum_{j=1}^n \frac{1}{\lambda_j} \right) \right).$$

*Proof of Theorem 2.1.* Let  $n \in \mathbb{N}$ . Let  $k \in \mathbb{N} \cup \{0\}$  be such that  $2^k \leq n < 2^{k+1}$  and put  $a_i = (i/2^k)$  for  $i = 0, 1, 2, 3, \dots, 2^k$ . Since  $\varphi_n$  takes the value 1 on one half of each of the intervals  $(a_{i-1}, a_i)$  and the value  $-1$  on the other half, we have

$$\int_{a_{i-1}}^{a_i} \varphi_n(x) dx = 0, \quad \text{for all } i = 1, 2, 3, \dots, 2^k.$$

Define a step function  $g$  by  $g(x) = f(a_{i-1})$  on  $[a_{i-1}, a_i), i = 1, 2, 3, \dots, 2^k$ . Then

$$\int_0^1 g(x) \varphi_n(x) dx = \sum_{i=1}^{2^k} f(a_{i-1}) \int_{a_{i-1}}^{a_i} \varphi_n(x) dx = 0.$$

Therefore,

$$\begin{aligned} |\hat{f}(n)| &= \left| \int_0^1 [f(x) - g(x)] \varphi_n(x) dx \right| \\ (2.1) \quad &\leq \int_0^1 |f(x) - g(x)| dx \\ &\leq \|f - g\|_p \|1\|_q \\ &= \left( \sum_{i=1}^{2^k} \int_{a_{i-1}}^{a_i} |f(x) - f(a_{i-1})|^p dx \right)^{\frac{1}{p}} \end{aligned}$$

by Hölder's inequality as  $f, g \in BV^{(p)}[0, 1]$  and  $BV^{(p)}[0, 1] \subset L^p[0, 1]$ .

Hence,

$$\begin{aligned} |\hat{f}(n)|^p &\leq \sum_{i=1}^{2^k} \int_{a_{i-1}}^{a_i} |f(x) - f(a_{i-1})|^p dx. \\ &\leq \sum_{i=1}^{2^k} \int_{a_{i-1}}^{a_i} (V(f, p, [a_{i-1}, a_i]))^p dx \\ &= \sum_{i=1}^{2^k} (V(f, p, [a_{i-1}, a_i]))^p \left( \frac{1}{2^k} \right) \end{aligned}$$

$$\begin{aligned} &\leq \left(\frac{1}{2^k}\right) (V(f, p, [0, 1]))^p \\ &\leq \left(\frac{2}{n}\right) (V(f, p, [0, 1]))^p, \end{aligned}$$

which completes the proof of Theorem 2.1.  $\square$

*Proof of Theorem 2.2.* Let  $c > 0$ . Using Jensen's inequality and proceeding as in Theorem 2.1, we get

$$\begin{aligned} \phi\left(c \int_0^1 |f(x) - g(x)| dx\right) &\leq \int_0^1 \phi(c|f(x) - g(x)|) dx \\ &= \sum_{i=1}^{2^k} \int_{a_{i-1}}^{a_i} \phi(c|f(x) - f(a_{i-1})|) dx \\ &\leq \sum_{i=1}^{2^k} \int_{a_{i-1}}^{a_i} V(cf, \phi, [a_{i-1}, a_i]) dx \\ &= \sum_{i=1}^{2^k} V(cf, \phi, [a_{i-1}, a_i]) \left(\frac{1}{2^k}\right) \\ &\leq \left(\frac{2}{n}\right) V(cf, \phi, [0, 1]). \end{aligned}$$

Since  $\phi$  is convex and  $\phi(0) = 0$ , for sufficiently small  $c \in (0, 1)$ ,  $V(cf, \phi, [0, 1]) < 1/2$ .

This completes the proof of Theorem 2.2 in view of (2.1).  $\square$

**Remark 1.** If  $\phi(x) = x^p$ ,  $p \geq 1$ , then the class  $\phi BV$  coincides with the class  $BV^{(p)}$  and Theorem 2.2 with Theorem 2.1.

**Remark 2.** Note that in the proof of Theorems 2.1 and 2.2, we have used the fact that if  $a = a_0 < a_1 < \dots < a_n = b$ , then

$$\sum_{i=1}^n (V(f, p, [a_{i-1}, a_i]))^p \leq (V(f, p, [a, b]))^p$$

and

$$\sum_{i=1}^n V(f, \phi, [a_{i-1}, a_i]) \leq V(f, \phi, [a, b]),$$

for any  $n \geq 2$  (see [2, 1.17, p. 15]). Such inequalities for functions of the class  $\Lambda BV^{(p)}$  ( $p \geq 1$ ) (resp.,  $\phi \Lambda BV$ ), which contain  $BV^{(p)}$  (resp.,  $\phi BV$ ) properly, do not hold true.

In fact, the following proposition shows that the validity of such inequalities for the class  $\Lambda BV^{(p)}$  (resp.,  $\phi \Lambda BV$ ) virtually reduces the class to  $BV^{(p)}$  (resp.,  $\phi BV$ ). Hence we prove Theorem 2.3 and Theorem 2.4 applying a technique different from the Taibleson-like technique [6] which we have applied in proving Theorem 2.1 and Theorem 2.2.

**Proposition 2.6.** Let  $f \in \phi \Lambda BV[a, b]$ . If there is a constant  $C$  such that

$$\sum_{i=1}^n V_{\Lambda}(f, \phi, [a_{i-1}, a_i]) \leq C V_{\Lambda}(f, \phi, [a, b]),$$

for any sequence of points  $\{a_i\}_{i=0}^n$  with  $a = a_0 < a_1 < \dots < a_n = b$ , then  $f \in \phi BV[a, b]$ .

*Proof.* For any partition  $a = x_0 < x_1 < \dots < x_n = b$  of  $[a, b]$ , we have

$$\begin{aligned} \sum_{i=1}^n \phi(|f(x_i) - f(x_{i-1})|) &= \lambda_1 \sum_{i=1}^n \frac{\phi(|f(x_i) - f(x_{i-1})|)}{\lambda_1} \\ &\leq \lambda_1 \sum_{i=1}^n V_\Lambda(f, \phi, [x_{i-1}, x_i]) \\ &\leq \lambda_1 CV_\Lambda(f, \phi, [a, b]), \end{aligned}$$

which shows that  $f \in \phi BV[a, b]$ . □

**Remark 3.**  $\phi(x) = x^p$  ( $p \geq 1$ ) in this proposition will give an analogous result for  $\Lambda BV^{(p)}$ .

To prove Theorem 2.3 and Theorem 2.4, we need the following lemma.

**Lemma 2.7.** For any  $n \in \mathbb{N}$ ,  $|\hat{f}(n)| \leq \omega_p(1/n; f)$ , where  $\omega_p(\delta; f)$  ( $\delta > 0$ ,  $p \geq 1$ ) denotes the integral modulus of continuity of order  $p$  of  $f$  given by

$$\omega_p(\delta; f) = \sup_{|h| \leq \delta} \left( \int_0^1 |f(x+h) - f(x)|^p dx \right)^{\frac{1}{p}}.$$

*Proof.* The inequality [1, Theorem IV, p. 382]  $|\hat{f}(n)| \leq \omega_1(1/n; f)$  and the fact that  $\omega_1(1/n; f) \leq \omega_p(1/n; f)$  for  $p \geq 1$  immediately proves the lemma. □

*Proof of Theorem 2.3.* For any  $n \in \mathbb{N}$ , put  $\theta_n = \sum_{j=1}^n 1/\lambda_j$ . Let  $f \in \Lambda BV^{(p)}[0, 1]$ . For  $0 < h \leq 1/n$ , put  $k = [1/h]$ . Then for a given  $x \in \mathbb{R}$ , all the points  $x + jh$ ,  $j = 0, 1, \dots, k$  lie in the interval  $[x, x + 1]$  of length 1 and

$$\int_0^1 |f(x) - f(x+h)|^p dx = \int_0^1 |f_j(x)|^p dx, \quad j = 1, 2, \dots, k,$$

where  $f_j(x) = f(x + (j-1)h) - f(x + jh)$ , for all  $j = 1, 2, \dots, k$ . Since the left hand side of this equation is independent of  $j$ , multiplying both sides by  $1/(\lambda_j \theta_k)$  and summing over  $j = 1, 2, \dots, k$ , we get

$$\begin{aligned} \int_0^1 |f(x) - f(x+h)|^p dx &\leq \left( \frac{1}{\theta_k} \right) \int_0^1 \sum_{j=1}^k \left( \frac{|f_j(x)|^p}{\lambda_j} \right) dx \\ &\leq \frac{(V_\Lambda(f, p, [0, 1]))^p}{\theta_k} \\ &\leq \frac{(V_\Lambda(f, p, [0, 1]))^p}{\theta_n}, \end{aligned}$$

because  $\{\lambda_j\}$  is non-decreasing and  $0 < h \leq 1/n$ . The case  $-1/n \leq h < 0$  is similar and we get using Lemma 2.7,

$$|\hat{f}(n)|^p \leq (\omega_p(1/n; f))^p \leq \frac{(V_\Lambda(f, p, [0, 1]))^p}{\theta_n}.$$

This proves Theorem 2.3. □

*Proof of Theorem 2.4.* Let  $f \in \phi\Lambda BV[0, 1]$ . Then for  $h, k$  and  $f_j(x)$  as in the proof of Theorem 2.3 and for  $c > 0$  by Jensen's inequality,

$$\begin{aligned} \phi\left(c \int_0^1 |f(x) - f(x+h)| dx\right) &\leq \int_0^1 \phi(c|f(x) - f(x+h)|) dx \\ &= \int_0^1 \phi(c|f_j(x)|) dx, \quad j = 1, 2, \dots, k. \end{aligned}$$

Multiplying both sides by  $1/(\lambda_j\theta_k)$  and summing over  $j = 1, 2, \dots, k$ , we get

$$\begin{aligned} \phi\left(c \int_0^1 |f(x) - f(x+h)| dx\right) &\leq \left(\frac{1}{\theta_k}\right) \int_0^1 \sum_{j=1}^k \left(\frac{\phi(c|f_j(x)|)}{\lambda_j}\right) dx \\ &\leq \frac{V_\Lambda(cf, \phi, [0, 1])}{\theta_k} \\ &\leq \frac{V_\Lambda(cf, \phi, [0, 1])}{\theta_n}. \end{aligned}$$

Since  $\phi$  is convex and  $\phi(0) = 0$ ,  $\phi(\alpha x) \leq \alpha\phi(x)$  for  $0 < \alpha < 1$ . So we may choose  $c$  sufficiently small so that  $V_\Lambda(cf, \phi, [0, 1]) \leq 1$ . But then we have

$$\int_0^1 |f(x) - f(x+h)| dx \leq \frac{1}{c} \phi^{-1}\left(\frac{1}{\theta_n}\right).$$

Thus it follows in view of Lemma 2.7 that

$$|\hat{f}(n)| \leq \omega_1(1/n; f) \leq \frac{1}{c} \phi^{-1}\left(\frac{1}{\theta_n}\right),$$

which proves Theorem 2.4.  $\square$

*Proof of Theorem 2.5.* It is known [3] that if  $\Gamma BV$  contains  $\Lambda BV$  properly with  $\Gamma = \{\gamma_n\}$  then  $\theta_n \neq O(\rho_n)$ , where  $\rho_n = \sum_{j=1}^n \frac{1}{\gamma_j}$  for each  $n$ . Also, if  $c_0 = 0$ ,  $c_{n+1} = 1$  and  $c_1 < c_2 < \dots < c_n$  denote all the  $n$  points of  $(0, 1)$  where the function  $\varphi_n$  changes its sign in  $(0, 1)$ ,  $n_0 \in \mathbb{N}$  is such that  $\rho_n \geq \frac{1}{2}$  for all  $n \geq n_0$  and  $E = \{n \in \mathbb{N} : n \geq n_0 \text{ is even}\}$ , then for each  $n \in E$ , for the function

$$f_n = \sum_{k=1}^{n+1} \frac{(-1)^{k-1}}{4\rho_n} \chi_{[c_{k-1}, c_k)}$$

extended 1-periodically on  $\mathbb{R}$ ,

$$V_\Gamma(f_n, [0, 1]) = \sum_{k=1}^{n+1} \frac{|f_n(c_k) - f_n(c_{k-1})|}{\gamma_k} = \sum_{k=1}^n \frac{1}{\gamma_k} \cdot \frac{1}{2\rho_n} = \frac{1}{2}$$

because

$$f_n(c_{n+1}) = f_n(1) = f_n(0) = \frac{1}{4\rho_n} = f_n(c_n)$$

as  $\varphi_n \equiv 1$  on  $[c_0, c_1)$ . Hence  $\|f_n\| = \frac{1}{4\rho_n} + \frac{1}{2} \leq 1$  for each  $n \in E$  in the Banach space  $\Gamma BV[0, 1]$  with  $\|f\| = |f(0)| + V_\Gamma(f, [0, 1])$ . Observe that for  $f \in \Gamma BV[0, 1]$

$$\|f\|_1 \leq \int_0^1 \left( \frac{|f(x) - f(0)|}{\gamma_1} \gamma_1 + |f(0)| \right) dx \leq C \|f\|, \quad C = \max\{1, \gamma_1\},$$

and hence, for each  $n \in \mathbb{N}$  the linear map  $T_n : \Gamma BV[0, 1] \rightarrow \mathbb{R}$  defined by  $T_n(f) = \theta_n \hat{f}(n)$  is bounded as

$$|T_n(f)| = \theta_n |\hat{f}(n)| \leq \theta_n \|f\|_1 \leq \theta_n C \|f\|, \quad \forall f \in \Gamma BV[0, 1].$$

Next, for each  $n \in E$  since  $f_n \cdot \varphi_n = \frac{1}{4\rho_n}$  on  $[0, 1)$ , we see that

$$T_n(f_n) = \theta_n \hat{f}_n(n) = \theta_n \int_0^1 f_n(x) \varphi_n(x) dx = \frac{1}{4} \left( \frac{\theta_n}{\rho_n} \right) \neq O(1)$$

and hence

$$\sup\{|T_n| : n \in \mathbb{N}\} \geq \sup\{|T_n| : n \in E\} \geq \sup\{|T_n(f_n)| : n \in E\} = \infty.$$

Therefore, an application of the Banach-Steinhaus theorem gives an  $f \in \Gamma BV[0, 1]$  such that  $\sup\{|T_n(f)| : n \in \mathbb{N}\} = \infty$ . It follows that  $\theta_n \hat{f}(n) = T_n(f) \neq O(1)$  and hence the theorem is proved.  $\square$

### REFERENCES

- [1] N.J. FINE, On the Walsh functions, *Trans. Amer. Math. Soc.*, **65** (1949), 372–414.
- [2] J. MUSIELAK AND W. ORLICZ, On generalized variations (I), *Studia Math.*, **18** (1959), 11–41.
- [3] S. PERLMAN AND D. WATERMAN, Some remarks on functions of  $\Lambda$ -bounded variation, *Proc. Amer. Math. Soc.*, **74**(1) (1979), 113–118.
- [4] M. SCHRAMM AND D. WATERMAN, On the magnitude of Fourier coefficients, *Proc. Amer. Math. Soc.*, **85** (1982), 407–410.
- [5] M. SHIBA, On absolute convergence of Fourier series of function of class  $\Lambda - BV^{(p)}$ , *Sci. Rep. Fac. Ed. Fukushima Univ.*, **30** (1980), 7–12.
- [6] M. TAIBLESON, Fourier coefficients of functions of bounded variation, *Proc. Amer. Math. Soc.*, **18** (1967), 766.
- [7] J.L. WALSH, A closed set of normal orthogonal functions, *Amer. J. Math.*, **55** (1923), 5–24.
- [8] D. WATERMAN, On convergence of Fourier series of functions of generalized bounded variation, *Studia Math.*, **44** (1972), 107–117.
- [9] N. WIENER, The quadratic variation of a function and its Fourier coefficients, *Mass. J. Math.*, **3** (1924), 72–94.