

GENERAL SYSTEM OF STRONGLY PSEUDOMONOTONE NONLINEAR VARIATIONAL INEQUALITIES BASED ON PROJECTION SYSTEMS

R. U. VERMA

Department of Mathematics, University of Toledo
Toledo, Ohio 43606, USA
EMail: verma99@msn.com

Received: 26 February, 2006

Accepted: 11 December, 2006

Communicated by: **R.N. Mohapatra**

2000 AMS Sub. Class.: 49J40, 65B05, 47H20.

Key words: Strongly pseudomonotone mappings, Approximation solvability, Projection methods, System of nonlinear variational inequalities.

Abstract: Let K_1 and K_2 , respectively, be non empty closed convex subsets of real Hilbert spaces H_1 and H_2 . The *Approximation – solvability* of a generalized system of nonlinear variational inequality (SNVI) problems based on the convergence of projection methods is discussed. The SNVI problem is stated as follows: find an element $(x^*, y^*) \in K_1 \times K_2$ such that

$$\langle \rho S(x^*, y^*), x - x^* \rangle \geq 0, \quad \forall x \in K_1 \text{ and for } \rho > 0,$$

$$\langle \eta T(x^*, y^*), y - y^* \rangle \geq 0, \quad \forall y \in K_2 \text{ and for } \eta > 0,$$

where $S : K_1 \times K_2 \rightarrow H_1$ and $T : K_1 \times K_2 \rightarrow H_2$ are nonlinear mappings.



Strongly Pseudomonotone
Nonlinear Variational
Inequalities

R. U. Verma

vol. 8, iss. 1, art. 6, 2007

[Title Page](#)

[Contents](#)



Page 1 of 19

[Go Back](#)

[Full Screen](#)


[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

Contents

1	Introduction	3
2	General Projection Methods	7





**Strongly Pseudomonotone
Nonlinear Variational
Inequalities**

R. U. Verma

vol. 8, iss. 1, art. 6, 2007

Title Page

Contents



Page 2 of 19

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

1. Introduction


Projection-like methods in general have been one of the most fundamental techniques for establishing the convergence analysis for solutions of problems arising from several fields, such as complementarity theory, convex quadratic programming, and variational problems. There exists a vast literature on approximation-solvability of several classes of variational/hemivariational inequalities in different space settings. The author [6, 7] introduced and studied a new system of nonlinear variational inequalities in Hilbert space settings. This class encompasses several classes of nonlinear variational inequality problems. In this paper we intend to explore, based on a general system of projection-like methods, the approximation-solvability of a system of nonlinear strongly pseudomonotone variational inequalities in Hilbert spaces. The obtained results extend/generalize the results in [1], [5] – [7] to the case of strongly pseudomonotone system of nonlinear variational inequalities. Approximation solvability of this system can also be established using the resolvent operator technique but in the more relaxed setting of Hilbert spaces. For more details, we refer the reader to [1] – [10].

Let H_1 and H_2 be two real Hilbert spaces with the inner product $\langle \cdot, \cdot \rangle$ and norm $\| \cdot \|$. Let $S : K_1 \times K_2 \rightarrow H_1$ and $T : K_1 \times K_2 \rightarrow H_2$ be any mappings on $K_1 \times K_2$, where K_1 and K_2 are nonempty closed convex subsets of H_1 and H_2 , respectively. We consider a system of nonlinear variational inequality (abbreviated as SNVI) problems: determine an element $(x^*, y^*) \in K_1 \times K_2$ such that

$$(1.1) \quad \langle \rho S(x^*, y^*), x - x^* \rangle \geq 0 \quad \forall x \in K_1$$


$$(1.2) \quad \langle \eta T(x^*, y^*), y - y^* \rangle \geq 0 \quad \forall y \in K_2,$$

where $\rho, \eta > 0$.



Title Page

Contents



Page 3 of 19

Go Back

Full Screen

Close



The SNVI (1.1)–(1.2) problem is equivalent to the following projection formulas

$$x^* = P_k[x^* - \rho S(x^*, y^*)] \quad \text{for } \rho > 0$$

$$y^* = Q_k[y^* - \eta T(x^*, y^*)] \quad \text{for } \eta > 0,$$

where P_k is the projection of H_1 onto K_1 and Q_k is the projection of H_2 onto K_2 .

We note that the SNVI (1.1)–(1.2) problem extends the NVI problem: determine an element $x^* \in K_1$ such that

$$(1.3) \quad \langle S(x^*), x - x^* \rangle \geq 0, \quad \forall x \in K_1.$$

Also, we note that the SNVI (1.1) – (1.2) problem is equivalent to a system of nonlinear complementarities (abbreviated as SNC): find an element $(x^*, y^*) \in K_1 \times K_2$ such that $S(x^*, y^*) \in K_1^*$, $T(x^*, y^*) \in K_2^*$, and

$$(1.4) \quad \langle \rho S(x^*, y^*), x^* \rangle = 0 \quad \text{for } \rho > 0,$$

$$(1.5) \quad \langle \eta T(x^*, y^*), y^* \rangle = 0 \quad \text{for } \eta > 0,$$

where K_1^* and K_2^* , respectively, are polar cones to K_1 and K_2 defined by

$$K_1^* = \{f \in H_1 : \langle f, x \rangle \geq 0, \quad \forall x \in K_1\}.$$

$$K_2^* = \{g \in H_2 : \langle g, y \rangle \geq 0, \quad \forall g \in K_2\}.$$

Now, we recall some auxiliary results and notions crucial to the problem on hand.

Lemma 1.1. *For an element $z \in H$, we have*

$$x \in K \quad \text{and} \quad \langle x - z, y - x \rangle \geq 0, \quad \forall y \in K \quad \text{if and only if} \quad x = P_k(z).$$



Title Page

Contents



Page 5 of 19

Go Back

Full Screen

Close

Lemma 1.2 ([3]). Let $\{\alpha^k\}$, $\{\beta^k\}$, and $\{\gamma^k\}$ be three nonnegative sequences such that

$$\alpha^{k+1} \leq (1 - t^k)\alpha^k + \beta^k + \gamma^k \quad \text{for } k = 0, 1, 2, \dots,$$

where $t^k \in [0, 1]$, $\sum_{k=0}^{\infty} t^k = \infty$, $\beta^k = o(t^k)$, and $\sum_{k=0}^{\infty} \gamma^k < \infty$. Then $\alpha^k \rightarrow 0$ as $k \rightarrow \infty$.

A mapping $T : H \rightarrow H$ from a Hilbert space H into H is called monotone if $\langle T(x) - T(y), x - y \rangle \geq 0$ for all $x, y \in H$. The mapping T is (r)-strongly monotone if for each $x, y \in H$, we have

$$\langle T(x) - T(y), x - y \rangle \geq r\|x - y\|^2 \quad \text{for a constant } r > 0.$$

This implies that $\|T(x) - T(y)\| \geq r\|x - y\|$, that is, T is (r)-expansive, and when $r = 1$, it is expansive. The mapping T is called (s)-Lipschitz continuous (or Lipschitzian) if there exists a constant $s \geq 0$ such that $\|T(x) - T(y)\| \leq s\|x - y\|$, $\forall x, y \in H$. T is called (μ)-cocoercive if for each $x, y \in H$, we have

$$\langle T(x) - T(y), x - y \rangle \geq \mu\|T(x) - T(y)\|^2 \quad \text{for a constant } \mu > 0.$$

Clearly, every (μ)-cocoercive mapping T is ($\frac{1}{\mu}$)-Lipschitz continuous. We can easily see that the following implications on monotonicity, strong monotonicity and expansiveness hold:


strong monotonicity \Rightarrow expansiveness



monotonicity


T is called relaxed (γ)-cocoercive if there exists a constant $\gamma > 0$ such that

$$\langle T(x) - T(y), x - y \rangle \geq (-\gamma)\|T(x) - T(y)\|^2, \quad \forall x, y \in H.$$



Title Page

Contents



Page 6 of 19

Go Back

Full Screen

Close

T is said to be (r) -strongly pseudomonotone if there exists a positive constant r such that

$$\langle T(y), x - y \rangle \geq 0 \Rightarrow \langle T(x), x - y \rangle \geq r \|x - y\|^2, \quad \forall x, y \in H.$$

T is said to be relaxed (γ, r) -cocoercive if there exist constants $\gamma, r > 0$ such that

$$\langle T(x) - T(y), x - y \rangle \geq (-\gamma) \|T(x) - T(y)\|^2 + r \|x - y\|^2.$$

Clearly, it implies that

$$\langle T(x) - T(y), x - y \rangle \geq (-\gamma) \|T(x) - T(y)\|^2,$$


that is, T is relaxed (γ) -cocoercive.

T is said to be relaxed (γ, r) -pseudococoercive if there exist positive constants γ and r such that

$$\langle T(y), x - y \rangle \geq 0 \Rightarrow \langle T(x), x - y \rangle \geq (-\gamma) \|T(x) - T(y)\|^2 + r \|x - y\|^2, \quad \forall x, y \in H.$$

Thus, we have following implications:

$$\begin{array}{l} (r)\text{-strong monotonicity} \quad \Rightarrow \quad \text{strong } (r)\text{-pseudomonotonicity} \\ \downarrow \\ \text{relaxed } (\gamma, r)\text{-cocoercivity} \\ \downarrow \\ \text{relaxed } (\gamma, r)\text{-pseudococoercivity} \end{array}$$



2. General Projection Methods

This section deals with the convergence of projection methods in the context of the approximation-solvability of the SNVI (1.1) – (1.2) problem.

Algorithm 1. For an arbitrarily chosen initial point $(x^0, y^0) \in K_1 \times K_2$, compute the sequences $\{x^k\}$ and $\{y^k\}$ such that

$$x^{k+1} = (1 - a^k - b^k)x^k + a^k P_k[x^k - \rho S(x^k, y^k)] + b^k u^k$$

$$y^{k+1} = (1 - \alpha^k - \beta^k)y^k + \alpha^k Q_K[y^k - \eta T(x^k, y^k)] + \beta^k v^k,$$

where P_K is the projection of H_1 onto K_1 , Q_K is the projection of H_2 onto K_2 , $\rho, \eta > 0$ are constants, $S : K_1 \times K_2 \rightarrow H_1$ and $T : K_1 \times K_2 \rightarrow H_2$ are any two mappings, and u^k and v^k , respectively, are bounded sequences in K_1 and K_2 . The sequences $\{a^k\}$, $\{b^k\}$, $\{\alpha^k\}$, and $\{\beta^k\}$ are in $[0, 1]$ with $(k \geq 0)$

$$0 \leq a^k + b^k \leq 1, \quad 0 \leq \alpha^k + \beta^k \leq 1.$$


Algorithm 2. For an arbitrarily chosen initial point $(x^0, y^0) \in K_1 \times K_2$, compute the sequences $\{x^k\}$ and $\{y^k\}$ such that

$$x^{k+1} = (1 - a^k - b^k)x^k + a^k P_k[x^k - \rho S(x^k, y^k)] + b^k u^k$$

$$y^{k+1} = (1 - a^k - b^k)y^k + a^k Q_K[y^k - \eta T(x^k, y^k)] + b^k v^k,$$

where P_K is the projection of H_1 onto K_1 , Q_K is the projection of H_2 onto K_2 , $\rho, \eta > 0$ are constants, $S : K_1 \times K_2 \rightarrow H_1$ and $T : K_1 \times K_2 \rightarrow H_2$ are any two mappings, and u^k and v^k , respectively, are bounded sequences in K_1 and K_2 . The sequences $\{a^k\}$ and $\{b^k\}$, are in $[0, 1]$ with $(k \geq 0)$

$$0 \leq a^k + b^k \leq 1.$$



[Title Page](#)

[Contents](#)



Page 8 of 19

[Go Back](#)

[Full Screen](#)

[Close](#)

Algorithm 3. For an arbitrarily chosen initial point $(x^0, y^0) \in K_1 \times K_2$, compute the sequences $\{x^k\}$ and $\{y^k\}$ such that

$$x^{k+1} = (1 - a^k)x^k + a^k P_k[x^k - \rho S(x^k, y^k)]$$

$$y^{k+1} = (1 - a^k)y^k + a^k Q_K[y^k - \eta T(x^k, y^k)],$$

where P_K is the projection of H_1 onto K_1 , Q_K is the projection of H_2 onto K_2 , $\rho, \eta > 0$ are constants, $S : K_1 \times K_2 \rightarrow H_1$ and $T : K_1 \times K_2 \rightarrow H_2$ are any two mappings. The sequence $\{a^k\} \in [0, 1]$ for $k \geq 0$.

We consider, based on Algorithm 2, the approximation solvability of the SNVI (1.1) – (1.2) problem involving strongly pseudomonotone and Lipschitz continuous mappings in Hilbert space settings.


Theorem 2.1. Let H_1 and H_2 be two real Hilbert spaces and, K_1 and K_2 , respectively, be nonempty closed convex subsets of H_1 and H_2 . Let $S : K_1 \times K_2 \rightarrow H_1$ be strongly (r) -pseudomonotone and (μ) -Lipschitz continuous in the first variable and let S be (ν) -Lipschitz continuous in the second variable. Let $T : K_1 \times K_2 \rightarrow H_2$ be strongly (s) -pseudomonotone and (β) -Lipschitz continuous in the second variable and let T be (τ) -Lipschitz continuous in the first variable. Let $\|\cdot\|^*$ denote the norm on $H_1 \times H_2$ defined by

$$\|(x, y)\|^* = (\|x\| + \|y\|) \forall (x, y) \in H_1 \times H_2.$$

In addition, let


$$\theta = \sqrt{1 - 2\rho r + \rho + \left(\frac{\rho\mu^2}{2}\right) + \rho^2\mu^2 + \eta\tau} < 1$$

$$\sigma = \sqrt{1 - 2\eta r + \eta + \left(\frac{\eta\beta^2}{2}\right) + \eta^2\beta^2 + \rho\nu} < 1,$$



Title Page

Contents



Page 9 of 19

Go Back

Full Screen

Close

let $(x^*, y^*) \in K_1 \times K_2$ form a solution to the SNVI (1.1) – (1.2) problem, and let sequences $\{x^k\}$, and $\{y^k\}$ be generated by Algorithm 2. Furthermore, let

$$(i) \langle S(x^*, y^k), x^k - x^* \rangle \geq 0;$$

$$(ii) \langle T(x^k, y^*), y^k - y^* \rangle \geq 0;$$

$$(iii) 0 \leq a^k + b^k \leq 1;$$

$$(iv) \sum_{k=0}^{\infty} a^k = \infty, \text{ and } \sum_{k=0}^{\infty} b^k < \infty;$$

$$(v) 0 < \rho < \frac{2r}{\mu^2} \text{ and } 0 < \eta < \frac{2s}{\beta^2}.$$


Then the sequence $\{x^k, y^k\}$ converges to (x^*, y^*) .

Proof. Since $(x^*, y^*) \in K_1 \times K_2$ forms a solution to the SNVI (1.1) – (1.2) problem, it follows that

$$x^* = P_K[x^* - \rho S(x^*, y^*)] \quad \text{and} \quad y^* = Q_K[x^* - \eta T(x^*, y^*)].$$


Applying Algorithm 2, we have

$$\begin{aligned} (2.1) \quad & \|x^{k+1} - x^*\| \\ &= \|(1 - a^k - b^k)x^k + a^k P_K[x^k - \rho S(x^k, y^k)] + b^k u^k \\ &\quad - (1 - a^k - b^k)x^* - a^k P_K[x^* - \rho S(x^*, y^*)] - b^k x^*\| \\ &\leq (1 - a^k - b^k)\|x^k - x^*\| \\ &\quad + a^k \|P_K[x^k - \rho S(x^k, y^k)] - P_K[x^* - \rho S(x^*, y^*)]\| + Mb^k \\ &\leq (1 - a^k)\|x^k - x^*\| + a^k \|x^k - x^* - \rho[S(x^k, y^k) - S(x^*, y^k)] \\ &\quad + S(x^*, y^k) - S(x^*, y^*)\| + Mb^k \\ &\leq (1 - a^k)\|x^k - x^*\| + a^k \|x^k - x^* - \rho[S(x^k, y^k) - S(x^*, y^k)]\| \\ &\quad + \rho \|S(x^*, y^k) - S(x^*, y^*)\| + Mb^k, \end{aligned}$$



Title Page

Contents



Page 10 of 19

Go Back

Full Screen

Close

where

$$M = \max\{\sup \|u^k - x^*\|, \sup \|v^k - y^*\|\} < \infty.$$

Since S is strongly (r) -pseudomonotone and (μ) -Lipschitz continuous in the first variable, and S is (ν) -Lipschitz continuous in the second variable, we have in light of (i) that


$$\begin{aligned} (2.2) \quad & \|x^k - x^* - \rho[S(x^k, y^k) - S(x^*, y^k)]\|^2 \\ &= \|x^k - x^*\|^2 - 2\rho\langle S(x^k, y^k) - S(x^*, y^k), x^k - x^* \rangle \\ &\quad + \rho^2\|S(x^k, y^k) - S(x^*, y^k)\|^2 \\ &= \|x^k - x^*\|^2 - 2\rho\langle S(x^k, y^k), x^k - x^* \rangle + 2\rho\langle S(x^*, y^k), x^k - x^* \rangle \\ &\quad + \rho^2\|S(x^k, y^k) - S(x^*, y^k)\|^2 \\ &\leq \|x^k - x^*\|^2 - 2\rho r\|x^k - x^*\|^2 + \rho^2\mu^2\|x^k - x^*\|^2 \\ &\quad + 2\rho\langle S(x^*, y^k), x^k - x^* \rangle \\ &\leq \|x^k - x^*\|^2 - 2\rho r\|x^k - x^*\|^2 + \rho^2\mu^2\|x^k - x^*\|^2 \\ &\quad + 2\rho\langle S(x^*, y^k), x^k - x^* \rangle \\ &= [1 - 2\rho r + \rho^2\mu^2]\|x^k - x^*\|^2 + 2\rho\langle S(x^*, y^k), x^k - x^* \rangle. \end{aligned}$$

On the other hand, we have

$$2\rho\langle S(x^*, y^k), x^k - x^* \rangle \leq \rho[\|S(x^*, y^k)\|^2 + \|x^k - x^*\|^2]$$



and

$$\begin{aligned} (2.3) \quad & \|S(x^*, y^k)\|^2 \\ &= \frac{1}{2} \left\{ \|S(x^*, y^k) - S(x^k, y^k)\|^2 - 2\|S(x^k, y^k)\|^2 \right\} \end{aligned}$$



Title Page

Contents



Page 11 of 19

Go Back

Full Screen

Close

$$\begin{aligned}
 & - \frac{1}{2} \|S(x^k, y^k) + S(x^*, y^k)\|^2 \} \\
 \leq & \frac{1}{2} \left\{ \|S(x^*, y^k) - S(x^k, y^k)\|^2 - 2[\|S(x^k, y^k)\|^2 \right. \\
 & \left. - \frac{1}{2} \| \|S(x^k, y^k) - S(x^*, y^k)\|^2 \right\} \\
 \leq & \frac{1}{2} \|S(x^*, y^k) - S(x^k, y^k)\|^2 \\
 \leq & \frac{\mu^2}{2} \|x^k - x^*\|^2,
 \end{aligned}$$

where

$$\|S(x^k, y^k)\|^2 - \frac{1}{2} \| \|S(x^k, y^k) - S(x^*, y^k)\|^2 > 0.$$

Therefore, we get

$$2\rho \langle S(x^*, y^k), x^k - x^* \rangle \leq \left[\rho + \left(\frac{\rho\mu^2}{2} \right) \right] \|x^k - x^*\|^2.$$


It follows that

$$\|x^k - x^* - \rho[S(x^k, y^k) - S(x^*, y^k)]\|^2 \leq \left[1 - 2\rho r + \rho + \left(\frac{\rho\mu^2}{2} \right) + (\rho\mu)^2 \right] \|x^k - x^*\|^2.$$

As a result, we have


$$(2.4) \quad \|x^{k+1} - x^*\| \leq (1 - a^k) \|x^k - x^*\| + a^k \theta \|x^k - x^*\| + a^k \rho \nu \|y^k - y^*\| + Mb^k,$$

$$\text{where } \theta = \sqrt{1 - 2\rho r + \rho + \left(\frac{\rho\mu^2}{2} \right) + \rho^2\mu^2}.$$



[Title Page](#)

[Contents](#)



Page 12 of 19

[Go Back](#)

[Full Screen](#)

[Close](#)

Similarly, we have

$$(2.5) \quad \|y^{k+1} - y^*\| \leq (1 - a^k)\|y^k - y^*\| + a^k\sigma\|y^k - y^*\| + a^k\eta\tau\|x^k - x^*\| + Mb^k,$$

$$\text{where } \sigma = \sqrt{1 - 2\eta r + \eta + \left(\frac{\eta\beta^2}{2}\right) + \eta^2\beta^2}.$$

It follows from (2.4) and (2.5) that

$$(2.6) \quad \begin{aligned} & \|x^{k+1} - x^*\| + \|y^{k+1} - y^*\| \\ & \leq (1 - a^k)\|x^k - x^*\| + a^k\theta\|x^k - x^*\| + a^k\eta\tau\|x^k - x^*\| + Mb^k \\ & \quad + (1 - a^k)\|y^k - y^*\| + a^k\sigma\|y^k - y^*\| + a^k\rho\nu\|y^k - y^*\| + Mb^k \\ & = [1 - (1 - \delta)a^k](\|x^k - x^*\| + \|y^k - y^*\|) + 2Mb^k, \end{aligned}$$

where $\delta = \max\{\theta + \eta\tau, \sigma + \rho\nu\}$ and $H_1 \times H_2$ is a Banach space under the norm $\|\cdot\|^*$.

If we set

$$\alpha^k = \|x^k - x^*\| + \|y^k - y^*\|, \quad t^k = (1 - \delta)a^k,$$

$$\beta^k = 2Mb^k \quad \text{for } k = 0, 1, 2, \dots,$$

in Lemma 1.2, and apply (iii) and (iv), we conclude that


$$\|x^k - x^*\| + \|y^k - y^*\| \rightarrow 0$$

as $k \rightarrow \infty$.

Hence,



$$\|x^{k+1} - x^*\| + \|y^{k+1} - y^*\| \rightarrow 0.$$

Consequently, the sequence $\{(x^k, y^k)\}$ converges strongly to (x^*, y^*) , a solution to the SNVI (1.1) – (1.2) problem. This completes the proof. ■



[Title Page](#)

[Contents](#)



Page 13 of 19

[Go Back](#)

[Full Screen](#)

[Close](#)

Note that the proof of the following theorem follows rather directly without using Lemma 1.2.

Theorem 2.2. Let H_1 and H_2 be two real Hilbert spaces and, K_1 and K_2 , respectively, be nonempty closed convex subsets of H_1 and H_2 . Let $S : K_1 \times K_2 \rightarrow H_1$ be strongly (r) -pseudomonotone and (μ) -Lipschitz continuous in the first variable and let S be (ν) -Lipschitz continuous in the second variable. Let $T : K_1 \times K_2 \rightarrow H_2$ be strongly (s) -pseudomonotone and (β) -Lipschitz continuous in the second variable and let T be (τ) -Lipschitz continuous in the first variable. Let $\|\cdot\|^*$ denote the norm on $H_1 \times H_2$ defined by

$$\|(x, y)\|^* = (\|x\| + \|y\|) \quad \forall (x, y) \in H_1 \times H_2.$$

In addition, let

$$\theta = \sqrt{1 - 2\rho r + \rho + \left(\frac{\rho\mu^2}{2}\right) + \rho^2\mu^2 + \eta\tau} < 1,$$

$$\sigma = \sqrt{1 - 2\eta r + \eta + \left(\frac{\eta\beta^2}{2}\right) + \eta^2\beta^2 + \rho\nu} < 1,$$

let $(x^*, y^*) \in K_1 \times K_2$ form a solution to the SNVI (1.1) – (1.2) problem, and let sequences $\{x^k\}$, and $\{y^k\}$ be generated by Algorithm 3. Furthermore, let

(i) $\langle S(x^*, y^*), x^k - x^* \rangle \geq 0$

(ii) $\langle T(x^k, y^*), y^k - y^* \rangle \geq 0$

(iii) $0 \leq a^k \leq 1$

(iv) $\sum_{k=0}^{\infty} a^k = \infty$

$$(v) \ 0 < \rho < \frac{2r}{\mu^2} \text{ and } 0 < \eta < \frac{2s}{\beta^2}.$$

Then the sequence $\{x^k, y^k\}$ converges strongly to (x^*, y^*) .

Proof. Since $(x^*, y^*) \in K_1 \times K_2$ forms a solution to the SNVI (1.1) – (1.2) problem, it follows that


$$x^* = P_K[x^* - \rho S(x^*, y^*)] \quad \text{and} \quad y^* = Q_K[x^* - \eta T(x^*, y^*)].$$

Applying Algorithm 3, we have

$$\begin{aligned} (2.7) \quad & \|x^{k+1} - x^*\| \\ &= \|(1 - a^k)x^k + a^k P_K[x^k - \rho S(x^k, y^k)] - (1 - a^k)x^* \\ &\quad - a^k P_K[x^* - \rho S(x^*, y^*)]\| \\ &\leq (1 - a^k)\|x^k - x^*\| + a^k \|P_K[x^k - \rho S(x^k, y^k)] - P_K[x^* - \rho S(x^*, y^*)]\| \\ &\leq (1 - a^k)\|x^k - x^*\| \\ &\quad + a^k \|x^k - x^* - \rho[S(x^k, y^k) - S(x^*, y^k)] + S(x^*, y^k) - S(x^*, y^*)\| \\ &\leq (1 - a^k)\|x^k - x^*\| + a^k \|x^k - x^* - \rho[S(x^k, y^k) - S(x^*, y^k)]\| \\ &\quad + \rho \| [S(x^*, y^k) - S(x^*, y^*)] \|. \end{aligned}$$


Since S is strongly (r) –pseudomonotone and (μ) –Lipschitz continuous in the first variable, and S is (ν) –Lipschitz continuous in the second variable, we have in light of (i) that

$$\begin{aligned} (2.8) \quad & \|x^k - x^* - \rho[S(x^k, y^k) - S(x^*, y^k)]\|^2 \\ &= \|x^k - x^*\|^2 - 2\rho \langle S(x^k, y^k) - S(x^*, y^k), x^k - x^* \rangle \\ &\quad + \rho^2 \|S(x^k, y^k) - S(x^*, y^k)\|^2 \\ &= \|x^k - x^*\|^2 - 2\rho \langle S(x^k, y^k), x^k - x^* \rangle + 2\rho \langle S(x^*, y^k), x^k - x^* \rangle \end{aligned}$$



Title Page

Contents




Page 14 of 19

Go Back


Full Screen

Close



Title Page

Contents



Page 15 of 19

Go Back

Full Screen

Close


$$\begin{aligned} & + \rho^2 \|S(x^k, y^k) - S(x^*, y^k)\|^2 \\ \leq & \|x^k - x^*\|^2 - 2\rho r \|x^k - x^*\|^2 + \rho^2 \mu^2 \|x^k - x^*\|^2 \\ & + 2\rho \langle S(x^*, y^k), x^k - x^* \rangle \\ \leq & \|x^k - x^*\|^2 - 2\rho r \|x^k - x^*\|^2 + \rho^2 \mu^2 \|x^k - x^*\|^2 \\ & + 2\rho \langle S(x^*, y^k), x^k - x^* \rangle \\ = & [1 - 2\rho r + \rho^2 \mu^2] \|x^k - x^*\|^2 + 2\rho \langle S(x^*, y^k), x^k - x^* \rangle. \end{aligned}$$

On the other hand, we have

$$2\rho \langle S(x^*, y^k), x^k - x^* \rangle \leq \rho [\|S(x^*, y^k)\|^2 + \|x^k - x^*\|^2]$$

and

$$\begin{aligned} (2.9) \quad & \|S(x^*, y^k)\|^2 \\ = & \frac{1}{2} \left\{ \|S(x^*, y^k) - S(x^k, y^k)\|^2 \right. \\ & \left. - 2 \left[\|S(x^k, y^k)\|^2 - \frac{1}{2} \|S(x^k, y^k) + S(x^*, y^k)\|^2 \right] \right\} \\ \leq & \frac{1}{2} \left\{ \|S(x^*, y^k) - S(x^k, y^k)\|^2 \right. \\ & \left. - 2 \left[\|S(x^k, y^k)\|^2 - \frac{1}{2} \|S(x^k, y^k) - S(x^*, y^k)\|^2 \right] \right\} \\ \leq & \frac{1}{2} \|S(x^*, y^k) - S(x^k, y^k)\|^2 \leq \frac{\mu^2}{2} \|x^k - x^*\|^2, \end{aligned}$$



Title Page

Contents



Page 16 of 19

Go Back

Full Screen

Close

where

$$\|S(x^k, y^k)\|^2 - \frac{1}{2} \|S(x^k, y^k) - S(x^*, y^k)\|^2 > 0.$$

Therefore, we get

$$2\rho \langle S(x^*, y^k), x^k - x^* \rangle \leq \left[\rho + \left(\frac{\rho\mu^2}{2} \right) \right] \|x^k - x^*\|^2.$$

It follows that

$$\|x^k - x^* - \rho[S(x^k, y^k) - S(x^*, y^k)]\|^2 \leq \left[1 - 2\rho r + \rho + \left(\frac{\rho\mu^2}{2} \right) + (\rho\mu)^2 \right] \|x^k - x^*\|^2.$$

As a result, we have

$$(2.10) \quad \|x^{k+1} - x^*\| \leq (1 - a^k) \|x^k - x^*\| + a^k \theta \|x^k - x^*\| + a^k \rho \nu \|y^k - y^*\|,$$

$$\text{where } \theta = \sqrt{1 - 2\rho r + \rho + \left(\frac{\rho\mu^2}{2} \right) + \rho^2\mu^2}.$$


Similarly, we have

$$(2.11) \quad \|y^{k+1} - y^*\| \leq (1 - a^k) \|y^k - y^*\| + a^k \sigma \|y^k - y^*\| + a^k \eta \tau \|x^k - x^*\|,$$

$$\text{where } \sigma = \sqrt{1 - 2\eta r + \eta + \left(\frac{\eta\beta^2}{2} \right) + \eta^2\beta^2}.$$


It follows from (2.9) and (2.10) that

$$(2.12) \quad \begin{aligned} & \|x^{k+1} - x^*\| + \|y^{k+1} - y^*\| \\ & \leq (1 - a^k) \|x^k - x^*\| + a^k \theta \|x^k - x^*\| + a^k \eta \tau \|x^k - x^*\| \\ & \quad + (1 - a^k) \|y^k - y^*\| + a^k \sigma \|y^k - y^*\| + a^k \rho \nu \|y^k - y^*\| \end{aligned}$$



Title Page

Contents



Page 17 of 19

Go Back

Full Screen

Close

$$\begin{aligned} &= [1 - (1 - \delta)a^k](\|x^k - x^*\| + \|y^k - y^*\|) \\ &\leq \prod_{j=0}^k [1 - (1 - \delta)a^j](\|x^0 - x^*\| + \|y^0 - y^*\|), \end{aligned}$$

where $\delta = \max\{\theta + \eta\tau, \sigma + \rho\nu\}$ and $H_1 \times H_2$ is a Banach space under the norm $\|\cdot\|^*$.

Since $\delta < 1$ and $\sum_{k=0}^{\infty} a^k$ is divergent, it follows that

$$\lim_{k \rightarrow \infty} \prod_{j=0}^k [1 - (1 - \delta)a^j] = 0 \quad \text{as } k \rightarrow \infty.$$


Therefore,

$$\|x^{k+1} - x^*\| + \|y^{k+1} - y^*\| \rightarrow 0,$$

and consequently, the sequence $\{(x^k, y^k)\}$ converges strongly to (x^*, y^*) , a solution to the *SNVI* (1.1) – (1.2) problem. This completes the proof. ■

References

- [1] S.S. CHANG, Y.J. CHO, AND J.K. KIM, On the two-step projection methods and applications to variational inequalities, *Mathematical Inequalities and Applications*, accepted.
- [2] Z. LIU, J.S. UME, AND S.M. KANG, Generalized nonlinear variational-like inequalities in reflexive Banach spaces, *Journal of Optimization Theory and Applications*, **126**(1) (2005), 157–174.
- [3] L. S. LIU, Ishikawa and Mann iterative process with errors for nonlinear strongly accretive mappings in Banach spaces, *Journal of Mathematical Analysis and Applications*, **194** (1995), 114–127.
- [4] H. NIE, Z. LIU, K. H. KIM AND S. M. KANG, A system of nonlinear variational inequalities involving strongly monotone and pseudocontractive mappings, *Advances in Nonlinear Variational Inequalities*, **6**(2) (2003), 91–99.
- [5] R.U. VERMA, Nonlinear variational and constrained hemivariational inequalities, *ZAMM: Z. Angew. Math. Mech.*, **77**(5) (1997), 387–391.
- [6] R.U. VERMA, Generalized convergence analysis for two-step projection methods and applications to variational problems, *Applied Mathematics Letters*, **18** (2005), 1286–1292.
- [7] R.U. VERMA, Projection methods, algorithms and a new system of nonlinear variational inequalities, *Computers and Mathematics with Applications*, **41** (2001), 1025–1031.
- [8] R. WITTMANN, Approximation of fixed points of nonexpansive mappings, *Archiv der Mathematik*, **58** (1992), 486–491.



Strongly Pseudomonotone
Nonlinear Variational
Inequalities

R. U. Verma

vol. 8, iss. 1, art. 6, 2007

Title Page

Contents



Page 18 of 19

Go Back


Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

- [9] E. ZEIDLER, *Nonlinear Functional Analysis and its Applications I*, Springer-Verlag, New York, New York, 1986.
- [10] E. ZEIDLER, *Nonlinear Functional Analysis and its Applications II/B*, Springer-Verlag, New York, New York, 1990.



**Strongly Pseudomonotone
Nonlinear Variational
Inequalities**

R. U. Verma

vol. 8, iss. 1, art. 6, 2007

[Title Page](#)

[Contents](#)



Page 19 of 19

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756