

Journal of Inequalities in Pure and Applied Mathematics

ON INTEGRAL FORMS OF GENERALISED MATHIEU SERIES

P. CERONE AND C.T. LENARD

School of Computer Sciences and Mathematics
Victoria University of Technology
PO Box 14428, MCMC 8001
VIC, Australia.

*E*Mail: pc@cs.vu.edu.au


*U*RL: <http://rgmia.vu.edu.au/cerone>

Department of Mathematics
LaTrobe University
PO Box 199, Bendigo,
Victoria, 3552, Australia.

*E*Mail: C.Lenard@bendigo.latrobe.edu.au

*U*RL: <http://www.bendigo.latrobe.edu.au/mte/mathstaff/lenard/>

©2000 Victoria University
ISSN (electronic): 1443-5756
080-03



volume 4, issue 5, article 100,
2003.

Received 16 June, 2003;
accepted 08 December, 2003.

Communicated by: F. Qi

Abstract

Contents



Home Page

Go Back

Close

Quit

Abstract


Integral representations for generalised Mathieu series are obtained which recapture the Mathieu series as a special case. Bounds are obtained through the use of the integral representations.

2000 Mathematics Subject Classification: Primary 26D15, 33E20; Secondary 26A42, 40A30.

Key words: Mathieu Series, Bounds, Identities.

Contents

1	Introduction	3
2	Integral Representation of the Generalised Mathieu Series $S_\mu(r)$	6
3	Bounds for $S_\mu(r)$	15
4	Further Integral Expressions for Generalised Mathieu Series	19
	References	



On Integral Forms of Generalised Mathieu Series

P. Cerone and C.T. Lenard

Title Page

Contents



Go Back

Close

Quit

Page 2 of 23

1. Introduction

The series

$$(1.1) \quad S(r) = \sum_{n=1}^{\infty} \frac{2n}{(n^2 + r^2)^2}, \quad r > 0$$

is well known in the literature as Mathieu's series. It has been extensively studied in the past since its introduction by Mathieu [12] in 1890, where it arose in connection with work on elasticity of solid bodies. The reader is directed to the references for further illustration.

One of the main questions addressed in relation (1.1) is to obtain sharp bounds. Alzer, Brenner and Ruehr [2] showed that the best constants a and b in

$$\frac{1}{x^2 + a} < S(x) < \frac{1}{x^2 + b}, \quad x \neq 0$$

are $a = \frac{1}{2\zeta(3)}$ and $b = \frac{1}{6}$ where $\zeta(\cdot)$ denotes the Riemann zeta function defined by

$$(1.2) \quad \zeta(p) = \sum_{n=1}^{\infty} \frac{1}{n^p}.$$

An integral representation for $S(r)$ as given in (1.1) was presented in [6] and [7] as

$$(1.3) \quad S(r) = \frac{1}{r} \int_0^{\infty} \frac{x}{e^x - 1} \sin(rx) dx.$$



On Integral Forms of
Generalised Mathieu Series

P. Cerone and C.T. Lenard

Title Page

Contents



Go Back

Close

Quit

Page 3 of 23

Guo [10] utilised (1.3) and a lemma [3, pp. 89–90] to obtain bounds on $S(r)$, namely,

$$(1.4) \quad \frac{\pi}{r^3} \sum_{k=0}^{\infty} \frac{(-1)^k \left(k + \frac{1}{2}\right)}{e^{(k+\frac{1}{2})\frac{\pi}{r}} - 1} < S(r) < \frac{1}{r^2} \left(1 + \frac{\pi}{r} \sum_{k=0}^{\infty} \frac{(-1)^k \left(k + \frac{1}{2}\right)}{e^{(k+\frac{1}{2})\frac{\pi}{r}} - 1}\right).$$

The following results were obtained by Qi and coworkers (see [4], [15] – [17])

$$(1.5) \quad \frac{4(1+r^2)(e^{-\frac{\pi}{r}} + e^{-\frac{\pi}{2r}}) - 4r^2 - 1}{(e^{-\frac{\pi}{r}} - 1)(1+r^2)(1+4r^2)} \leq S(r) \leq \frac{(1+4r^2)(e^{-\frac{\pi}{r}} - e^{-\frac{\pi}{2r}}) - 4(1+r^2)}{(e^{-\frac{\pi}{r}} - 1)(1+r^2)(1+4r^2)}$$


$$S(r) < \frac{1}{r} \int_0^{\frac{\pi}{r}} \frac{x}{e^x - 1} \sin(rx) dx < \frac{1 + e^{-\frac{\pi}{2r}}}{r^2 + \frac{1}{4}},$$

and

$$S(r) \geq \frac{1}{8r(1+r^2)^3} \left[16r(r^2 - 3) + \pi^3(r^2 + 1)^3 \operatorname{sech}^2\left(\frac{\pi r}{2}\right) \tanh\left(\frac{\pi r}{2}\right) \right].$$

Guo in [10] poses the interesting problem as to whether there is an integral representation of the generalised Mathieu series

$$(1.6) \quad S_{\mu}(r) = \sum_{n=1}^{\infty} \frac{2n}{(n^2 + r^2)^{1+\mu}}, \quad r > 0, \mu > 0.$$



On Integral Forms of Generalised Mathieu Series

P. Cerone and C.T. Lenard

Title Page

Contents



Go Back

Close

Quit

Page 4 of 23

This is resolved in Section 2.

Recently in [18] an integral representation was obtained for $S_m(r)$, where $m \in \mathbb{N}$, namely

$$(1.7) \quad S_m(r) = \frac{2}{(2r)^m m!} \int_0^\infty \frac{t^m}{e^t - 1} \cos\left(\frac{m\pi}{2} - rt\right) dt \\ - 2 \sum_{k=1}^m \left[\frac{(k-1)(2r)^{k-2m-1}}{k!(m-k+1)} \binom{-(m+1)}{m-k} \right. \\ \left. \times \int_0^\infty \frac{t^k \cos\left[\frac{\pi}{2}(2m-k+1) - rt\right]}{e^t - 1} dt \right].$$

Bounds were obtained by Tomovski and Trenčevski [18] using (1.3).

It is the intention of the current paper to investigate further integral representations of the generalised Mathieu series (1.6).

Bounds are obtained in Section 3 for $S_\mu(r)$. In Section 4 the open problem of obtaining an integral representation for


$$S(r; \mu, \gamma) = \sum_{n=1}^{\infty} \frac{2n^\gamma}{(n^{2\gamma} + r^2)^{\mu+1}}$$

posed by Qi [15] is addressed.

We notice that

$$S(r; 1, 1) = S_1(r) = S(r),$$

the Mathieu series.



On Integral Forms of
Generalised Mathieu Series

P. Cerone and C.T. Lenard

Title Page

Contents



Go Back

Close

Quit

Page 5 of 23

2. Integral Representation of the Generalised Mathieu Series $S_\mu(r)$

Before proceeding to obtain an integral representation for $S_\mu(r)$ as given by (1.6), it is instructive to present an alternative representation in terms of the zeta function $\zeta(p)$ presented in (1.2). Namely, a straightforward series expansion gives

$$(2.1) \quad S_\mu(r) = 2 \sum_{k=0}^{\infty} r^{2k} (-1)^k \binom{\mu+k}{k} \zeta(2\mu+2k+1)$$

on using the result $\binom{\alpha}{k} = (-1)^k \binom{k-\alpha-1}{k}$ with $\alpha = -(\mu+1)$.

Theorem 2.1. *The generalised Mathieu series $S_\mu(r)$ defined by (1.6) may be represented in the integral form*

$$(2.2) \quad S_\mu(r) = C_\mu(r) \int_0^\infty \frac{x^{\mu+\frac{1}{2}}}{e^x - 1} J_{\mu-\frac{1}{2}}(rx) dx, \quad \mu > 0,$$


where

$$(2.3) \quad C_\mu(r) = \frac{\sqrt{\pi}}{(2r)^{\mu-\frac{1}{2}} \Gamma(\mu+1)}$$

and $J_\nu(z)$ is the ν^{th} order Bessel function of the first kind.

Proof (A). Consider

$$(2.4) \quad T_\mu(r) = \int_0^\infty \frac{x^{\mu+\frac{1}{2}}}{e^x - 1} J_{\mu-\frac{1}{2}}(rx) dx.$$



On Integral Forms of
Generalised Mathieu Series

P. Cerone and C.T. Lenard

Title Page

Contents



Go Back

Close

Quit

Page 6 of 23

Then using the series definition for $J_\nu(z)$ (Gradshteyn and Ryzhik [9]),

$$J_\nu(z) = \sum_{k=0}^{\infty} \frac{(-1)^k \left(\frac{z}{2}\right)^{\nu+2k}}{k! \Gamma(\nu+k+1)}$$

in (2.4) produces after the permissible interchange of summation and integral,

$$(2.5) \quad T_\mu(r) = \sum_{k=0}^{\infty} \frac{(-1)^k \left(\frac{r}{2}\right)^{\mu+2k-\frac{1}{2}}}{k! \Gamma\left(\mu+k+\frac{1}{2}\right)} \int_0^\infty \frac{x^{2(\mu+k)}}{e^x - 1} dx.$$

Now, the well known representation [9]

$$(2.6) \quad \int_0^\infty \frac{x^p}{e^x - 1} dx = \Gamma(p+1) \zeta(p+1)$$

gives from (2.5) with $p = 2(\mu+k)$


$$(2.7) \quad T_\mu(r) = \sum_{k=0}^{\infty} \frac{(-1)^k \left(\frac{r}{2}\right)^{\mu+2k-\frac{1}{2}} \Gamma(2\mu+2k+1) \zeta(2\mu+2k+1)}{k! \Gamma\left(\mu+k+\frac{1}{2}\right)}.$$

An application of the duplication identity for the gamma function

$$\sqrt{\pi} \Gamma(2z) = 2^{2z-1} \Gamma(z) \Gamma\left(z + \frac{1}{2}\right),$$

with $z = \mu+k + \frac{1}{2}$ simplifies the expression in (2.7) to

$$(2.8) \quad T_\mu(r) = \frac{(2r)^{\mu-\frac{1}{2}}}{\sqrt{\pi}} \sum_{k=0}^{\infty} (-1)^k r^{2k} \frac{\Gamma(\mu+k+1)}{k!} \zeta(2\mu+2k+1).$$



On Integral Forms of
Generalised Mathieu Series

P. Cerone and C.T. Lenard

Title Page

Contents



Go Back

Close

Quit

Page 7 of 23

Repeated use of the identity $\Gamma(z + 1) = z\Gamma(z)$ gives

$$\frac{\Gamma(\mu + k + 1)}{k!} = \binom{\mu + k}{k} \Gamma(\mu + 1)$$

and so from (2.8)

$$T_\mu(r) = \frac{(2r)^{\mu - \frac{1}{2}} \Gamma(\mu + 1)}{\sqrt{\pi}} 2 \sum_{k=0}^{\infty} (-1)^k r^{2k} \binom{\mu + k}{k} \zeta(2\mu + 2k + 1)$$

produces the result (2.2) on reference to (2.1), (2.3) and (2.4).

Proof (A) is now complete. □

Proof (B). From (2.4) we have


$$\begin{aligned} (2.9) \quad T_\mu(r) &= \int_0^\infty \frac{e^{-x}}{1 - e^{-x}} x^{\mu + \frac{1}{2}} J_{\mu - \frac{1}{2}}(rx) dx \\ &= \sum_{k=1}^{\infty} \int_0^\infty e^{-nx} x^{\mu + \frac{1}{2}} J_{\mu - \frac{1}{2}}(rx) dx. \end{aligned}$$

Now Gradshteyn and Ryzhik [9] on page 712 has the result

$$(2.10) \quad \int_0^\infty e^{-\alpha x} x^{\nu+1} J_\nu(\beta x) dx = \frac{2\alpha(2\beta)^\nu \Gamma(\nu + \frac{3}{2})}{\sqrt{\pi} [\alpha^2 + \beta^2]^{\nu + \frac{3}{2}}},$$

$$\operatorname{Re}(\nu) > -1, \quad \operatorname{Re}(\alpha) > |\operatorname{Im} \beta|,$$

which is referred to in Watson [20] whom in turn attributes the result to an 1875 result of Gegenbauer.



**On Integral Forms of
Generalised Mathieu Series**

P. Cerone and C.T. Lenard

Title Page

Contents



Go Back

Close

Quit

Page 8 of 23

Taking $\alpha = n$, $\nu = \mu - \frac{1}{2}$ and $\beta = r$, all real, in (2.10) and substituting in (2.9) readily produces

$$T_\mu(r) = \frac{(2r)^{\mu-\frac{1}{2}} \Gamma(\mu+1)}{\sqrt{\pi}} \sum_{n=1}^{\infty} \frac{2n}{[n^2+r^2]^{\mu+1}},$$

giving from (1.6), (2.4) and (2.3) the result (2.2).

We note that the more restrictive condition of $\mu > 0$ needs to be imposed for the convergence of the series although (2.10) requires $\text{Re}(\nu) = \mu - \frac{1}{2} > -1$. \square

Remark 2.1. *If we take $\mu = 1$ in (1.6) and (2.2) – (2.3) then $S_1(r) \equiv S(r)$, the Mathieu series given by (1.1) and its integral representation (1.3). This is easily seen to be the case since $J_{\frac{1}{2}}(z) = \sqrt{\frac{2}{\pi z}} \sin z$ and taking $\mu = 1$ in (2.2) – (2.3) produces (1.3).*


Remark 2.2. *Gradshtein and Ryzhik [9] on page 712 also quote the result*

$$(2.11) \quad \int_0^\infty e^{-\alpha x} x^\nu J_\nu(\beta x) dx = \frac{(2\beta)^\nu \Gamma(\nu + \frac{1}{2})}{\sqrt{\pi} (\alpha^2 + \beta^2)^{\nu + \frac{1}{2}}}$$

$$\text{Re}(\nu) > -\frac{1}{2}, \quad \text{Re}(\alpha) > |\text{Im}(\beta)|,$$

which Watson [20] again attributes to an 1875 result by Gegenbauer.

We note that formal differentiation of (2.11) with respect to α produces the result (2.10).



On Integral Forms of
Generalised Mathieu Series

P. Cerone and C.T. Lenard

Title Page

Contents



Go Back

Close

Quit

Page 9 of 23

Following a similar process as in Proof (B) above, we may show that

$$(2.12) \quad \int_0^\infty \frac{x^{\mu-\frac{1}{2}}}{e^x - 1} J_{\mu-\frac{1}{2}}(rx) dx = \frac{(2r)^{\mu-\frac{1}{2}} \Gamma(\mu)}{\sqrt{\pi}} \sum_{n=1}^\infty \frac{1}{(n^2 + r^2)^\mu}.$$

Gradshteyn and Ryzhik [9] have an explicit expression which can be transformed by a simple change of variables to (2.12). Namely,

$$(2.13) \quad \int_0^\infty \frac{x^\nu J_\nu(bx)}{e^{\pi x} - 1} dx = \frac{(2b)^\nu \Gamma(\nu + \frac{1}{2})}{\sqrt{\pi}} \sum_{n=1}^\infty \frac{1}{(n^2 \pi^2 + b^2)^{\nu+\frac{1}{2}}},$$

$\text{Re}(\nu) > 0$, $|\text{Im}(b)| < \pi$, which is attributed by Watson [20] to a 1906 result by Kapteyn.

An explicit integral expression for $S_\mu(r)$ of the current form does not seem to have been available previously.


Finally, we note that (2.10) or (2.11) may be looked upon as an integral transform such as the Laplace or Hankel transform and the results may be found in tables of such.

Remark 2.3. $S_\mu(r)$ as given in (2.2) – (2.3) may be written in the alternate form

$$(2.14) \quad S_\mu(r) = \frac{\sqrt{\pi}}{2^{\mu-\frac{1}{2}} r^{2\mu-1} \Gamma(\mu+1)} \int_0^\infty \frac{x}{e^x - 1} \left[(rx)^{\mu-\frac{1}{2}} J_{\mu-\frac{1}{2}}(rx) \right] dx,$$

which, for $\mu = m$, a positive integer

$$(2.15) \quad S_m(r) = \frac{1}{2^{m-1} r^{2m-1} m!} \sqrt{\frac{\pi}{2}} \int_0^\infty \frac{x}{e^x - 1} R_m(rx) dx,$$



On Integral Forms of
Generalised Mathieu Series

P. Cerone and C.T. Lenard

Title Page

Contents



Go Back

Close

Quit

Page 10 of 23

where

$$(2.16) \quad \sqrt{\frac{\pi}{2}} R_m(z) = \sqrt{\frac{\pi}{2}} z^{m-\frac{1}{2}} J_{m-\frac{1}{2}}(z).$$

For $m = 1, 2, 3, 4$ we have

$$\sqrt{\frac{\pi}{2}} R_m(z) = \sin z, \quad \sin z - z \cos z, \quad 3 \sin z - 3z \cos z - z^2 \sin z,$$

and

$$15 \sin z - 15z \cos z - 6z^2 \sin z + z^3 \cos z,$$

respectively.

Thus, for example,


$$S_1(r) = \frac{1}{r} \int_0^\infty \frac{x}{e^x - 1} \sin(rx) dx,$$

$$S_2(r) = \frac{1}{4r^3} \int_0^\infty \frac{x}{e^x - 1} [\sin(rx) - (rx) \cos(rx)] dx,$$

$$S_3(r) = \frac{1}{24r^5} \int_0^\infty \frac{x}{e^x - 1} [3 \sin(rx) - 3(rx) \cos(rx) - (rx)^2 \sin(rx)] dx,$$

and

$$S_4(r) = \frac{1}{192r^7} \int_0^\infty \frac{x}{e^x - 1} [15 \sin(rx) - 15(rx) \cos(rx) - 6(rx)^2 \sin(rx) + (rx) \cos(rx)] dx.$$



On Integral Forms of
Generalised Mathieu Series

P. Cerone and C.T. Lenard

Title Page

Contents



Go Back

Close

Quit

Page 11 of 23

The above results for integer m can also be obtained using the relationship from (1.1) and (1.3)

$$(2.17) \quad S_1(r) = S(r) = \sum_{n=1}^{\infty} \frac{2n}{(n^2 + r^2)^2} = \frac{1}{r} \int_0^{\infty} \frac{x}{e^x - 1} \sin(rx) dx.$$

Formal differentiation with respect to r of (2.17) gives

$$\begin{aligned} (-4r) S_2(r) &= \int_0^{\infty} \frac{x}{e^x - 1} \left[\frac{x \cos rx}{r} - \frac{\sin rx}{r^2} \right] dx \\ &= -\frac{1}{r^2} \int_0^{\infty} \frac{x}{e^x - 1} (\sin rx - rx \cos rx) dx \end{aligned}$$

producing the result above. Continuing in this manner would produce further representations for $S_m(r)$.


The following theorem gives an explicit representation for $S_m(r)$, $m \in \mathbb{N}$.

Theorem 2.2. *For m a positive integer we have*

$$(2.18) \quad S_m(r) = \frac{1}{2^{m-1}} \cdot \frac{1}{r^{2m-1}} \times \frac{1}{m} \sum_{k=0}^{m-1} \frac{(-1)^{\lfloor \frac{3k}{2} \rfloor}}{k!} r^k [\delta_{k \text{ even}} A_k(r) + \delta_{k \text{ odd}} B_k(r)],$$

where

$$(2.19) \quad A_k(r) = \int_0^{\infty} \frac{x^{k+1}}{e^x - 1} \sin(rx) dx, \quad B_k(r) = \int_0^{\infty} \frac{x^{k+1}}{e^x - 1} \cos(rx) dx,$$



On Integral Forms of
Generalised Mathieu Series

P. Cerone and C.T. Lenard

Title Page

Contents



Go Back

Close

Quit

Page 12 of 23

with $\delta_{\text{condition}} = 1$ if condition holds and zero otherwise and $\lfloor x \rfloor$ is the greatest integer less than or equal to x .

Proof. From (2.17) we may differentiate $m - 1$ times with respect to r to produce

$$(2.20) \quad S_1^{(m-1)}(r) = (-1)^{m-1} m! (2r)^{m-1} S_m(r) \\ = \int_0^\infty \frac{x}{e^x - 1} \cdot \frac{d^{m-1}}{dr^{m-1}} \left(\frac{\sin rx}{r} \right) dx.$$

Now,

$$(2.21) \quad \frac{d^{m-1}}{dr^{m-1}} \left(\frac{\sin rx}{r} \right) = \sum_{k=0}^{m-1} \binom{m-1}{k} \frac{d^{m-1-k}}{dr^{m-1-k}} (r^{-1}) \cdot \frac{d^k}{dr^k} (\sin rx)$$


and

$$\frac{d^l}{dr^l} (r^{-1}) = (-1)^l l! r^{-(l+1)}, \\ \frac{d^k}{dr^k} (\sin rx) = (-1)^{\lfloor \frac{k}{2} \rfloor} x^k [\delta_{k \text{ even}} \sin(rx) + \delta_{k \text{ odd}} \cos(rx)]$$

where $\delta_{\text{condition}} = 1$ if condition is true and zero otherwise.

Thus from (2.21)

$$(2.22) \quad \frac{d^{m-1}}{dr^{m-1}} \left(\frac{\sin(rx)}{r} \right) = \frac{1}{r^m} \sum_{k=0}^{m-1} \binom{m-1}{k} (-1)^{m-1-k+\lfloor \frac{k}{2} \rfloor} \\ \times (m-1-k)! r^k x^k [\delta_{k \text{ even}} \sin(rx) + \delta_{k \text{ odd}} \cos(rx)]$$



On Integral Forms of
Generalised Mathieu Series

P. Cerone and C.T. Lenard

Title Page

Contents



Go Back

Close


Quit

Page 13 of 23

$$= \frac{(-1)^{m-1}}{r^m} (m-1)! \sum_{k=0}^{m-1} \frac{(-1)^{\lfloor \frac{3k}{2} \rfloor}}{k!} r^k x^k [\delta_{k \text{ even}} \sin(rx) + \delta_{k \text{ odd}} \cos(rx)].$$

Substitution of (2.22) into (2.20) and simplifying produces the stated result (2.18). \square

Remark 2.4. *The integral representation for $S_m(r)$ given in Theorem 2.2 is simpler than that obtained in [18] as given by (1.7). Further, the derivation here is much more straight forward.*



On Integral Forms of
Generalised Mathieu Series

P. Cerone and C.T. Lenard

Title Page

Contents



Go Back

Close

Quit

Page 14 of 23

3. Bounds for $S_\mu(r)$

It was stated in the introduction that considerable effort has been expended in determining bounds for the generalised Mathieu series. More recently, bounds for the generalised Mathieu series (1.6) has been investigated in particular by Qi and coworkers and by Tomovski and Trenčevski [18].

In a recent article Landau [11] obtained the best possible uniform bounds for Bessel functions using monotonicity arguments. Of particular interest to us here is that he showed that

$$(3.1) \quad |J_\nu(x)| < \frac{b_L}{\nu^{\frac{1}{3}}}$$

uniformly in the argument x and is best possible in the exponent $\frac{1}{3}$ and constant

$$(3.2) \quad b_L = 2^{\frac{1}{3}} \sup_x Ai(x) = 0.674885 \dots,$$

where $Ai(x)$ is the Airy function satisfying

$$w'' - xw = 0.$$


Landau also showed that

$$(3.3) \quad |J_\nu(x)| \leq \frac{c_L}{x^{\frac{1}{3}}}$$

uniformly in the order $\nu > 0$ and the exponent $\frac{1}{3}$ is best possible with

$$(3.4) \quad \begin{aligned} c_L &= \sup_x x^{\frac{1}{3}} J_0(x) \\ &= 0.78574687 \dots \end{aligned}$$

The following theorem is based on the Landau bounds (3.1) – (3.4).



On Integral Forms of
Generalised Mathieu Series

P. Cerone and C.T. Lenard

Title Page

Contents



Go Back

Close

Quit

Page 15 of 23

Theorem 3.1. *The generalised Mathieu series $S_\mu(r)$ satisfies the bounds for $\mu > \frac{1}{2}$ and $r > 0$*

$$(3.5) \quad S_\mu(r) \leq b_L \frac{\sqrt{\pi}}{(2r)^{\mu-\frac{1}{2}}} \cdot \frac{1}{\left(\mu - \frac{1}{2}\right)^{\frac{1}{3}}} \cdot \frac{\Gamma\left(\mu + \frac{3}{2}\right)}{\Gamma(\mu + 1)} \zeta\left(\mu + \frac{3}{2}\right),$$

and

$$(3.6) \quad S_\mu(r) \leq c_L \cdot \frac{\sqrt{\pi}}{2^{\mu-\frac{1}{2}} r^{\mu-\frac{1}{6}}} \cdot \Gamma\left(\mu + \frac{7}{6}\right) \zeta\left(\mu + \frac{7}{6}\right),$$

where b_L and c_L are given by (4.2) and (4.4) respectively.

Proof. From (2.2) and (2.3) we have

$$(3.7) \quad S_\mu(r) \leq C_\mu(r) \int_0^\infty \frac{x^{\mu+\frac{1}{2}}}{e^x - 1} \left| J_{\mu-\frac{1}{2}}(rx) \right| dx, \quad r > 0$$


and so from (3.1) we obtain, on utilising (2.6)

$$S_\mu(r) \leq C_\mu(r) \cdot \frac{b_L}{\left(\mu - \frac{1}{2}\right)^{\frac{1}{3}}} \Gamma\left(\mu + \frac{3}{2}\right) \zeta\left(\mu + \frac{3}{2}\right),$$

which simplifies down to (3.5).

Further, using (3.3) into (3.7) gives

$$\begin{aligned} S_\mu(r) &\leq C_\mu(r) \cdot c_L \cdot \int_0^\infty \frac{x^{\mu+\frac{1}{2}}}{e^x - 1} \cdot \frac{1}{|rx|^{\frac{1}{3}}} dx \\ &= C_\mu(r) \cdot \frac{c_L}{r^{\frac{1}{3}}} \int_0^\infty \frac{x^{\mu+\frac{1}{6}}}{e^x - 1} dx \end{aligned}$$



On Integral Forms of
Generalised Mathieu Series

P. Cerone and C.T. Lenard

Title Page

Contents



Go Back

Close

Quit

Page 16 of 23

which upon using (2.6) produces

$$(3.8) \quad S_\mu(r) \leq C_\mu(r) \cdot \frac{c_L}{r^{\frac{1}{3}}} \cdot \Gamma\left(\mu + \frac{7}{6}\right) \zeta\left(\mu + \frac{7}{6}\right).$$

Simplifying (3.8) and using (2.3) gives the stated result (3.6). \square

Corollary 3.2. *The Mathieu series $S(r)$ satisfies the following bounds*

$$(3.9) \quad S(r) \leq \frac{3\pi}{2^{\frac{11}{12}}} b_L \zeta\left(\frac{5}{2}\right)$$

and

$$(3.10) \quad S(r) \leq \frac{7c_L}{36} \cdot \sqrt{\frac{\pi}{2}} \cdot \Gamma\left(\frac{1}{6}\right) \zeta\left(\frac{13}{6}\right) \cdot r^{-\frac{5}{6}},$$


where b_L and c_L are given by (3.2) and (3.4) respectively.

Proof. Taking $\mu = 1$ in (3.5) and (3.6), noting that $S(r) = S_1(r)$ gives the stated results after some simplification. \square

The following corollary gives coarser bounds than Theorem 3.1 without the presence of the zeta function.

Corollary 3.3. *The generalised Mathieu series $S_\mu(r)$ satisfies the bounds for $\mu > \frac{1}{2}$ and $r > 0$*

$$(3.11) \quad S_\mu(r) \leq 2\sqrt{\pi} \cdot \frac{b_L}{\left(\mu - \frac{1}{2}\right)^{\frac{1}{3}}} \cdot \frac{1}{r^{\mu - \frac{1}{2}}} \cdot \frac{\Gamma\left(\mu + \frac{1}{2}\right)}{\Gamma(\mu + 1)}$$



On Integral Forms of
Generalised Mathieu Series

P. Cerone and C.T. Lenard

Title Page

Contents



Go Back

Close

Quit

Page 17 of 23

and

$$(3.12) \quad S_\mu(r) \leq 2^{\frac{2}{3}} \sqrt{\pi} \cdot \frac{c_L}{r^{\mu-\frac{1}{6}}} \cdot \frac{\Gamma(\mu + \frac{1}{6})}{\Gamma(\mu + 1)}$$

with b_L and c_L given by (3.2) and (3.4).

Proof. We use the well known inequality

$$e^{-x} < \frac{x}{e^x - 1} < e^{-\frac{x}{2}}$$


to produce from (3.7)

$$(3.13) \quad S_\mu(r) \leq C_\mu(r) \int_0^\infty e^{-\frac{x}{2}} x^{\mu-\frac{1}{2}} \left| J_{\mu-\frac{1}{2}}(rx) \right| dx.$$

We know from Laplace transforms or the definition of the gamma function that

$$(3.14) \quad \int_0^\infty e^{-\alpha x} x^s dx = \frac{\Gamma(s+1)}{\alpha^{s+1}}.$$

Hence, placing (3.1) into (3.13) and utilising (3.14) we obtain (3.11) after simplification. A similar approach produces (3.12) starting from (3.3) rather than (3.1). \square



On Integral Forms of
Generalised Mathieu Series

P. Cerone and C.T. Lenard

Title Page

Contents



Go Back

Close

Quit

Page 18 of 23

4. Further Integral Expressions for Generalised Mathieu Series

In [18], Tomovski and Trenčevski gave the integral representation

$$(4.1) \quad S_{\mu}(r) = \frac{2}{\Gamma(\mu+1)} \int_0^{\infty} x^{\mu} e^{-r^2 x} f(x) dx,$$

where

$$(4.2) \quad f(x) = \sum_{n=1}^{\infty} n e^{-n^2 x}, \quad \text{convergent for finite } x > 0,$$

by effectively utilising the result (3.14).

They leave the summation of the series in (4.2) as an open problem.

If we place $\nu = \mu - \frac{1}{2}$ and $\beta = r$, all real in (2.9) then we obtain the identity


$$(4.3) \quad C_{\mu}(r) \int_0^{\infty} e^{-\alpha x} x^{\mu+\frac{1}{2}} J_{\mu-\frac{1}{2}}(rx) dx = \frac{2\alpha}{[\alpha^2 + r^2]^{\mu+1}},$$

where $C_{\mu}(r)$ is as given by (2.3).

Proof B of Theorem 2.1 takes $\alpha = n$ and sums to produce the identity (2.1) – (2.2).

If we take $\alpha = n^{\gamma}$ then we have from (4.3) on summing

$$(4.4) \quad \begin{aligned} S(r; \mu, \gamma) &= \sum_{n=1}^{\infty} \frac{2n^{\gamma}}{(n^{2\gamma} + r^2)^{\mu+1}} \\ &= C_{\mu}(r) \int_0^{\infty} \left(\sum_{n=1}^{\infty} e^{-n^{\gamma} x} \right) x^{\mu+\frac{1}{2}} J_{\mu-\frac{1}{2}}(rx) dx, \end{aligned}$$



On Integral Forms of
Generalised Mathieu Series

P. Cerone and C.T. Lenard

Title Page

Contents



Go Back

Close

Quit

Page 19 of 23

giving an integral representation that was left as an open problem by Qi [15].

As a matter of fact, if we take $\alpha = a_n$ where $\mathbf{a} = (a_1, a_2, \dots, a_n, \dots)$ is a positive sequence, then

$$(4.5) \quad \mathcal{S}(r; \mu; \mathbf{a}) = \sum_{n=1}^{\infty} \frac{2a_n}{(a_n^2 + r^2)^{\mu+1}}$$

$$= C_{\mu}(r) \int_0^{\infty} \left(\sum_{n=1}^{\infty} e^{-a_n x} \right) x^{\mu+\frac{1}{2}} J_{\mu-\frac{1}{2}}(rx) dx.$$

We note that for $\mathbf{a}^+ = (1^{\gamma}, 2^{\gamma}, \dots)$ then

$$\mathcal{S}(r; \mu; \mathbf{a}^+) = S(r; \mu, \gamma).$$

The series

$$\sum_{n=1}^{\infty} \frac{2a_n}{(a_n^2 + r^2)^2}$$

has been investigated in [16].


A closed form expression for

$$F(\mathbf{a}) = \sum_{n=1}^{\infty} e^{-a_n x}, \quad x > 0$$

where a_n is a positive sequence, remains an open problem.

If $\mathbf{a}^* = (1, 2, 3, \dots, n, \dots)$, then

$$F(\mathbf{a}^*) = \frac{1}{e^x - 1}.$$



On Integral Forms of
Generalised Mathieu Series

P. Cerone and C.T. Lenard

Title Page

Contents



Go Back


Close

Quit

Page 20 of 23

References

- [1] M. ABRAMOWITZ AND I.A. STEGUN (Eds.), *Handbook of Mathematical Functions with Formulas, Graphs and Mathematical Tables*, National Bureau of Standards, Applied Mathematics Series **55**, 4th printing, Washington, 1965.
- [2] H. ALZER, J.L. BRENNER AND O.G. RUEHR, On Mathieu's inequality, *J. Math. Anal. Appl.*, **218** (1998), 607–610.
- [3] P.S. BULLEN, *A Dictionary of Inequalities*, Pitman Monographs and Surveys in Pure and Applied Mathematics **97**, Addison Wesley Longman Limited, 1998.
- [4] Ch.-P. CHEN AND F. QI, On an evaluation of upper bound of Mathieu series, *Studies in College Mathematics* (Gāoděng Shùxué Yánjiū), **6**(1) (2003), 48–49. (Chinese).
- [5] P.H. DIANANDA, Some inequalities related to an inequality of Mathieu, *Math. Ann.*, **250** (1980), 95–98.
- [6] A. ELBERT, Asymptotic expansion and continued fraction for Mathieu's series, *Period. Math. Hungar.*, **13**(1) (1982), 1–8.
- [7] O.E. EMERSLEBEN, Über die Reihe $\sum_{k=1}^{\infty} k(k^2+c^2)^{-2}$, *Math. Ann.*, **125** (1952), 165–171.
- [8] I. GAVREA, Some remarks on Mathieu's series, *Mathematical Analysis and Approximation Theory*, 113–117, Burg Verlag, 2002.



On Integral Forms of
Generalised Mathieu Series

P. Cerone and C.T. Lenard

Title Page

Contents




Go Back

Close

Quit

Page **21** of **23**

- [9] I.S. GRADSHTEIN AND I.M. RYZHIK, *Table in Integrals, Series and Products*, (1980), Academic Press, New York.
- [10] B.-N. GUO, Note on Mathieu's inequality, *RGMA Res. Rep. Coll.*, **3**(3) (2000), Art. 5. Available online at <http://rgmia.vu.edu.au/v3n3.html>.
- [11] L. LANDAU, Monotonicity and bounds on Bessel functions, *Electronic J. of Differential Equations*, (2002), 147–154.
- [12] E. MATHIEU, *Traité de physique mathématique, VI–VII: Théorie de l'élasticité des corps solides*, Gauthier-Villars, Paris, 1890.
- [13] D.S. MITRINOVIĆ, J.E. PEČARIĆ AND A.M. FINK, *Classical and New Inequalities in Analysis*, Kluwer Academic Publishers, Dordrecht, 1993.
- [14] F. QI, An integral expression and some inequalities of Mathieu type series, *Rostock Math. Kolloq.*, (2003), 389–392.
- [15] F. QI, Inequalities for Mathieu's series, *RGMA Res. Rep. Coll.*, **4**(2) (2001), Art. 3, 187–193. Available online at <http://rgmia.vu.edu.au/v4n2.html>
- [16] F. QI, Integral expressions and inequalities of Mathieu type series, *RGMA Res. Rep. Coll.*, **6**(2) (2003), Art. 10. Available online at <http://rgmia.vu.edu.au/v6n2.html>
- [17] F. QI AND Ch.-P. CHEN, Notes on double inequalities of Mathieu's series, *Internat. J. Pure Appl. Math.* (2003), accepted.



**On Integral Forms of
Generalised Mathieu Series**

P. Cerone and C.T. Lenard

Title Page

Contents




Go Back

Close

Quit

Page 22 of 23

- [18] Z. TOMOVSKI AND K. TRENČEVSKI, On an open problem of Bai-Ni Gou and Feng Qi, *J. Inequal. Pure Appl. Math.*, **4**(2) (2003), Art. 19. Available online at http://jipam.vu.edu.au/v4n2/102_02.html.
- [19] Ch.-L. WANG AND X.-H. WANG, A refinement of the Mathieu inequality, *Univ. Beograd. Publ. Elektroteh. Fak. Ser. Mat.*, No. **716-734** (1981), 22–24.
- [20] J.E. WILKINS, Jr., Solution of Problem 97-1, *Siam Rev.*
- [21] G.N. WATSON, A treatise on the theory of Bessel functions, 2nd Edn., Cambridge University Press (1966).



**On Integral Forms of
Generalised Mathieu Series**

P. Cerone and C.T. Lenard

Title Page

Contents



Go Back

Close

Quit

Page 23 of 23