


CERTAIN INEQUALITIES CONCERNING SOME KINDS OF CHORDAL POLYGONS

MIRKO RADIĆ

University of Rijeka,
Faculty of Philosophy,
Department of Mathematics,
51000 Rijeka, Omladinska 14, Croatia.
EMail: mradic@pefri.hr



volume 5, issue 1, article 1,
2004.

*Received 09 July, 2003;
accepted 25 November, 2003.*

Communicated by: J. Sándor

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)



Abstract

This paper deals with certain inequalities concerning some kinds of chordal polygons (Definition 1.2). The main part of the article concerns the inequality

$$\sum_{j=1}^n \cos \beta_j > 2k,$$

where

$$\sum_{j=1}^n \beta_j = (n - 2k) \frac{\pi}{2}, \quad n - 2k > 0, \quad 0 < \beta_j < \frac{\pi}{2}, \quad j = \overline{1, n}.$$

This inequality is considered and proved in [5, Theorem 1, pp.143-145]. Here we have obtained some new results. Among others we found some chordal polygons with the property that $\sum_{j=1}^n \cos^2 \beta_j = 2k$, where $n = 4k$ (Theorem 2.14). Also it could be mentioned that Theorem 2.16 is a modest generalization of the Pythagorean theorem.

2000 Mathematics Subject Classification: Primary: 51E12.

Key words: Inequality, k -chordal polygon, k -inscribed chordal polygon, index of k -inscribed chordal polygon, characteristic of k -chordal polygon.

Contents

1	Introduction	3
2	Certain Inequalities Concerning k -Chordal Polygons	11
3	Inequalities Concerning k -Inscribed Polygons	35
	References	

Certain Inequalities Concerning Some Kinds Of Chordal Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 2 of 40

1. Introduction

To begin, we will quote some results given in [5], [6].

A polygon with vertices A_1, \dots, A_n (in this order) will be denoted by $\mathcal{A} \equiv A_1 \cdots A_n$ and the lengths of its sides we will denote by a_1, \dots, a_n . The interior angle at the vertex A_j will be signed by α_j or $\sphericalangle A_j$. Thus

$$\sphericalangle A_j = \sphericalangle A_{j-1}A_jA_{j+1}, \quad j = \overline{1, n},$$

where $A_0 = A_n$ and $A_{n+1} = A_1$.

A polygon \mathcal{A} is called a chordal polygon if there exists a circle \mathcal{K} such that $A_j \in \mathcal{K}$, $j = \overline{1, n}$.

Remark 1.1. *We shall assume that the considered chordal polygon has the property that no two of its consecutive vertices are the same.*

For \mathcal{A} chordal, by C and r we denote its centre and the radius of its circum-circle \mathcal{K} respectively.

A very important role will be played by the angles

$$(1.1) \quad \beta_j = \sphericalangle CA_jA_{j+1},$$

$$(1.2) \quad \varphi_j = \sphericalangle A_jCA_{j+1}, \quad j = \overline{1, n}.$$

We shall use oriented angles, as it is known, an angle $\sphericalangle PQR$ is positively or negatively oriented if it is going from QP to QR counter-clockwise or clockwise. It is very important to emphasize that the angles β_j , φ_j have opposite orientations, see e.g. Fig. 1. Of course, the measure of an oriented angle will be taken with $+$ or $-$ depending on whether the angle is positively or negatively oriented. The measure of an angle will usually be expressed by radians.



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 3 of 40




Figure 1:


Remark 1.2. For the sake of simplicity, we shall also write the measures of the oriented angles in (1.1) and (1.2) as β_j , φ_j . Obviously, for all β_j , φ_j the following is valid

$$0 \leq |\beta_j| < \frac{\pi}{2}, \quad 0 < |\varphi_j| \leq \pi,$$

since no two consecutive vertices in $A_1 \cdots A_n$ are the same, compare Remark 1.1.

Remark 1.3. In the sequel, unless specified otherwise, we shall suppose that no $\beta_j = 0$, i.e.

$$0 < |\beta_j| < \frac{\pi}{2}, \quad j = \overline{1, n}.$$



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 4 of 40

Accordingly, in the sequel when we refer to chordal polygons, it will be meant (by Remark 1.1 and Remark 1.2) that the polygon has no two consecutive overlapping vertices and no one of its sides is its diameter.

Definition 1.1. Let \mathcal{A} be a chordal polygon. We say that \mathcal{A} is of the first kind if inside of \mathcal{A} there is a point O such that all oriented angles $\sphericalangle A_j O A_{j+1}$, $j = \overline{1, n}$ have the same orientation. If such a point O does not exist, we say that \mathcal{A} is of the second kind.

Definition 1.2. Let \mathcal{A} be a chordal polygon and let $O \in \text{Int}(\mathcal{A})$, such that

$$|\psi_1 + \cdots + \psi_n| = 2k\pi,$$

where $\psi_j = \text{measure of the oriented angle } \sphericalangle A_j O A_{j+1}$ and k is a positive integer. Then \mathcal{A} is called a k -inscribed chordal polygon or, for brevity, k -inscribed polygon if O is such a point that k is maximal, i.e. no other interior point P exists such that $k < m$ and at the same time the following is valid

$$|\psi_1 + \cdots + \psi_n| = 2m\pi,$$

where now $\psi_j = \text{measure of the oriented angle } \sphericalangle A_j P A_{j+1}$.

For example, the heptagon $A_1 \cdots A_7$ drawn in Fig. 2 is 2-inscribed chordal, since $|\psi_1 + \cdots + \psi_7| = 4\pi$. This heptagon is, according to Definition 1.1, of the first kind – all angles ψ_j have the same, negative orientation.

Of course, a k -inscribed polygon is of the second kind if not all angles ψ_j have the same orientation.



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 5 of 40




Figure 2:


Definition 1.3. Let \mathcal{A} be a k -inscribed chordal n -gon and let

$$|\varphi_1 + \dots + \varphi_n| = 2m\pi, \quad m \in \{0, 1, 2, \dots, k\}$$

and φ_j is given by (1.2). Then m is the index of \mathcal{A} , denoted as $\text{Ind}(\mathcal{A})$.

For example, the heptagon on Fig. 2 has index equal to 1, since $|\varphi_1 + \dots + \varphi_7| = 2\pi$. (See Figure 3. Let us remark that φ_4 is positively and all other angles are negatively oriented.)

Definition 1.4. A k -inscribed polygon \mathcal{A} will be called a k -chordal polygon if it is of the first kind and $\text{Ind}(\mathcal{A}) = k$.



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 6 of 40




Figure 3:


Theorem A. Let A be a k -chordal polygon and let β_j be given by (1.1). Then we have

$$|\beta_1 + \dots + \beta_n| = (n - 2k) \frac{\pi}{2}.$$

Proof. Since every k -chordal polygon is of the first kind (Definition 1.4), then either $\beta_j > 0$, $j = \overline{1, n}$ or $\beta_j < 0$, $j = \overline{1, n}$. If $\beta_j > 0$, then $\varphi_j < 0$ and the following holds

$$\varphi_1 + \dots + \varphi_n = -2k\pi.$$

In this case, because $2\beta_j + |\varphi_j| = \pi$ or $\varphi_j = 2\beta_j - \pi$, the above equality can be



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 7 of 40

written as

$$\sum_{j=1}^n (2\beta_j - \pi) = -2k\pi,$$

or equivalently

$$\sum_{j=1}^n \beta_j = (n - 2k)\frac{\pi}{2}.$$

If $\beta_j < 0$, then $\varphi_j > 0$ and it holds $\varphi_1 + \dots + \varphi_n = 2k\pi$. In this case we have

$$\sum_{j=1}^n \beta_j = -(n - 2k)\frac{\pi}{2}.$$

□


If \mathcal{A} is a k -chordal polygon, then each β_j , $j = \overline{1, n}$, is negative if \mathcal{A} is positively oriented and vice versa. But in the case when \mathcal{A} is a k -inscribed polygon of the second kind, then some of the β_j are negative and some are positive.

Remark 1.4. *In the sequel, for the sake of simplicity, we shall assume that the considered polygon is negatively oriented. Thus, in the case when a k -inscribed polygon \mathcal{A} is negatively oriented, then*

$$\varphi_1 + \dots + \varphi_n \leq 0 \quad \text{but} \quad \beta_1 + \dots + \beta_n \geq 0.$$

Finally, let us point out that for $\text{Ind}(\mathcal{A}) = 0$, the following holds

$$\varphi_1 + \dots + \varphi_n = 0 = \beta_1 + \dots + \beta_n.$$



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 8 of 40

Theorem B. Let \mathcal{A} be a k -inscribed polygon. Then

$$(1.3) \quad |\beta_1 + \cdots + \beta_n| = [n - 2(m + \nu)] \frac{\pi}{2},$$

where $\text{Ind}(\mathcal{A}) = m$ and ν is number of all negative β_j 's.

Proof. As $\varphi_j = -\pi + 2\beta_j$ if $\beta_j > 0$ and $\varphi_j = \pi + 2\beta_j$ if $\beta_j < 0$, the equality $\varphi_1 + \cdots + \varphi_n = -2m\pi$ can be written as

$$2\beta_1 + \cdots + 2\beta_n + \nu\pi - (n - \nu)\pi = -2m\pi,$$

from which (1.3) follows.

If $\beta_{j_1}, \dots, \beta_{j_\nu}$ are the negative angles in (1.3), then we have

$$(1.4) \quad |\beta_1| + \cdots + |\beta_n| = [n - 2(m + \nu)] \frac{\pi}{2} + 2\tau,$$

where $\tau = -(\beta_{j_1} + \cdots + \beta_{j_\nu})$. □


The greatest part of this article is in some way connected to the following theorem, see [5, Theorem 1] as well.

Theorem C. Let \mathcal{A} be a k -chordal polygon. Then

$$(1.5) \quad \sum_{j=1}^n \cos \beta_j > 2k,$$

where

$$\sum_{j=1}^n \beta_j = (n - 2k) \frac{\pi}{2}, \quad 0 < \beta_j < \frac{\pi}{2}, j = \overline{1, n}.$$



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 9 of 40

Proof. Since $\cos \pi x > 1 - 2x$ if $x \in (0, 1/2)$, putting $\alpha = \pi x$ we obtain

$$(1.6) \quad \cos \alpha > 1 - \frac{2}{\pi} \alpha, \quad 0 < \alpha < \frac{\pi}{2}.$$

Thus, we deduce

$$\sum_{j=1}^n \cos \beta_j > n - \frac{2}{\pi} \sum_{j=1}^n \beta_j = n - \frac{2}{\pi} (n - 2k) \frac{\pi}{2} = 2k.$$


□

Remark 1.5. After this paper had been written, J. Sándor informed me that the inequality (1.6) follows from Jordan's inequality

$$\sin x > \frac{2}{\pi} x, \quad x \in \left(0, \frac{\pi}{2}\right),$$

putting $x = \pi/2 - \alpha$.

At this point let us remark that we can consult the articles [1], [2], [3], [4] and [8] for further information and generalizations of certain inequalities concerning plane and space polygons.



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 10 of 40

2. Certain Inequalities Concerning k -Chordal Polygons

In this section we deal with k -chordal polygons. By Remark 1.4 and Definition 1.4, all angles β_j are positive. First of all we give the following remark.

Remark 2.1. *By the relation $\beta_j \approx 0$ we mean that β_j is near to zero, but it is positive. Similarly, $\beta_j \approx \pi/2$ denotes the case, when β_j is close to $\pi/2$, but it is less than $\pi/2$.*

Theorem 2.1. *Let k, n be positive integers such that $n - 2k > 0$ and let β_1, \dots, β_n be angles such that*

$$(2.1) \quad \sum_{j=1}^n \beta_j = (n - 2k) \frac{\pi}{2}, \quad 0 < \beta_j < \frac{\pi}{2}, j = \overline{1, n}.$$


Then there exists a positive number h such that

$$(2.2) \quad \sum_{j=1}^n \cos^h \beta_j = 2k,$$

where

$$(2.3) \quad 1 < h < \frac{\log \frac{2k}{2k+1}}{\log \cos \frac{\pi}{4k+2}}.$$

Proof. From (1.5) it follows that there is a positive h for which (2.2) holds as well. Now, we only need to prove that this h satisfies (2.3). For this purpose we will first prove the following lemma.



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 11 of 40

Lemma 2.2. Let $h \geq 1$ be fixed. Then the function $y = \cosh^h x$ is concave in the interval $(0, \arctan(1/\sqrt{h-1}))$.

Proof. As

$$y'' = h \cosh^{h-2} x [(h-1) \sin^2 x - \cos^2 x],$$

it follows that

$$\begin{aligned} y'' < 0 & \quad \text{if} \quad (h-1) \tan^2 x < 1, \\ y'' > 0 & \quad \text{if} \quad (h-1) \tan^2 x > 1. \end{aligned}$$

Thus, the function $y = \cosh^h x$ is concave in $(0, \arctan(1/\sqrt{h-1}))$ and convex in the interval $(\arctan(1/\sqrt{h-1}), \pi/2)$. This proves Lemma 2.2. \square


Now, assume that (2.1) is fulfilled. Then it is easy to see that the sum $\sum_{j=1}^n \cos^h \beta_j$ has the following properties.

(i₁) If $(n-2k)\frac{\pi}{2n} < \arctan(1/\sqrt{h-1})$, then the sum $\sum_{j=1}^n \cos^h \beta_j$ attains its maximum for $\beta_1 = \dots = \beta_n = (n-2k)\frac{\pi}{2n}$.

(i₂) If $(n-2k)\frac{\pi}{2n} > \arctan(1/\sqrt{h-1})$, then the sum $\sum_{j=1}^n \cos^h \beta_j$ attains its minimum for $\beta_1 = \dots = \beta_n = (n-2k)\frac{\pi}{2n}$.

(i₃) If $\beta_1 = \dots = \beta_{2k} \approx 0$, $\beta_{2k+1} = \dots = \beta_n \approx \frac{\pi}{2}$, then

$$\sum_{j=1}^n \cos^h \beta_j \approx 2k.$$



**Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)**

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 12 of 40

(i₄) For h sufficiently large the following result holds:

$$n \cos^h(n - 2k) \frac{\pi}{2n} < 2k.$$

(i₅) There are $h_1 \geq 1$, $h_2 > 1$ such that

$$n \cos^{h_1}(n - 2k) \frac{\pi}{2n} > 2k, \quad n \cos^{h_2}(n - 2k) \frac{\pi}{2n} < 2k,$$

and the equality $n \cos^{h_0}(n - 2k) \frac{\pi}{2n} = 2k$ is obtained for

$$(2.4) \quad h_0 = h(n, k) = \frac{\log \frac{2k}{n}}{\log \cos(n - 2k) \frac{\pi}{2n}}.$$

Lemma 2.3. Let $h(k)$, $k \in \mathbb{N}$ be given by


$$(2.5) \quad h(k) = \frac{\log \frac{2k}{2k+1}}{\log \cos \frac{\pi}{4k+2}}.$$

Then the sum $\sum_{j=1}^n \cos^{h(k)} \beta_j$ attains its maximum for

$$(2.6) \quad \beta_1 = \dots = \beta_{2k+1} \approx \frac{\pi}{4k+2}, \quad \beta_{2k+2} = \dots = \beta_n \approx \frac{\pi}{2}.$$

Proof. Firstly let us remark that $\frac{\pi}{4k+2} = \frac{\pi}{2} : (2k+1)$ and this practically means that

$$\beta_{2k+2} + \dots + \beta_n = (n - (2k+1)) \frac{\pi}{2},$$



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 13 of 40

so, from

$$(2.7) \quad (2k + 1) \cos^{h(k)} \frac{\pi}{4k + 2} + (n - (2k + 1)) \cos^{h(k)} \frac{\pi}{2} = 2k$$

we get (2.5). To prove Lemma 2.3 we have to prove the inequality

$$(2.8) \quad \arctan \frac{1}{\sqrt{h(k)} - 1} > \frac{\pi}{4k + 2}.$$

Starting from (2.6), we can write

$$\sqrt{h(k)} - 1 < \cot \frac{\pi}{4k + 2},$$

i.e.

$$h(k) < 1 + \cot^2 \frac{\pi}{4k + 2},$$

so

$$\frac{\log \frac{2k}{2k+1}}{\log \cos \frac{\pi}{4k+2}} < 1 + \cot^2 \frac{\pi}{4k + 2},$$


implying

$$\log \frac{2k}{2k + 1} > \log \left(\cos \frac{\pi}{4k + 2} \right)^{1/\sin^2 \frac{\pi}{4k+2}},$$

thus

$$\frac{2k}{2k + 1} > \frac{1}{\sqrt{\left(1 - \sin^2 \frac{\pi}{4k+2}\right)^{-1/\sin^2 \frac{\pi}{4k+2}}}.$$

Letting $k \rightarrow \infty$ in the last relation we get a valid result since the expression on the left-hand side tends to 1, while the right-hand side tends to $1/\sqrt{e}$. This finishes the proof of Lemma 2.3. \square



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 14 of 40

Finally we have to show that

$$(2k + 1) \cos^{h(k)} \frac{\pi}{4k + 2} > 2k,$$

$$(2k + 1) \cos^{h(k)} \frac{\pi}{4k + 2} > (2k + 2) \cos^{h(k)} \frac{\pi}{2k + 2},$$

where one can write $2k = 2k \cos^{h(k)} 0$, $\frac{\pi}{2k+2} = (\frac{\pi}{2} + \frac{\pi}{2}) : (2k + 2)$. For this purpose it is sufficient to check that the above relations hold, e.g. for $k = 1, 2, 3$. Thus, we have

$$3 \cos^{h(1)} \frac{\pi}{6} = 2.000000001,$$

$$5 \cos^{h(2)} \frac{\pi}{10} = 4,$$

$$7 \cos^{h(3)} \frac{\pi}{14} = 6.000000006,$$


$$4 \cos^{h(1)} \frac{\pi}{4} = 1.50585114 < 3 \cos^{h(1)} \frac{\pi}{6},$$

$$6 \cos^{h(2)} \frac{\pi}{6} = 3.164961846 < 5 \cos^{h(2)} \frac{\pi}{10},$$

$$8 \cos^{h(3)} \frac{\pi}{8} = 4.947027176 < 7 \cos^{h(3)} \frac{\pi}{14}.$$

Let us remark that $\frac{\pi}{4k+2} \approx \frac{1}{2} \frac{\pi}{2k+2}$ for sufficiently large k . This completes the proof of the Theorem 2.1. \square

As an interesting illustrative example we provide the following table.



**Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)**

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 15 of 40

k	$h(k)$	$\arctan 1/\sqrt{h(k) - 1}$	$\frac{\pi}{4k+2}$
1	2.818841678	36.55639173^0	30^0
2	4.446703708	28.30865018^0	18^0
3	6.070896923	23.94487335^0	12.85714286^0
4	7.693796543	21.13214916^0	10^0
5	9.316082999	19.12497372^0	8.18181812^0
10	17.42431500	13.86082784^0	4.28571428^0
100	163.3293834	04.48781187^0	0.44776119^0

Table 1.

Example 2.1. We give an illustrative example with respect to $h(2)$. The function $y = \cos^{h(2)} x$ is shown in Fig. 4 for $x \in [0, \frac{\pi}{2}]$. The point $x_0 = \arctan 1/\sqrt{h(2) - 1} = 28.30865018$ is its inflection point. For $n = 11$, under the constraint (2.1),





Figure 4:



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 16 of 40

the sum $\sum_{j=1}^{11} \cos^{h(2)} \beta_j$ takes its maximum for

$$\beta_1 = \dots = \beta_5 = \frac{\pi}{10}, \quad \beta_6 = \dots = \beta_{11} \approx \frac{\pi}{2}.$$

Here we point out that $y = \cos^{h(2)} x$ is concave in $(0, x_0)$ and

$$5 \cos^{h(2)} \frac{\pi}{10} \geq \sum_{j=1}^5 \cos^{h(2)} x_j,$$


holds true for every x_1, \dots, x_5 such that $x_1 + \dots + x_5 = \frac{\pi}{2}$, $0 < x_j < \frac{\pi}{2}$, $j = \overline{1, 5}$.

Also,

$$\begin{aligned} 5 \cos^{h(2)} \frac{\pi}{10} &> 6 \cos^{h(2)} \frac{2\pi}{12} = 3.164961846 \\ &> 7 \cos^{h(2)} \frac{3\pi}{14} = 2.343170592 \\ &> 8 \cos^{h(2)} \frac{4\pi}{16} = 1.713146048 \\ &\dots\dots\dots \\ &> 11 \cos^{h(2)} \frac{7\pi}{22} = 0.714031536, \end{aligned}$$

holds, where

$$\frac{2\pi}{12} = \left(\frac{\pi}{2} + \frac{\pi}{2} \right) : 6, \quad \beta_7 = \dots = \beta_{11} \approx \frac{\pi}{2},$$



**Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)**

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 17 of 40

$$\frac{3\pi}{14} = \left(\frac{\pi}{2} + \frac{\pi}{2} + \frac{\pi}{2}\right) : 7, \quad \beta_8 = \dots = \beta_{11} \approx \frac{\pi}{2},$$

etc.

These relations can be clearly explained by the convexity of $\cos^{h(2)} x$ on $(x_0, \frac{\pi}{2})$ and by $x_0 < \frac{2\pi}{12} < \frac{3\pi}{14} < \dots < \frac{7\pi}{22}$.

Now, we shall state and prove some corollaries of Theorem 2.1.

Corollary 2.4. *One has $h(k) \rightarrow \infty$ when $k \rightarrow \infty$.*

Proof. It can be found that

$$\frac{\frac{d}{dk} \left(\log \frac{2k}{2k+1}\right)}{\frac{d}{dk} \left(\log \cos \frac{\pi}{4k+2}\right)} = \frac{2k+1}{4} \cot \frac{\pi}{4k+2}.$$

□


For example, $h(500) = 811.78$, $h(10^3) = 1622.38$, $h(10^4) = 16233.22$, etc.

Corollary 2.5. *$h(k)$ is the same for all $n > 2k$.*

Proof. This is a consequence of (2.7).

□

Corollary 2.6. *Let k be a fixed positive integer and $h(n, k)$ be given by (2.4). Then $h(n, k) \rightarrow 1$ when $n \rightarrow \infty$.*



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 18 of 40

Proof. It can be easily seen that

$$\frac{\frac{d}{dn} \left(\log \frac{2k}{n} \right)}{\frac{d}{dn} \left(\log \sin \frac{k\pi}{n} \right)} = \frac{n}{k\pi} \tan \frac{k\pi}{n}.$$

Now, obvious transformations give the assertion. □

For example, we have

$$h(5, 1) = 1.72432, \quad h(6, 1) = 1.58496, \quad h(7, 1) = 1.50035,$$

$$h(5, 2) = 4.44670, \quad h(6, 2) = 2.81884, \quad h(7, 2) = 2.27279.$$

Corollary 2.7. *Let n_1, k_1, n_2, k_2 be any given positive integers, such that $n_j > 2k_j, j = 1, 2$. If*

$$(2.9) \quad \frac{k_1}{n_1} = \frac{k_2}{n_2},$$


then $h(n_1, k_1) = h(n_2, k_2)$.

Proof. Suppose that (2.9) holds. Then we can write

$$\frac{k_1\pi}{n_1} = \frac{k_2\pi}{n_2} \implies (n_1 - 2k_1)\frac{\pi}{2n_1} = (n_2 - 2k_2)\frac{\pi}{2n_2}.$$

From this we easily deduce the assertion. □

Corollary 2.8. *Let $k \in \mathbb{N}$ be fixed. Then $h(n, k) \leq h(k)$ for any integer $n > 2k$. The equality $h(n, k) = h(k)$ holds for $n = 2k + 1$.*



**Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)**

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 19 of 40

Proof. This follows from the Corollary 2.4 and Corollary 2.6. The asserted inequality is the straightforward consequence of (2.4) and (2.5). \square

As an example we give the following numerical results (see Table 1 and the previous example):

$$h(5, 1) = 1.72432 < h(1) = 2.81884$$

$$h(5, 2) = 4.44670 = h(2) = 4.44670 \quad (\text{since } 5 = 2 \cdot 2 + 1)$$

$$h(6, 2) = 2.81884 < h(2).$$

Theorem 2.9. *Let \mathcal{A} be a given k -chordal n -gon and let a_1, \dots, a_n be the lengths of its sides. Then*


$$(2.10) \quad \left(\frac{a_1^{h(k)} + \dots + a_n^{h(k)}}{2k} \right)^{1/h(k)} \leq 2r < \frac{a_1 + \dots + a_n}{2k},$$

where r denotes the radius of the circumcircle of \mathcal{A} .

Proof. From (2.2) and (2.3) it follows that

$$(2.11) \quad \sum_{j=1}^n \cos^{h(k)} \beta_j < 2k < \sum_{j=1}^n \cos \beta_j.$$

Since $a_j = 2r \cos \beta_j$, $j = \overline{1, n}$, the above inequalities can be written as in (2.10). Thus, Theorem 2.9 is proved. \square



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 20 of 40

Corollary 2.10. *The following equality holds:*

$$(2.12) \quad a_1^{m(k)} + \dots + a_n^{m(k)} = 2k(2r)^{m(k)},$$

where $1 < m(k) \leq h(k)$.

Corollary 2.11. *Let a_1, \dots, a_n be given lengths. Then there exists a k -chordal n -gon with radius r whose sides have given lengths, if there is an $m(k)$ satisfying (2.12). (In this connection Example 2.6 may be interesting.)*

Corollary 2.12. *Let $a_1 = \dots = a_n = a$. Then*

$$(2.13) \quad r = \frac{a}{2} \left(\frac{n}{2k} \right)^{1/h(n,k)}.$$

Proof. The relation (2.13) follows from (2.2) if $\beta_1 = \dots = \beta_n$. □


Corollary 2.13. *The following equality holds:*

$$\sin \frac{k\pi}{n} = \left(\frac{2k}{n} \right)^{1/h(n,k)}.$$

Proof. As $a = 2r \cos(n - 2k) \frac{\pi}{2n} = 2r \sin \frac{k\pi}{n}$, we have $a/(2r) = \sin \frac{k\pi}{n}$. From (2.13) it follows that

$$\frac{a}{2r} = \left(\frac{2k}{n} \right)^{1/h(n,k)}.$$

□



**Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)**

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 21 of 40

Example 2.2. Let $\beta_1 = 20^\circ$, $\beta_2 = 30^\circ$, $\beta_3 = 40^\circ$, $r = 5$. By the well-known relation $a_j = 2r \cos \beta_j$ we get

$$a_1 = 9.396926208, \quad a_2 = 8.660254038, \quad a_3 = 7.660444431.$$

From $\beta_1 + \beta_2 + \beta_3 = (3 - 2 \cdot 1)\frac{\pi}{2}$, it is clear that $k = 1$. It can be found that

$$\cos^m \beta_1 + \cos^m \beta_2 + \cos^m \beta_3 = 1.999999783 \quad \text{for} \quad m = 2.737684,$$

$$\cos^m \beta_1 + \cos^m \beta_2 + \cos^m \beta_3 = 2.000000061 \quad \text{for} \quad m = 2.737683.$$

Thus, we have the approximative equality

$$a_1^m + a_2^m + a_3^m = 2k(2r)^m,$$

where $k = 1$ and $m = 2.737683$. We see immediately that $2.737683 < h(1) = 2.81884$. But it follows from the fact that β_j are not equal to each other, i.e. $\beta_j \neq \pi/6$. Therefore


$$\cos^{h(1)} 20^\circ + \cos^{h(1)} 30^\circ + \cos^{h(1)} 40^\circ = 1.97761 < 2;$$

in the case of equal β_j 's we have $3 \cos^{h(1)} \pi/6 = 2$.

Example 2.3. Let $\beta_1 = 10^\circ$, $\beta_2 = 15^\circ$, $\beta_3 = 18^\circ$, $\beta_4 = 22^\circ$, $\beta_5 = 25^\circ$, $r = 4$. With the help of $a_j = 2r \cos \beta_j$ we derive

$$a_1 = 7.87846202, \quad a_2 = 7.72740661, \quad a_3 = 7.60845213,$$

$$a_4 = 7.41747084, \quad a_5 = 7.25046230.$$



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 22 of 40

From $\beta_1 + \dots + \beta_5 = (5 - 2 \cdot 2) \frac{\pi}{2}$ we conclude that $k = 2$. The corresponding pentagon is shown in Fig. 5. Let us remark that

$$\sum_{j=1}^5 \text{measure of } \sphericalangle A_j C A_{j+1} = 4\pi.$$

It can be easily computed that





Figure 5:

$$\sum_{j=1}^5 \cos^m \beta_j = 3.999977021 \quad \text{for} \quad m = 4.2082782$$



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 23 of 40

$$\sum_{j=1}^5 \cos^m \beta_j = 4.000022422 \quad \text{for} \quad m = 4.2082781.$$

Finally the approximate equality $\sum_{j=1}^5 a_j^m = 2k(2r)^m$ holds for $k = 2$ and $m = 4.2082782$, where $m < h(2) = 4.446703708$.

Example 2.4. There is a 1-chordal pentagon \mathcal{B} such that $b_j = |B_j B_{j+1}| = |A_j A_{j+1}| = a_j$, $j = \overline{1, 5}$, where \mathcal{A} is the 2-chordal pentagon shown in Fig. 5. It can be found that

$$\sum_{j=1}^5 \arccos \frac{a_j}{12.90} = 270.011718^0 > 270^0,$$

$$\sum_{j=1}^5 \arccos \frac{a_j}{12.89} = 269.955703^0 < 270^0.$$

Thus, the radius of the circumcircle of \mathcal{B} satisfies the relation


$$12.89 < 2r_B < 12.90$$

and for the angles of \mathcal{B} we have $\beta_1 + \dots + \beta_5 = (5 - 2 \cdot 1) \frac{\pi}{2}$, since, here $k = 1$.

Thus, besides the equality in Example 2.3 there is the equality

$$a_1^m + \dots + a_5^m = 2k(2r_B)^m,$$

for $k = 1$ and $m < h(1) = 2.81884$.



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 24 of 40

Example 2.5. Let $\beta_1 = 9^\circ$, $\beta_2 = 63^\circ$, $\beta_3 = 65^\circ$, $\beta_4 = 66^\circ$, $\beta_5 = 67^\circ$, $r = 3$. Then there is 1-chordal pentagon such that

$$a_1^m + \cdots + a_5^m = 2 \cdot 6^m, \quad 1 < m < h(1).$$

But there is no 2-chordal pentagon $\mathcal{B} \equiv B_1 \cdots B_5$ such that $a_j = |B_j B_{j+1}|$. Indeed, it is easy to show this by

$$a_1^m + \cdots + a_5^m < 4(2r_{\mathcal{B}})^m$$

for all $m \geq 1$, and for all $r_{\mathcal{B}} \geq 3 \cos \beta_1$ when

$$a_1 = 5.92613, \quad a_2 = 2.72394, \quad a_3 = 2.53571, \quad a_4 = 2.44042, \quad a_5 = 2.34439.$$

Finally, we can show that for $m = 1$ and $m = h(2)$ we have

$$a_1^m + \cdots + a_5^m < 4a_1^m.$$


Definition 2.1. Let \mathcal{A} be a k -chordal n -gon. Then the number $m > 1$ for which we obtain

$$\sum_{j=1}^n a_j^m = 2k(2r)^m$$

is the characteristic of \mathcal{A} in the notation $\text{Char}(\mathcal{A})$. Here r is the radius of the circumcircle of \mathcal{A} and $a_j = |A_j A_{j+1}|$, $j = \overline{1, n}$.

Remark 2.2. By Theorem 2.1, we have

$$(2.14) \quad 1 < \text{Char}(\mathcal{A}) < h(k),$$



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 25 of 40

where $h(k)$ is given by (2.5).

In particular, if $\beta_1 = \dots = \beta_n = (n - 2k) \frac{\pi}{2n}$, then

$$\text{Char}(\mathcal{A}) = h(n, k),$$

where $h(n, k)$ is given by (2.4).

Remark 2.3. Since there are situations when certain angles β_j are close to 0, and other angles are close to $\pi/2$, it is clear that instead of constraint (2.6) in the proving procedure of Theorem 2.1, we can take

$$\beta_1 = \dots = \beta_{2k+1} = \frac{\pi}{4k+2}, \quad \beta_{2k+2} = \dots = \beta_n = \frac{\pi}{2}.$$

Thus, instead of (2.14) it can be written

$$1 < \text{Char}(\mathcal{A}) \lesssim h(k),$$

where $\text{Char}(\mathcal{A}) \lesssim h(k)$ means that $\text{Char}(\mathcal{A}) < h(k)$ and that $\text{Char}(\mathcal{A})$ may be close to $h(k)$.

For example, let \mathcal{A} be a 1-chordal pentagon such that


$$\beta_1 = \beta_2 = \beta_3 = \frac{90.000002^0}{3}, \quad \beta_4 = \beta_5 = 89.999999^0.$$

Then

$$\text{Char}(\mathcal{A}) \approx h(1) = 2.818841678,$$

because

$$3 \cos^{h(1)} \frac{90.000002^0}{3} + 2 \cos^{h(1)} 89.999999^0 = 1.999999962 \approx 2.$$



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 26 of 40

Example 2.6. Let \mathcal{A} be 1-chordal quadrilateral such that

$$\beta_1 + \beta_3 = \frac{\pi}{2} = \beta_2 + \beta_4.$$

Then $\text{Char}(\mathcal{A}) = 2$. This is clear, since

$$\cos^2 \beta_1 + \cos^2 \beta_3 = 1 = \cos^2 \beta_2 + \cos^2 \beta_4.$$

Thus

$$a_1^2 + a_2^2 + a_3^2 + a_4^2 = 2(2r)^2.$$

Of course this property is not true for every 1-chordal quadrilateral; there are chordal quadrilaterals where $\beta_1 + \beta_3 \neq \frac{\pi}{2}$, compare Fig 6.





Figure 6:



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 27 of 40

Example 2.7. Let \mathcal{A} be a 2-chordal octagon such that

$$(2.15) \quad \beta_1 + \beta_5 = \beta_2 + \beta_6 = \beta_3 + \beta_7 = \beta_4 + \beta_8 = \frac{\pi}{2}.$$

As an illustration, see Fig. 7





Figure 7:

As

$$\cos^2 \beta_j + \cos^2 \beta_{j+4} = 1, \quad j = 1, 2, 3, 4,$$

then $\text{Char}(\mathcal{A}) = 2$. Thus we clearly deduce by $\sum_{j=1}^8 \cos^2 \beta_j = 4$ that

$$\sum_{j=1}^8 a_j^2 = 4(2r)^2.$$



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 28 of 40

Of course, instead of (2.15) we can assume that

$$\beta_{i_1} + \beta_{i_2} = \beta_{i_3} + \beta_{i_4} = \beta_{i_5} + \beta_{i_6} = \beta_{i_7} + \beta_{i_8} = \frac{\pi}{2}.$$

Here $i_j \in \{1, 2, \dots, 8\}$.

Theorem 2.14. Let \mathcal{A} be a k -chordal n -gon, where $n = 4k$, and let

$$\beta_{i_1} + \beta_{i_2} = \dots = \beta_{i_{n-1}} + \beta_{i_n} = \frac{\pi}{2}.$$

Then $\text{Char}(\mathcal{A}) = 2$.

Proof. Since $\frac{4k}{2} = 2k$, we have

$$\sum_{j=1}^n \cos^2 \beta_{i_j} = 2k,$$

and this proves Theorem 2.14. □


Corollary 2.15. We have

$$\sum_{j=1}^n a_j^2 = 2k(2r)^2.$$

Theorem 2.16. Let \mathcal{A} be a chordal n -gon such that

$$(2.16) \quad \sum_{j=1}^{n-1} \beta_j = (n-2)\frac{\pi}{2}, \quad \beta_n = 0, \quad 0 < \beta_j < \frac{\pi}{2}, \quad j = \overline{1, n-1}.$$

Then $\text{Char}(\mathcal{A}) \leq 2$.



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 29 of 40

Proof. As it will be seen, this theorem is a corollary of Theorem 2.1. First we point out that (2.16) is obtained by putting $k = 1$, $\beta_n = 0$ into (2.1). Also, let us remark that in (2.1) we can take $\beta_n \approx 0$ as well. Therefore the proof of Theorem 2.16 is a straightforward consequence of Theorem 2.1 where, instead of (2.6), we write

$$\beta_1 = \dots = \beta_{2k} \approx \frac{\pi}{4k}, \quad \beta_{2k+1} = \dots = \beta_{n-1} \approx \frac{\pi}{2}; \quad \beta_n = 0,$$

or, because $k = 1$, we put

$$\beta_1 = \beta_2 \approx \frac{\pi}{4}, \quad \beta_3 = \dots = \beta_{n-1} \approx \frac{\pi}{2}, \quad \beta_n = 0.$$

For these specified values of β_1, \dots, β_n we obtain

$$\sum_{j=1}^n \cos^2 \beta_j \approx \cos^2 0 + 2 \cos^2 \frac{\pi}{4} = 2,$$


and

$$\sum_{j=1}^n \cos^m \beta_j \approx \cos^m 0 + 2 \cos^m \frac{\pi}{4} < 2$$

when $m > 2$. Theorem 2.16 is thus proved. □

Corollary 2.17. *Let the situation be the same as in Theorem 2.16. Then there is a m such that*

$$a_1^m + \dots + a_{n-1}^m = a_n^m, \quad 1 < m \leq 2.$$



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 30 of 40

Proof. The assertion immediately follows from $\sum_{j=1}^n a_j^m = 2(2r)^m$ because $a_n = 2r$. \square

Corollary 2.18. *Under conditions of Theorem 2.16, $\text{Char}(\mathcal{A}) = 2$ only if $n = 3$.*

Proof. Without loss of generality we can take $n = 5$ and consider the pentagon in Fig. 8. By Theorem 2.16, we have $\sum_{j=1}^5 \cos^2 \beta_j \leq 2$. But if $A_3 \rightarrow A_5$ and





Figure 8:

$A_4 \rightarrow A_5$, then

$$\beta_1 + \beta_2 \rightarrow \frac{\pi}{2}, \quad \beta_3, \beta_4 \rightarrow \frac{\pi}{2}, \quad \beta_5 = 0$$

and $\sum_{j=1}^5 \cos^2 \beta_j \rightarrow 2$. \square



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 31 of 40

Example 2.8. Specify $\beta_1 = 62^\circ, \beta_2 = 65^\circ, \beta_3 = 68^\circ, \beta_4 = 75^\circ$, then

$$\sum_{j=1}^5 \cos^{3/2} \beta_j = 1.957416 < 2,$$

$$\sum_{j=1}^5 \cos^{7/5} \beta_j = 2.050053 > 2.$$

When $\beta_1 = 44.1^\circ, \beta_2 = 46.9^\circ, \beta_3 = 89.4^\circ, \beta_4 = 89.6^\circ$, then

$$\sum_{j=1}^5 \cos^2 \beta_j = 1.9827268 < 2,$$

$$\sum_{j=1}^5 \cos^{19/10} \beta_j = 2.0183067 > 2.$$


Remark 2.4. As we can see, Theorem 2.16 may be considered as a generalization of the Pythagorean theorem. For example, all positive solutions of the equation

$$x_1^{3/2} + \dots + x_{n-1}^{3/2} = x_n^{3/2}$$

are related to chordal n -gons whose characteristic is $3/2$.

Thus, the problem "find all positive solutions of the above equation" is in fact the problem "find all angles β_1, \dots, β_n such that (2.16) is satisfied under the constraint

$$\sum_{j=1}^n \cos^{3/2} \beta_j = 2."$$



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 32 of 40

This problem is obvious when $n = 3$ since then $\beta_1 + \beta_2 = \frac{\pi}{2}$. But the case $n > 3$ could be very difficult.

Theorem 2.19. Let \mathcal{A} be a k -chordal n -gon. Then for every real $p > 1$, we have

$$(2.17) \quad \sum_{j=1}^n \cos^p \beta_j > n \left(\frac{2k}{n} \right)^p,$$

where β_1, \dots, β_n satisfy (2.1).

Proof. In [6, Theorem 2] it was proved that (2.17) holds for every positive integer p . Here we give an abbreviated and simplified proof of this result by which we deduce the assertion of Theorem 2.19.

By the Jordan-type inequality (1.6), i.e. by

$$\cos \beta_j > 1 - \frac{2}{\pi} \beta_j,$$


using the properties of the arithmetical mean, we can write

$$\sum_{j=1}^n \cos^p \beta_j \geq n \left(\frac{1}{n} \sum_{j=1}^n \cos \beta_j \right)^p > n \left(\frac{1}{n} \sum_{j=1}^n \left(1 - \frac{2}{\pi} \beta_j \right) \right)^p = n \left(\frac{2k}{n} \right)^p.$$

Indeed, here we have

$$\sum_{j=1}^n \left(1 - \frac{2}{\pi} \beta_j \right) = n - \frac{2}{\pi} \sum_{j=1}^n \beta_j = n - \frac{2}{\pi} (n - 2k) \frac{\pi}{2} = 2k.$$

This completes the proof of Theorem 2.19. □



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 33 of 40


Corollary 2.20. *Under the same assumptions as in Theorem 2.19, the following holds*

$$a_1^p + \cdots + a_n^p > n \left(\frac{4kr}{n} \right)^p.$$

For $p = 1$ one obtains an interesting relation:

$$a_1 + \cdots + a_n > 4kr.$$

For example if $n = 7$, $k = 3$, then $\sum_{j=1}^7 a_j > 12r$.



**Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)**

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 34 of 40

3. Inequalities Concerning k -Inscribed Polygons

In this section we start with the equality (1.4):

$$(1.4) \quad |\beta_1| + \cdots + |\beta_n| = [n - 2(m + \nu)] \frac{\pi}{2} + 2\tau$$

where $\tau = -(\beta_{j_1} + \cdots + \beta_{j_\nu})$ and $\beta_{j_1}, \dots, \beta_{j_\nu}$ are the negative angles, while $\text{Ind}(\mathcal{A}) = m$.

Let λ be defined by $2\tau = \lambda\pi$. Then (1.4) becomes

$$(3.1) \quad |\beta_1| + \cdots + |\beta_n| = [n - 2(m + \nu - \lambda)] \frac{\pi}{2}.$$

Using the inequality (1.6) we can write

$$\sum_{j=1}^n \cos \beta_j > \sum_{j=1}^n \left(1 - \frac{2}{\pi} |\beta_j|\right) = n - \frac{2}{\pi} \sum_{j=1}^n |\beta_j| = 2(m + \nu - \lambda).$$


So, by (3.1) it follows that

$$(3.2) \quad \sum_{j=1}^n \cos \beta_j > 2(m + \nu - \lambda).$$

In the case $\text{Ind}(\mathcal{A}) = k$, $\nu = 0$, we get the inequality (1.5).

It can be easily seen that

$$(3.3) \quad \nu - \lambda > 0.$$



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 35 of 40

Let us remark that

$$2\tau = 2 \sum_{j=1}^{\nu} |\beta_{i_j}| < 2 \left(\nu \cdot \frac{\pi}{2} \right),$$

from which it follows that $2\tau < \nu\pi$.

For the sake of brevity, we denote in the sequel

$$(3.4) \quad w = \text{Ind}(\mathcal{A}) + \nu - \lambda.$$

Now, we have the following theorem which is in fact a corollary of Theorem 2.1.

Theorem 3.1. *Let \mathcal{A} be a k -inscribed n -gon and let*


$$(3.5) \quad \sum_{j=1}^n |\beta_j| = (n - 2w) \frac{\pi}{2}, \quad 0 < |\beta_j| < \frac{\pi}{2}.$$

Then there is a q such that

$$(3.6) \quad \sum_{j=1}^n \cos^q \beta_j = 2w, \quad 1 < q \leq h(w),$$

where

$$(3.7) \quad h(w) = \frac{\log \frac{2w}{2w+1}}{\log \cos \frac{\pi}{4w+2}} \quad \text{if } w \text{ is an integer,}$$



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 36 of 40

but if w is not an integer, that is, when $\lambda = z + u$, where $z \geq 0$ is an integer and u is a positive number such that $0 < u < 1$, then

$$(3.8) \quad h(w) = \frac{\log \frac{\text{Ind}(\mathcal{A}) + \nu - \lambda}{\text{Ind}(\mathcal{A}) + \nu - z}}{\log \cos \frac{u\pi}{2(\text{Ind}(\mathcal{A}) + \nu - z)}}.$$

Proof. If w is an integer then the proof is quite analogous to the proof of Theorem 2.1.

In the case when w is not an integer, then instead of (2.6), we have the expressions

$$|\beta_1| = \cdots = |\beta_{2(\text{Ind}(\mathcal{A}) + \nu - z)}| \approx \frac{u\pi}{2(\text{Ind}(\mathcal{A}) + \nu - z)},$$

$$|\beta_{2(\text{Ind}(\mathcal{A}) + \nu - z) + 1}| = \cdots = |\beta_n| \approx \frac{\pi}{2}.$$

Let us remark that now the equality from (3.5) can be written as

$$\sum_{j=1}^n |\beta_j| = [n - 2(\text{Ind}(\mathcal{A}) + \nu - z)] \frac{\pi}{2} + u\pi$$

since


$$[n - 2(\text{Ind}(\mathcal{A}) + \nu - z - u)] \frac{\pi}{2} = [n - 2(\text{Ind}(\mathcal{A}) + \nu - z)] \frac{\pi}{2} + u\pi$$

Thus, from

$$2(\text{Ind}(\mathcal{A}) + \nu - z) \cos^{h(w)} \frac{u\pi}{2(\text{Ind}(\mathcal{A}) + \nu - z)} = 2w$$

we get (3.8).

Theorem 3.1 is thus proved. □



Certain Inequalities Concerning Some Kinds Of Chordal Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 37 of 40

Corollary 3.2. Let $a_i = |A_i A_{i+1}|$, $i = 1, \dots, n$. Then

$$\sum_{j=1}^n a_j^m = 2w(2r)^m, \quad 1 < m \lesssim h(w).$$

Example 3.1. In Fig. 9 we have drawn an 1-inscribed pentagon \mathcal{A} , with $r = 3$ and $\beta_1 = -6^\circ, \beta_2 = 56^\circ, \beta_3 = 28^\circ, \beta_4 = -58^\circ, \beta_5 = 70^\circ$. In this case we have





Figure 9:

$k = 1, \text{Ind}(\mathcal{A}) = 0, \nu = 2$ and

$$\sum_{j=1}^5 \beta_j = 90^\circ = \frac{\pi}{2} \text{ radian}, \quad \sum_{j=1}^5 |\beta_j| = 218^\circ = \left(\frac{\pi}{2} + 2 \cdot \frac{64}{90} \cdot \frac{\pi}{2} \right) \text{ radian}.$$



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 38 of 40

Thus we can write

$$\sum_{j=1}^5 |\beta_j| = (5 - 2\nu) \frac{\pi}{2} + 2 \cdot \frac{64}{90} \cdot \frac{\pi}{2} = (5 - 2(\nu - 0.71111111)) \frac{\pi}{2}.$$

Since $\nu = 2$, we have


$$w = \nu - 0.71111111 = 1.28888888, \quad 2w = 2.57777776$$

$$\lambda = 0.71111111, \quad z = 0, \quad u = \lambda \text{ since } \lambda < 1.$$

Finally, it can be written that

$$\sum_{j=1}^5 \cos^m \beta_j = 2.571645882 < 2w \quad \text{for } m = 1.85,$$

$$\sum_{j=1}^5 \cos^m \beta_j = 2.578128456 > 2w \quad \text{for } m = 1.84.$$



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back


Close

Quit

Page 39 of 40

References

- [1] D.S. MITRINOVIĆ, J.E. PEČARIĆ and V. VOLENEC, *Recent Advances in Geometric Inequalities*, Kluwer Acad. Publ., Dordrecht/Boston/London, 1989.
- [2] B.H. NEUMANN, Some remarks on polygons, *J. London. Math. Soc.*, **16** (1941), 230–245.
- [3] P. PECH, Inequality between sides and diagonals of a space n -gon and its integral analog, *Čas. Pro. Pěst. Mat.*, **115** (1990), 343–350.
- [4] P. PECH, Relations between inequalities for polygons, *Rad HAZU*, [472]13 (1997), 69–75.
- [5] M. RADIĆ, Some inequalities and properties concerning chordal polygons, *Math. Ineq. Appl.*, **2** (1998), 141–150.
- [6] M. RADIĆ, Some inequalities and properties concerning chordal semi-polygons, *Math. Ineq. Appl.*, **4**(2) (2001), 301–322.
- [7] M. RADIĆ, Some relations concerning k -chordal and k -tangential polygons, *Math. Commun.*, **7** (2002), 21–34.
- [8] J. SÁNDOR, Certain trigonometric inequalities, *Octagon M. M.*, **9**(1A) (2001), 331–336.



Certain Inequalities Concerning
Some Kinds Of Chordal
Polygons)

Mirko Radić

Title Page

Contents



Go Back

Close

Quit

Page 40 of 40