

# Journal of Inequalities in Pure and Applied Mathematics

## INEQUALITIES APPLICABLE TO CERTAIN PARTIAL DIFFERENTIAL EQUATIONS

B.G. PACHPATTE

57 Shri Niketan Colony  
Near Abhinay Talkies  
Aurangabad 431 001  
(Maharashtra) India.  
EMail: [bgpachpatte@hotmail.com](mailto:bgpachpatte@hotmail.com)

©2000 Victoria University  
ISSN (electronic): 1443-5756  
157-03



---

volume 5, issue 2, article 27,  
2004.

*Received 24 August, 2003;  
accepted 06 November, 2003.*

*Communicated by: J. Sándor*

---

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

## Abstract


In this paper explicit bounds on certain retarded integral inequalities involving functions of two independent variables are established. Some applications are also given to illustrate the usefulness of one of our results.

*2000 Mathematics Subject Classification:* 26D15, 26D20

*Key words:* Explicit bounds, retarded integral inequalities, two independent variables, non-self-adjoint, Hyperbolic partial differential equations, partial derivatives.

## Contents

1	Introduction .....	3
2	Statement of Results .....	4
3	Proofs of Theorems 2.1 – 2.4 .....	11
4	Some Applications .....	17
	References	



---

### Inequalities Applicable To Certain Partial Differential Equations

B.G. Pachpatte

---

Title Page

Contents



Go Back


Close

Quit

Page 2 of 27

# 1. Introduction

The integral inequalities which furnish explicit bounds on unknown functions has become a rich source of inspiration in the development of the theory of differential and integral equations. Over the years a great deal of attention has been given to such inequalities and their applications. A detailed account related to such inequalities can be found in [1] – [6] and the references given therein. However, in certain situations the bounds provided by such inequalities available in the literature are inadequate and we need bounds on some new integral inequalities in order to achieve a diversity of desired goals. In this paper, we offer some basic integral inequalities in two independent variables which can be used more conveniently in specific applications. Some applications are also given to study the behavior of solutions of non-self-adjoint hyperbolic partial differential equations with several retarded arguments.



---

**Inequalities Applicable To  
Certain Partial Differential  
Equations**

B.G. Pachpatte

---

Title Page

Contents



Go Back

Close

Quit

Page 3 of 27

## 2. Statement of Results

In what follows  $\mathbb{R}$  denotes the set of real numbers,  $\mathbb{R}_+ = [0, \infty)$ ,  $I_1 = [x_0, X)$ ,  $I_2 = [y_0, Y)$  are the subsets of  $\mathbb{R}$  and  $\Delta = I_1 \times I_2$ . The partial derivatives of a function  $z(x, y)$ ,  $x, y \in \mathbb{R}$  with respect to  $x, y$  and  $xy$  are denoted by  $D_1z(x, y)$ ,  $D_2z(x, y)$  and  $D_1D_2z(x, y)$  (or  $z_{xy}$ ) respectively.

Our main results are established in the following theorems.

**Theorem 2.1.** *Let  $u, a, b_i \in C(\Delta, \mathbb{R}_+)$  and  $\alpha_i \in C^1(I_1, I_1)$ ,  $\beta_i \in C^1(I_2, I_2)$  be nondecreasing with  $\alpha_i(x) \leq x$  on  $I_1$ ,  $\beta_i(y) \leq y$  on  $I_2$  for  $i = 1, \dots, n$  and  $k \geq 0$  be a constant.*

(A<sub>1</sub>) If

$$(2.1) \quad u(x, y) \leq k + \int_{x_0}^x a(s, y) u(s, y) ds + \sum_{i=1}^n \int_{\alpha_i(x_0)}^{\alpha_i(x)} \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(s, t) u(s, t) dt ds,$$


for  $x \in I_1, y \in I_2$ , then

$$(2.2) \quad u(x, y) \leq kq(x, y) \exp \left( \sum_{i=1}^n \int_{\alpha_i(x_0)}^{\alpha_i(x)} \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(s, t) q(s, t) dt ds \right),$$

for  $x \in I_1, y \in I_2$ , where

$$(2.3) \quad q(x, y) = \exp \left( \int_{x_0}^x a(\xi, y) d\xi \right),$$

for  $x \in I_1, y \in I_2$ .



Inequalities Applicable To  
Certain Partial Differential  
Equations

B.G. Pachpatte

Title Page

Contents




Go Back

Close

Quit

Page 4 of 27



**Inequalities Applicable To  
Certain Partial Differential  
Equations**

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 5 of 27

(A<sub>2</sub>) If

$$(2.4) \quad u(x, y) \leq k + \int_{y_0}^y a(x, t) u(x, t) dt + \sum_{i=1}^n \int_{\alpha_i(x_0)}^{\alpha_i(x)} \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(s, t) u(s, t) dt ds,$$

for  $x \in I_1, y \in I_2$ , then

$$(2.5) \quad u(x, y) \leq kr(x, y) \exp \left( \sum_{i=1}^n \int_{\alpha_i(x_0)}^{\alpha_i(x)} \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(s, t) r(s, t) dt ds \right),$$

for  $x \in I_1, y \in I_2$ , where


$$(2.6) \quad r(x, y) = \exp \left( \int_{y_0}^y a(x, \eta) d\eta \right),$$

for  $x \in I_1, y \in I_2$ .

**Theorem 2.2.** Let  $u, a, b_i, \alpha_i, \beta_i, k$  be as in Theorem 2.1. Let  $g \in C(\mathbb{R}_+, \mathbb{R}_+)$  be nondecreasing and submultiplicative function with  $g(u) > 0$  for  $u > 0$ .

(B<sub>1</sub>) If

$$(2.7) \quad u(x, y) \leq k + \int_{x_0}^x a(s, y) u(s, y) ds + \sum_{i=1}^n \int_{\alpha_i(x_0)}^{\alpha_i(x)} \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(s, t) g(u(s, t)) dt ds,$$



Title Page

Contents



Go Back

Close

Quit

Page 6 of 27

for  $x \in I_1, y \in I_2$ ; then for  $x_0 \leq x \leq x_1, y_0 \leq y \leq y_1; x, x_1 \in I_1, y, y_1 \in I_2$ ,

$$(2.8) \quad u(x, y) \leq q(x, y) \times G^{-1} \left[ G(k) + \sum_{i=1}^n \int_{\alpha_i(x_0)}^{\alpha_i(x)} \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(s, t) g(q(s, t)) dt ds \right],$$

where  $q(x, y)$  is given by (2.3) and  $G^{-1}$  is the inverse function of

$$(2.9) \quad G(r) = \int_{r_0}^r \frac{ds}{g(s)}, r > 0,$$


$r_0 > 0$  is arbitrary and  $x_1 \in I_1, y_1 \in I_2$  are chosen so that

$$G(k) + \sum_{i=1}^n \int_{\alpha_i(x_0)}^{\alpha_i(x)} \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(s, t) g(q(s, t)) dt ds \in \text{Dom}(G^{-1}),$$

for all  $x$  and  $y$  lying in  $[x_0, x_1]$  and  $[y_0, y_1]$  respectively.

(B<sub>2</sub>) If

$$(2.10) \quad u(x, y) \leq k + \int_{y_0}^y a(x, t) u(x, t) dt + \sum_{i=1}^n \int_{\alpha_i(x_0)}^{\alpha_i(x)} \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(s, t) g(u(s, t)) dt ds,$$



Title Page

Contents



Go Back

Close

Quit

Page 7 of 27

for  $x \in I_1, y \in I_2$ ; then for  $x_0 \leq x \leq x_2, y_0 \leq y \leq y_2; x, x_2 \in I_1, y, y_2 \in I_2$ ,

$$(2.11) \quad u(x, y) \leq r(x, y) \times G^{-1} \left[ G(k) + \sum_{i=1}^n \int_{\alpha_i(x_0)}^{\alpha_i(x)} \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(s, t) g(r(s, t)) dt ds \right],$$

where  $G, G^{-1}$  are as in part  $(B_1)$ ,  $r(x, y)$  is given by (2.6) and  $x_2 \in I_1, y_2 \in I_2$  are chosen so that

$$G(k) + \sum_{i=1}^n \int_{\alpha_i(x_0)}^{\alpha_i(x)} \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(s, t) g(r(s, t)) dt ds \in \text{Dom}(G^{-1}),$$


for all  $x$  and  $y$  lying in  $[x_0, x_2]$  and  $[y_0, y_2]$  respectively.

The inequalities in the following theorems can be used in the qualitative analysis of certain partial integrodifferential equations involving several retarded arguments.

**Theorem 2.3.** Let  $u, a, b_i, \alpha_i, \beta_i, k$  be as in Theorem 2.1.

$(C_1)$  If  $c \in C(\Delta, \mathbb{R}_+)$  and

$$(2.12) \quad u(x, y) \leq k + \int_{x_0}^x a(s, y) \left( u(s, y) + \int_{x_0}^s c(\sigma, y) u(\sigma, y) d\sigma \right) ds + \sum_{i=1}^n \int_{\alpha_i(x_0)}^{\alpha_i(x)} \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(s, t) u(s, t) dt ds,$$



**Inequalities Applicable To  
Certain Partial Differential  
Equations**

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 8 of 27

for  $x \in I_1, y \in I_2$ , then

$$(2.13) \quad u(x, y) \leq kp(x, y) \times \exp \left( \sum_{i=1}^n \int_{\alpha_i(x_0)}^{\alpha_i(x)} \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(s, t) p(s, t) dt ds \right),$$

for  $x \in I_1, y \in I_2$ , where

$$(2.14) \quad p(x, y) = 1 + \int_{x_0}^x a(\xi, y) \exp \left( \int_{x_0}^{\xi} [a(\sigma, y) + c(\sigma, y)] d\sigma \right) d\xi,$$

for  $x \in I_1, y \in I_2$ .


(C<sub>2</sub>) If  $c \in C(\Delta, \mathbb{R}_+)$  and

$$(2.15) \quad u(x, y) \leq k + \int_{y_0}^y a(x, t) \left( u(x, t) + \int_{y_0}^t c(x, \tau) u(x, \tau) d\tau \right) dt + \sum_{i=1}^n \int_{\alpha_i(x_0)}^{\alpha_i(x)} \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(s, t) u(s, t) dt ds,$$

for  $x \in I_1, y \in I_2$ , then

$$(2.16) \quad u(x, y) \leq kw(x, y) \times \exp \left( \sum_{i=1}^n \int_{\alpha_i(x_0)}^{\alpha_i(x)} \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(s, t) w(s, t) dt ds \right),$$





for  $x \in I_1, y \in I_2$ , where

$$(2.17) \quad w(x, y) = 1 + \int_{y_0}^y a(x, \eta) \exp \left( \int_{y_0}^{\eta} [a(x, \tau) + c(x, \tau)] d\tau \right) d\eta,$$

for  $x \in I_1, y \in I_2$ .

**Theorem 2.4.** Let  $u, a, b_i, \alpha_i, \beta_i, k$  be as in Theorem 2.1 and  $g$  be as in Theorem 2.2.

(D<sub>1</sub>) If  $c \in C(\Delta, \mathbb{R}_+)$  and


$$(2.18) \quad u(x, y) \leq k + \int_{x_0}^x a(s, y) \left( u(s, y) + \int_{x_0}^s c(\sigma, y) u(\sigma, y) d\sigma \right) ds \\ + \sum_{i=1}^n \int_{\alpha_i(x_0)}^{\alpha_i(x)} \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(s, t) g(u(s, t)) dt ds,$$

for  $x \in I_1, y \in I_2$ ; then for  $x_0 \leq x \leq x_3, y_0 \leq y \leq y_3; x, x_3 \in I_1, y, y_3 \in I_2$ ,

$$(2.19) \quad u(x, y) \leq p(x, y) \\ \times G^{-1} \left[ G(k) + \sum_{i=1}^n \int_{\alpha_i(x_0)}^{\alpha_i(x)} \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(s, t) g(p(s, t)) dt ds \right],$$

where  $p(x, y)$  is given by (2.14),  $G, G^{-1}$  are as in part (B<sub>1</sub>) in Theorem 2.2 and  $x_3 \in I_1, y_3 \in I_2$  are chosen so that

$$G(k) + \sum_{i=1}^n \int_{\alpha_i(x_0)}^{\alpha_i(x)} \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(s, t) g(p(s, t)) dt ds \in \text{Dom}(G^{-1}),$$



**Inequalities Applicable To  
Certain Partial Differential  
Equations**

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 10 of 27

for all  $x$  and  $y$  lying in  $[x_0, x_3]$  and  $[y_0, y_3]$  respectively.

(D<sub>2</sub>) If  $c \in C(\Delta, \mathbb{R}_+)$  and

$$(2.20) \quad u(x, y) \leq k + \int_{y_0}^y a(x, t) \left( u(x, t) + \int_{y_0}^t c(x, \tau) u(x, \tau) d\tau \right) dt \\ + \sum_{i=1}^n \int_{\alpha_i(x_0)}^{\alpha_i(x)} \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(s, t) g(u(s, t)) dt ds,$$

for  $x \in I_1, y \in I_2$ ; then for  $x_0 \leq x \leq x_4, y_0 \leq y \leq y_4; x, x_4 \in I_1, y, y_4 \in I_2$ ,

$$(2.21) \quad u(x, y) \leq w(x, y) \\ \times G^{-1} \left[ G(k) + \sum_{i=1}^n \int_{\alpha_i(x_0)}^{\alpha_i(x)} \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(s, t) g(w(s, t)) dt ds \right],$$

where  $w(x, y)$  is given by (2.17),  $G, G^{-1}$  are as in part (B<sub>1</sub>) in Theorem 2.2 and  $x_4 \in I_1, y_4 \in I_2$  are chosen so that

$$G(k) + \sum_{i=1}^n \int_{\alpha_i(x_0)}^{\alpha_i(x)} \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(s, t) g(w(s, t)) dt ds \in \text{Dom}(G^{-1}),$$

for all  $x$  and  $y$  lying in  $[x_0, x_4]$  and  $[y_0, y_4]$  respectively.

### 3. Proofs of Theorems 2.1 – 2.4

We give the details of the proofs of  $(A_1)$ ,  $(B_1)$  and  $(C_1)$  only. The proofs of the remaining inequalities can be completed by closely looking at the proofs of the above mentioned inequalities with suitable modifications.

$(A_1)$  Define a function  $z(x, y)$  by

$$(3.1) \quad z(x, y) = k + \sum_{i=1}^n \int_{\alpha_i(x_0)}^{\alpha_i(x)} \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(s, t) u(s, t) dt ds.$$

Then (2.1) can be restated as


$$(3.2) \quad u(x, y) \leq z(x, y) + \int_{x_0}^x a(s, y) u(s, y) ds.$$

It is easy to observe that  $z(x, y)$  is a nonnegative, continuous and nondecreasing function for  $x \in I_1, y \in I_2$ . Treating  $y, y \in I_2$  fixed in (3.2) and using Lemma 2.1 in [4] (see also [3, Theorem 1.3.1]) to (3.2), we get

$$(3.3) \quad u(x, y) \leq q(x, y) z(x, y),$$

for  $x \in I_1, y \in I_2$ , where  $q(x, y)$  is defined by (2.3). From (3.1) and (3.3) we have

$$(3.4) \quad z(x, y) \leq k + \sum_{i=1}^n \int_{\alpha_i(x_0)}^{\alpha_i(x)} \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(s, t) q(s, t) z(s, t) dt ds.$$



---

Inequalities Applicable To  
Certain Partial Differential  
Equations

B.G. Pachpatte

---

Title Page

Contents



Go Back

Close

Quit

Page 11 of 27

Let  $k > 0$  and define a function  $v(x, y)$  by the right hand side of (3.4). Then it is easy to observe that

$$v(x, y) > 0, \quad v(x_0, y) = v(x, y_0) = k, \quad z(x, y) \leq v(x, y)$$

and


$$\begin{aligned} D_1 v(x, y) &= \sum_{i=1}^n \left( \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(\alpha_i(x), t) q(\alpha_i(x), t) z(\alpha_i(x), t) dt \right) \alpha_i'(x) \\ &\leq \sum_{i=1}^n \left( \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(\alpha_i(x), t) q(\alpha_i(x), t) v(\alpha_i(x), t) dt \right) \alpha_i'(x) \\ &\leq v(x, y) \sum_{i=1}^n \left( \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(\alpha_i(x), t) q(\alpha_i(x), t) dt \right) \alpha_i'(x) \end{aligned}$$

i.e.

$$(3.5) \quad \frac{D_1 v(x, y)}{v(x, y)} \leq \sum_{i=1}^n \left( \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(\alpha_i(x), t) q(\alpha_i(x), t) dt \right) \alpha_i'(x).$$

Keeping  $y$  fixed in (3.5), setting  $x = \sigma$  and integrating it with respect to  $\sigma$  from  $x_0$  to  $x$ ,  $x \in I_1$ , and making the change of variables we get

$$(3.6) \quad v(x, y) \leq k \exp \left( \sum_{i=1}^n \int_{\alpha_i(x_0)}^{\alpha_i(x)} \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(s, t) q(s, t) dt ds \right),$$



**Inequalities Applicable To  
Certain Partial Differential  
Equations**

B.G. Pachpatte

Title Page

Contents




Go Back

Close

Quit

Page 12 of 27




---

**Inequalities Applicable To  
Certain Partial Differential  
Equations**

B.G. Pachpatte

---

Title Page

Contents



Go Back

Close

Quit

Page 13 of 27

for  $x \in I_1, y \in I_2$ . Using (3.6) in  $z(x, y) \leq v(x, y)$  we get

$$(3.7) \quad z(x, y) \leq k \exp \left( \sum_{i=1}^n \int_{\alpha_i(x_0)}^{\alpha_i(x)} \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(s, t) q(s, t) dt ds \right).$$

Using (3.7) in (3.3) we get the required inequality in (2.5).

If  $k \geq 0$  we carry out the above procedure with  $k + \varepsilon$  instead of  $k$ , where  $\varepsilon > 0$  is an arbitrary small constant, and subsequently pass the limit  $\varepsilon \rightarrow 0$  to obtain (2.5).

(B<sub>1</sub>) Define a function  $z(x, y)$  by

$$(3.8) \quad z(x, y) = k + \sum_{i=1}^n \int_{\alpha_i(x_0)}^{\alpha_i(x)} \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(s, t) g(u(s, t)) dt ds.$$


Then (2.7) can be stated as

$$(3.9) \quad u(x, y) \leq z(x, y) + \int_{x_0}^x a(s, y) u(s, y) ds.$$

As in the proof of part (A<sub>1</sub>), using Lemma 2.1 in [4] to (3.9) we have

$$(3.10) \quad u(x, y) \leq q(x, y) z(x, y),$$

for  $x \in I_1, y \in I_2$ , where  $q(x, y)$  and  $z(x, y)$  are defined by (2.3) and



(3.8). From (3.8) and (3.10) and the hypotheses on  $g$  we have

$$\begin{aligned}
 (3.11) \quad z(x, y) &\leq k + \sum_{i=1}^n \int_{\alpha_i(x_0)}^{\alpha_i(x)} \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(s, t) g(q(s, t) z(s, t)) dt ds \\
 &\leq k + \sum_{i=1}^n \int_{\alpha_i(x_0)}^{\alpha_i(x)} \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(s, t) g(q(s, t)) g(z(s, t)) dt ds.
 \end{aligned}$$

Let  $k > 0$  and define a function  $v(x, y)$  by the right hand side of (3.11). Then, it is easy to observe that  $v(x, y) > 0$ ,  $v(x_0, y) = v(x, y_0) = k$ ,  $z(x, y) \leq v(x, y)$  and

$$\begin{aligned}
 (3.12) \quad D_1 v(x, y) &= \sum_{i=1}^n \left( \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(\alpha_i(x), t) g(q(\alpha_i(x), t)) g(z(\alpha_i(x), t)) dt \right) \alpha_i'(x) \\
 &\leq \sum_{i=1}^n \left( \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(\alpha_i(x), t) g(q(\alpha_i(x), t)) g(v(\alpha_i(x), t)) dt \right) \alpha_i'(x) \\
 &\leq g(v(x, y)) \sum_{i=1}^n \left( \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(\alpha_i(x), t) g(q(\alpha_i(x), t)) dt \right) \alpha_i'(x).
 \end{aligned}$$

From (2.9) and (3.12) we have

$$(3.13) \quad D_1 G(v(x, y)) = \frac{D_1 v(x, y)}{g(v(x, y))}$$

$$\leq \sum_{i=1}^n \left( \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(\alpha_i(x), t) g(q(\alpha_i(x), t)) dt \right) \alpha_i'(x).$$

Keeping  $y$  fixed in (3.13), setting  $x = \sigma$  and integrating it with respect to  $\sigma$  from  $x_0$  to  $x$ ,  $x \in I_1$  and making the change of variables we get

$$(3.14) \quad G(v(x, y)) \leq G(k) + \sum_{i=1}^n \int_{\alpha_i(x_0)}^{\alpha_i(x)} \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(s, t) g(q(s, t)) dt ds.$$


Since  $G^{-1}(v)$  is increasing, from (3.14) we have

$$(3.15) \quad v(x, y) \leq G^{-1} \left[ G(k) + \sum_{i=1}^n \int_{\alpha_i(x_0)}^{\alpha_i(x)} \int_{\beta_i(y_0)}^{\beta_i(y)} b_i(s, t) g(q(s, t)) dt ds \right].$$

Using (3.15) in  $z(x, y) \leq v(x, y)$  and then the bound on  $z(x, y)$  in (3.10) we get the required inequality in (2.8). The case  $k \geq 0$  can be completed as mentioned in the proof of  $(A_1)$ .

$(C_1)$  Define a function  $z(x, y)$  by (3.1). Then (2.12) can be stated as

$$(3.16) \quad u(x, y) \leq z(x, y) + \int_{x_0}^x a(s, y) \left( u(s, y) + \int_{x_0}^s c(\sigma, y) u(\sigma, y) d\sigma \right) ds.$$



**Inequalities Applicable To  
Certain Partial Differential  
Equations**

B.G. Pachpatte

Title Page

Contents



Go Back

Close


Quit

Page 15 of 27

Clearly,  $z(x, y)$  is nonnegative, continuous and nondecreasing function for  $x \in I_1, y \in I_2$ . Treating  $y, y \in I_2$  fixed in (3.16) and applying Theorem 1.7.4 given in [3, p. 39] to (3.16) yields

$$u(x, y) \leq p(x, y) z(x, y),$$

where  $p(x, y)$  and  $z(x, y)$  are defined by (2.14) and (3.1). Now by following the proof of  $(A_1)$  with suitable changes we get the desired inequality in (2.13).



---

**Inequalities Applicable To  
Certain Partial Differential  
Equations**

B.G. Pachpatte

---

Title Page

Contents



Go Back

Close

Quit

Page 16 of 27



## 4. Some Applications

In this section, we present applications of the inequality  $(A_1)$  given in Theorem 2.1 which display the importance of our results to the literature. Consider the following retarded non-self-adjoint hyperbolic partial differential equation

$$(4.1) \quad z_{xy}(x, y) = D_2(a(x, y)z(x, y)) \\ + f(x, y, z(x - h_1(x), y - g_1(y)), \\ \dots, z(x - h_n(x), y - g_n(y))),$$

with the given initial boundary conditions

$$(4.2) \quad z(x, y_0) = a_1(x), z(x_0, y) = a_2(y), a_1(x_0) = a_2(y_0) = 0,$$


where  $f \in C(\Delta \times \mathbb{R}^n, \mathbb{R})$ ,  $a_1 \in C^1(I_1, \mathbb{R})$ ,  $a_2 \in C^1(I_2, \mathbb{R})$ , and  $a \in C(\Delta, \mathbb{R})$  is differentiable with respect to  $\mathbf{y}$ ;  $h_i \in C(I_1, \mathbb{R}_+)$ ,  $g_i \in C(I_2, \mathbb{R}_+)$  are nonincreasing, and such that  $x - h_i(x) \geq 0$ ,  $x - h_i(x) \in C^1(I_1, I_1)$ ,  $y - g_i(y) \geq 0$ ,  $y - g_i(y) \in C^1(I_2, I_2)$ ,  $h'_i(x) < 1$ ,  $g'_i(y) < 1$ ,  $h_i(x_0) = g_i(y_0) = 0$  for  $i = 1, \dots, n$ ;  $x \in I_1, y \in I_2$  and

$$(4.3) \quad M_i = \max_{x \in I_1} \frac{1}{1 - h'_i(x)}, \quad N_i = \max_{y \in I_2} \frac{1}{1 - g'_i(y)}.$$

Our first result gives the bound on the solution of the problem (4.1) – (4.2).

**Theorem 4.1.** *Suppose that*

$$(4.4) \quad |f(x, y, u_1, \dots, u_n)| \leq \sum_{i=1}^n b_i(x, y) |u_i|,$$



Inequalities Applicable To  
Certain Partial Differential  
Equations

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 17 of 27

$$(4.5) \quad |e(x, y)| \leq k,$$

where  $b_i(x, y)$ ,  $k$  are as in Theorem 2.1 and

$$(4.6) \quad e(x, y) = a_1(x) + a_2(y) - \int_{x_0}^x a(s, y_0) a_1(s) ds.$$

If  $z(x, y)$  is any solution of (4.1) – (4.2), then

$$(4.7) \quad |z(x, y)| \leq k\bar{q}(x, y) \exp\left(\sum_{i=1}^n \int_{\phi_i(x_0)}^{\phi_i(x)} \int_{\psi_i(y_0)}^{\psi_i(y)} \bar{b}_i(\sigma, \tau) \bar{q}(\sigma, \tau) d\tau d\sigma\right),$$


for  $x \in I_1, y \in I_2$ , where  $\phi_i(x) = x - h_i(x)$ ,  $x \in I_1$ ,  $\psi_i(y) = y - g_i(y)$ ,  $y \in I_2$ ,  $\bar{b}_i(\sigma, \tau) = M_i N_i b_i(\sigma + h_i(s), \tau + g_i(t))$  for  $\sigma, s \in I_1, \tau, t \in I_2$  and

$$(4.8) \quad \bar{q}(x, y) = \exp\left(\int_{x_0}^x |a(\xi, y)| d\xi\right),$$

for  $x \in I_1, y \in I_2$ .

*Proof.* It is easy to see that, the solution  $z(x, y)$  of the problem (4.1) – (4.2) satisfies the equivalent integral equation

$$(4.9) \quad z(x, y) = e(x, y) + \int_{x_0}^x a(s, y) z(s, y) ds + \int_{x_0}^x \int_{y_0}^y f(s, t, z(s - h_1(s), t - g_1(t)), \dots, z(s - h_n(s), t - g_n(t))) dt ds,$$



## Inequalities Applicable To Certain Partial Differential Equations

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 18 of 27

where  $e(x, y)$  is given by (4.6). From (4.9), (4.4), (4.5), (4.3) and making the change of variables we have

$$\begin{aligned}
 (4.10) \quad |z(x, y)| &\leq k + \int_{x_0}^x |a(s, y)| |z(s, y)| ds \\
 &\quad + \int_{x_0}^x \int_{y_0}^y \sum_{i=1}^n b_i(s, t) |z(s - h_i(s), t - g_i(t))| dt ds \\
 &\leq k + \int_{x_0}^x |a(s, y)| |z(s, y)| ds \\
 &\quad + \sum_{i=1}^n \int_{\phi_i(x_0)}^{\phi_i(x)} \int_{\psi_i(y_0)}^{\psi_i(y)} \bar{b}_i(\sigma, \tau) |z(\sigma, \tau)| d\tau d\sigma.
 \end{aligned}$$


Now a suitable application of the inequality  $(A_1)$  given in Theorem 2.1 to (4.10) yields (4.7).  $\square$

The next theorem deals with the uniqueness of solutions of (4.1) – (4.2).

**Theorem 4.2.** . Suppose that the function  $f$  in (4.1) satisfies the condition

$$(4.11) \quad |f(x, y, u_1, \dots, u_n) - f(x, y, v_1, \dots, v_n)| \leq \sum_{i=1}^n b_i(x, y) |u_i - v_i|,$$

where  $b_i(x, y)$  are as in Theorem 2.1. Let  $M_i, N_i, \phi_i, \psi_i, \bar{b}_i$  be as in Theorem 4.1. Then the problem (4.1) – (4.2) has at most one solution on  $\Delta$ .



**Inequalities Applicable To  
Certain Partial Differential  
Equations**

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit


Page 19 of 27

*Proof.* Let  $u(x, y)$  and  $v(x, y)$  be two solutions of (4.1) – (4.2) on  $\Delta$ , then

$$(4.12) \quad u(x, y) - v(x, y) = \int_{x_0}^x a(s, y) \{u(s, y) - v(s, y)\} ds \\ + \int_{x_0}^x \int_{y_0}^y \{f(s, t, u(s - h_1(s), t - g_1(t)), \\ \dots, u(s - h_n(s), t - g_n(t))) \\ - f(s, t, v(s - h_1(s), t - g_1(t)), \\ \dots, v(s - h_n(s), t - g_n(t)))\} dt ds.$$

From (4.12), (4.11), making the change of variables and in view of (4.3) we have

$$(4.13) \quad |u(x, y) - v(x, y)| \\ \leq \int_{x_0}^x |a(s, y)| |u(s, y) - v(s, y)| ds \\ + \int_{x_0}^x \int_{y_0}^y \sum_{i=1}^n b_i(s, t) |u(s - h_i(s), t - g_i(t)) \\ - v(s - h_i(s), t - g_i(t))| dt ds \\ \leq \int_{x_0}^x |a(s, y)| |u(s, y) - v(s, y)| ds \\ + \sum_{i=1}^n \int_{\phi_i(x_0)}^{\phi_i(x)} \int_{\psi(y_0)}^{\psi(y)} \bar{b}_i(\sigma, \tau) |u(\sigma, \tau) - v(\sigma, \tau)| d\tau d\sigma.$$



**Inequalities Applicable To  
Certain Partial Differential  
Equations**

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 20 of 27

A suitable application of the inequality ( $A_1$ ) in Theorem 2.1 to (4.13) yields

$$|u(x, y) - v(x, y)| \leq 0.$$

Therefore  $u(x, y) = v(x, y)$  i.e. there is at most one solution of the problem (4.1) – (4.2).  $\square$

The following theorem shows the dependency of solutions of equation (4.1) on given initial boundary data.

**Theorem 4.3.** *Let  $u(x, y)$  and  $v(x, y)$  be the solutions of (4.1) with the given initial boundary data*

$$(4.14) \quad u(x, y_0) = c_1(x), u(x_0, y) = c_2(y), c_1(x_0) = c_2(y_0) = 0,$$


and

$$(4.15) \quad v(x, y_0) = d_1(x), v(x_0, y) = d_2(y), d_1(x_0) = d_2(y_0) = 0,$$

respectively, where  $c_1, d_1 \in C^1(I_1, \mathbb{R})$ ,  $c_2, d_2 \in C^1(I_2, \mathbb{R})$ . Suppose that the function  $f$  satisfies the condition (4.11) in Theorem 4.2. Let

$$(4.16) \quad e_1(x, y) = c_1(x) + c_2(y) - \int_{x_0}^x a(s, y_0) c_1(s) ds,$$

$$(4.17) \quad e_2(x, y) = d_1(x) + d_2(y) - \int_{x_0}^x a(s, y_0) d_1(s) ds,$$



Inequalities Applicable To  
Certain Partial Differential  
Equations

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 21 of 27



**Inequalities Applicable To  
Certain Partial Differential  
Equations**

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 22 of 27

for  $x \in I_1, y \in I_2$  and

$$(4.18) \quad |e_1(x, y) - e_2(x, y)| \leq k,$$

where  $k$  is as in Theorem 2.1. Let  $M_i, N_i, \phi_i, \psi_i, \bar{b}_i, \bar{q}$  be as in Theorem 4.1. Then

$$(4.19) \quad |u(x, y) - v(x, y)| \leq k\bar{q}(x, y) \exp \left( \sum_{i=1}^n \int_{\phi_i(x_0)}^{\phi_i(x)} \int_{\psi(y_0)}^{\psi(y)} \bar{b}_i(\sigma, \tau) \bar{q}(\sigma, \tau) d\tau d\sigma \right),$$

for  $x \in I_1, y \in I_2$ .

*Proof.* Since  $u(x, y)$  and  $v(x, y)$  are the solutions of (4.1) – (4.14) and (4.1) – (4.15) respectively, we have

$$(4.20) \quad u(x, y) - v(x, y) = e_1(x, y) - e_2(x, y) + \int_{x_0}^x a(s, y) \{u(s, y) - v(s, y)\} ds + \int_{x_0}^x \int_{y_0}^y \{f(s, t, u(s - h_1(s), t - g_1(t)), \dots, u(s - h_n(s), t - g_n(t))) - f(s, t, v(s - h_1(s), t - g_1(t)), \dots, v(s - h_n(s), t - g_n(t)))\} dt ds,$$

for  $x \in I_1, y \in I_2$ . From (4.20), (4.18), (4.11), making the change of variables

and in view of (4.3) we have

$$(4.21) \quad |u(x, y) - v(x, y)| \leq k + \int_{x_0}^x |a(s, y)| |u(s, y) - v(s, y)| ds \\ + \sum_{i=1}^n \int_{\phi_i(x_0)}^{\phi_i(x)} \int_{\psi(y_0)}^{\psi(y)} \bar{b}_i(\sigma, \tau) |u(\sigma, \tau) - v(\sigma, \tau)| d\tau d\sigma,$$

for  $x \in I_1, y \in I_2$ . Now a suitable application of the inequality  $(A_1)$  in Theorem 2.1 to (4.21) yields the required estimate in (4.19), which shows the dependency of solutions of (4.1) on given initial boundary data.  $\square$


We next consider the following retarded non-self-adjoint hyperbolic partial differential equations

$$(4.22) \quad z_{xy}(x, y) = D_2(a(x, y)z(x, y)) \\ + f(x, y, z(x - h_1(x), y - g_1(y)), \\ \dots, z(x - h_n(x), y - g_n(y)), \mu),$$

$$(4.23) \quad z_{xy}(x, y) = D_2(a(x, y)z(x, y)) \\ + f(x, y, z(x - h_1(x), y - g_1(y)), \\ \dots, z(x - h_n(x), y - g_n(y)), \mu_0),$$

with the given initial boundary conditions (4.2), where  $f \in C(\Delta \times \mathbb{R}^n \times \mathbb{R}, \mathbb{R})$ ,  $h_i, g_i$  are as in (4.1) and  $\mu, \mu_0$  are real parameters.

The following theorem shows the dependency of solutions of problems (4.22) – (4.2) and (4.23) – (4.2) on parameters.



**Inequalities Applicable To  
Certain Partial Differential  
Equations**

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 23 of 27

**Theorem 4.4.** *Suppose that*

$$(4.24) \quad |f(x, y, u_1, \dots, u_n, \mu) - f(x, y, v_1, \dots, v_n, \mu)| \\ \leq \sum_{i=1}^n b_i(x, y) |u_i - v_i|,$$

$$(4.25) \quad |f(x, y, u_1, \dots, u_n, \mu) - f(x, y, u_1, \dots, u_n, \mu_0)| \leq m(x, y) |\mu - \mu_0|,$$


where  $b_i(x, y)$  are as in Theorem 2.1 and  $m : \Delta \rightarrow \mathbb{R}$  is a continuous function such that

$$(4.26) \quad \int_{x_0}^x \int_{y_0}^y m(s, t) dt ds \leq M,$$

where  $M \geq 0$  is a real constant. Let  $M_i, N_i, \phi_i, \psi_i, \bar{b}_i$  be as in Theorem 4.1. If  $z_1(x, y)$  and  $z_2(x, y)$  are the solutions of (4.22) – (4.2) and (4.23) =- (4.2), then

$$(4.27) \quad |z_1(x, y) - z_2(x, y)| \\ \leq \bar{k}\bar{q}(x, y) \exp\left(\sum_{i=1}^n \int_{\phi_i(x_0)}^{\phi_i(x)} \int_{\psi(y_0)}^{\psi(y)} \bar{b}_i(\sigma, \tau) \bar{q}(\sigma, \tau) d\tau d\sigma\right),$$

for  $x \in I_1, y \in I_2$ , where  $\bar{k} = |\mu - \mu_0| M$  and  $\bar{q}(x, y)$  is defined by (4.8).



**Inequalities Applicable To  
Certain Partial Differential  
Equations**

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 24 of 27




*Proof.* Let  $z(x, y) = z_1(x, y) - z_2(x, y)$  for  $x \in I_1, y \in I_2$ . As in the proof of Theorem 4.2, from the hypotheses we have

$$(4.28) \quad z(x, y) = \int_{x_0}^x a(s, y) z(s, y) ds + \int_{x_0}^x \int_{y_0}^y \{f(s, t, z_1(s - h_1(s), t - g_1(t)), \dots, z_1(s - h_n(s), t - g_n(t)), \mu) - f(s, t, z_2(s - h_1(s), t - g_1(t)), \dots, z_2(s - h_n(s), t - g_n(t)), \mu) + f(s, t, z_2(s - h_1(s), t - g_1(t)), \dots, z_2(s - h_n(s), t - g_n(t)), \mu) - f(s, t, z_2(s - h_1(s), t - g_1(t)), \dots, z_2(s - h_n(s), t - g_n(t)), \mu_0)\} dt ds.$$

From (4.28), (4.24) – (4.26), making the change of variables and in view of (4.3) we have

$$(4.29) \quad |z(x, y)| \leq \int_{x_0}^x |a(s, y)| |z(s, y)| ds + \int_{x_0}^x \int_{y_0}^y \sum_{i=1}^n b_i(s, t) |z_1(s - h_i(s), t - g_i(t)) - z_2(s - h_i(s), t - g_i(t))| dt ds + \int_{x_0}^x \int_{y_0}^y m(s, t) |\mu - \mu_0| dt ds \leq \bar{k} + \int_{x_0}^x |a(s, y)| |z(s, y)| ds$$



**Inequalities Applicable To  
Certain Partial Differential  
Equations**

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 25 of 27

$$+ \sum_{i=1}^n \int_{\phi_i(x_0)}^{\phi_i(x)} \int_{\psi_i(y_0)}^{\psi_i(y)} \bar{b}_i(\sigma, \tau) |z(\sigma, \tau)| d\tau d\sigma.$$

A suitable application of the inequality ( $A_1$ ) in Theorem 2.1 to (4.29) yields (4.27), which shows the dependency of solutions of problems (4.22) – (4.2) and (4.23) – (4.2) on parameters  $\mu$  and  $\mu_0$ .  $\square$

We note that the inequality given in Theorem 2.1 part ( $A_2$ ) can be used to study the similar properties as in Theorems 4.1 – 4.4 by replacing  $D_2(a(x, y)z(x, y))$  by  $D_1(a(x, y)z(x, y))$  in the equations (4.1), (4.22), (4.23) with the corresponding given initial-boundary conditions, under some suitable conditions on the functions involved therein. We also note that the inequalities given in Theorem 2.3 can be used to establish similar results as in Theorems 4.1 – 4.4 by replacing  $D_2(a(x, y)z(x, y))$  by


$$D_2 \left( Q_1 \left( x, y, z(x, y), \int_{x_0}^x k_1(\sigma, y, z(\sigma, y)) d\sigma \right) \right)$$

or

$$D_1 \left( Q_2 \left( x, y, z(x, y), \int_{y_0}^y k_2(x, \tau, z(x, \tau)) d\tau \right) \right)$$

in the equations (4.1), (4.22), (4.23) with the corresponding given initial-boundary conditions and under some suitable conditions on the functions involved therein.

Further it is to be noted that the inequalities and their applications given here can be extended very easily to functions involving many independent variables.



### Inequalities Applicable To Certain Partial Differential Equations

B.G. Pachpatte

Title Page

Contents



Go Back


Close

Quit

Page 26 of 27

## References

- [1] D. BAINOV AND P. SIMENOV, *Integral Inequalities and Applications*, Kluwer Academic Publishers, Dordrecht, 1992 .
- [2] O. LIPOVAN, A retarded Gronwall-like inequality and its applications, *J. Math. Anal. Appl.*, **252** (2000), 389–401.
- [3] B.G. PACHPATTE, *Inequalities for Differential and Integral Equations*, Academic Press, New York, 1998.
- [4] B.G. PACHPATTE, On some fundamental integral inequalities and their discrete analogues, *J. Inequal. Pure and Appl. Math.*, **2**(2) (2001), Art.15. [ONLINE <http://jipam.vu.edu.au>]
- [5] B.G. PACHPATTE, Explicit bounds on certain integral inequalities, *J. Math. Anal. Appl.*, **267** (2002), 48–61.
- [6] B.G. PACHPATTE, On some retarded integral inequalities and applications, *J. Inequal. Pure and Appl. Math.*, **3**(2) (2002), Art. 18. [ONLINE <http://jipam.vu.edu.au>]



---

Inequalities Applicable To  
Certain Partial Differential  
Equations

B.G. Pachpatte

---

Title Page

Contents



Go Back

Close

Quit

Page 27 of 27