

A GENERALIZED RESIDUAL INFORMATION MEASURE AND ITS PROPERTIES

M.A.K. BAIG AND JAVID GANI DAR

P. G. Department of Statistics

University of Kashmir

Srinagar - 190006 (INDIA)

Email: baigmak@yahoo.co.in javinfo.stat@yahoo.co.in

Received: 15 April, 2008


Accepted: 26 June, 200

Communicated by: N.S. Barnett

2000 AMS Sub. Class.: 60E15, 62N05, 90B25, 94A17, 94A24.

Key words: Shannon entropy, Renyi entropy, Residual entropy, Generalized residual entropy, Life time distributions.

Abstract: Ebrahim and Pellery [7] and Ebrahim [4] proposed the Shannon residual entropy function as a dynamic measure of uncertainty. In this paper we introduce and study a generalized information measure for residual lifetime distributions. It is shown that the proposed measure uniquely determines the distribution function. Also, characterization results for some lifetime distributions are discussed. Some discrete distribution results are also addressed.





Generalized Residual
Information Measure

M.A.K. Baig and Javid Gani Dar

vol. 10, iss. 3, art. 84, 2009

[Title Page](#)

[Contents](#)



Page 1 of 19

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

Contents

1	Introduction	3
2	Characterization of Distributions	6
2.1	Continuous Case	6
2.2	Discrete Case	11
3	A New Class of Life Time Distribution	14



**Generalized Residual
Information Measure**

M.A.K. Baig and Javid Gani Dar

vol. 10, iss. 3, art. 84, 2009

Title Page

Contents



Page 2 of 19


Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



Title Page

Contents



Page 3 of 19

Go Back

Full Screen

Close

1. Introduction

Let X be an absolutely continuous non-negative variable describing the random life-time of a component. Let $f(x)$ be the probability density function, $F(x)$ be the cumulative distribution and $R(x)$ be the survival function of the random variable X . A classical measure of uncertainty for X is the differential entropy, also known as the Shannon information measure, defined as

$$(1.1) \quad H(X) = - \int_0^{\infty} f(x) \log f(x) dx.$$

If X is a discrete random variable taking values x_1, x_2, \dots, x_n with respective probabilities p_1, p_2, \dots, p_n , then Shannon's entropy is defined as

$$(1.2) \quad H(P) = H(p_1, p_2, \dots, p_n) = - \sum_{k=1}^n p_k \log(p_k).$$

Renyi [11] generalized (1.1) and defined the measure

$$(1.3) \quad H_{\alpha}(X) = \frac{1}{\alpha(1-\alpha)} \log \int_0^{\infty} f^{\alpha}(x) dx, \quad \alpha > 1$$

and in the discrete case

$$(1.4) \quad H_{\alpha}(X) = \frac{1}{\alpha(1-\alpha)} \log \sum_{k=1}^n p_k^{\alpha}, \quad \alpha > 1.$$

Furthermore, in the continuous case

$$(1.5) \quad \lim_{\alpha \rightarrow 1} H_{\alpha}(X) = - \int_0^{\infty} f(x) \log f(x) dx = H(X)$$



Title Page

Contents



Page 4 of 19

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

© 2007 Victoria University. All rights reserved.

and in discrete case

$$(1.6) \quad \lim_{\alpha \rightarrow 1} H_{\alpha}(X) = - \sum_{k=1}^n p_k \log(p_k) = H(P),$$

which is Shannon's entropy in both cases.

The role of differential entropy as a measure of uncertainty in residual lifetime distributions has attracted increasing attention in recent years. As stated by Ebrahimi [4], the residual entropy at a time t of a random life time X is defined as the differential entropy of $(X/X > t)$. Formally, for all $t > 0$, the residual entropy of X is given by

$$(1.7) \quad H(X; t) = - \int_t^{\infty} \frac{f(x)}{R(t)} \log \frac{f(x)}{R(t)} dx$$

or

$$H(X; t) = 1 - \frac{1}{R(t)} \int_t^{\infty} f(x) \log h(x) dx,$$

where $h(t) = \frac{f(t)}{R(t)}$ is the hazard function or failure rate of the random variable X . Given that an item has survived up to t , $H(X; t)$ measures the uncertainty of the remaining lifetime of the component.

In the case of a discrete random variable, we have

$$(1.8) \quad H(t_j) = - \sum_{k=j}^n \frac{P(t_k)}{R(t_j)} \log \frac{P(t_k)}{R(t_j)},$$

where $R(t)$ is the reliability function of the random variable X .


Nair and Rajesh [9] studied the characterization of lifetime distributions by using the residual entropy function corresponding to the Shannon's entropy. In this sequel,

we investigate the problem of the characterization of a lifetime distribution using the following generalized residual entropy function:

$$(1.9) \quad H_\alpha(X; t) = \frac{1}{\alpha(1-\alpha)} \log \left(\frac{\int_t^\infty f^\alpha(x) dx}{R^\alpha(t)} \right), \quad \alpha > 1.$$

As $\alpha \rightarrow 1$, (1.9) reduces to (1.7).

The measure (1.9) is the residual life entropy corresponding to (1.3).



**Generalized Residual
Information Measure**

M.A.K. Baig and Javid Gani Dar

vol. 10, iss. 3, art. 84, 2009

Title Page

Contents



Page 5 of 19


Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



2. Characterization of Distributions

2.1. Continuous Case

Let X be a continuous non-negative random variable representing component failure time with failure distribution $F(t) = P(X \leq t)$ and survival function $R(t) = 1 - F(t)$ with $R(0) = 1$. We define the generalized entropy for residual life as

$$(2.1) \quad H_\alpha(X; t) = \frac{1}{\alpha(1 - \alpha)} \log \left(\frac{\int_t^\infty f^\alpha(x) dx}{R^\alpha(t)} \right), \quad \alpha > 1$$

and so

$$(2.2) \quad \int_t^\infty f^\alpha(x) dx = R^\alpha(t) \exp(\alpha(1 - \alpha)H_\alpha(X; t)), \quad \alpha > 1.$$

We now show that $H_\alpha(X; t)$ uniquely determines $R(t)$.

Theorem 2.1. *If X has an absolutely continuous distribution $F(t)$ with reliability function $R(t)$ and an increasing residual entropy $H_\alpha(X; t)$, then $H_\alpha(X; t)$ uniquely determines $R(t)$.*

Proof. Differentiating (2.2) with respect to t , we have

$$(2.3) \quad h^\alpha(t) = \alpha h(t) \exp(\alpha(1 - \alpha)H_\alpha(X; t)) - (\alpha)(1 - \alpha) \exp(\alpha(1 - \alpha)H_\alpha(X; t)) H'_\alpha(X; t),$$

where $h(t) = \frac{f(t)}{R(t)}$ is the failure rate function.

Hence for a fixed $t > 0$, $h(t)$ is a solution of

$$(2.4) \quad g(x) = (x)^\alpha - \alpha x \exp(\alpha(1 - \alpha)H_\alpha(X; t)) + \alpha(1 - \alpha) \exp(\alpha(1 - \alpha)H_\alpha(X; t)) H'_\alpha(X; t) = 0.$$

Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 6 of 19

Go Back

Full Screen

Close



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 7 of 19

[Go Back](#)

[Full Screen](#)

[Close](#)

Differentiating both sides with respect to x , we have

$$(2.5) \quad g'(x) = \alpha(x)^{\alpha-1} - \alpha \exp(\alpha(1-\alpha)H_\alpha(X;t)).$$

Now for $\alpha > 1$, $g(0) \leq 0$, $g(\infty) = \infty$, $g(x)$ first decreases and then increases with minimum at $x_t = \exp(-\alpha H_\alpha(X;t))$.

So, the unique solution to $g(x) = 0$ is given by $x = h(t)$. Thus $H_\alpha(X;t)$ determines $h(t)$ uniquely and hence determines $R(t)$ uniquely. \square

Theorem 2.2. *The uniform distribution over (a, b) , $a < b$ can be characterized by a decreasing generalized residual entropy $H_\alpha(X;t) = \frac{1}{\alpha} \log(b-t)$, $b > t$.*

Proof. For the case of uniform distribution over (a, b) , $a < b$, we have

$$(2.6) \quad H_\alpha(X;t) = \frac{1}{\alpha} \log(b-t), \quad b > t$$

which is decreasing in t .

Also, $x_t = \exp(-\alpha H_\alpha(X;t))$, therefore,

$$\begin{aligned} g(x_t) &= (x_t)^\alpha - \alpha x_t \exp(\alpha(1-\alpha)H_\alpha(X;t)) \\ &\quad + \alpha(1-\alpha) \exp(\alpha(1-\alpha)H_\alpha(X;t)) H'_\alpha(X;t) \\ &= 0. \end{aligned}$$

Hence $H_\alpha(X;t) = \frac{1}{\alpha} \log(b-t)$ is the unique solution to $g(x_t) = 0$, which proves the theorem. \square

Theorem 2.3. *Let X be a random variable having a generalized residual entropy of the form*

$$(2.7) \quad H_\alpha(X;t) = \frac{1}{\alpha(1-\alpha)} \log k - \frac{1}{\alpha} \log h(t),$$

where $h(t)$ is the failure rate function of X . Then X has



Title Page

Contents



Page 8 of 19

Go Back

Full Screen

Close

(i) an exponential distribution iff $k = \frac{1}{\alpha}$,

(ii) a Pareto distribution iff $k < \frac{1}{\alpha}$ and

(iii) a finite range distribution iff $k > \frac{1}{\alpha}$.

Proof. (i) Let X have the exponential distribution,

$$f(t) = \frac{1}{\theta} \exp\left(-\left(\frac{t}{\theta}\right)\right), \quad t > 0, \theta > 0.$$

The reliability function is given by

$$R(t) = \exp\left(-\frac{t}{\theta}\right)$$

and the failure rate function by

$$h(t) = \frac{1}{\theta}.$$

Therefore, after simplification, using (2.1),

$$(2.8) \quad H_{\alpha}(X; t) = \frac{1}{\alpha(1-\alpha)} \log k - \frac{1}{\alpha} \log h(t),$$

where $k = \frac{1}{\alpha}$ and $h(t) = \frac{1}{\theta}$.

Thus (2.7) holds.

Conversely, suppose that $k = \frac{1}{\alpha}$, then

$$\frac{1}{\alpha(1-\alpha)} \log k - \frac{1}{\alpha} \log h(t) = \frac{1}{\alpha(1-\alpha)} \log\left(\frac{\int_t^{\infty} f^{\alpha}(x) dx}{R^{\alpha}(t)}\right), \quad \alpha > 1$$



Title Page

Contents



Page 9 of 19

Go Back

Full Screen

Close

which gives,

$$(2.9) \quad h(t) = \left(\frac{1 - k\alpha}{k(\alpha - 1)}t + \frac{1}{h(0)} \right)^{-1} = (at + b)^{-1},$$

where $a = \left(\frac{1 - k\alpha}{k(\alpha - 1)} \right) = 0$, since $k = \frac{1}{\alpha}$ and $b = \frac{1}{h(0)}$.

Clearly (2.9) is the failure rate function of the exponential distribution.

(ii) The density function of the Pareto distribution is given by

$$f(t) = \frac{(b)^{\frac{1}{a}}}{(at + b)^{1 + \frac{1}{a}}}, \quad t \geq 0, a > 0, b > 0.$$

The reliability function is given by

$$R(t) = \frac{(b)^{\frac{1}{a}}}{(at + b)^{\frac{1}{a}}}, \quad t \geq 0, a > 0, b > 0$$

and failure rate is given by

$$(2.10) \quad h(t) = (at + b)^{-1}.$$

After simplification, (2.1) yields

$$(2.11) \quad H_{\alpha}(X; t) = \frac{1}{\alpha(1 - \alpha)} \log k - \frac{1}{\alpha} \log h(t),$$

where $k = \frac{1}{a\alpha + \alpha - a} < \frac{1}{\alpha}$, since $\alpha > 1$ and $h(t) = (at + b)^{-1}$.

Thus (2.7) holds.



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 10 of 19

Go Back

Full Screen

Close

Conversely, suppose that $k < \frac{1}{\alpha}$. Proceeding as in (i), (2.9) gives

$$(2.12) \quad h(t) = \left(\frac{1 - k\alpha}{k(\alpha - 1)}t + \frac{1}{h(0)} \right)^{-1} = (at + b)^{-1},$$

where $a = \left(\frac{1 - k\alpha}{k(\alpha - 1)} \right) > 0$, since $k < \frac{1}{\alpha}$, $\alpha > 1$ and $b = \frac{1}{h(0)}$.

Clearly, (2.12) is the failure rate function of the Pareto distribution given in (2.10).

(iii) The density function of the finite range distribution is given by

$$f(t) = \frac{\beta_1}{\nu} \left(1 - \frac{t}{\nu} \right)^{\beta_1 - 1}, \quad \beta_1 > 0, 0 \leq t \leq \nu < \infty.$$

The reliability function is given by

$$R(t) = \left(1 - \frac{t}{\nu} \right)^{\beta_1}, \quad \beta_1 > 0, 0 \leq t \leq \nu < \infty$$

and the failure rate function by

$$(2.13) \quad h(t) = \left(\frac{\beta_1}{\nu} \right) \left(1 - \frac{t}{\nu} \right)^{-1}.$$

It follows that


$$H_\alpha(X; t) = \frac{1}{\alpha(1 - \alpha)} \log k - \frac{1}{\alpha} \log h(t),$$

where $k = \frac{\beta_1}{\alpha\beta_1 - \alpha + 1} > \frac{1}{\alpha}$, since $\alpha > 1$ and $h(t) = \left(\frac{\beta_1}{\nu} \right) \left(1 - \frac{t}{\nu} \right)^{-1}$.

Thus (2.7) holds.

Conversely, suppose $k > \frac{1}{\alpha}$. Proceeding as in (i), (2.9) gives

$$(2.14) \quad h(t) = h(0) \left(1 - \frac{k\alpha - 1}{k(\alpha - 1)} h(0)t \right)^{-1},$$



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 11 of 19

Go Back

Full Screen

Close

which is the failure rate function of the distribution given by (2.13), if $k > \frac{1}{\alpha}$. \square

2.2. Discrete Case

Let X be a discrete random variable taking values x_1, x_2, \dots, x_n with respective probabilities p_1, p_2, \dots, p_n . The discrete residual entropy is defined as

$$(2.15) \quad H(p; j) = - \sum_{k=j}^n \frac{p_k}{R(j)} \log \left(\frac{p_k}{R(j)} \right).$$

The generalized residual entropy for the discrete case is defined as

$$(2.16) \quad H_\alpha(p; j) = \frac{1}{\alpha(1-\alpha)} \log \sum_{k=j}^n \left(\frac{p_k}{R(j)} \right)^\alpha.$$

For $\alpha \rightarrow 1$, (2.16) reduces to (2.15).

Theorem 2.4. *If X has a discrete distribution $F(t)$ with support $(t_j : t_j < t_{j+1})$ and an increasing generalized residual entropy $H_\alpha(X; t)$ then $H_\alpha(X; t)$ uniquely determines $F(t)$.*

Proof. We have

$$H_\alpha(p; j) = \frac{1}{\alpha(1-\alpha)} \log \sum_{k=j}^n \left(\frac{p_k}{R(j)} \right)^\alpha$$

and so

$$(2.17) \quad \sum_{k=j}^n p_k^\alpha = R^\alpha(j) \exp(\alpha(1-\alpha)H_\alpha(p; j)).$$



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 12 of 19

Go Back

Full Screen

Close

For $j + 1$, we have

$$(2.18) \quad \sum_{k=j+1}^n p_k^\alpha = R^\alpha(j+1) \exp(\alpha(1-\alpha)H_\alpha(p; j+1)).$$

Subtracting (2.18) from (2.17), using $p_j = R(j) - R(j+1)$ and $\lambda_j = \frac{R(j+1)}{R(j)}$, we have

$$\exp(\alpha(1-\alpha)H_\alpha(p; j)) = (1-\lambda_j)^\alpha + (\lambda_j)^\alpha \exp(\alpha(1-\alpha)H_\alpha(p; j+1)).$$

Hence, λ_j is a number in $(0, 1)$ which is a solution of

$$(2.19) \quad \phi(x) = \exp(\alpha(1-\alpha)H_\alpha(p; j)) - (1-x)^\alpha - (x)^\alpha \exp(\alpha(1-\alpha)H_\alpha(p; j+1)).$$

Differentiating both sides with respect to x , we have

$$(2.20) \quad \phi'(x) = \alpha(1-x)^{\alpha-1} - \alpha(x)^{\alpha-1} \exp(\alpha(1-\alpha)H_\alpha(p; j+1)).$$

Note that $\phi'(x) = 0$ gives

$$x = [1 + \exp(-\alpha H_\alpha(p; j+1))]^{-1} = x_j.$$

Now for $\alpha > 1$, $\phi(0) \leq 0$ and $\phi(1) \leq 0$, $\phi(x)$ first increases and then decreases in $(0, 1)$ with a maximum at $x_j = [1 + \exp(-\alpha H_\alpha(p; j+1))]^{-1}$.

So the unique solution to $\phi(x) = 0$ is given by $x = x_j$.

Thus $H_\alpha(X; t)$ uniquely determines $F(t)$. □

Theorem 2.5. *A discrete uniform distribution with support $(1, 2, \dots, n)$ is characterized by the decreasing generalized discrete residual entropy*

$$H_\alpha(p; j) = \frac{1}{\alpha} \log(n - j + 1), \quad j = 1, 2, \dots, n.$$



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 13 of 19

Go Back

Full Screen

Close

Proof. In the case of a discrete uniform distribution with support $(1, 2, \dots, n)$,

$$H_{\alpha}(p; j) = \frac{1}{\alpha} \log(n - j + 1), \quad j = 1, 2, \dots, n$$

which is decreasing in j .


Also,

$$x_j = [1 + \exp(-\alpha H_{\alpha}(p; j + 1))]^{-1}.$$

Therefore,

$$\begin{aligned} \phi(x_j) &= \exp(\alpha(1 - \alpha)H_{\alpha}(p; j)) - (1 - x_j)^{\alpha} - (x_j)^{\alpha} \exp(\alpha(1 - \alpha)H_{\alpha}(p; j + 1)) \\ &= 0 \end{aligned}$$

which proves the theorem. \square



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 14 of 19

[Go Back](#)

[Full Screen](#)

[Close](#)

3. A New Class of Life Time Distribution

Ebrahimi [4] defined two nonparametric classes of distribution based on the measure $H(X; t)$ as follows:

Definition 3.1. A random variable X is said to have decreasing (increasing) uncertainty in residual life DURL (IURL) if $H(X; t)$ is decreasing (increasing) in $t \geq 0$.

Definition 3.2. A non-negative random variable X is said to have decreasing (increasing) uncertainty in a generalized residual entropy of order α DUGRL(IUGRL) if $H_\alpha(X; t)$ is decreasing (increasing) in t , $t > 0$.

This implies that the random variable X has DUGRL(IUGRL),

$$H'_\alpha(X; t) \leq 0,$$

$$H'_\alpha(X; t) \geq 0.$$

Now we present a relationship between the new classes and the decreasing(increasing) failure rate class of lifetime distributions.

Remark 1. R is said to be an IFR(DFR) if $h(t)$ is increasing(decreasing) in t .

Theorem 3.3. If R has an increasing(decreasing) failure rate, IFR(DFR) then it is also a DUGRL(IUGRL).

Proof. We have,

$$(3.1) \quad H'_\alpha(X; t) = \frac{1}{1-\alpha} [h(t) - h^\alpha(t) \exp(-\alpha(1-\alpha)H_\alpha(X; t))].$$

Since R is IFR, by (3.1) and Remark 1, we have

$$H'_\alpha(X; t) \leq 0,$$



Title Page

Contents



Page 15 of 19

Go Back

Full Screen

Close

which means that $H_\alpha(X; t)$ is decreasing in t , i.e, R is DUGRL. The proof for IUGRL is similar. \square

Theorem 3.4. *If a distribution is DUGRL as well as IUGRL for some constant, then it must be exponential.*

Proof. Since the random variable X is both DUGRL and IUGRL, then,

$$H_\alpha(X; t) = \text{constant.}$$

Differentiating both sides with respect to t , we get

$$h(t) = \text{constant,}$$

which means that the distribution is exponential. \square

The following lemma which gives the value of the function $H_\alpha(X; t)$ under linear transformation will be used in proving the upcoming theorem.

Lemma 3.5. *For any absolutely continuous random variable X , define $Z = ax + b$, where $a > 0, b \geq 0$ are constants, then*

$$H_\alpha(Z; t) = \frac{\log a}{\alpha} + H_\alpha\left(X; \frac{t-b}{a}\right).$$

Proof. We have, $H_\alpha(X; t)$ from (2.1) and $Z = ax + b$, therefore,

$$H_\alpha(Z; t) = \frac{\log a}{\alpha} + H_\alpha\left(X; \frac{t-b}{a}\right),$$

which proves the lemma. \square



Title Page

Contents



Page 16 of 19

Go Back

Full Screen

Close

Theorem 3.6. Let X be an absolutely continuous random variable and $X \in DUGRL(IUGRL)$. Define $Z = aX + b$, where $a > 0$ and $b \geq 0$ are constants, then $Z \in DUGRL(IUGRL)$.

Proof. Since $X \in DUGRL(IUGRL)$, then,

$$H'_\alpha(X; t) \leq 0,$$

$$H'_\alpha(X; t) \geq 0.$$

By applying Lemma 3.5, it follows that $Z \in DUGRL(IUGRL)$, which proves the theorem. \square

The next theorem gives upper(lower) bounds for the failure rate function.

Theorem 3.7. If X is $DUGRL(IUGRL)$, then

$$h(t) \geq (\leq) (\alpha)^{\frac{1}{\alpha-1}} \exp(-\alpha H_\alpha(X; t)).$$

Proof. If X is $DUGRL$, then

$$H'_\alpha(X; t) \leq 0$$

which gives,

$$(3.2) \quad h(t) \geq (\alpha)^{\frac{1}{\alpha-1}} \exp(-\alpha H_\alpha(X; t)).$$

Similarly, if X is $IUGRL$, then


$$(3.3) \quad h(t) \leq (\alpha)^{\frac{1}{\alpha-1}} \exp(-\alpha H_\alpha(X; t)).$$

\square

Corollary 3.8. *Let $R(t)$ be a DUGRL(IUGRL), then*

$$R(t) \leq (\geq) \exp \left(- \int_0^t (\alpha)^{\frac{1}{\alpha-1}} \exp(-\alpha H_\alpha(X; u) du) \right)$$

for all $t \geq 0$.



**Generalized Residual
Information Measure**

M.A.K. Baig and Javid Gani Dar

vol. 10, iss. 3, art. 84, 2009

Title Page

Contents



Page 17 of 19

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

References

- [1] M. ASAID AND N. EBRAHIMI, Residual entropy and its characterizations in terms of hazard function and mean residual life time function, *Statist. Prob. Lett.*, **49** (2000), 263–269.
- [2] A.M. AWAD, A statistical information measure, *Dirasat XIV*, **12** (1987), 7–20.
- [3] A.D. CRESCENZO AND M. LONGOBARDI, entropy based measure of uncertainty in past life time distributions, *J. of Applied Probability*, **39** (2002), 434–440.
- [4] N. EBRAHIMI, How to measure uncertainty in the life time distributions, *Sankhya*, **58** (1996), 48–57.
- [5] N. EBRAHIMI, Testing whether life time distribution is decreasing uncertainty, *J. Statist. Plann. Infer.*, **64** (1997), 9–19.
- [6] N. EBRAHIMI AND S.KIRMANI, Some results on ordering of survival function through uncertainty, *Statist. Prob. Lett.*, **29** (1996), 167–176.
- [7] N. EBRAHIMI AND F. PELLERY, New partial ordering of survival function based on the notion of uncertainty, *Journal of Applied Probability*, **32** (1995), 202–211.
- [8] F. BELZUNCE, J. NAVARROR, J.M. RUIZ AND Y. AGUILA, Some results on residual entropy function, *Metrika*, **59** (2004), 147–161.
- [9] K.R.M. NAIR AND G. RAJESH, Characterization of the probability distributions using the residual entropy function, *J. Indian Statist. Assoc.*, **36** (1998), 157–166.





Generalized Residual
Information Measure

M.A.K. Baig and Javid Gani Dar
vol. 10, iss. 3, art. 84, 2009

Title Page

Contents



Page 18 of 19

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



- [10] R.D. GUPTA AND A.K. NANDA, α and β - entropies and relative entropies of distributions, *J. of Statistical Theory and Applications*, **3** (2002), 177–190.
- [11] A. RENYI, On measure of entropy and information, *Proc. of the Fourth Berkeley Symposium on Math. Statist. Prob.*, University of California Press, Berkely, 1 (1961), 547–561.
- [12] P.G. SANKARAN AND R.P GUPTA, Characterization of the life time distributions using measure of uncertainty, *Calcutta Statistical Association Bulletin*, **49** (1999), 154–166.
- [13] C.E. SHANNON, A mathematical theory of communication, *Bell System Technical J.*, **27** (1948), 379–423.

Title Page

Contents



Page 19 of 19

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756