


Journal of Inequalities in Pure and Applied Mathematics

A UNIFIED TREATMENT OF SOME SHARP INEQUALITIES

ZHENG LIU

Institute of Applied Mathematics, Faculty of Science
Anshan University of Science and Technology
Anshan 114044, Liaoning, China
EMail: lewzheng@163.net

©2000 Victoria University
ISSN (electronic): 1443-5756
195-06



volume 7, issue 5, article 172,
2006.

*Received 24 July, 2006;
accepted 11 December, 2006.*

Communicated by: S.S. Dragomir

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

Abstract

A generalization of some recent sharp inequalities by N. Ujević is established. Applications in numerical integration are also considered.

2000 Mathematics Subject Classification: Primary 26D15

Key words: Quadrature formula, Sharp error bounds, Generalization, Numerical integration.

Contents

1	Introduction	3
2	Main Results	6
3	Applications in Numerical Integration	13
	References	



A Unified Treatment of Some Sharp Inequalities

Zheng Liu

Title Page

Contents



Go Back

Close

Quit

Page 2 of 16

1. Introduction

In [1] we can find a generalization of the pre-Grüss inequality as:

Lemma 1.1. *Let $f, g, \Psi \in L_2(a, b)$. Then we have*

$$(1.1) \quad S_{\Psi}(f, g)^2 \leq S_{\Psi}(f, f)S_{\Psi}(g, g),$$

where

$$(1.2) \quad S_{\Psi}(f, g) = \int_a^b f(t)g(t) dt - \frac{1}{b-a} \int_a^b f(t) dt \int_a^b g(t) dt \\ - \frac{1}{\|\Psi\|_2^2} \int_a^b f(t)\Psi(t) dt \int_a^b g(t)\Psi(t) dt$$


and Ψ satisfies

$$(1.3) \quad \int_a^b \Psi(t) dt = 0,$$

while as usual, $\|\cdot\|_2$ is the norm in $L_2(a, b)$. i.e.,

$$\|\Psi\|_2^2 = \int_a^b \Psi^2(t) dt.$$

Using the above inequality, Ujević in [1] obtained the following interesting results:



A Unified Treatment of Some Sharp Inequalities

Zheng Liu

Title Page

Contents



Go Back

Close

Quit

Page 3 of 16

Theorem 1.2. Let $f : [a, b] \rightarrow \mathbb{R}$ be an absolutely continuous function whose derivative $f' \in L_2(a, b)$. Then

$$(1.4) \quad \left| f\left(\frac{a+b}{2}\right)(b-a) - \int_a^b f(t) dt \right| \leq \frac{(b-a)^{\frac{3}{2}}}{2\sqrt{3}} C_1$$

where

$$(1.5) \quad C_1 = \left\{ \|f'\|_2^2 - \frac{[f(b) - f(a)]^2}{b-a} - [Q(f; a, b)]^2 \right\}^{\frac{1}{2}}$$

and

$$(1.6) \quad Q(f; a, b) = \frac{2}{\sqrt{b-a}} \left[f(a) + f\left(\frac{a+b}{2}\right) + f(b) - \frac{3}{b-a} \int_a^b f(t) dt \right].$$

Theorem 1.3. Let the assumptions of Theorem 1.2 hold. Then


$$(1.7) \quad \left| \left(\frac{f(a) + f(b)}{2} \right) (b-a) - \int_a^b f(t) dt \right| \leq \frac{(b-a)^{\frac{3}{2}}}{2\sqrt{3}} C_2,$$

where

$$(1.8) \quad C_2 = \left\{ \|f'\|_2^2 - \frac{[f(b) - f(a)]^2}{b-a} - [P(f; a, b)]^2 \right\}^{\frac{1}{2}}$$

and

$$(1.9) \quad P(f; a, b) = \frac{1}{\sqrt{b-a}} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) - \frac{6}{b-a} \int_a^b f(t) dt \right].$$



A Unified Treatment of Some Sharp Inequalities

Zheng Liu

Title Page

Contents



Go Back

Close

Quit

Page 4 of 16

Theorem 1.4. Let the assumptions of Theorem 1.2 hold. Then

$$(1.10) \quad \left| \frac{f(a) + 2f\left(\frac{a+b}{2}\right) + f(b)}{4}(b-a) - \int_a^b f(t) dt \right| \leq \frac{(b-a)^{\frac{3}{2}}}{4\sqrt{3}} C_3,$$


where

$$(1.11) \quad C_3 = \left\{ \|f'\|_2^2 - \frac{[f(b) - f(a)]^2}{b-a} - \frac{1}{b-a} \left(f(a) - 2f\left(\frac{a+b}{2}\right) + f(b) \right)^2 \right\}^{\frac{1}{2}}$$

$$= \left\{ \|f'\|_2^2 - \frac{2}{b-a} \left[f\left(\frac{a+b}{2}\right) - f(a) \right]^2 - \frac{2}{b-a} \left[f(b) - f\left(\frac{a+b}{2}\right) \right]^2 \right\}^{\frac{1}{2}}.$$

In [2], Ujević further proved that the above all inequalities are sharp.

In this paper, we will derive a new sharp inequality with a parameter for absolutely continuous functions with derivatives belonging to $L_2(a, b)$, which not only provides a unified treatment of all the above sharp inequalities, but also gives some other interesting results as special cases. Applications in numerical integration are also considered.



A Unified Treatment of Some Sharp Inequalities

Zheng Liu

Title Page

Contents



Go Back

Close

Quit

Page 5 of 16

2. Main Results

Theorem 2.1. *Let the assumptions of Theorem 1.2 hold. Then for any $\theta \in [0, 1]$ we have*

$$(2.1) \quad \left| (b-a) \left[(1-\theta)f\left(\frac{a+b}{2}\right) + \theta \frac{f(a)+f(b)}{2} \right] - \int_a^b f(t) dt \right| \leq \frac{(b-a)^{\frac{3}{2}}}{2\sqrt{3}} \sqrt{1-3\theta+3\theta^2} C(\theta),$$

where

$$(2.2) \quad C(\theta) = \left\{ \|f'\|_2^2 - \frac{[f(b)-f(a)]^2}{b-a} - [N(f; a, b; \theta)]^2 \right\}^{\frac{1}{2}}$$


and

$$(2.3) \quad N(f; a, b; \theta) = \frac{2}{\sqrt{(1-3\theta+3\theta^2)(b-a)}} \times \left| (1-3\theta)f\left(\frac{a+b}{2}\right) + (2-3\theta)\frac{f(a)+f(b)}{2} - \frac{3-6\theta}{b-a} \int_a^b f(t) dt \right|.$$

The inequality (2.1) with (2.2) and (2.3) is sharp in the sense that the constant $\frac{1}{2\sqrt{3}}$ cannot be replaced by a smaller one.

Proof. Let us define the functions

$$p(t) = \begin{cases} t-a, & t \in [a, \frac{a+b}{2}], \\ t-b, & t \in (\frac{a+b}{2}, b], \end{cases}$$



A Unified Treatment of Some Sharp Inequalities

Zheng Liu

Title Page

Contents



Go Back

Close

Quit

Page 6 of 16

and

$$\Psi(t) = \begin{cases} t - (a + \theta \frac{b-a}{2}), & t \in [a, \frac{a+b}{2}], \\ t - (b - \theta \frac{b-a}{2}), & t \in (\frac{a+b}{2}, b], \end{cases}$$

where $\theta \in [0, 1]$.

It is not difficult to verify that

$$(2.4) \quad \int_a^b p(t) dt = \int_a^b \Psi(t) dt = 0.$$

i.e., Ψ satisfies the condition (1.3).

We also have


$$(2.5) \quad \|p\|_2^2 = \int_a^b p^2(t) dt = \frac{(b-a)^3}{12}$$

and

$$(2.6) \quad \|\Psi\|_2^2 = \int_a^b \Psi^2(t) dt = \frac{(b-a)^3}{12} (1 - 3\theta + 3\theta^2).$$

We now calculate

$$(2.7) \quad \begin{aligned} \int_a^b p(t)\Psi(t) dt &= \int_a^{\frac{a+b}{2}} (t-a) \left(t-a - \theta \frac{b-a}{2} \right) dt \\ &\quad + \int_{\frac{a+b}{2}}^b (t-b) \left(t-b + \theta \frac{b-a}{2} \right) dt \\ &= \left(\frac{1}{12} - \frac{\theta}{8} \right) (b-a)^3. \end{aligned}$$



A Unified Treatment of Some Sharp Inequalities

Zheng Liu

Title Page

Contents



Go Back

Close

Quit

Page 7 of 16

Integrating by parts, we have

$$(2.8) \quad \int_a^b f'(t)p(t) dt = f\left(\frac{a+b}{2}\right)(b-a) - \int_a^b f(t) dt$$

and

$$(2.9) \quad \int_a^b f'(t)\Psi(t) dt \\ = \int_a^{\frac{a+b}{2}} \left(t - a - \theta\frac{b-a}{2}\right) f'(t) dt + \int_{\frac{a+b}{2}}^b \left(t - b + \theta\frac{b-a}{2}\right) f'(t) dt \\ = (b-a) \left[(1-\theta)f\left(\frac{a+b}{2}\right) + \theta\frac{f(a)+f(b)}{2} \right] - \int_a^b f(t) dt.$$

From (2.4), (2.6) – (2.9) and (1.2) we get

$$(2.10) \quad S_{\Psi}(f', p) \\ = \int_a^b f'(t)p(t) dt - \frac{1}{b-a} \int_a^b f'(t) dt \int_a^b p(t) dt \\ - \frac{1}{\|\Psi\|_2^2} \int_a^b f'(t)\Psi(t) dt \int_a^b p(t)\Psi(t) dt \\ = f\left(\frac{a+b}{2}\right)(b-a) - \int_a^b f(t) dt - \frac{2-3\theta}{2(1-3\theta+3\theta^2)} \\ \times \left\{ (b-a) \left[(1-\theta)f\left(\frac{a+b}{2}\right) + \theta\frac{f(a)+f(b)}{2} \right] - \int_a^b f(t) dt \right\}$$



A Unified Treatment of Some Sharp Inequalities

Zheng Liu

Title Page

Contents



Go Back

Close

Quit

Page 8 of 16

$$= \frac{\theta}{2(1-3\theta+3\theta^2)} \left\{ (b-a) \left[(1-3\theta)f\left(\frac{a+b}{2}\right) + (2-3\theta)\frac{f(a)+f(b)}{2} \right] - (3-6\theta) \int_a^b f(t) dt \right\}.$$

From (2.4) – (2.7) and (1.2) we also have


$$(2.11) \quad S_{\Psi}(p, p) = \|p\|_2^2 - \frac{1}{b-a} \left(\int_a^b p(t) dt \right)^2 - \frac{1}{\|\Psi\|_2^2} \left(\int_a^b p(t)\Psi(t) dt \right)^2 \\ = \frac{\theta^2(b-a)^3}{16(1-3\theta+3\theta^2)}$$

and

$$(2.12) \quad S_{\Psi}(f', f') \\ = \|f'\|_2^2 - \frac{1}{b-a} \left(\int_a^b f'(t) dt \right)^2 - \frac{1}{\|\Psi\|_2^2} \left(\int_a^b f'(t)\Psi(t) dt \right)^2 \\ = \|f'\|_2^2 - \frac{[f(b)-f(a)]^2}{b-a} - \frac{12}{(1-3\theta+3\theta^2)(b-a)} \\ \times \left[(1-\theta)f\left(\frac{a+b}{2}\right) + \theta\frac{f(a)+f(b)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right]^2$$

Thus from (2.10) – (2.12) and (1.1) we can easily get

$$(2.13) \quad \left| (b-a) \left[(1-3\theta)f\left(\frac{a+b}{2}\right) + (2-3\theta)\frac{f(a)+f(b)}{2} \right] \right|$$



A Unified Treatment of Some Sharp Inequalities

Zheng Liu

Title Page

Contents



Go Back

Close

Quit

Page 9 of 16

$$\begin{aligned}
& -\left(3 - 6\theta\right) \left| \int_a^b f(t) dt \right|^2 \leq \frac{(1 - 3\theta + 3\theta^2)(b - a)^3}{4} \\
& \times \left\{ \|f'\|_2^2 - \frac{[f(b) - f(a)]^2}{b - a} - \frac{12}{(1 - 3\theta + 3\theta^2)(b - a)} \right. \\
& \left. \times \left[(1 - \theta)f\left(\frac{a + b}{2}\right) + \theta \frac{f(a) + f(b)}{2} - \frac{1}{b - a} \int_a^b f(t) dt \right]^2 \right\}.
\end{aligned}$$

It is equivalent to


$$\begin{aligned}
(2.14) \quad & 3(b - a)^2 \left[(1 - \theta)f\left(\frac{a + b}{2}\right) + \theta \frac{f(a) + f(b)}{2} - \frac{1}{b - a} \int_a^b f(t) dt \right]^2 \\
& \leq \frac{1 - 3\theta + 3\theta^2}{4} (b - a)^3 \left\{ \|f'\|_2^2 - \frac{[f(b) - f(a)]^2}{b - a} - \frac{4}{(1 - 3\theta + 3\theta^2)(b - a)} \right. \\
& \left. \times \left| (1 - 3\theta)f\left(\frac{a + b}{2}\right) + (2 - 3\theta) \frac{f(a) + f(b)}{2} - \frac{3 - 6\theta}{b - a} \int_a^b f(t) dt \right|^2 \right\}.
\end{aligned}$$

Consequently, inequality (2.1) with (2.2) and (2.3) follow from (2.14).

In order to prove that the inequality (2.1) with (2.2) and (2.3) is sharp for any $\theta \in [0, 1]$, we define the function

$$(2.15) \quad f(t) = \begin{cases} \frac{1}{2}t^2 - \frac{\theta}{2}t, & t \in [0, \frac{1}{2}], \\ \frac{1}{2}t^2 - (1 - \frac{\theta}{2})t + \frac{1 - \theta}{2}, & t \in (\frac{1}{2}, 1] \end{cases}$$

The function given in (2.15) is absolutely continuous since it is a continuous piecewise polynomial function.



A Unified Treatment of Some Sharp Inequalities

Zheng Liu

Title Page

Contents



Go Back

Close

Quit

Page 10 of 16

We now suppose that (2.1) holds with a constant $C > 0$ as

$$(2.16) \quad \left| (b-a) \left[(1-\theta)f\left(\frac{a+b}{2}\right) + \theta \frac{f(a)+f(b)}{2} \right] - \int_a^b f(t) dt \right| \leq C(b-a)^{\frac{3}{2}} \sqrt{1-3\theta+3\theta^2} C(\theta),$$

where $C(\theta)$ is as defined in (2.2) and (2.3).

Choosing $a = 0$, $b = 1$, and f defined in (2.15), we get

$$\int_0^1 f(t) dt = \frac{1}{24} - \frac{\theta}{8},$$

$$f(0) = f(1) = 0, \quad f\left(\frac{1}{2}\right) = \frac{1}{8} - \frac{\theta}{4},$$

$$\int_0^1 (f'(t))^2 dt = \frac{1-3\theta+3\theta^2}{12}$$

and


$$N(f; a, b; \theta) = 0$$

such that the left-hand side becomes

$$(2.17) \quad L.H.S.(2.16) = \frac{1-3\theta+3\theta^2}{12}.$$

We also find that the right-hand side is

$$(2.18) \quad R.H.S.(2.16) = \frac{C(1-3\theta+3\theta^2)}{2\sqrt{3}}.$$



A Unified Treatment of Some Sharp Inequalities

Zheng Liu

Title Page

Contents



Go Back

Close

Quit

Page 11 of 16

From (2.16) – (2.18), we find that $C \geq \frac{1}{2\sqrt{3}}$, proving that the constant $\frac{1}{2\sqrt{3}}$ is the best possible in (2.1). \square

Remark 1. If we take $\theta = 0$, $\theta = 1$ and $\theta = \frac{1}{2}$ in (2.1) with (2.2) and (2.3), we recapture the sharp midpoint type inequality (1.4) with (1.5) and (1.6), the sharp trapezoid type inequality (1.7) with (1.8) and (1.9) and the sharp averaged midpoint-trapezoid type inequality (1.10) with (1.11), respectively. Thus Theorem 2.1 may be regarded as a generalization of Theorem 1.2, Theorem 1.3 and Theorem 1.4.

Remark 2. If we take $\theta = \frac{1}{3}$, we get a sharp Simpson type inequality as

$$(2.19) \quad \left| \frac{b-a}{6} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] - \int_a^b f(t) dt \right| \leq \frac{(b-a)^{\frac{3}{2}}}{6} C_4,$$

where

$$(2.20) \quad C_4 = \left\{ \|f'\|_2^2 - \frac{[f(b) - f(a)]^2}{b-a} - [R(f; a, b)]^2 \right\}^{\frac{1}{2}}$$

and

$$(2.21) \quad R(f; a, b) = N \left(f; a, b; \frac{1}{3} \right) \\ = \frac{2\sqrt{3}}{\sqrt{b-a}} \left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(t) dt \right|.$$



A Unified Treatment of Some Sharp Inequalities

Zheng Liu

Title Page

Contents



Go Back

Close

Quit

Page 12 of 16

3. Applications in Numerical Integration

We restrict further considerations to the averaged midpoint-trapezoid quadrature rule. We also emphasize that similar considerations may be made for all the quadrature rules considered in the previous section.

Theorem 3.1. *Let $\pi = \{x_0 = a < x_1 < \dots < x_n = b\}$ be a given subdivision of the interval $[a, b]$ such that $h_i = x_{i+1} - x_i = h = \frac{b-a}{n}$ and let the assumptions of Theorem 1.4 hold. Then we have*


$$(3.1) \quad \left| \int_a^b f(t) dt - \frac{h}{4} \sum_{i=0}^{n-1} \left[f(x_i) + 2f\left(\frac{x_i + x_{i+1}}{2}\right) + f(x_{i+1}) \right] \right| \leq \frac{(b-a)^{\frac{3}{2}}}{4\sqrt{3}n} \delta_n(f) \leq \frac{(b-a)^{\frac{3}{2}}}{4\sqrt{3}n} \lambda_n(f).$$

where

$$(3.2) \quad \delta_n(f) = \left\{ \|f'\|_2^2 - \frac{[f(b) - f(a)]^2}{b-a} - \frac{1}{b-a} \left[f(x_0) + f(x_n) + 2 \sum_{i=1}^{n-1} f(x_i) - 2 \sum_{i=0}^{n-1} f\left(\frac{x_i + x_{i+1}}{2}\right) \right]^2 \right\}^{\frac{1}{2}}$$

and

$$(3.3) \quad \lambda_n(f) = \left\{ \|f'\|_2^2 - \frac{[f(b) - f(a)]^2}{b-a} \right\}^{\frac{1}{2}}.$$



A Unified Treatment of Some Sharp Inequalities

Zheng Liu

Title Page

Contents



Go Back

Close

Quit

Page 13 of 16

Proof. From (1.10) and (1.11) in Theorem 1.4 we obtain

$$(3.4) \quad \left| \frac{h}{4} \left[f(x_i) + 2f\left(\frac{x_i + x_{i+1}}{2}\right) + f(x_{i+1}) \right] - \int_{x_i}^{x_{i+1}} f(t) dt \right|$$

$$\leq \frac{h^{\frac{3}{2}}}{4\sqrt{3}} \left\{ \int_{x_i}^{x_{i+1}} (f'(t))^2 dt - \frac{1}{h} [f(x_{i+1}) - f(x_i)]^2 \right.$$

$$\left. - \frac{1}{h} \left[f(x_i) - 2f\left(\frac{x_i + x_{i+1}}{2}\right) + f(x_{i+1}) \right]^2 \right\}^{\frac{1}{2}}.$$

By summing (3.4) over i from 0 to $n - 1$ and using the generalized triangle inequality, we get


$$(3.5) \quad \left| \int_a^b f(t) dt - \frac{h}{4} \sum_{i=0}^{n-1} \left[f(x_i) + 2f\left(\frac{x_i + x_{i+1}}{2}\right) + f(x_{i+1}) \right] \right|$$

$$\leq \frac{h^{\frac{3}{2}}}{4\sqrt{3}} \sum_{i=0}^{n-1} \left\{ \int_{x_i}^{x_{i+1}} (f'(t))^2 dt - \frac{1}{h} [f(x_{i+1}) - f(x_i)]^2 \right.$$

$$\left. - \frac{1}{h} \left[f(x_i) - 2f\left(\frac{x_i + x_{i+1}}{2}\right) + f(x_{i+1}) \right]^2 \right\}^{\frac{1}{2}}.$$

By using the Cauchy inequality twice, we can easily obtain

$$(3.6) \quad \sum_{i=0}^{n-1} \left\{ \int_{x_i}^{x_{i+1}} (f'(t))^2 dt - \frac{1}{h} [f(x_{i+1}) - f(x_i)]^2 \right.$$



A Unified Treatment of Some Sharp Inequalities

Zheng Liu

Title Page

Contents




Go Back

Close

Quit

Page 14 of 16



A Unified Treatment of Some Sharp Inequalities

Zheng Liu

Title Page

Contents



Go Back

Close

Quit

Page 15 of 16


$$\begin{aligned}
 & -\frac{1}{h} \left[f(x_i) - 2f\left(\frac{x_i + x_{i+1}}{2}\right) + f(x_{i+1}) \right]^2 \Bigg\}^{\frac{1}{2}} \\
 \leq & \sqrt{n} \left\{ \|f'\|_2^2 - \frac{n}{b-a} \sum_{i=0}^{n-1} [f(x_{i+1}) - f(x_i)]^2 \right. \\
 & \left. - \frac{n}{b-a} \sum_{i=0}^{n-1} \left[f(x_i) - 2f\left(\frac{x_i + x_{i+1}}{2}\right) + f(x_{i+1}) \right]^2 \right\}^{\frac{1}{2}} \\
 \leq & \sqrt{n} \left\{ \|f'\|_2^2 - \frac{[f(b) - f(a)]^2}{b-a} \right. \\
 & \left. - \frac{1}{b-a} \left[f(x_0) + f(x_n) + 2 \sum_{i=1}^{n-1} f(x_i) - 2 \sum_{i=0}^{n-1} f\left(\frac{x_i + x_{i+1}}{2}\right) \right]^2 \right\}^{\frac{1}{2}} .
 \end{aligned}$$

Consequently, the inequality (3.1) with (3.2) and (3.3) follow from (3.5) and (3.6). \square

Remark 3. It should be noticed that Theorem 3.1 seems to be a revision and an improvement of the corresponding result in [2, Theorem 6.1].

References

- [1] N. UJEVIĆ, A generalization of the pre-Grüss inequality and applications to some quadrature formulae, *J. Inequal. Pure Appl. Math.*, **3**(1) (2002), Art. 13. [ONLINE: <http://jipam.vu.edu.au/article.php?sid=165>].
- [2] N. UJEVIĆ, Sharp error bounds for some quadrature formulae and applications, *J. Inequal. Pure Appl. Math.*, **7**(1) (2006), Art. 8. [ONLINE: <http://jipam.vu.edu.au/article.php?sid=622>].



A Unified Treatment of Some Sharp Inequalities

Zheng Liu

Title Page

Contents



Go Back

Close

Quit

Page 16 of 16