


CAUCHY'S MEANS OF LEVINSON TYPE





Cauchy's Means of
Levinson Type

M. Anwar and J. Pečarić

vol. 9, iss. 4, art. 120, 2008

[Title Page](#)

[Contents](#)



Page 1 of 16

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

© 2007 Victoria University. All rights reserved.

MATLOOB ANWAR

1- Abdus Salam School of Mathematical Sciences
GC University, Lahore,
Pakistan
EMail: matloob_t@yahoo.com

J. PEČARIĆ

University Of Zagreb
Faculty Of Textile Technology
Croatia
EMail: pecaric@mahazu.hazu.hr

Received: 14 July, 2008

Accepted: 12 October, 2008

Communicated by: [S. Abramovich](#)

2000 AMS Sub. Class.: Primary 26A51; Secondary 26A46, 26A48.

Key words: Convex function, Ky-Fan inequality, Weighted mean

Abstract: In this paper we introduce Levinson means of Cauchy's type. We show that these means are monotonic.

Acknowledgements: The research of the second author was supported by the Croatian Ministry of Science, Education and Sports under the Research Grants 117-1170889-0888.

Contents

1	Introduction and Preliminaries	3
2	Main Results	6
3	Related Results	12



**Cauchy's Means of
Levinson Type**

M. Anwar and J. Pečarić

vol. 9, iss. 4, art. 120, 2008

Title Page

Contents



Page 2 of 16


Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 3 of 16

[Go Back](#)

[Full Screen](#)

[Close](#)

1. Introduction and Preliminaries

Let x_1, x_2, \dots, x_n and p_1, p_2, \dots, p_n be real numbers such that $x_i \in [0, \frac{1}{2}]$, $p_i > 0$ with $P_n = \sum_{i=1}^n p_i$. Let G_n and A_n be the weighted geometric mean and arithmetic mean respectively defined by

$$G_n = \left(\prod_{i=1}^n x_i^{p_i} \right)^{\frac{1}{P_n}} \quad \text{and} \quad A_n = \frac{1}{P_n} \sum_{i=1}^n p_i x_i = \bar{x}.$$

In particular, consider the means

$$G'_n = \left(\prod_{i=1}^n (1 - x_i)^{p_i} \right)^{\frac{1}{P_n}} \quad \text{and} \quad A'_n = \frac{1}{P_n} \sum_{i=1}^n p_i (1 - x_i).$$

The well known Levinson inequality is the following ([1, 2] see also [6, p. 71]).

Theorem 1.1. *Let f be a real valued 3-convex function on $[0, 2a]$. Then for $0 < x_i < a$, $p_i > 0$ we have*

$$(1.1) \quad \frac{1}{P_n} \sum_{i=1}^n p_i f(x_i) - f\left(\frac{1}{P_n} \sum_{i=1}^n p_i x_i\right) \\ \leq \frac{1}{P_n} \sum_{i=1}^n p_i f(2a - x_i) - f\left(\frac{1}{P_n} \sum_{i=1}^n p_i (2a - x_i)\right).$$

In [4], the second author proved the following similar result.

Theorem 1.2. *Let f be a real valued 3-convex function on $[0, 2a]$ and x_i ($1 \leq i \leq n$) n points on $[0, a]$. Then*



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 4 of 16

Go Back

Full Screen

Close

$$(1.2) \quad \frac{1}{P_n} \sum_{i=1}^n p_i f(x_i) - f\left(\frac{1}{P_n} \sum_{i=1}^n p_i x_i\right) \\ \leq \frac{1}{P_n} \sum_{i=1}^n p_i f(a + x_i) - f\left(\frac{1}{P_n} \sum_{i=1}^n p_i (a + x_i)\right).$$

Lemma 1.3. *Let f be a log-convex function. If, $x_1 \leq y_1$, $x_2 \leq y_2$, $x_1 \neq x_2$, $y_1 \neq y_2$, then the following inequality is valid:*

$$(1.3) \quad \left[\frac{f(x_2)}{f(x_1)} \right]^{\frac{1}{x_2 - x_1}} \leq \left[\frac{f(y_2)}{f(y_1)} \right]^{\frac{1}{y_2 - y_1}}.$$

Lemma 1.4. *Let $f \in C^3(I)$ for some interval $I \subseteq \mathbb{R}$, such that f''' is bounded and $m = \min f'''$ and $M = \max f'''$. Consider the functions ϕ_1, ϕ_2 defined as,*

$$(1.4) \quad \phi_1(t) = \frac{M}{6} t^3 - f(t),$$

$$(1.5) \quad \phi_2(t) = f(t) - \frac{m}{6} t^3,$$

then ϕ_1 and ϕ_2 are 3-convex functions.

Lemma 1.5. *Let us define the function*

$$(1.6) \quad \varphi_s(x) = \begin{cases} \frac{x^s}{s(s-1)(s-2)}, & s \neq 0, 1, 2; \\ \frac{1}{2} \log x, & s=0; \\ -x \log x, & s=1; \\ \frac{1}{2} x^2 \log x, & s=2. \end{cases}$$

Then $\varphi_s'''(x) = x^{s-3}$, that is $\varphi_s(x)$ is 3-convex for $x > 0$.



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 5 of 16

Go Back

Full Screen

Close

Let $I \subseteq \mathbb{R}$ be an interval and let \mathcal{F} be some appropriately chosen vector space of real valued functions defined on I . Let Ψ be a functional on \mathcal{F} and let $A : \mathcal{F} \rightarrow \mathcal{R}$ be a linear operator, where \mathcal{R} is the vector of all real valued functions defined on I .

Suppose that for each $f \in \mathcal{F}$, there is a $\xi \in I$ such that

$$(1.7) \quad \Psi(f) = A(f)(\xi).$$

J. Pečarić, I. Perić and H. Srivastava in [5] proved the following important result for ϕ and A defined above.

Theorem 1.6. *For every $f, g \in \mathcal{F}$, there is a $\xi \in I$ such that*

$$(1.8) \quad A(g)(\xi)\Psi(f) = A(f)(\xi)\Psi(g).$$



Title Page

Contents

◀ ▶

◀ ▶

Page 6 of 16

Go Back

Full Screen

Close


2. Main Results

Theorem 2.1. Let $f \in C^3(I)$. Then for $x_i > 0$ and $p_i > 0$ there exist $\xi \in I$ such that the following equality holds true,

$$(2.1) \quad f\left(\frac{1}{P_n} \sum_{i=1}^n p_i x_i\right) - \frac{1}{P_n} \sum_{i=1}^n p_i f(x_i) \\ + \frac{1}{P_n} \sum_{i=1}^n p_i f(2a - x_i) - f\left(\frac{1}{P_n} \sum_{i=1}^n p_i (2a - x_i)\right) \\ = \frac{f'''(\xi)}{6} \left[\left(\frac{1}{P_n} \sum_{i=1}^n p_i x_i\right)^3 - \frac{1}{P_n} \sum_{i=1}^n p_i x_i^3 \right. \\ \left. + \frac{1}{P_n} \sum_{i=1}^n p_i (2a - x_i)^3 - \left(\frac{1}{P_n} \sum_{i=1}^n p_i (2a - x_i)\right)^3 \right].$$

Proof. Suppose that f''' is bounded, that is, $\min f''' = m$, $\max f''' = M$. By applying the Levinson inequality (1.1) to the functions ϕ_1 and ϕ_2 defined in Lemma 1.4, we get the following inequalities,

$$(2.2) \quad f\left(\frac{1}{P_n} \sum_{i=1}^n p_i x_i\right) - \frac{1}{P_n} \sum_{i=1}^n p_i f(x_i) \\ + \frac{1}{P_n} \sum_{i=1}^n p_i f(2a - x_i) - f\left(\frac{1}{P_n} \sum_{i=1}^n p_i (2a - x_i)\right)$$



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 7 of 16

[Go Back](#)

[Full Screen](#)

[Close](#)

$$\leq \frac{M}{6} \left[\left(\frac{1}{P_n} \sum_{i=1}^n p_i x_i \right)^3 - \frac{1}{P_n} \sum_{i=1}^n p_i x_i^3 \right. \\ \left. + \frac{1}{P_n} \sum_{i=1}^n p_i (2a - x_i)^3 - \left(\frac{1}{P_n} \sum_{i=1}^n p_i (2a - x_i) \right)^3 \right]$$


and

$$(2.3) \quad \frac{m}{6} \left[\left(\frac{1}{P_n} \sum_{i=1}^n p_i x_i \right)^3 - \frac{1}{P_n} \sum_{i=1}^n p_i x_i^3 \right. \\ \left. + \frac{1}{P_n} \sum_{i=1}^n p_i (2a - x_i)^3 - \left(\frac{1}{P_n} \sum_{i=1}^n p_i (2a - x_i) \right)^3 \right] \\ \leq f \left(\frac{1}{P_n} \sum_{i=1}^n p_i x_i \right) - \frac{1}{P_n} \sum_{i=1}^n p_i f(x_i) \\ + \frac{1}{P_n} \sum_{i=1}^n p_i f(2a - x_i) - f \left(\frac{1}{P_n} \sum_{i=1}^n p_i (2a - x_i) \right).$$

By combining both inequalities and using the fact that for $m \leq \rho \leq M$ there exist $\xi \in I$ such that $f'''(\xi) = \rho$, we get (2.1). Moreover, if f''' is (for example) bounded from above we have that (2.2) is valid and again (2.1) is valid.

Of course (2.1) is obvious if f''' is not bounded from above and below. \square

Theorem 2.2. Let $f, g \in C^3(I)$. Then for $x_i > 0$ and $p_i > 0$, $i = 1, \dots, n$ there



Title Page

Contents



Page 8 of 16

Go Back

Full Screen

Close

exist $\xi \in I$ such that the following equality holds true,

$$(2.4) \quad \frac{f'''(\xi)}{g'''(\xi)} = \frac{f\left(\frac{1}{P_n} \sum_{i=1}^n p_i x_i\right) - \frac{1}{P_n} \sum_{i=1}^n p_i f(x_i) + \frac{1}{P_n} \sum_{i=1}^n p_i f(2a - x_i) - f\left(\frac{1}{P_n} \sum_{i=1}^n p_i (2a - x_i)\right)}{g\left(\frac{1}{P_n} \sum_{i=1}^n p_i x_i\right) - \frac{1}{P_n} \sum_{i=1}^n p_i g(x_i) + \frac{1}{P_n} \sum_{i=1}^n p_i g(2a - x_i) - g\left(\frac{1}{P_n} \sum_{i=1}^n p_i (2a - x_i)\right)}.$$

Proof. Consider the linear functionals Ψ and A as in (1.7) for $\mathcal{F} = C^3(I)$ and \mathcal{R} the vector space of real valued functions such that $\Psi(k) = A(k)(\xi)$ for some function k . Let A be defined as:

$$(2.5) \quad A(f)(\xi) = \frac{f'''(\xi)}{6} \left[\left(\frac{1}{P_n} \sum_{i=1}^n p_i x_i \right)^3 - \frac{1}{P_n} \sum_{i=1}^n p_i x_i^3 + \frac{1}{P_n} \sum_{i=1}^n p_i (2a - x_i)^3 - \left(\frac{1}{P_n} \sum_{i=1}^n p_i (2a - x_i) \right)^3 \right].$$

Also, consider the linear combination $k = c_1 f - c_2 g$, where $f, g \in C^3(I)$ and c_1, c_2 are defined by

$$(2.6) \quad c_1 = \Psi(g) = g\left(\frac{1}{P_n} \sum_{i=1}^n p_i x_i\right) - \frac{1}{P_n} \sum_{i=1}^n p_i g(x_i) + \frac{1}{P_n} \sum_{i=1}^n p_i g(2a - x_i) - g\left(\frac{1}{P_n} \sum_{i=1}^n p_i (2a - x_i)\right),$$



[Title Page](#)

[Contents](#)

◀ ▶

◀ ▶

Page 9 of 16

[Go Back](#)

[Full Screen](#)

[Close](#)

$$(2.7) \quad c_2 = \Psi(f) = f \left(\frac{1}{P_n} \sum_{i=1}^n p_i x_i \right) - \frac{1}{P_n} \sum_{i=1}^n p_i f(x_i) \\ + \frac{1}{P_n} \sum_{i=1}^n p_i f(2a - x_i) - f \left(\frac{1}{P_n} \sum_{i=1}^n p_i (2a - x_i) \right).$$

Obviously, we have $\Psi(k) = 0$. This implies that (as in Theorem 1.6):

$$(2.8) \quad \Psi(g)A(f)(\xi) = \Psi(f)A(g)(\xi).$$

Now since $\Psi(g) \neq 0$ and $A(g)(\xi) \neq 0$ we have from the last equation

$$(2.9) \quad \frac{\Psi(f)}{\Psi(g)} = \frac{A(f)(\xi)}{A(g)(\xi)}.$$

After putting in the values we get (2.4). □

Corollary 2.3. Let $\frac{f'''}{g'''}$ be invertible then (2.4) suggests new means. That is,

$$(2.10) \quad \xi = \left(\frac{f'''}{g'''} \right)^{-1} \left[\frac{f \left(\frac{1}{P_n} \sum_{i=1}^n p_i x_i \right) - \frac{1}{P_n} \sum_{i=1}^n p_i f(x_i) + \frac{1}{P_n} \sum_{i=1}^n p_i f(2a - x_i) - f \left(\frac{1}{P_n} \sum_{i=1}^n p_i (2a - x_i) \right)}{g \left(\frac{1}{P_n} \sum_{i=1}^n p_i x_i \right) - \frac{1}{P_n} \sum_{i=1}^n p_i g(x_i) + \frac{1}{P_n} \sum_{i=1}^n p_i g(2a - x_i) - g \left(\frac{1}{P_n} \sum_{i=1}^n p_i (2a - x_i) \right)} \right]$$

is a new mean.

Definition 2.4. Define the function

$$(2.11) \quad \xi_s = \frac{1}{P_n} \sum_{i=1}^n p_i \left(\varphi_s(2a - x_i) - \varphi_s(x_i) \right) - \varphi_s(2a - \bar{x}) + \varphi_s(\bar{x}),$$

when $s \neq 0, 1, 2$. $s = 0, 1, 2$ are limiting cases defined by

$$\xi_0 = \frac{1}{2} \ln \left(\frac{G_n^a A_n}{G_n A_n^a} \right),$$



[Title Page](#)

[Contents](#)

◀◀ ▶▶

◀ ▶

Page 10 of 16

[Go Back](#)

[Full Screen](#)

[Close](#)

where

$$G_n^a = \left[\prod_{i=1}^n (2a - x_i)^{p_i} \right]^{\frac{1}{P_n}} \quad \text{and} \quad A_n^a = \frac{1}{P_n} \sum_{i=1}^n p_i (2a - x_i),$$

$$\xi_1 = \frac{1}{P_n} \sum_{i=1}^n p_i \left(x_i \ln x_i - (2a - x_i) \ln(2a - x_i) \right) + (2a - \bar{x}) \ln(2a - \bar{x}) - \bar{x} \ln \bar{x},$$

$$\xi_2 = \frac{1}{2} \left[\frac{1}{P_n} \sum_{i=1}^n p_i \left((2a - x_i)^2 \ln(2a - x_i) - x_i^2 \ln x_i \right) - (2a - \bar{x})^2 \ln(2a - \bar{x}) + \bar{x}^2 \ln \bar{x} \right].$$

Then we define the new means $M_{s,t}$ as:

Definition 2.5. Let us denote:

$$(2.12) \quad M_{s,t} = \left(\frac{\xi_s}{\xi_t} \right)^{\frac{1}{s-t}}$$

for $s \neq t \neq 0, 1, 2$. We define these limiting cases as

$$M_{s,s} = \exp \left[\frac{\eta}{\frac{1}{P_n} \sum_{i=1}^n p_i \left((2a - x_i)^s - x_i^s \right) - (2a - \bar{x})^s + \bar{x}^s} - \frac{3s^2 - 6s + 2}{s(s-1)(s-2)} \right],$$

where

$$\eta = \frac{1}{P_n} \sum_{i=1}^n p_i \left((2a - x_i)^s \log(2a - x_i) - x_i^s \log x_i \right) - (2a - \bar{x})^s \log(2a - \bar{x}) + \bar{x}^s \log \bar{x}$$

for $s \neq 0, 1, 2$

$$M_{0,0} = \exp \left[\frac{2 \left(\frac{1}{P_n} \sum_{i=1}^n p_i [(\log(2a - x_i))^2 - (\log x_i)^2] \right)}{4\xi_0} - \frac{(\log(2a - \bar{x}))^2 - (\log \bar{x})^2 6\xi_0}{4\xi_0} \right],$$

$$M_{1,1} = \exp \left[\frac{\frac{1}{P_n} \sum_{i=1}^n p_i [(2a - x_i)(\log(2a - x_i))^2 - x_i(\log x_i)^2]}{2\xi_1} - \frac{(2a - \bar{x})(\log(2a - \bar{x}))^2 - \bar{x}(\log \bar{x})^2}{2\xi_1} \right],$$


$$M_{2,2} = \exp \left[\frac{\frac{1}{P_n} \sum_{i=1}^n p_i [(2a - x_i)^2(\log(2a - x_i))^2 - x_i^2(\log x_i)^2]}{3\xi_2} - \frac{(2a - \bar{x})^2(\log(2a - \bar{x}))^2 - \bar{x}^2(\log \bar{x})^2}{3\xi_2} - 1 \right].$$

In our next result we prove that this new mean is monotonic.

Theorem 2.6. *Let $r \leq s, t \leq u, r \neq t, s \neq u$, then the following inequality is valid:*

$$(2.13) \quad M_{r,t} \leq M_{s,u}.$$

Proof. Since ξ_s is log convex as proved in [3, Theorem 2.2], then applying Lemma 1.4 for $r \leq s, t \leq u, r \neq t, s \neq u$ we get our required result. \square





Cauchy's Means of
Levinson Type

M. Anwar and J. Pečarić

vol. 9, iss. 4, art. 120, 2008

Title Page

Contents



Page 11 of 16


Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 12 of 16

Go Back

Full Screen

Close

3. Related Results

Theorem 3.1. Let $f \in C^3(I)$. For $x_i > 0$ and $p_i > 0$, $i = 1, \dots, n$ there exist $\xi \in I$ such that the following equality holds true,

$$\begin{aligned}
 (3.1) \quad & f\left(\frac{1}{P_n} \sum_{i=1}^n p_i x_i\right) - \frac{1}{P_n} \sum_{i=1}^n p_i f(x_i) \\
 & + \frac{1}{P_n} \sum_{i=1}^n p_i f(a + x_i) - f\left(\frac{1}{P_n} \sum_{i=1}^n p_i (a + x_i)\right) \\
 & = \frac{f'''(\xi)}{6} \left[\left(\frac{1}{P_n} \sum_{i=1}^n p_i x_i\right)^3 - \frac{1}{P_n} \sum_{i=1}^n p_i x_i^3 \right. \\
 & \quad \left. + \frac{1}{P_n} \sum_{i=1}^n p_i (a + x_i)^3 - \left(\frac{1}{P_n} \sum_{i=1}^n p_i (a + x_i)\right)^3 \right].
 \end{aligned}$$

Proof. Similar to proof of Theorem 2.1. □

Theorem 3.2. Let $f, g \in C^3(I)$. Then for $x_i > 0$ and $p_i > 0$, $i = 1, \dots, n$ there exist $\xi \in I$ such that the following equality holds true,

$$\begin{aligned}
 (3.2) \quad & \frac{f'''(\xi)}{g'''(\xi)} \\
 & = \frac{f\left(\frac{1}{P_n} \sum_{i=1}^n p_i x_i\right) - \frac{1}{P_n} \sum_{i=1}^n p_i f(x_i) + \frac{1}{P_n} \sum_{i=1}^n p_i f(a + x_i) - f\left(\frac{1}{P_n} \sum_{i=1}^n p_i (a + x_i)\right)}{g\left(\frac{1}{P_n} \sum_{i=1}^n p_i x_i\right) - \frac{1}{P_n} \sum_{i=1}^n p_i g(x_i) + \frac{1}{P_n} \sum_{i=1}^n p_i g(a + x_i) - g\left(\frac{1}{P_n} \sum_{i=1}^n p_i (a + x_i)\right)}
 \end{aligned}$$



[Title Page](#)

[Contents](#)

◀◀ ▶▶

◀ ▶

Page 13 of 16

[Go Back](#)

[Full Screen](#)

[Close](#)

Proof. Similar to proof of Theorem 2.2. □

Corollary 3.3. Let $\frac{f'''}{g'''}$ be invertible. Then (3.2) suggests new means. That is,

$$(3.3) \quad \xi = \left(\frac{f'''}{g'''} \right)^{-1} \left[\frac{f\left(\frac{1}{P_n} \sum_{i=1}^n p_i x_i\right) - \frac{1}{P_n} \sum_{i=1}^n p_i f(x_i) + \frac{1}{P_n} \sum_{i=1}^n p_i f(a+x_i) - f\left(\frac{1}{P_n} \sum_{i=1}^n p_i (a+x_i)\right)}{g\left(\frac{1}{P_n} \sum_{i=1}^n p_i x_i\right) - \frac{1}{P_n} \sum_{i=1}^n p_i g(x_i) + \frac{1}{P_n} \sum_{i=1}^n p_i g(a+x_i) - g\left(\frac{1}{P_n} \sum_{i=1}^n p_i (a+x_i)\right)} \right]$$

is a new mean.

Definition 3.4. Define the function

$$(3.4) \quad \bar{\xi}_s = \frac{1}{P_n} \sum_{i=1}^n p_i \left(\varphi_s(a+x_i) - \varphi_s(x_i) \right) - \varphi_s(a+\bar{x}) + \varphi_s(\bar{x}),$$

when $s \neq 0, 1, 2$. $s = 0, 1, 2$ are limiting cases defined by

$$\bar{\xi}_0 = \frac{1}{2} \ln \left(\frac{\bar{G}_n^a A_n}{G_n \bar{A}_n^a} \right),$$

where

$$\bar{G}_n^a = \left(\prod_{i=1}^n (a+x_i)^{p_i} \right)^{\frac{1}{P_n}}, \quad \text{and} \quad \bar{A}_n^a = \frac{1}{P_n} \sum_{i=1}^n p_i (a+x_i),$$

$$\bar{\xi}_1 = \frac{1}{P_n} \sum_{i=1}^n p_i \left(x_i \ln x_i - (a+x_i) \ln(2a-x_i) \right) + (a+\bar{x}) \ln(a+\bar{x}) - \bar{x} \ln \bar{x},$$

$$\bar{\xi}_2 = \frac{1}{2} \left[\frac{1}{P_n} \sum_{i=1}^n p_i \left((a+x_i)^2 \ln(a+x_i) - x_i^2 \ln x_i \right) - (a+\bar{x})^2 \ln(a+\bar{x}) + \bar{x}^2 \ln \bar{x} \right].$$



[Title Page](#)

[Contents](#)

◀◀ ▶▶

◀ ▶

Page 14 of 16

[Go Back](#)

[Full Screen](#)

[Close](#)

We now define new means $\overline{M}_{s,t}$ as:

Definition 3.5. Let us denote:

$$(3.5) \quad \overline{M}_{s,t} = \left(\frac{\bar{\xi}_s}{\bar{\xi}_t} \right)^{\frac{1}{s-t}}$$

for $s \neq t \neq 0, 1, 2$. We define these limiting cases as

$$\overline{M}_{s,s} = \exp \left(\frac{\bar{\eta}}{\frac{1}{P_n} \sum_{i=1}^n p_i ((a+x_i)^s - x_i^s) - (a+\bar{x})^s + \bar{x}^s} - \frac{3s^2 - 6s + 2}{s(s-1)(s-2)} \right),$$


where

$$\bar{\eta} = \frac{1}{P_n} \sum_{i=1}^n p_i ((a+x_i)^s \log(a+x_i) - x_i^s \log x_i) - (a+\bar{x})^s \log(a+\bar{x}) + \bar{x}^s \log \bar{x}$$

for $s \neq 0, 1, 2$

$$\overline{M}_{0,0} = \exp \left[\frac{2 \left(\frac{1}{P_n} \sum_{i=1}^n p_i [(\log(a+x_i))^2 - (\log x_i)^2] \right)}{4\bar{\xi}_0} - \frac{(\log(a+\bar{x}))^2 - (\log \bar{x})^2}{4\bar{\xi}_0} \right],$$

$$\overline{M}_{1,1} = \exp \left[\frac{\frac{1}{P_n} \sum_{i=1}^n p_i ((a+x_i)(\log(a+x_i))^2 - x_i(\log x_i)^2)}{2\bar{\xi}_1} - \frac{(a+\bar{x})(\log(a+\bar{x}))^2 - \bar{x}(\log \bar{x})^2}{2\bar{\xi}_1} \right],$$



Title Page

Contents



Page 15 of 16

Go Back

Full Screen

Close

$$\overline{M}_{2,2} = \exp \left[\frac{\frac{1}{P_n} \sum_{i=1}^n p_i ((a + x_i)^2 (\log(a + x_i))^2 - x_i^2 (\log x_i)^2)}{3\bar{\xi}_2} - \frac{(a + \bar{x})^2 (\log(a + \bar{x}))^2 - \bar{x}^2 (\log \bar{x})^2}{3\bar{\xi}_2} - 1 \right].$$

In our next result we prove that this new mean is monotonic.


Theorem 3.6. *Let $r \leq s, t \leq u, r \neq t, s \neq u$, then the following inequality is valid:*

$$(3.6) \quad \overline{M}_{r,t} \leq \overline{M}_{s,u}.$$

Proof. Since $\bar{\xi}_s$ is log convex as proved in [3, Theorem 2.5] ($\bar{\xi}_s = \rho_s$), then applying Lemma 1.3 for $r \leq s, t \leq u, r \neq t, s \neq u$ we get our required result. \square

References

- [1] S. LAWRENCE AND D. SEGALMAN, A generalization of two inequalities involving means, *Proc. Amer. Math. Soc.*, **35**(1) (1972), 96–100.
- [2] N. LEVINSON, Generalization of an inequality of Ky Fan, *J. Math. Anal. Appl.*, **8** (1964), 133–134.
- [3] M. ANWAR AND J.E. PEČARIĆ, On logarithmic convexity for Ky-Fan inequality, *J. Inequal. and Appl.*, (2008), Article ID 870950.
- [4] J.E. PEČARIĆ, An inequality for 3-convex functions, *J. Math. Anal. Appl.*, **19** (1982), 213–218.
- [5] J.E. PEČARIĆ, I. PERIĆ AND H. SRIVASTAVA A family of the Cauchy type mean-value theorems, *J. Math. Anal. Appl.*, **306** (2005), 730–739.
- [6] J.E. PEČARIĆ, F. PROSCHAN AND Y.C. TONG, *Convex Functions, Partial Orderings and Statistical Applications*, Academic Press, New York, 1992.



Cauchy's Means of
Levinson Type

M. Anwar and J. Pečarić

vol. 9, iss. 4, art. 120, 2008

Title Page

Contents



Page 16 of 16

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756