

Journal of Inequalities in Pure and Applied Mathematics


ON THE ABSOLUTE CONVERGENCE OF SMALL GAPS FOURIER SERIES OF FUNCTIONS OF $\varphi \wedge BV$

R. G. VYAS

Department of Mathematics
Faculty of Science
The Maharaja Sayajirao University of Baroda
Vadodara-390002, Gujarat, India.

EMail: drvgvyas@yahoo.com

©2000 Victoria University
ISSN (electronic): 1443-5756
205-05



volume 6, issue 4, article 94,
2005.

*Received 06 July, 2005;
accepted 29 July, 2005.*

Communicated by: L. Leindler

Abstract

Contents



Home Page

Go Back

Close

Quit



Abstract

Let f be a 2π periodic function in $L^1[0, 2\pi]$ and $\sum_{k=-\infty}^{\infty} \widehat{f}(n_k) e^{in_k x}$ be its Fourier series with 'small' gaps $n_{k+1} - n_k \geq q \geq 1$. Here we obtain a sufficiency condition for the convergence of the series $\sum_{k \in \mathbb{Z}} |\widehat{f}(n_k)|^\beta$ ($0 < \beta \leq 2$) if f is of $\varphi \wedge BV$ locally. We also obtain beautiful interconnections between the types of lacunarity in Fourier series and the localness of the hypothesis to be satisfied by the generic function allows us to interpolate results concerning lacunary Fourier series and non-lacunary Fourier series.

2000 Mathematics Subject Classification: 42A16, 42A28, 26A45.

Key words: Fourier series with small gaps, Absolute convergence of Fourier series and $\varphi \wedge$ -bounded variation.

Contents

1	Introduction	3
	References	

**On The Absolute Convergence
Of Small Gaps Fourier Series Of
Functions Of $\varphi \wedge BV$**

R. G. Vyas

Title Page

Contents



Go Back

Close

Quit

Page 2 of 11

1. Introduction

Let f be a 2π periodic function in $L^1[0, 2\pi]$ and $\widehat{f}(n)$, $n \in \mathbb{Z}$, be its Fourier coefficients. The series

$$(1.1) \quad \sum_{k \in \mathbb{Z}} \widehat{f}(n_k) e^{in_k x},$$

wherein $\{n_k\}_1^\infty$ is a strictly increasing sequence of natural numbers and $n_{-k} = -n_k$, for all k , satisfies an inequality

$$(1.2) \quad (n_{k+1} - n_k) \geq q \geq 1 \quad \text{for all } k = 0, 1, 2, \dots,$$

is called the Fourier series of f with ‘small’ gaps.


Obviously, if $n_k = k$, for all k , (i.e. $n_{k+1} - n_k = q = 1$, for all k), then we get non-lacunary Fourier series and if $\{n_k\}$ is such that

$$(1.3) \quad (n_{k+1} - n_k) \rightarrow \infty \quad \text{as } k \rightarrow \infty$$

then (1.1) is said to be the lacunary Fourier series.

In 1982 M. Schramm and D. Waterman [3] have introduced the class $\varphi \wedge BV(I)$ of functions of $\varphi \wedge$ -bounded variation over I and have studied sufficiency conditions for the absolute convergence of Fourier series of functions of $\wedge BV^{(p)}$ and $\varphi \wedge BV$.

Definition 1.1. Given a nonnegative convex function φ , defined on $[0, \infty)$ such that $\frac{\varphi(x)}{x} \rightarrow 0$ as $x \rightarrow 0$, for some constant $d \geq 2$, $\varphi(2x) \leq d\varphi(x)$ for all $x \geq 0$ and given a sequence of non-decreasing positive real numbers $\wedge = \{\lambda_m\}$



On The Absolute Convergence
Of Small Gaps Fourier Series Of
Functions Of $\varphi \wedge BV$

R. G. Vyas

Title Page

Contents



Go Back

Close

Quit

Page 3 of 11

($m = 1, 2, \dots$) such that $\sum_m \frac{1}{\lambda_m}$ diverges we say that $f \in \varphi \wedge BV$ (that is f is a function of $\varphi \wedge$ -bounded variation over (I)) if

$$V_{\Lambda_\varphi}(f, I) = \sup_{\{I_m\}} \{V_{\Lambda_\varphi}(\{I_m\}, f, I)\} < \infty,$$

where

$$V_{\Lambda_\varphi}(\{I_m\}, f, I) = \left(\sum_m \frac{\varphi |f(b_m) - f(a_m)|}{\lambda_m} \right),$$


and $\{I_m\}$ is a sequence of non-overlapping subintervals $I_m = [a_m, b_m] \subset I = [a, b]$.

Definition 1.2. For $p \geq 1$, the p -integral modulus of continuity $\omega^{(p)}(\delta, f, I)$ of f over I is defined as

$$\omega^{(p)}(\delta, f, I) = \sup_{0 \leq h \leq \delta} \|(T_h f - f)(x)\|_{p, I},$$

where $T_h f(x) = f(x + h)$ for all x and $\|(\cdot)\|_{p, I} = \|(\cdot)\chi_I\|_p$ in which χ_I is the characteristic function of I and $\|(\cdot)\|_p$ denotes the L^p -norm. $p = \infty$ gives the modulus of continuity $\omega(\delta, f, I)$.

By applying the Wiener-Ingham result [1, Vol. I, p. 222] for the finite trigonometric sums with ‘small’ gap (1.2) we have already studied the sufficiency conditions for the convergence of the series $\sum_{k \in \mathbb{Z}} \left| \widehat{f}(n_k) \right|^\beta$ ($0 < \beta \leq 2$) for the functions of $\wedge BV$ and $\wedge BV^{(p)}$ in terms of the modulus of continuity [6]. Here we obtain a sufficiency condition if function f is of $\varphi \wedge BV$. We prove the following theorem.



On The Absolute Convergence
Of Small Gaps Fourier Series Of
Functions Of $\varphi \wedge BV$

R. G. Vyas

Title Page

Contents



Go Back

Close

Quit

Page 4 of 11

Theorem 1.1. Let $f \in L[-\pi, \pi]$ possess a Fourier series with ‘small’ gaps (1.2) and I be a subinterval of length $\delta_1 > \frac{2\pi}{q}$. If $f \in \varphi \wedge BV(I)$, $1 \leq p < 2r$, $1 \leq r < \infty$, and

$$\sum_{k=1}^{\infty} \left(\left[\varphi^{-1} \left(\frac{\left(\omega^{((2-p)s+p)} \left(\frac{1}{n_k}, f, I \right) \right)^{2r-p}}{\sum_{j=1}^{n_k} \frac{1}{\lambda_j}} \right)^{\frac{1}{r}} \right] / k \right)^{\frac{\beta}{2}} < \infty,$$

where $\frac{1}{r} + \frac{1}{s} = 1$, then

$$(1.4) \quad \sum_{k \in \mathbb{Z}} \left| \widehat{f}(n_k) \right|^{\beta} < \infty \quad (0 < \beta \leq 2).$$


Theorem 1.1 with $\beta = 1$ is a ‘small’ gaps analogue of the Schramm and Waterman result [3, Theorem 2]. Observe that the interval I considered in the theorem for the gap condition (1.2) is of length $> \frac{2\pi}{q}$, so that when $n_k = k$, for all k , I is of length 2π . Hence for non-lacunary Fourier series (equality throughout in (1.2)) the theorem with $\beta = 1$ gives the Schramm and Waterman result [3, Theorem 2] as a particular case.

We need the following lemmas to prove the theorem.

Lemma 1.2 ([2, Lemma 2]). Let f and I be as in Theorem 1.1. If $f \in L^2(I)$ then

$$(1.5) \quad \sum_{k \in \mathbb{Z}} \left| \widehat{f}(n_k) \right|^2 \leq A_{\delta} |I|^{-1} \|f\|_{2,I}^2,$$

where A_{δ} depends only on δ .



On The Absolute Convergence
Of Small Gaps Fourier Series Of
Functions Of $\varphi \wedge BV$

R. G. Vyas

Title Page

Contents



Go Back

Close

Quit

Page 5 of 11

Lemma 1.3. If $|n_k| > p$ then for $t \in \mathbb{N}$ one has

$$\int_0^{\frac{\pi}{p}} \sin^{2t} |n_k| h dh \geq \frac{\pi}{2^{t+1} p}.$$

Proof. Obvious. □

Lemma 1.4 (Stechkin, refer to [5]). If $u_n \geq 0$ for $n \in \mathbb{N}$, $u_n \neq 0$ and a function $F(u)$ is concave, increasing, and $F(0) = 0$, then

$$\sum_1^{\infty} F(u_n) \leq 2 \sum_1^{\infty} F\left(\frac{u_n + u_{n+1} + \dots}{n}\right).$$

Proof of Theorem 1.1. Let $I = [x_0 - \frac{\delta_1}{2}, x_0 + \frac{\delta_1}{2}]$ for some x_0 and δ_2 be such that $0 < \frac{2\pi}{q} < \delta_2 < \delta_1$. Put $\delta_3 = \delta_1 - \delta_2$ and $J = [x_0 - \frac{\delta_2}{2}, x_0 + \frac{\delta_2}{2}]$. Suppose integers T and j satisfy


$$(1.6) \quad |n_T| > \frac{4\pi}{\delta_3} \quad \text{and} \quad 0 \leq j \leq \frac{\delta_3 |n_T|}{4\pi}.$$

$f \in \varphi \wedge BV(I)$ implies

$$|f(x)| \leq |f(a)| + |f(x) - f(a)| \leq |f(a)| + C\varphi^{-1}(V_{\wedge\varphi}(f, I)) \quad \text{for all } x \in I.$$

Since f is bounded over I , we have $f \in L^2(I)$, so that (1.5) holds and $f \in L^2[-\pi, \pi]$. If we put $f_j = (T_{2^j h} f - T_{2^{j-1} h} f)$ then $f_j \in L^2(I)$ and the Fourier series of f_j also possess gaps (1.2). Hence by Lemma 1.2 we get

$$(1.7) \quad \sum_{k \in \mathbb{Z}} \left| \hat{f}(n_k) \right|^2 \sin^2 \left(\frac{n_k h}{2} \right) = O \left(\|f_j\|_{2,J}^2 \right)$$



On The Absolute Convergence
Of Small Gaps Fourier Series Of
Functions Of $\varphi \wedge BV$

R. G. Vyas

Title Page

Contents



Go Back

Close

Quit

Page 6 of 11

because

$$\hat{f}_j(n_k) = 2i\hat{f}(n_k)e^{in_k(2j-\frac{1}{2}h)} \sin\left(\frac{n_k h}{2}\right).$$

Integrating both the sides of (1.7) over $(0, \frac{\pi}{n_T})$ with respect to h and using Lemma 1.3, we get

$$(1.8) \quad R_{n_T} = \sum_{|n_k| \geq n_T}^{\infty} \left| \hat{f}(n_k) \right|^2 = O(n_T) \int_0^{\frac{\pi}{n_T}} \left(\|f_j\|_{2,J}^2 \right) dh.$$


Since $2 = \frac{(2-p)s+p}{s} + \frac{p}{r}$, by using Hölder's inequality, we get from (1.8)

$$\begin{aligned} B &= \int_J |f_j(x)|^2 dx \\ &\leq \left(\int_J |f_j(x)|^{(2-p)s+p} dx \right)^{\frac{1}{s}} \left(\int_J |f_j(x)|^p dx \right)^{\frac{1}{r}} \\ &\leq \Omega_{h,J}^{1/r} \left(\int_J |f_j(x)|^p dx \right)^{\frac{1}{r}}, \end{aligned}$$

where $\Omega_{h,J} = (\omega^{(2-p)s+p}(h, f, J))^{2r-p}$. Thus

$$(1.9) \quad B^r \leq \Omega_{h,J} \int_J |f_j(x)|^p dx.$$

Since f is bounded over I , there exists some positive constant $M \geq \frac{1}{2}$ such that $|f(x)| \leq M$ for all $x \in I$. Dividing f by the positive constant M alters



On The Absolute Convergence
Of Small Gaps Fourier Series Of
Functions Of $\varphi \wedge BV$

R. G. Vyas

Title Page

Contents



Go Back

Close

Quit

Page 7 of 11

$\omega_p(h, f, J)$ by the same constant M and $\varphi(2|f(x)|) \leq d\varphi(|f(x)|)$ for all x , we may assume that $|f(x)| \leq 1$ for all $x \in I$. Hence from (1.9) we get


$$B^r \leq \Omega_{h,J} \int_J |f_j(x)| dx.$$

Since $\varphi(2x) \leq d\varphi(x)$, we have $\varphi(ax) \leq d^{\log_2 a} \varphi(x)$, so

$$\begin{aligned} \varphi\left(\frac{B^r}{\delta_2}\right) &\leq d^{\log_2 C\Omega_{h,J}} \varphi\left(\frac{\int_J |f_j(x)| dx}{\delta_2}\right) \\ &= C\Omega_{h,J}^{\log_2 d} \varphi\left(\frac{\int_J |f_j(x)| dx}{\delta_2}\right) \\ &= C\Omega_{h,J}^{\log_2 d-1} \Omega_{h,J} \varphi\left(\frac{\int_J |f_j(x)| dx}{\int_J 1 dx}\right) \\ &\leq C\Omega_{h,J} \left(\frac{\int_J \varphi(|f_j(x)|) dx}{\delta_2}\right) \quad (\text{by Jensen's inequality for integrals}) \\ &= C\Omega_{h,J} \left(\int_J \varphi(|f_j(x)|) dx\right). \end{aligned}$$

Multiplying both the sides of the equation by $\frac{1}{\lambda_j}$ and then taking the summation over $j = 1$ to n_T ($T \in \mathbb{N}$) we get

$$(1.10) \quad \varphi\left(\frac{B^r}{\delta_2}\right) \leq C \left(\frac{\Omega_{h,J}}{\sum_{j=1}^{n_T} \left(\frac{1}{\lambda_j}\right)}\right) \left(\int_J \left(\sum_{j=1}^{n_T} \frac{\varphi(|f_j(x)|)}{\lambda_j}\right) dx\right).$$



On The Absolute Convergence
Of Small Gaps Fourier Series Of
Functions Of $\varphi \wedge BV$

R. G. Vyas

Title Page

Contents



Go Back

Close

Quit

Page 8 of 11

Observe that for x in J , h in $(0, \frac{\pi}{n_T})$ and for each j of the summation the points $x + 2jh$ and $x + (2j - 1)h$ lie in I ; moreover $f \in \varphi \wedge BV(I)$ implies

$$\sum_{j=1}^{n_T} \frac{\varphi |f_j(x)|}{\lambda_j} = O(1).$$

Therefore, it follows from (1.10) that


$$B = O \left(\left[\varphi^{-1} \left(\frac{\Omega_{1/n_T, I}}{\sum_{j=1}^{n_T} \left(\frac{1}{\lambda_j} \right)} \right) \right]^{\frac{1}{r}} \right).$$

Substituting back the value of B in the equation (1.8), we get

$$R_{n_T} = \sum_{|n_k| \geq n_T}^{\infty} |\hat{f}(n_k)|^2 = O \left(\left[\varphi^{-1} \left(\frac{\Omega_{1/n_T, I}}{\sum_{j=1}^{n_T} \left(\frac{1}{\lambda_j} \right)} \right) \right]^{\frac{1}{r}} \right).$$

Thus

$$R_{n_T} = O \left(\left[\varphi^{-1} \left(\frac{\left(\omega^{(2-p)s+p} \left(\frac{1}{n_T}, f, I \right) \right)^{2r-p}}{\sum_{j=1}^{n_T} \left(\frac{1}{\lambda_j} \right)} \right) \right]^{\frac{1}{r}} \right).$$



On The Absolute Convergence
Of Small Gaps Fourier Series Of
Functions Of $\varphi \wedge BV$

R. G. Vyas

Title Page

Contents



Go Back

Close


Quit

Page 9 of 11

Finally, Lemma 1.4 with $u_k = \left| \hat{f}(n_k) \right|^2$ ($k \in \mathbb{Z}$) and $F(u) = u^{\beta/2}$ gives

$$\begin{aligned} \sum_{|k|=1}^{\infty} \left| \hat{f}(n_k) \right|^{\beta} &= 2 \sum_{k=1}^{\infty} F \left(\left| \hat{f}(n_k) \right|^2 \right) \\ &\leq 4 \sum_{k=1}^{\infty} F \left(\frac{R_{n_k}}{k} \right) \\ &= 4 \sum_{k=1}^{\infty} \left(\frac{R_{n_k}}{k} \right)^{\frac{\beta}{2}} \\ &= O(1) \left(\sum_{k=1}^{\infty} \left[\left[\varphi^{-1} \left(\frac{(\omega^{(2-p)s+p}(\frac{1}{n_k}, f, I))^{2r-p}}{\sum_{j=1}^{n_k} \frac{1}{\lambda_j}} \right) \right]^{\frac{1}{r}} / k \right]^{\frac{\beta}{2}} \right). \end{aligned}$$

This proves the theorem. \square



**On The Absolute Convergence
Of Small Gaps Fourier Series Of
Functions Of $\varphi \wedge BV$**

R. G. Vyas

Title Page

Contents



Go Back

Close

Quit

Page 10 of 11

References

- [1] A. ZYMUND, *Trigonometric Series*, 2nd ed., Cambridge Univ. Press, Cambridge, 1979 (reprint).
- [2] J.R. PATADIA AND R.G. VYAS, Fourier series with small gaps and functions of generalized variations, *J. Math. Anal. and Appl.*, **182**(1) (1994), 113–126.
- [3] M. SCHRAMM AND D. WATERMAN, Absolute convergence of Fourier series of functions of $\wedge BV^{(p)}$ and $\Phi \wedge BV$, *Acta. Math. Hungar.*, **40** (1982), 273–276.
- [4] N.K. BARRY, *A Treatise on Trigonometric Series*, Pergamon, New York, 1964.
- [5] N.V. PATEL AND V.M. SHAH, A note on the absolute convergence of lacunary Fourier series, *Proc. Amer. Math. Soc.*, **93** (1985), 433–439.
- [6] R.G. VYAS, On the Absolute convergence of small gaps Fourier series of functions of $\wedge BV^{(p)}$, *J. Inequal. Pure and Appl. Math.*, **6**(1) (2005), Art. 23, 1–6. [ONLINE: <http://jipam.vu.edu.au/article.php?sid=492>]



On The Absolute Convergence
Of Small Gaps Fourier Series Of
Functions Of $\varphi \wedge BV$

R. G. Vyas

Title Page

Contents



Go Back

Close

Quit

Page 11 of 11