

AN ELEMENTARY PROOF OF BLUNDON'S INEQUALITY

GABRIEL DOSPINESCU

École Normale Supérieure, Paris, France.

E-Mail: gdospi2002@yahoo.com

MIRCEA LASCU

GIL Publishing House, Zalău, Romania.

E-Mail: gil1993@zalau.astral.ro

COSMIN POHOATA

13 Pridvorului Street, Bucharest 010014, Romania.

E-Mail: pohoata_cosmin2000@yahoo.com

MARIAN TETIVA

"Gheorghe Roșca Codreanu" High-School,

Bârlad 731183, Romania.

E-Mail: rianamro@yahoo.com

Received: 05 August, 2008


Accepted: 11 October, 2008

Communicated by: **K.B. Stolarsky**

2000 AMS Sub. Class.: Primary 52A40; Secondary 52C05.

Key words: Blundon's Inequality, Geometric Inequality, Arithmetic-Geometric Mean Inequality.



Abstract: In this note, we give an elementary proof of Blundon's Inequality. We make use of a simple auxiliary result, provable by only using the Arithmetic Mean - Geometric Mean Inequality.



Blundon's Inequality
G. Dospinescu, M. Lascu,
C. Pohoata and M. Tetiva
vol. 9, iss. 4, art. 100, 2008

[Title Page](#)

[Contents](#)



Page 1 of 6

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



Blundon's Inequality
G. Dospinescu, M. Lascu,
C. Pohoata and M. Tetiva
vol. 9, iss. 4, art. 100, 2008

For a given triangle ABC we shall consider that A, B, C denote the magnitudes of its angles, and a, b, c denote the lengths of its corresponding sides. Let R, r and s be the circumradius, the inradius and the semi-perimeter of the triangle, respectively. In addition, we will occasionally make use of the symbols \sum (cyclic sum) and \prod (cyclic product), where

$$\sum f(a) = f(a) + f(b) + f(c), \quad \prod f(a) = f(a)f(b)f(c).$$

In the AMERICAN MATHEMATICAL MONTHLY, W. J. Blundon [1] asked for the proof of the inequality

$$s \leq 2R + (3\sqrt{3} - 4)r$$

which holds in any triangle ABC . The solution given by the editors was in fact a comment made by A. Makowski [3], who refers the reader to [2], where Blundon originally published this inequality, and where he actually proves more, namely that this is the best such inequality in the following sense: if, for the numbers k and h the inequality

$$s \leq kR + hr$$

is valid in any triangle, with the equality occurring when the triangle is equilateral, then

$$2R + (3\sqrt{3} - 4)r \leq kR + hr.$$

In this note we give a new proof of Blundon's inequality by making use of the following preliminary result:

Lemma 1. *Any positive real numbers x, y, z such that*

$$x + y + z = xyz$$

Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 2 of 6

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

satisfy the inequality

$$(x - 1)(y - 1)(z - 1) \leq 6\sqrt{3} - 10.$$

Proof. Since the numbers are positive, from the given condition it follows immediately that $x < xyz \Leftrightarrow yz > 1$, and similarly $xz > 1$ and $yz > 1$, which shows that it is not possible for two of the numbers to be less than or equal to 1 (neither can all the numbers be less than 1). Because if a number is less than 1 and two are greater than 1 the inequality is obviously true (the product from the left-hand side being negative), we still have to consider the case when $x > 1, y > 1, z > 1$. Then the numbers $u = x - 1, v = y - 1$ and $w = z - 1$ are positive and, replacing $x = u + 1, y = v + 1, z = w + 1$ in the condition from the hypothesis, one gets

$$uvw + uv + uw + vw = 2.$$

By the Arithmetic Mean - Geometric Mean inequality

$$uvw + 3\sqrt[3]{u^2v^2w^2} \leq uvw + uv + uw + vw = 2,$$

and hence for $t = \sqrt[3]{uvw}$ we have

$$t^3 + 3t^2 - 2 \leq 0 \Leftrightarrow (t + 1)(t + 1 + \sqrt{3})(t + 1 - \sqrt{3}) \leq 0.$$

We conclude that $t \leq \sqrt{3} - 1$ and thus,

$$(x - 1)(y - 1)(z - 1) \leq 6\sqrt{3} - 10.$$

The equality occurs when $x = y = z = \sqrt{3}$. This proves Lemma 1. \square

We now proceed to prove Blundon's Inequality.





Blundon's Inequality
G. Dospinescu, M. Lascu,
C. Pohoata and M. Tetiva

vol. 9, iss. 4, art. 100, 2008

Title Page

Contents



Page 3 of 6


Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



Theorem 2. In any triangle ABC , we have that

$$s \leq 2R + (3\sqrt{3} - 4)r.$$

The equality occurs if and only if ABC is equilateral.

Proof. According to the well-known formulae

$$\cot \frac{A}{2} = \sqrt{\frac{s(s-a)}{(s-b)(s-c)}}, \quad \cot \frac{B}{2} = \sqrt{\frac{s(s-b)}{(s-c)(s-a)}}, \quad \cot \frac{C}{2} = \sqrt{\frac{s(s-c)}{(s-a)(s-b)}},$$

we deduce that

$$\sum \cot \frac{A}{2} = \prod \cot \frac{A}{2} = \frac{s}{r},$$

and

$$\sum \cot \frac{A}{2} \cot \frac{B}{2} = \sum \frac{s}{s-a} = \frac{4R+r}{r}.$$

In this case, by applying Lemma 1 to the positive numbers $x = \cot \frac{A}{2}$, $y = \cot \frac{B}{2}$ and $z = \cot \frac{C}{2}$, it follows that

$$\left(\cot \frac{A}{2} - 1\right) \left(\cot \frac{B}{2} - 1\right) \left(\cot \frac{C}{2} - 1\right) \leq 6\sqrt{3} - 10,$$

and therefore

$$2 \prod \cot \frac{A}{2} - \left(\sum \cot \frac{A}{2} \cot \frac{B}{2}\right) \leq 6\sqrt{3} - 9.$$

This can be rewritten as

$$\frac{2s}{r} - \frac{4R+r}{r} \leq 6\sqrt{3} - 9,$$

Title Page

Contents



Page 4 of 6

Go Back

Full Screen

Close


journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

and thus

$$s \leq 2R + (3\sqrt{3} - 4)r.$$

The equality occurs if and only if $\cot \frac{A}{2} = \cot \frac{B}{2} = \cot \frac{C}{2}$, i.e. when the triangle ABC is equilateral. This completes the proof of Blundon's Inequality. \square



Blundon's Inequality
G. Dospinescu, M. Lascu,
C. Pohoata and M. Tetiva

vol. 9, iss. 4, art. 100, 2008

Title Page

Contents



Page 5 of 6

Go Back

Full Screen


Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

References

- [1] W.J. BLUNDON, Problem E1935, *The Amer. Math. Monthly*, **73** (1966), 1122.
- [2] W.J. BLUNDON, Inequalities associated with the triangle, *Canad. Math. Bull.*, **8** (1965), 615–626.
- [3] A. MAKOWSKI, Solution of the Problem E1935, *The Amer. Math. Monthly*, **75** (1968), 404.



Blundon's Inequality
G. Dospinescu, M. Lascu,
C. Pohoata and M. Tetiva

vol. 9, iss. 4, art. 100, 2008

Title Page

Contents



Page 6 of 6

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756