

MAXIMAL OPERATORS OF FEJÉR MEANS OF VILENKIN-FOURIER SERIES

ISTVÁN BLAHOTA, GYÖRGY GÁT AND USHANGI GOGINAVA

Institute of Mathematics and Computer Science
College of Nyíregyháza
P.O. Box 166, Nyíregyháza
H-4400 Hungary

EMail: blahota@nyf.hu

EMail: gatgy@nyf.hu

Department of Mechanics and Mathematics
Tbilisi State University
Chavchavadze str. 1
Tbilisi 0128, Georgia
EMail: z_goginava@hotmail.com



volume 7, issue 4, article 149,
2006.

*Received 12 June, 2006;
accepted 22 November, 2006.*

Communicated by: Zs. Páles

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)



Abstract

The main aim of this paper is to prove that the maximal operator $\sigma^* := \sup_n |\sigma_n|$ of the Fejér means of the Vilenkin-Fourier series is not bounded from the Hardy space $H_{1/2}$ to the space $L_{1/2}$.

2000 Mathematics Subject Classification: 42C10.

Key words: Vilenkin system, Hardy space, Maximal operator.


The first author is supported by the Békésy Postdoctoral fellowship of the Hungarian Ministry of Education Bö 91/2003, the second author is supported by the Hungarian National Foundation for Scientific Research (OTKA), grant no. M 36511/2001., T 048780 and by the Széchenyi fellowship of the Hungarian Ministry of Education Szö 184/2003.

Let \mathbb{N}_+ denote the set of positive integers, $\mathbb{N} := \mathbb{N}_+ \cup \{0\}$. Let $m := (m_0, m_1, \dots)$ denote a sequence of positive integers not less than 2. Denote by $Z_{m_k} := \{0, 1, \dots, m_k - 1\}$ the additive group of integers modulo m_k .

Define the group G_m as the complete direct product of the groups Z_{m_j} , with the product of the discrete topologies of Z_{m_j} 's.

The direct product μ of the measures

$$\mu_k(\{j\}) := \frac{1}{m_k} \quad (j \in Z_{m_k})$$



Maximal Operators of Fejér Means of Vilenkin-Fourier Series

István Blahota, György Gát and Ushangi Goginava

Title Page

Contents



Go Back

Close

Quit

Page 2 of 13

is the Haar measure on G_m with $\mu(G_m) = 1$.

If the sequence m is bounded, then G_m is called a bounded Vilenkin group, else it is called an unbounded one. The elements of G_m can be represented by sequences $x := (x_0, x_1, \dots, x_j, \dots)$ ($x_j \in Z_{m_j}$). It is easy to give a base for the neighborhoods of G_m :

$$I_0(x) := G_m,$$

$$I_n(x) := \{y \in G_m | y_0 = x_0, \dots, y_{n-1} = x_{n-1}\}$$

for $x \in G_m$, $n \in \mathbb{N}$. Define $I_n := I_n(0)$ for $n \in \mathbb{N}_+$.

If we define the so-called generalized number system based on m in the following way:


$$M_0 := 1, \quad M_{k+1} := m_k M_k (k \in \mathbb{N}),$$

then every $n \in \mathbb{N}$ can be uniquely expressed as $n = \sum_{j=0}^{\infty} n_j M_j$, where $n_j \in Z_{m_j}$ ($j \in \mathbb{N}_+$) and only a finite number of n_j 's differ from zero. We use the following notations. Let (for $n > 0$) $|n| := \max\{k \in \mathbb{N} : n_k \neq 0\}$ (that is, $M_{|n|} \leq n < M_{|n|+1}$), $n^{(k)} = \sum_{j=k}^{\infty} n_j M_j$ and $n_{(k)} := n - n^{(k)}$.

Denote by $L_p(G_m)$ the usual (one dimensional) Lebesgue spaces ($\|\cdot\|_p$ the corresponding norms) ($1 \leq p \leq \infty$).

Next, we introduce on G_m an orthonormal system which is called the Vilenkin system. At first define the complex valued functions $r_k(x) : G_m \rightarrow \mathbb{C}$, the generalized Rademacher functions as

$$r_k(x) := \exp \frac{2\pi i x_k}{m_k} \quad (i^2 = -1, x \in G_m, k \in \mathbb{N}).$$



Maximal Operators of Fejér Means of Vilenkin-Fourier Series

István Blahota, György Gát and Ushangi Goginava

Title Page

Contents



Go Back

Close

Quit

Page 3 of 13

Now define the Vilenkin system $\psi := (\psi_n : n \in \mathbb{N})$ on G_m as:

$$\psi_n(x) := \prod_{k=0}^{\infty} r_k^{n_k}(x) \quad (n \in \mathbb{N}).$$

Specifically, we call this system the Walsh-Paley one if $m \equiv 2$.

The Vilenkin system is orthonormal and complete in $L_1(G_m)$ [9].


Now, we introduce analogues of the usual definitions in Fourier-analysis. If $f \in L_1(G_m)$ we can establish the following definitions in the usual manner:

(Fourier coefficients)
$$\widehat{f}(k) := \int_{G_m} f \overline{\psi_k} d\mu \quad (k \in \mathbb{N}),$$

(Partial sums)
$$S_n f := \sum_{k=0}^{n-1} \widehat{f}(k) \psi_k \quad (n \in \mathbb{N}_+, S_0 f := 0),$$

(Fejér means)
$$\sigma_n f := \frac{1}{n} \sum_{k=0}^{n-1} S_k f \quad (n \in \mathbb{N}_+),$$

(Dirichlet kernels)
$$D_n := \sum_{k=0}^{n-1} \psi_k \quad (n \in \mathbb{N}_+).$$



Maximal Operators of Fejér Means of Vilenkin-Fourier Series

István Blahota, György Gát and Ushangi Goginava

Title Page

Contents



Go Back

Close

Quit

Page 4 of 13

Recall that

$$(1) \quad D_{M_n}(x) = \begin{cases} M_n, & \text{if } x \in I_n, \\ 0, & \text{if } x \in G_m \setminus I_n. \end{cases}$$

The norm (or quasinorm) of the space $L_p(G_m)$ is defined by

$$\|f\|_p := \left(\int_{G_m} |f(x)|^p \mu(x) \right)^{\frac{1}{p}} \quad (0 < p < +\infty).$$

The space weak- $L_p(G_m)$ consists of all measurable functions f for which

$$\|f\|_{\text{weak-}L_p(G_m)} := \sup_{\lambda > 0} \lambda \mu(|f| > \lambda)^{\frac{1}{p}} < +\infty.$$

The σ -algebra generated by the intervals $\{I_n(x) : (x) \in G_m\}$ will be denoted by \mathcal{F}_n ($n \in \mathbb{N}$).

Denote by $f = (f^{(n)}, n \in \mathbb{N})$ a martingale with respect to $(\mathcal{F}_n, n \in \mathbb{N})$ (for details see, e. g. [10, 14]).

The maximal function of a martingale f is defined by

$$f^* = \sup_{n \in \mathbb{N}} |f^{(n)}|,$$

respectively.

In case $f \in L_1(G_m)$, the maximal functions are also be given by

$$f^*(x) = \sup_{n \in \mathbb{N}} \frac{1}{\mu(I_n(x))} \left| \int_{I_n(x)} f(u) \mu(u) \right|.$$



Maximal Operators of Fejér Means of Vilenkin-Fourier Series

István Blahota, György Gát and Ushangi Goginava

Title Page

Contents



Go Back

Close

Quit

Page 5 of 13

For $0 < p < \infty$ the Hardy martingale spaces $H_p(G_m)$ consist of all martingales for which

$$\|f\|_{H_p} := \|f^*\|_p < \infty.$$

If $f \in L_1(G_m)$, then it is easy to show that the sequence $(S_{M_n}(f) : n \in \mathbb{N})$ is a martingale. If f is a martingale, that is $f = (f^{(n)} : n \in \mathbb{N})$, then the Vilenkin-Fourier coefficients must be defined in a slightly different manner:

$$\widehat{f}(i) = \lim_{k \rightarrow \infty} \int_{G_m} f^{(k)}(x) \overline{\psi}_i(x) \mu(x).$$

The Vilenkin-Fourier coefficients of $f \in L_1(G_m)$ are the same as those of the martingale $(S_{M_n}(f) : n \in \mathbb{N})$ obtained from f .


For a martingale f the maximal operators of the Fejér means are defined by

$$\sigma^* f(x) = \sup_{n \in \mathbb{N}} |\sigma_n(f; x)|.$$

In this one-dimensional case the weak type inequality

$$\mu(\sigma^* f > \lambda) \leq \frac{c}{\lambda} \|f\|_1 \quad (\lambda > 0)$$

can be found in Zygmund [16] for the trigonometric series, in Schipp [6] for Walsh series and in Pál, Simon [5] for bounded Vilenkin series. Again in one-dimension, Fujji [3] and Simon [8] verified that σ^* is bounded from H_1 to L_1 . Weisz [11, 13] generalized this result and proved the boundedness of σ^* from the martingale Hardy space H_p to the space L_p for $p > 1/2$. Simon [7] gave a counterexample, which shows that this boundedness does not hold for $0 < p < 1/2$. In the endpoint case $p = 1/2$ Weisz [15] proved that σ^* is bounded from



Maximal Operators of Fejér Means of Vilenkin-Fourier Series

István Blahota, György Gát and Ushangi Goginava

Title Page

Contents



Go Back

Close

Quit

Page 6 of 13

the Hardy space $H_{1/2}$ to the space weak- $L_{1/2}$. By interpolation it follows that σ^* is not bounded from H_p to the space weak- L_p for all $0 < p < 1/2$.

Theorem 1. *For any bounded Vilenkin system the maximal operator σ^* of the Fejér means is not bounded from the Hardy space $H_{1/2}$ to the space $L_{1/2}$.*

The Fejér kernel of order n of the Vilenkin-Fourier series is defined by

$$K_n(x) := \frac{1}{n} \sum_{k=0}^{n-1} D_k(x).$$

In order to prove the theorem we need the following lemmas.

Lemma 2 ([4]). *Suppose that $s, t, n \in \mathbb{N}$ and $x \in I_t \setminus I_{t+1}$. If $t \leq s \leq |n|$, then*


$$\begin{aligned} & (n^{(s+1)} + M_s)K_{n^{(s+1)}+M_s}(x) - n^{(s+1)}K_{n^{(s+1)}} \\ &= \begin{cases} M_t M_s \psi_{n^{(s+1)}}(x) \frac{1}{1-r_t(x)}, & \text{if } x - x_t e_t \in I_s, \\ 0, & \text{otherwise.} \end{cases} \end{aligned}$$

Lemma 3 ([2]). *Let $2 < A \in \mathbb{N}_+$, $k \leq s < A$ and $n_A^* := M_{2A} + M_{2A-2} + \dots + M_2 + M_0$. Then*

$$n_{A-1}^* \left| K_{n_{A-1}^*}(x) \right| \geq \frac{M_{2k} M_{2s}}{4}$$

for

$$\begin{aligned} x \in I_{2A}(0, \dots, 0, x_{2k} \neq 0, 0, \dots, 0, x_{2s} \neq 0, x_{2s+1}, \dots, x_{2A-1}), \\ k = 0, 1, \dots, A-3, \quad s = k+2, k+3, \dots, A-1. \end{aligned}$$



Maximal Operators of Fejér Means of Vilenkin-Fourier Series

István Blahota, György Gát and Ushangi Goginava

Title Page

Contents



Go Back

Close

Quit

Page 7 of 13

Proof of Theorem 1. Let $A \in \mathbb{N}_+$ and

$$f_A(x) := D_{M_{2A+1}}(x) - D_{M_{2A}}(x).$$

In the sequel we are going to prove for the function f_A that

$$\frac{\|\sigma^* f_A\|_{1/2}}{\|f_A\|_{H_{1/2}}} \geq c \log_q^2 A,$$

where $q = \sup \{m_0, m_1, \dots\}$ and constant c depends only on q . This inequality obviously would show the unboundedness of σ^* .

It is evident that


$$\widehat{f}_A(i) = \begin{cases} 1, & \text{if } i = M_{2A}, \dots, M_{2A+1} - 1, \\ 0, & \text{otherwise.} \end{cases}$$

Then we can write

$$(2) \quad S_i(f_A; x) = \begin{cases} D_i(x) - D_{M_{2A}}(x), & \text{if } i = M_{2A} + 1, \dots, M_{2A+1} - 1, \\ f_A(x), & \text{if } i \geq M_{2A+1}, \\ 0, & \text{otherwise.} \end{cases}$$

Since

$$f_A^*(x) = \sup_{n \in \mathbb{N}} |S_{M_n}(f_A; x)| = |f_A(x)|,$$



Maximal Operators of Fejér Means of Vilenkin-Fourier Series

István Blahota, György Gát and Ushangi Goginava

Title Page

Contents



Go Back

Close

Quit

Page 8 of 13

from (1) we get


$$\begin{aligned}
 (3) \quad \|f_A\|_{H_{1/2}} &= \|f_A^*\|_{1/2} = \|D_{M_{2A+1}} - D_{M_{2A}}\|_{1/2} \\
 &= \left(\int_{I_{2A} \setminus I_{2A+1}} M_{2A}^{\frac{1}{2}} + \int_{I_{2A+1}} |M_{2A+1} - M_{2A}|^{\frac{1}{2}} \right)^2 \\
 &= \left(\frac{m_{2A} - 1}{M_{2A+1}} M_{2A}^{\frac{1}{2}} + \frac{(m_{2A} - 1)^{\frac{1}{2}}}{M_{2A+1}} M_{2A}^{\frac{1}{2}} \right)^2 \\
 &\leq 2^2 m_{2A} M_{2A}^{-1} \\
 &\leq c M_{2A}^{-1}.
 \end{aligned}$$

Since

$$D_{k+M_{2A}} - D_{M_{2A}} = \psi_{M_{2A}} D_k, \quad k = 1, 2, \dots, M_{2A},$$

from (2) we obtain

$$\begin{aligned}
 (4) \quad \sigma^* f_A(x) &= \sup_{n \in \mathbb{N}} |\sigma_n(f_A; x)| \\
 &\geq |\sigma_{n_A^*}(f_A; x)| \\
 &= \frac{1}{n_A^*} \left| \sum_{i=0}^{n_A^*-1} S_i(f_A; x) \right| \\
 &= \frac{1}{n_A^*} \left| \sum_{i=M_{2A}+1}^{n_A^*-1} (D_i(x) - D_{M_{2A}}(x)) \right|
 \end{aligned}$$



Maximal Operators of Fejér Means of Vilenkin-Fourier Series

István Blahota, György Gát and Ushangi Goginava

Title Page

Contents



Go Back

Close

Quit

Page 9 of 13

$$\begin{aligned}
&= \frac{1}{n_A^*} \left| \sum_{i=1}^{n_{A-1}^*-1} (D_{i+M_{2A}}(x) - D_{M_{2A}}(x)) \right| \\
&= \frac{n_{A-1}^*}{n_A^*} \left| K_{n_{A-1}^*}(x) \right|.
\end{aligned}$$

Let $q := \sup\{m_i : i \in \mathbb{N}\}$. For every $l = 1, \dots, \left[\frac{1}{4} \log_q \sqrt{A}\right] - 1$ (A is supposed to be large enough) let k_l be the smallest natural numbers, for which

$$M_{2A} \sqrt{A} \frac{1}{q^{4l}} \leq M_{2k_l}^2 < M_{2A} \sqrt{A} \frac{1}{q^{4l-4}}$$

hold.

Denote

$$I_{2A}^{k,s}(x) := I_{2A}(0, \dots, 0, x_{2k} \neq 0, 0, \dots, 0, x_{2s} \neq 0, x_{2s+1}, \dots, x_{2A-1})$$

and let


$$x \in I_{2A}^{k_l, k_l+1}(z)$$

Then from Lemma 3 and (4) we obtain that

$$\sigma^* f_A(x) \geq c \frac{M_{2k_l}^2}{M_{2A}} \geq c \frac{\sqrt{A}}{q^{4l}}$$

On the other hand,

$$\sqrt{\|\sigma^* f_A\|_{1/2}} \geq c \sum_{l=1}^{\left[\frac{1}{4} \log_q \sqrt{A}\right]} \sum_{x_{2k_l+3}=0}^{m_{2k_l+3}-1} \cdots \sum_{x_{2A-1}=0}^{m_{2A-1}-1} \frac{\sqrt{A}}{q^{2l}} \mu \left(I_{2A}^{k_l, k_l+1}(x) \right)$$



Maximal Operators of Fejér Means of Vilenkin-Fourier Series

István Blahota, György Gát and Ushangi Goginava

Title Page

Contents



Go Back

Close

Quit


Page 10 of 13

$$\begin{aligned}
&\geq c\sqrt[4]{A} \sum_{l=1}^{\lfloor \frac{1}{4} \log_q \sqrt{A} \rfloor} \frac{m_{2k_l+3} \cdots m_{2A-1}}{q^{2l} M_{2A}} \\
&= c\sqrt[4]{A} \sum_{l=1}^{\lfloor \frac{1}{4} \log_q \sqrt{A} \rfloor} \frac{1}{q^{2l} M_{2k_l+2}} \\
&\geq c\sqrt[4]{A} \sum_{l=1}^{\lfloor \frac{1}{4} \log_q \sqrt{A} \rfloor} \frac{1}{q^{2l} M_{2k_l}} \\
&\geq c\sqrt[4]{A} \sum_{l=1}^{\lfloor \frac{1}{4} \log_q \sqrt{A} \rfloor} \frac{1}{q^{2l} \sqrt{M_{2A} \sqrt{A} q^{-4l+4}}} \\
&\geq c \frac{\log_q A}{\sqrt{M_{2A}}}.
\end{aligned}$$

Combining this with (3) we obtain

$$\frac{\|\sigma^* f_A\|_{1/2}}{\|f_A\|_{H_{1/2}}} \geq \frac{c \log_q^2 A}{M_{2A}} M_{2A} = c \log_q^2 A \rightarrow \infty \quad \text{as } A \rightarrow \infty.$$

Thus, the theorem is proved. \square



Maximal Operators of Fejér Means of Vilenkin-Fourier Series

István Blahota, György Gát and Ushangi Goginava

Title Page

Contents



Go Back


Close

Quit

Page 11 of 13

References

- [1] G.N. AGAEV, N.Ya. VILENKIN, G.M. DZHAFARLI AND A.I. RUBIN-SHTEJN, Multiplicative systems of functions and harmonic analysis on zero-dimensional groups, Baku, Ehlms, 1981 (in Russian).
- [2] I. BLAHOTA, G. GÁT AND U. GOGINAVA, Maximal operators of Fejér means of double Vilenkin-Fourier series, *Colloq. Math.*, to appear.
- [3] N.J. FUJII, Cesàro summability of Walsh-Fourier series, *Proc. Amer. Math. Soc.*, **77** (1979), 111–116.
- [4] G. GÁT, Pointwise convergence of the Fejér means of functions on unbounded Vilenkin groups, *J. of Approximation Theory*, **101**(1) (1999), 1–36.
- [5] J. PÁL AND P. SIMON, On a generalization of the concept of derivate, *Acta Math. Hung.*, **29** (1977), 155–164.
- [6] F. SCHIPP, Certain rearrangements of series in the Walsh series, *Mat. Zametki*, **18** (1975), 193–201.
- [7] P. SIMON, Cesaro summability with respect to two-parameter Walsh system, *Monatsh. Math.*, **131** (2000), 321–334.
- [8] P. SIMON, Investigations with respect to the Vilenkin system, *Annales Univ. Sci. Budapest Eötv., Sect. Math.*, **28** (1985), 87–101.
- [9] N. Ya. VILENKIN, A class of complete orthonormal systems, *Izv. Akad. Nauk. U.S.S.R., Ser. Mat.*, **11** (1947), 363–400



Maximal Operators of Fejér Means of Vilenkin-Fourier Series

István Blahota, György Gát and Ushangi Goginava

Title Page

Contents




Go Back

Close

Quit

Page 12 of 13

- [10] F. WEISZ, *Martingale Hardy Spaces and their Applications in Fourier Analysis*, Springer, Berlin - Heidelberg - New York, 1994.
- [11] F. WEISZ, Cesàro summability of one and two-dimensional Walsh-Fourier series, *Anal. Math.*, **22** (1996), 229–242.
- [12] F. WEISZ, Hardy spaces and Cesàro means of two-dimensional Fourier series, *Bolyai Soc. Math. Studies*, **5** (1996), 353–367.
- [13] F. WEISZ, Bounded operators on Weak Hardy spaces and applications, *Acta Math. Hungar.*, **80** (1998), 249–264.
- [14] F. WEISZ, *Summability of Multi-dimensional Fourier Series and Hardy Space*, Kluwer Academic, Dordrecht, 2002.
- [15] F. WEISZ, ϑ -summability of Fourier series, *Acta Math. Hungar.*, **103**(1-2) (2004), 139–176.
- [16] A. ZYGMUND, *Trigonometric Series*, Vol. 1, Cambridge Univ. Press, 1959.



Maximal Operators of Fejér Means of Vilenkin-Fourier Series

István Blahota, György Gát and Ushangi Goginava

Title Page

Contents



Go Back

Close

Quit

Page 13 of 13