

Journal of Inequalities in Pure and Applied Mathematics

A NOTE ON ABSOLUTE NÖRLUND SUMMABILITY

HÜSEYİN BOR

Department of Mathematics
Erciyes University
38039 Kayseri, Turkey

EMail: bor@erciyes.edu.tr

©2000 Victoria University
ISSN (electronic): 1443-5756
287-05



volume 6, issue 4, article 115,
2005.

*Received 26 September, 2005;
accepted 17 October, 2005.*

Communicated by: L. Leindler

Abstract

Contents



Home Page

Go Back

Close

Quit

Abstract

In this paper a main theorem on $|N, p_n|_k$ summability factors, which generalizes a result of Bor [2] on $|N, p_n|$ summability factors, has been proved.

2000 Mathematics Subject Classification: 40D15, 40F05, 40G05.

Key words: Nörlund summability, summability factors, power increasing sequences.

The author is grateful to the referee for his valuable suggestions for the improvement of this paper.

Contents

1	Introduction	3
2	Main Result	6
3	Proof of Theorem 2.1	7
	References	



A Note on Absolute Nörlund Summability

Hüseyin Bor

Title Page

Contents



Go Back

Close

Quit

Page 2 of 10

1. Introduction

A positive sequence (b_n) is said to be almost increasing if there exist a positive increasing sequence (c_n) and two positive constants A and B such that $Ac_n \leq b_n \leq Bc_n$ (see [1]). A positive sequence (γ_n) is said to be a quasi β -power increasing sequence if there exists a constant $K = K(\beta, \gamma) \geq 1$ such that

$$(1.1) \quad Kn^\beta \gamma_n \geq m^\beta \gamma_m$$

holds for all $n \geq m \geq 1$. It should be noted that every almost increasing sequence is a quasi β -power increasing sequence for any nonnegative β , but the converse need not be true as can be seen by taking the example, say $\gamma_n = n^{-\beta}$ for $\beta > 0$. We denote by \mathcal{BV}_O the $\mathcal{BV} \cap \mathcal{C}_O$, where \mathcal{C}_O and \mathcal{BV} are the null sequences and sequences with bounded variation, respectively.

Let $\sum a_n$ be a given infinite series with the sequence of partial sums (s_n) and $w_n = na_n$. By u_n^α and t_n^α we denote the n -th Cesàro means of order α , with $\alpha > -1$, of the sequences (s_n) and (w_n) , respectively.

The series $\sum a_n$ is said to be summable $|C, \alpha|_k$, $k \geq 1$, if (see [4])


$$(1.2) \quad \sum_{n=1}^{\infty} n^{k-1} |u_n^\alpha - u_{n-1}^\alpha|^k = \sum_{n=1}^{\infty} \frac{1}{n} |t_n^\alpha|^k < \infty.$$

Let (p_n) be a sequence of constants, real or complex, and let us write

$$(1.3) \quad P_n = p_0 + p_1 + p_2 + \cdots + p_n \neq 0, \quad (n \geq 0).$$

The sequence-to-sequence transformation

$$(1.4) \quad \sigma_n = \frac{1}{P_n} \sum_{v=0}^n p_{n-v} s_v$$



A Note on Absolute Nörlund Summability

Hüseyin Bor

Title Page

Contents



Go Back

Close

Quit

Page 3 of 10

defines the sequence (σ_n) of the Nörlund mean of the sequence (s_n) , generated by the sequence of coefficients (p_n) . The series $\sum a_n$ is said to be summable $|N, p_n|_k, k \geq 1$, if (see [3])

$$(1.5) \quad \sum_{n=1}^{\infty} n^{k-1} |\sigma_n - \sigma_{n-1}|^k < \infty.$$

In the special case when

$$(1.6) \quad p_n = \frac{\Gamma(n + \alpha)}{\Gamma(\alpha)\Gamma(n + 1)}, \quad \alpha \geq 0$$

the Nörlund mean reduces to the (C, α) mean and $|N, p_n|_k$ summability becomes $|C, \alpha|_k$ summability. For $p_n = 1$ and $P_n = n$, we get the $(C, 1)$ mean and then $|N, p_n|_k$ summability becomes $|C, 1|_k$ summability. For any sequence (λ_n) , we write $\Delta\lambda_n = \lambda_n - \lambda_{n+1}$.


The known results. Concerning the $|C, 1|_k$ and $|N, p_n|_k$ summabilities Varma [6] has proved the following theorem.

Theorem A. *Let $p_0 > 0, p_n \geq 0$ and (p_n) be a non-increasing sequence. If $\sum a_n$ is summable $|C, 1|_k$, then the series $\sum a_n P_n (n + 1)^{-1}$ is summable $|N, p_n|_k, k \geq 1$.*

Quite recently Bor [2] has proved the following theorem.

Theorem B. *Let (p_n) be as in Theorem A, and let (X_n) be a quasi β -power increasing sequence with some $0 < \beta < 1$. If*

$$(1.7) \quad \sum_{v=1}^n \frac{1}{v} |t_v| = O(X_n) \quad \text{as } n \rightarrow \infty,$$



A Note on Absolute Nörlund Summability

Hüseyin Bor

Title Page

Contents



Go Back

Close

Quit

Page 4 of 10

and the sequences (λ_n) and (β_n) satisfy the following conditions


$$(1.8) \quad X_n \lambda_n = O(1),$$

$$(1.9) \quad |\Delta \lambda_n| \leq \beta_n,$$

$$(1.10) \quad \beta_n \rightarrow 0,$$

$$(1.11) \quad \sum n X_n |\Delta \beta_n| < \infty,$$

then the series $\sum a_n P_n \lambda_n (n+1)^{-1}$ is summable $|N, p_n|$.



**A Note on Absolute Nörlund
Summability**

Hüseyin Bor

Title Page

Contents



Go Back

Close

Quit

Page 5 of 10

2. Main Result

The aim of this paper is to generalize Theorem B for $|N, p_n|_k$ summability. Now we shall prove the following theorem.

Theorem 2.1. *Let (p_n) be as in Theorem A, and let (X_n) be a quasi β -power increasing sequence with some $0 < \beta < 1$. If*

$$(2.1) \quad \sum_{v=1}^n \frac{1}{v} |t_v|^k = O(X_n) \quad \text{as } n \rightarrow \infty,$$

and the sequences (λ_n) and (β_n) satisfy the conditions from (1.8) to (1.11) of Theorem B; further suppose that

$$(2.2) \quad (\lambda_n) \in \mathcal{BV}_O,$$

then the series $\sum a_n P_n \lambda_n (n+1)^{-1}$ is summable $|N, p_n|_k$, $k \geq 1$.


Remark 1. *It should be noted that if we take $k = 1$, then we get Theorem B. In this case condition (2.2) is not needed.*

We need the following lemma for the proof of our theorem.

Lemma 2.2 ([5]). *Except for the condition (2.2), under the conditions on (X_n) , (λ_n) and (β_n) as taken in the statement of the theorem, the following conditions hold when (1.11) is satisfied:*

$$(2.3) \quad n\beta_n X_n = O(1) \text{ as } n \rightarrow \infty,$$

$$(2.4) \quad \sum_{n=1}^{\infty} \beta_n X_n < \infty.$$



A Note on Absolute Nörlund Summability

Hüseyin Bor

Title Page

Contents



Go Back

Close

Quit

Page 6 of 10

3. Proof of Theorem 2.1

In order to prove the theorem, we need consider only the special case in which (N, p_n) is $(C, 1)$, that is, we shall prove that $\sum a_n \lambda_n$ is summable $|C, 1|_k$. Our theorem will then follow by means of Theorem A. Let T_n be the n -th $(C, 1)$ mean of the sequence $(na_n \lambda_n)$, that is,

$$(3.1) \quad T_n = \frac{1}{n+1} \sum_{v=1}^n va_v \lambda_v.$$

Using Abel's transformation, we have


$$\begin{aligned} T_n &= \frac{1}{n+1} \sum_{v=1}^n va_v \lambda_v = \frac{1}{n+1} \sum_{v=1}^{n-1} \Delta \lambda_v (v+1)t_v + \lambda_n t_n \\ &= T_{n,1} + T_{n,2}, \text{ say.} \end{aligned}$$

To complete the proof of the theorem, it is sufficient to show that

$$(3.2) \quad \sum_{n=1}^{\infty} \frac{1}{n} |T_{n,r}|^k < \infty \quad \text{for } r = 1, 2, \text{ by (1.2).}$$

Now, we have that

$$\begin{aligned} \sum_{n=2}^{m+1} \frac{1}{n} |T_{n,1}|^k &\leq \sum_{n=2}^{m+1} \frac{1}{n(n+1)^k} \left\{ \sum_{v=1}^{n-1} \frac{v+1}{v} v |\Delta \lambda_v| |t_v| \right\}^k \\ &= O(1) \sum_{n=2}^{m+1} \frac{1}{n^{k+1}} \left\{ \sum_{v=1}^{n-1} v |\Delta \lambda_v| |t_v| \right\}^k \end{aligned}$$



A Note on Absolute Nörlund
Summability

Hüseyin Bor

Title Page

Contents



Go Back


Close

Quit

Page 7 of 10

$$\begin{aligned}
&= O(1) \sum_{n=2}^{m+1} \frac{1}{n^2} \left\{ \sum_{v=1}^{n-1} v |\Delta\lambda_v| |t_v|^k \right\} \times \left\{ \frac{1}{n} \sum_{v=1}^{n-1} v |\Delta\lambda_v| \right\}^{k-1} \\
&= O(1) \sum_{n=2}^{m+1} \frac{1}{n^2} \sum_{v=1}^{n-1} v |\Delta\lambda_v| |t_v|^k \quad (\text{by (2.2)}) \\
&= O(1) \sum_{n=2}^{m+1} \frac{1}{n^2} \left\{ \sum_{v=1}^{n-1} v \beta_v |t_v|^k \right\} \quad (\text{by (1.9)}) \\
&= O(1) \sum_{v=1}^m v \beta_v |t_v|^k \sum_{n=v+1}^{m+1} \frac{1}{n^2} = O(1) \sum_{v=1}^m v \beta_v \frac{|t_v|^k}{v} \\
&= O(1) \sum_{v=1}^{m-1} \Delta(v\beta_v) \sum_{r=1}^v \frac{|t_r|^k}{r} + O(1) m \beta_m \sum_{v=1}^m \frac{|t_v|^k}{v} \\
&= O(1) \sum_{v=1}^{m-1} |\Delta(v\beta_v)| X_v + O(1) m \beta_m X_m \quad (\text{by (2.1)}) \\
&= O(1) \sum_{v=1}^{m-1} |(v+1)\Delta\beta_v - \beta_v| X_v + O(1) m \beta_m X_m \\
&= O(1) \sum_{v=1}^{m-1} v |\Delta\beta_v| X_v + O(1) \sum_{v=1}^{m-1} |\beta_v| X_v + O(1) m \beta_m X_m \\
&= O(1) \quad \text{as } m \rightarrow \infty,
\end{aligned}$$

in view of (1.11), (2.3) and (2.4).



A Note on Absolute Nörlund Summability

Hüseyin Bor

Title Page

Contents



Go Back

Close


Quit

Page 8 of 10

Again

$$\begin{aligned}
 \sum_{n=1}^m \frac{1}{n} |T_{n,2}|^k &= \sum_{n=1}^m |\lambda_n|^k \frac{|t_n|^k}{n} \\
 &= \sum_{n=1}^m |\lambda_n|^{k-1} |\lambda_n| \frac{|t_n|^k}{n} = O(1) \sum_{n=1}^m |\lambda_n| \frac{|t_n|^k}{n} \quad (\text{by (2.2)}) \\
 &= O(1) \sum_{n=1}^{m-1} \Delta |\lambda_n| \sum_{v=1}^n \frac{|t_v|^k}{v} + O(1) |\lambda_m| \sum_{n=1}^m \frac{|t_n|^k}{n} \\
 &= O(1) \sum_{n=1}^{m-1} |\Delta \lambda_n| X_n + O(1) |\lambda_m| X_m \quad (\text{by (2.1)}) \\
 &= O(1) \sum_{n=1}^{m-1} \beta_n X_n + O(1) |\lambda_m| X_m = O(1) \quad \text{as } m \rightarrow \infty,
 \end{aligned}$$

by virtue of (1.8) and (2.4). This completes the proof of the theorem. □



**A Note on Absolute Nörlund
Summability**

Hüseyin Bor

Title Page

Contents



Go Back


Close

Quit

Page 9 of 10

References

- [1] S. ALJANČIĆ AND D. ARANDELOVIĆ, O -regularly varying functions, *Publ. Inst. Math.*, **22** (1977), 5–22.
- [2] H. BOR, Absolute Nörlund summability factors, *J. Inequal. Pure Appl. Math.*, **6**(3) (2005), Art. 62. [ONLINE <http://jipam.vu.edu.au/article.php?sid=535>].
- [3] D. BORWEIN AND F.P. CASS, Strong Nörlund summability, *Math. Zeith.*, **103** (1968), 94–111.
- [4] T.M. FLETT, On an extension of absolute summability and some theorems of Littlewood and Paley, *Proc. London Math. Soc.*, **7** (1957), 113–141.
- [5] L. LEINDLER, A new application of quasi power increasing sequences, *Publ. Math. Debrecen*, **58** (2001), 791–796.
- [6] R.S. VARMA, On the absolute Nörlund summability factors, *Riv. Math. Univ. Parma* (4), **3** (1977), 27–33.



A Note on Absolute Nörlund Summability

Hüseyin Bor

Title Page

Contents



Go Back

Close

Quit

Page 10 of 10