

# ON A QUESTION OF MILMAN AND ALESKER CONCERNING THE MONOTONICITY OF A DOMAIN FUNCTIONAL

G. KEADY

School of Mathematics and Statistics,  
University of Western Australia  
6009, Australia  
EMail: [keady@maths.uwa.edu.au](mailto:keady@maths.uwa.edu.au)

*Received:* 08 November, 2006


*Accepted:* 15 July, 2007

*Communicated by:* S.S. Dragomir

*2000 AMS Sub. Class.:* 26D20, 52A20, 52A40.

*Key words:* Moment of inertia, Convex set.

*Abstract:* We answer affirmatively the special case of  $q = 1$ ,  $n = 2$ ,  $j = 2$  of Question 3 on page 1004 of Alesker, *Annals of Mathematics*, **149** (1999).



Monotonicity of a  
Domain Functional

G. Keady

vol. 8, iss. 3, art. 76, 2007

[Title Page](#)

[Contents](#)



Page 1 of 8

[Go Back](#)

[Full Screen](#)


[Close](#)

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-5756

# Contents

1	Introduction	3
2	Support Functions	4
3	A Partial Answer to Alesker's Question 3	7



---

**Monotonicity of a  
Domain Functional**

G. Keady

vol. 8, iss. 3, art. 76, 2007

---

Title Page

Contents



Page 2 of 8


Go Back

Full Screen

Close

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-5756



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 3 of 8

Go Back

Full Screen

Close

## 1. Introduction

Let  $D_0$  be a bounded convex domain in  $\mathbb{R}^2$  containing the origin. Let  $B(0, \rho_1)$  be the disk, centred at 0, of radius  $\rho_1$ , and consider the Minkowski sum

$$D(t) = (1 - t)D_0 + tB(0, \rho_1).$$

Define  $I_O(\cdot, q)$  by

$$I_O(D(t), q) = \int_{D(t)} |x|^{2q},$$

and note that at  $q = 1$  this is the polar moment of inertia about the origin of the domain  $D(t)$ . When  $q = 1$  we will simply write  $I_O(D(t))$  and omit the second argument. The derivatives of  $I_O(D(t), q)$  with respect to  $\rho_1$  are shown to be positive: see Alesker [1, Theorem 6.1]. Alesker [1, p. 1004, Question 3] asks about the domain-monotonicity of the derivatives with respect to  $\rho_1$  evaluated at  $\rho_1 = 0$ . Our answer to the special case of this question will be given in Theorem 3.1 in §3.

Alesker considers one-dimensional sets. These suffice to show that domain monotonicity will not hold true in general unless the origin is in the domain.

The notation in this paper is the same as in [7].

When the convex sets are centrally-symmetric, several of the proofs in [7] simplify, and there are additional results. Alesker asked the question for centrally-symmetric convex sets, but here we can, for  $n = 2$ ,  $q = 1$ , answer it more generally, merely requiring the set to contain the origin.



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 4 of 8

[Go Back](#)

[Full Screen](#)

[Close](#)

## 2. Support Functions

We use the letter  $p$  for the support function as in [4] and Santalo's book [9]. An adequate description of  $p$  is as 'the perpendicular distance from the origin to the tangent'. The radius of curvature is then

$$\rho = p + \ddot{p}, \quad \text{where } \dot{f} = \frac{df}{d\varphi}, \quad ds = \rho d\varphi.$$

For a diagram, see page 2 of [9]. Santalo's  $\phi$  is the angle between a line *normal* to the tangent and the  $x$ -axis. Following Santalo's notation, let  $H$  be a point on the tangent line such that  $OH$  is perpendicular to the tangent line.  $|OH| = p$ . The boundaries of our convex sets  $D$  can be determined from the functions  $p(\phi)$  through formulae (1.3) of [9]. Our  $\varphi$  is the angle between the *tangent* line (through  $H$ ) and the  $x$ -axis. We have

$$\varphi = \phi + \frac{\pi}{2}.$$

Then the area and perimeter are given by

$$(2.1) \quad A = \text{Area}(D) = \frac{1}{2} \int_0^{2\pi} p\rho \, d\varphi = \frac{1}{2} \int_0^{2\pi} (p^2 - \dot{p}^2) d\varphi,$$

$$(2.2) \quad L = \int_0^{2\pi} \rho \, d\varphi = \int_0^{2\pi} p \, d\varphi.$$

$D$  is convex iff  $\rho \geq 0$ . In the case of a polygon, for example, we might interpret  $\rho$  as a nonnegative measure. The set  $\mathcal{S}$  of support functions forms a cone:  $\mathcal{S}$  is convex, and if  $t > 0$  and  $p \in \mathcal{S}$ , then  $tp \in \mathcal{S}$ .

We now suppose that we have two convex domains  $D_0$  and  $D_1$ . We denote  $\text{Area}(D_0) = A_0$  and  $\text{Area}(D_1) = A_1$ . We have the following pretty, and very well-known, result:



Title Page

Contents



Page 5 of 8

Go Back

Full Screen

Close

**Lemma 2.1.** For convex sets  $D_0, D_1$ , the support function for  $D(t)$  is given by

$$(2.3) \quad p_t = (1 - t)p_0 + tp_1.$$

In particular, the preceding lemma yields that

$$(2.4) \quad L(t) := L(D(t)) = (1 - t)L_0 + tL_1,$$

$$(2.5) \quad A(t) := \text{Area}(D(t)) = (1 - t)^2 A_0 + 2t(1 - t)A_{0,1} + t^2 A_1,$$

where the *mixed-area*  $A_{0,1}$  satisfies

$$(2.6) \quad A_{0,1} := A(D_0, D_1) = \frac{1}{2} \int_0^{2\pi} (p_0 p_1 - \dot{p}_0 \dot{p}_1) d\varphi.$$

(All of these functionals are monotonic under domain inclusion for convex sets.)

There are many nice properties of support functions. Here is one. See [10, p. 37], or the first page of [8].

**Theorem 2.2.** If  $0 \in D \subseteq \hat{D}$  then  $0 \leq p \leq \hat{p}$ .

We do not use, but state:

**Theorem 2.3** ([3, p. 56]). Let  $C$  denote the convex hull of the union of the convex domains  $D_0$  and  $D_1$ . Then, the support functions satisfy

$$p_C = \max(p_{D_0}, p_{D_1}).$$

(Further general references on convex domains and their support functions include [2], [4], [6], [8], [10].)

Starting from the expressions for  $x(\varphi)$  and  $y(\varphi)$  for boundary points, expressing the coordinates in terms of  $p$  and  $\dot{p} = dp/d\varphi$ , in [7], using Lemma 2.1, the following



Title Page

Contents



Page 6 of 8

Go Back

Full Screen

Close


expression for  $I_O(D(t))$  when  $D_1 = B(0, \rho_1)$  is derived:

$$I_O(D(t)) = (1-t)^4 I_O(D_0) + \rho_1 t (1-t)^3 I(\partial D_0) + (\rho_1 t (1-t))^2 Z \\ + (\rho_1 t)^3 (1-t) L + \frac{\pi}{2} (\rho_1 t)^4.$$

Here

$$(2.7) \quad Z := \frac{1}{2} \int_0^{2\pi} (3p^2 - \dot{p}^2) d\varphi.$$

In the two preceding equations,  $p$  is the support function for  $D_0$ , and  $L$  and  $Z$  are evaluated for  $D_0$ .



Title Page

Contents



Page 7 of 8

Go Back

Full Screen

Close

### 3. A Partial Answer to Alesker's Question 3

**Theorem 3.1.** *Let  $D, \hat{D}$  be convex domains containing the origin with smooth boundaries. Define*

$$\alpha(j) = \frac{d^j}{d\rho_1^j} I_O(D(t))|_{\rho_1=0}$$

and define  $\hat{\alpha}$  similarly. Then (for  $n = 2, q = 1$ ),

$$\alpha(2) = 2t^2(1-t)^2 Z,$$

and

$$Z = A + \int_0^{2\pi} p(\varphi)^2 d\varphi.$$

The function  $Z$  is nonnegative and is monotonic under domain inclusion, i.e. if  $0 \in D \subseteq \hat{D}$ , then  $0 \leq Z(D) \leq Z(\hat{D})$ .

*Proof.* The area is monotonic under domain inclusion. Using Theorem 2.2, so is  $\int_0^{2\pi} p(\varphi)^2 d\varphi$ . Hence we have the required monotonicity of  $Z$ .


The restriction that the boundaries be smooth can be removed by taking limits.

Alesker states that  $\alpha(1)$  can be shown to be monotonic under domain inclusion. Returning to  $n = 2, q = 1$ , as  $\alpha(3) = 6t^3(1-t)L$ , we also have that  $\alpha(3)$  is monotonic under domain inclusion.  $\alpha(4)$  is independent of domain.  $\square$

For centrally symmetric domains, other properties of the second derivatives can be deduced from the results in [5].

## References

- [1] S. ALESKER, Continuous rotation invariant valuations on convex sets, *Annals of Mathematics*, **149** (1999), 977–1005.
- [2] T. BONNESEN AND W. FENCHEL, *Theory of Convex Bodies*, German, 1934; English translation, 1987.
- [3] H.G. EGGLESTON, *Convexity*, Cambridge Univ. Press, 1969.
- [4] H. FLANDERS, A proof of Minkowski's inequality for convex curves, *Amer. Math. Monthly*, **75** (1968), 581–593.
- [5] H. HADWIGER, Konkave eikerperfunktionale und hoher tragheitsmomente, *Comment Math. Helv.*, **30** (1956), 285–296.
- [6] M. KALLAY, Reconstruction of a plane convex body from the curvature of its boundary, *Israel J. Math.*, **17** (1974), 149–161.
- [7] G. KEADY, On Hadwiger's results concerning Minkowski sums and isoperimetric inequalities for moments of inertia, *RGMIA Research Report Collection*, **9**(4) (2006), Art. 16. [ONLINE: <http://rgmia.vu.edu.au/v9n4.html>].
- [8] D.A. KLAIN, The Brunn-Minkowski inequality in the plane, *Preprint*, (2002).
- [9] L.A. SANTALO, *Integral Geometry and Geometric Probability*, Addison-Wesley, 1976.
- [10] R. SCHNEIDER, *Convex Bodies: The Brunn-Minkowski Theory*, Cambridge University Press, 1993.




Monotonicity of a  
Domain Functional

G. Keady

vol. 8, iss. 3, art. 76, 2007

Title Page

Contents



Page 8 of 8

Go Back

Full Screen

Close

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-5756