

Journal of Inequalities in Pure and Applied Mathematics

HYERS-ULAM STABILITY OF THE GENERALIZED TRIGONOMETRIC FORMULAS

¹AHMED REDOUANI, ¹ELHOUCIEN ELQORACHI AND
²BELAID BOUIKHALENE


¹Laboratory LAMA
Harmonic Analysis and Functional Equations Team
Department of Mathematics
Faculty of Sciences, University of Ibn Zohr
Agadir, Morocco

EMail: redouani_ahmed@yahoo.fr

EMail: elqorachi@hotmail.com

²Laboratory LAMA, Department of Mathematics
Faculty of Sciences, University of Ibn Tofail
Kenitra, Morocco

EMail: bbouikhalene@yahoo.fr



volume 7, issue 2, article 74,
2006.

*Received 14 September, 2005;
accepted 14 October, 2005.*

Communicated by: Th.M. Rassias

Abstract

Contents



Home Page

Go Back

Close

Quit

Abstract

In this paper, we will investigate the Hyers-Ulam stability of the following functional equations

$$\int_G \int_K f(xtk \cdot y) dk d\mu(t) = f(x)g(y) + g(x)f(y), \quad x, y \in G$$

and

$$\int_G \int_K f(xtk \cdot y) dk d\mu(t) = f(x)f(y) - g(x)g(y), \quad x, y \in G,$$


where K is a compact subgroup of morphisms of G , dk is a normalized Haar measure of K , μ is a complex K -invariant measure with compact support, the functions f, g are continuous on G and f is assumed to satisfies the Kannappan type condition $K(\mu)$

$$\int_G \int_G f(ztxsy) d\mu(t) d\mu(s) = \int_G \int_G f(ztysx) d\mu(t) d\mu(s), \quad x, y, z \in G.$$

The paper of Székelyhidi [30] is the essential motivation for the present work and the methods used here are closely related to and inspired by those in [30]. The concept of the generalized Hyers-Ulam stability of mappings was introduced in the subject of functional equations by Th. M. Rassias in [20].

2000 Mathematics Subject Classification: 39B42, 39B32.

Key words: Locally compact group, Functional equation, Hyers-Ulam stability, Superstability.



Hyers-Ulam stability of the Generalized Trigonometric Formulas

Ahmed Redouani,
Elhoucien Elqorachi and
Belaid Bouikhalene

Title Page

Contents



Go Back


Close

Quit

Page 2 of 29

Contents

1	Introduction	4
2	Notation and Preliminary Results	8
3	Stability of Equation (1.3)	13
4	Stability of Equation (1.4)	18
5	Superstability of Equation (1.5)	23
References		



Hyers-Ulam stability of the Generalized Trigonometric Formulas

Ahmed Redouani,
Elhoucien Elqorachi and
Belaid Bouikhalene

Title Page

Contents



Go Back

Close

Quit

Page 3 of 29

1. Introduction

The Hyers-Ulam stability problem for functional equation has its origin in the following question posed by S. Ulam [41] in 1940.

Given a group G and a metric group (G', d) and given a number $\varepsilon > 0$, does there exist a $\delta > 0$ such that, if $f : G \rightarrow G'$ satisfies the inequality

$$d(f(xy), f(x)f(y)) < \delta, \text{ for all } x, y \in G,$$

then a homomorphism $a : G \rightarrow G'$ exists such that

$$d(f(x), a(x)) < \varepsilon, \text{ for all } x \in G?$$


The first affirmative answer to Ulam's question for linear mappings came within a year when D. H. Hyers [8] proved the following result.

Theorem 1.1 ([8]). Let B and B' be Banach spaces and let $f : B \rightarrow B'$ be a function such that for some $\delta > 0$

$$\| f(x + y) - f(x) - f(y) \| \leq \delta, \text{ for all } x, y \in B.$$

Then there exists a unique additive function $\varphi : B \rightarrow B'$ such that $\| f(x) - \varphi(x) \| \leq \delta$, for all $x \in B$.

Furthermore, the continuity of f at a point $y \in B$ implies the continuity of φ on B . The continuity, for each $x \in B$, of the function $t \rightarrow f(tx)$, $t \in \mathbb{R}$, implies the homogeneity of φ .



Hyers-Ulam stability of the
Generalized Trigonometric
Formulas

Ahmed Redouani,
Elhoucien Elqorachi and
Belaid Bouikhalene

Title Page

Contents



Go Back

Close

Quit

Page 4 of 29

After Hyers's result a great number of papers on the subject have been published, generalizing Ulam's problem and Hyers's theorem in various directions. In 1951 D.G. Bourgin [3] treated this problem for additive mappings. In 1978, Th. M. Rassias [20] provided a remarkable generalization of Hyers's theorem, a fact which rekindled interest in the field of functional equations.

Theorem 1.2 ([20]). *Let $f : V \longrightarrow X$ be a mapping between Banach spaces and let $p < 1$ be fixed. If f satisfies the inequality*

$$\|f(x + y) - f(x) - f(y)\| \leq \theta(\|x\|^p + \|y\|^p)$$

for some $\theta \geq 0$ and for all $x, y \in V$ ($x, y \in V \setminus \{0\}$ if $p < 0$). Then there exists a unique additive mapping $T : V \longrightarrow X$ such that


$$\|f(x) - T(x)\| \leq \frac{2\theta}{|2 - 2^p|} \|x\|^p$$

for all $x \in V$ ($x \in V \setminus \{0\}$ if $p < 0$).

If, in addition, $f(tx)$ is continuous in t for each fixed x , then T is linear.

This theorem of Th. M. Rassias stimulated several mathematicians working in the theory of functional equations to investigate this kind of stability for a variety of significant functional equations. By taking into consideration the influence of S. M. Ulam, D. H. Hyers and Th. M. Rassias on the study of stability problems of functional equations in mathematical analysis, the stability phenomenon that was proved by Th. M. Rassias is called the Hyers-Ulam-Rassias stability.

The Hyers-Ulam-Rassias stability was taken up by a number of mathematicians and the study of this area has the grown to be one of the central subjects in



Hyers-Ulam stability of the Generalized Trigonometric Formulas

Ahmed Redouani,
Elhoucien Elqorachi and
Belaid Bouikhalene

Title Page

Contents



Go Back

Close

Quit

Page 5 of 29

the mathematical analysis area. For more information, we can see for examples ([3], [7], [8], [10], [12], ..., [40]) and the monographs [4], [9], [11] by D. H. Hyers, G. Isac and Th. M. Rassias, by S.-M. Jung and by S. Czerwik (ed.).

L. Székelyhidi in [30], studied the stability property of two well known functional equations: The sine and cosine functional equations

$$(1.1) \quad f(xy) = f(x)g(y) + f(y)g(x), \quad x, y \in G$$

and

$$(1.2) \quad f(xy) = f(x)f(y) - g(x)g(y), \quad x, y \in G,$$


where f, g are complex-valued functions on an amenable group G . More precisely, he proved that if $f, g : G \rightarrow \mathbb{C}$ are given functions, G is an amenable group, and the function $(x, y) \rightarrow f(xy) - f(x)g(y) - f(y)g(x)$ is bounded, then there exists a solution (f_0, g_0) of (1.1) such that $f - f_0$ and $g - g_0$ are bounded. An analogous result holds for equation (1.2).

The aim of the present paper is to extend the Székelyhidi's results [30] to the functional equations

$$(1.3) \quad \int_G \int_K f(xtk \cdot y) dk d\mu(t) = f(x)g(y) + g(x)f(y), \quad x, y \in G$$

and

$$(1.4) \quad \int_G \int_K f(xtk \cdot y) dk d\mu(t) = f(x)f(y) + g(x)g(y), \quad x, y \in G,$$



Hyers-Ulam stability of the Generalized Trigonometric Formulas

Ahmed Redouani,
Elhoucien Elqorachi and
Belaid Bouikhalene

Title Page

Contents



Go Back

Close

Quit

Page 6 of 29


where K is a compact subgroup of $Mor(G)$, μ is a complex K -invariant measure with compact support, f, g are continuous functions on G and f is assumed to satisfy the Kannappan type condition $K(\mu)$

$$\int_G \int_G f(ztxsy)d\mu(t)d\mu(s) = \int_G \int_G f(ztysx)d\mu(t)d\mu(s), \quad x, y, z \in G.$$

Furthermore, in the last subsection we study a superstability result of the generalized quadratical functional equation

$$(1.5) \quad \int_G \int_K f(xtk \cdot y)dkd\mu(t) = f(x) + f(y), \quad x, y \in G.$$

The result can be viewed as a generalization of the ones obtained by G. Maksa and Z. Páles in [12].



Hyers-Ulam stability of the Generalized Trigonometric Formulas

Ahmed Redouani,
Elhoucien Elqorachi and
Belaid Bouikhalene

Title Page

Contents



Go Back

Close

Quit

Page 7 of 29

2. Notation and Preliminary Results

Our notation is described in the following Set Up and it will be used throughout the paper.

Set-Up. We let G be a locally compact group, $C(G)$ (resp. $C_b(G)$) the complex algebra of all continuous (resp. continuous and bounded) complex valued functions on G . $M(G)$ denotes the topological dual of $C_0(G)$: the Banach space of continuous functions vanishing at infinity. We let K be a compact subgroup of the group $\text{Mor}(G)$ of all mappings k of G onto itself that are either automorphisms and homeomorphisms (i.e. $k \in K^+$), or anti-automorphisms and homeomorphisms (i.e. $k \in K^-$). The action of $k \in K$ on $x \in G$ will be denoted by $k \cdot x$ and the normalized Haar measure on K by dk .


For any function f on G , we put $(k \cdot f)(x) = f(k^{-1} \cdot x)$. For any $\mu \in M(G)$, $k \in K$ and any $f \in C_b(G)$, we put $\langle k \cdot \mu, f \rangle = \langle \mu, k \cdot f \rangle$, and we say that μ is K -invariant if $k \cdot \mu = \mu$, for all $k \in K$.

A non-zero function $\phi \in C_b(G)$ is said to be a solution of Badora's functional equation if it satisfies

$$(2.1) \quad \int_K \int_G \phi(xtk \cdot y) d\mu(t) dk = \phi(x)\phi(y), \quad x, y \in G.$$

Recently the functional equation (2.1) was completely solved in abelian groups by Badora [2] and by E. Elqorachi, M. Akkouchi, A. Bakali, and B. Bouikhalene [6] in non-abelian groups and the Hyers-Ulam-Rassias stability of this equation was investigated in [1] and [5].

In the following, we prove some lemmas that we need later.



**Hyers-Ulam stability of the
Generalized Trigonometric
Formulas**

Ahmed Redouani,
Elhoucien Elqorachi and
Belaid Bouikhalene

Title Page

Contents



Go Back

Close

Quit

Page 8 of 29

Lemma 2.1. Let K be a compact subgroup of $\text{Mor}(G)$. Let μ be a K -invariant bounded measure on G . If $f \in C_b(G)$ satisfies the Kannappan condition $K(\mu)$:

$$\int_G \int_G f(ztxsy) d\mu(t) d\mu(s) = \int_G \int_G f(ztysx) d\mu(t) d\mu(s), \quad x, y, z \in G,$$

then we have

$$\begin{aligned} \int_K \int_K \int_G \int_G f(zsk \cdot (xtk' \cdot y)) dk dk' d\mu(s) d\mu(t) \\ = \int_K \int_K \int_G \int_G f(zsk \cdot xtk' \cdot y) dk dk' d\mu(s) d\mu(t), \end{aligned}$$


for all $x, y, z \in G$.

Proof. Let $x, y, z \in G$. Let $f \in C_b(G)$ be a complex function such that f satisfies $K(\mu)$. Then

$$\begin{aligned} \int_K \int_K \int_G \int_G f(zsk \cdot (xtk' \cdot y)) dk dk' d\mu(s) d\mu(t) \\ = \int_{K^+} \int_K \int_G \int_G f(zsk \cdot xk \cdot t(kk') \cdot y) dk dk' d\mu(s) d\mu(t) \\ + \int_{K^-} \int_K \int_G \int_G f(zs(kk') \cdot yk \cdot tk \cdot x) dk dk' d\mu(s) d\mu(t). \end{aligned}$$

Since μ is K -invariant and dk' is invariant by translation, then we get

$$\int_{K^+} \int_K \int_G \int_G f(zsk \cdot xk \cdot t(kk') \cdot y) dk dk' d\mu(s) d\mu(t)$$



Hyers-Ulam stability of the Generalized Trigonometric Formulas

Ahmed Redouani,
Elhoucien Elqorachi and
Belaid Bouikhalene

Title Page

Contents



Go Back

Close

Quit

Page 9 of 29

$$= \int_{K^+} \int_K \int_G \int_G f(zsk \cdot xtk' \cdot y) dkdk' d\mu(s) d\mu(t),$$

$$\begin{aligned} & \int_{K^-} \int_K \int_G \int_G f(zs(kk') \cdot yk \cdot tk \cdot x) dkdk' d\mu(s) d\mu(t) \\ &= \int_{K^-} \int_K \int_G \int_G f(zsk' \cdot ytk \cdot x) dkdk' d\mu(s) d\mu(t) \\ &= \int_{K^-} \int_K \int_G \int_G f(zsk \cdot xtk' \cdot y) dkdk' d\mu(s) d\mu(t), \end{aligned}$$


because f satisfies $K(\mu)$.

Consequently,

$$\begin{aligned} & \int_K \int_K \int_G \int_G f(zsk \cdot (xtk' \cdot y)) dkdk' d\mu(s) d\mu(t) \\ &= \int_{K^+} \int_K \int_G \int_G f(zsk \cdot xtk' \cdot y) dkdk' d\mu(s) d\mu(t) \\ &\quad + \int_{K^-} \int_K \int_G \int_G f(zsk \cdot xtk' \cdot y) dkdk' d\mu(s) d\mu(t) \\ &= \int_K \int_K \int_G \int_G f(zsk \cdot xtk' \cdot y) dkdk' d\mu(s) d\mu(t). \end{aligned}$$

This completes the proof. \square

The following result is a generalization of the lemma obtained by G. Maksa and Z. Páles in [12].



Hyers-Ulam stability of the Generalized Trigonometric Formulas

Ahmed Redouani,
Elhoucien Elqorachi and
Belaid Bouikhalene

Title Page

Contents



Go Back

Close

Quit

Page 10 of 29

Lemma 2.2. Let K be a compact subgroup of $\text{Mor}(G)$. Let μ be a K -invariant bounded measure on G such that $\langle \mu, 1_G \rangle = 1$. Let $f \in C_b(G)$ be a complex function which satisfies $K(\mu)$, then the continuous and bounded function


$$(2.2) \quad L(x, y) = f(x) + f(y) - \int_G \int_K f(xtk \cdot y) dk d\mu(t), \quad x, y \in G$$

satisfies the functional equation

$$(2.3) \quad L(x, y) + \int_G \int_K L((xtk \cdot y), z) dk d\mu(t) \\ = L(y, z) + \int_G \int_K L(x, (ytk \cdot z)) dk d\mu(t), \quad x, y, z \in G.$$

Proof. The proof is closely related to the computation in ([12, Section 2, Lemma]), where K is a finite subgroup of $\text{Aut}(G)$ and $\mu = \delta_e$. Let f be a bounded and continuous function on G which satisfies the Kannappan condition $K(\mu)$ and let $L(x, y)$ be the function defined by (2.2), then we have

$$L(x, y) + \int_K \int_G L((xtk \cdot y), z) dk d\mu(t) \\ = f(x) + f(y) - \int_K \int_G f(xtk \cdot y) dk d\mu(t) \\ + \int_K \int_G f(xtk \cdot y) dk d\mu(t) + \langle \mu, 1_G \rangle \langle dk, 1_K \rangle f(z) \\ - \int_K \int_G \int_K \int_G f(xtk \cdot ysk' \cdot z) dk dk' d\mu(s) d\mu(t) \\ = f(x) + f(y) + f(z) - \int_K \int_G \int_K \int_G f(xtk \cdot ysk' \cdot z) dk dk' d\mu(s) d\mu(t).$$



Hyers-Ulam stability of the Generalized Trigonometric Formulas

Ahmed Redouani,
Elhoucien Elqorachi and
Belaid Bouikhalene

Title Page

Contents



Go Back

Close


Quit

Page 11 of 29

On the other hand, we have

$$\begin{aligned}
 & L(y, z) + \int_K \int_G L(x, (ytk \cdot z)) dk d\mu(t) \\
 &= f(y) + f(z) - \int_K \int_G f(ytk \cdot z) dk d\mu(t) \\
 &\quad + \langle \mu, 1_G \rangle \langle dk, 1_K \rangle f(x) + \int_K \int_G f(ytk \cdot z) dk d\mu(t) \\
 &\quad - \int_K \int_G \int_K \int_G f(xsk' \cdot (ytk \cdot z)) dk dk' d\mu(s) d\mu(t) \\
 &= f(y) + f(z) + f(x) - \int_K \int_G \int_K \int_G f(xsk' \cdot ytk \cdot z) dk dk' d\mu(s) d\mu(t).
 \end{aligned}$$

This ends the proof of Lemma 2.2. □



**Hyers-Ulam stability of the
Generalized Trigonometric
Formulas**

Ahmed Redouani,
Elhoucien Elqorachi and
Belaid Bouikhalene

Title Page

Contents



Go Back

Close

Quit

Page 12 of 29

3. Stability of Equation (1.3)

In this section, we investigate the stability properties of the functional equation (1.3), it's a generalization of the stability of equation (1.1) proved by Székelyhidi in [30].


Theorem 3.1. *Let K be a compact subgroup of $\text{Mor}(G)$, and let μ be a K -invariant measure with compact support. Let f, g be continuous complex-valued functions such that f satisfies $K(\mu)$ and the following function*

$$(3.1) \quad (G, G) \ni (x, y) \longrightarrow \int_K \int_G f(xtk \cdot y) dk d\mu(t) - f(x)g(y) - f(y)g(x)$$

is bounded. Then

- i) $f = 0$, g arbitrary in $C(G)$ or
- ii) f, g are bounded or
- iii) f is unbounded, g is a bounded solution of Badora's equation or
- iv) There exists φ a solution of Badora's equation, there exists b a continuous bounded function on G and $\gamma \in \mathbb{C}$ such that $f = \gamma(\varphi - b)$ and $g = \frac{\varphi + b}{2}$ or
- v) f, g are solutions of (1.3).

Proof. If $f = 0$, then g can be chosen arbitrarily in $C(G)$. This is case (i). If $f \neq 0$ is bounded, then the function $G \ni x \longmapsto f(x)g(y) + f(y)g(x)$ is



Hyers-Ulam stability of the Generalized Trigonometric Formulas

Ahmed Redouani,
Elhoucien Elqorachi and
Belaid Bouikhalene

Title Page

Contents



Go Back

Close

Quit

Page 13 of 29

bounded for all $y \in G$, so g is bounded. This is case (ii). If f is unbounded and g is bounded, the function

$$G \ni x \longmapsto \int_K \int_G f(xtk \cdot y) dk d\mu(t) - f(x)g(y)$$

is bounded, for all $y \in G$. In view of [5, Theorem 3.1], we get that g is a solution of Badora's equation. This is case (iii). If f, g are unbounded functions, we distinguish two cases:

First case. We assume that there exist $\alpha, \beta \in \mathbb{C} \setminus \{0\}$ such that $\alpha f + \beta g$ is bounded, then g can be written as $g = \frac{f}{2^\gamma} + b$, where b is a bounded function and $\gamma \in \mathbb{C} \setminus \{0\}$. Consequently, the function

$$G \ni x \longmapsto \int_K \int_G f(xtk \cdot y) dk d\mu(t) - \left(\frac{f(y)}{\gamma} + b(y) \right) f(x)$$


is bounded, for all $y \in G$. Hence by [5, Theorem 3.1], it follows that $\varphi(y) = \frac{f(y)}{\gamma} + b(y)$ is a solution of Badora's equation. This is case (iv).

Second case. For all $\alpha, \beta \in \mathbb{C} \setminus \{0\}$, $\alpha f + \beta g$ is an unbounded function on G . In this case we shall prove that f, g are solutions of (1.3). The idea of the proof is closely inspired by some good computations used in [30, Lemma 2.2]. We define the mapping

$$F(x, y) = \int_K \int_G f(xtk \cdot y) dk d\mu(t) - f(x)g(y) - f(y)g(x), \quad x, y \in G$$

and we will prove that $F(x, y) = 0$, for all $x, y \in G$. By assumption, there exist $\gamma, \delta, \lambda \in \mathbb{C}$ and $a \in G$ such that

$$(3.2) \quad g(x) = \gamma f(x) + \delta \int_K \int_G f(xtk \cdot a) dk d\mu(t) + \lambda F(x, a), \quad x \in G.$$



Hyers-Ulam stability of the Generalized Trigonometric Formulas

Ahmed Redouani,
Elhoucien Elqorachi and
Belaid Bouikhalene

Title Page

Contents



Go Back

Close

Quit


Page 14 of 29

For all $x, y, z \in G$, we have

$$\begin{aligned}
 & \int_K \int_K \int_G \int_G f((xtk \cdot y)sk' \cdot z)dkdk'd\mu(t)d\mu(s) \\
 &= g(z) \int_K \int_G f(xtk \cdot y)dkd\mu(t) + f(z) \int_K \int_G g(xtk \cdot y)dkd\mu(t) \\
 & \quad + \int_K \int_G F((xtk \cdot y), z)dkd\mu(t) \\
 &= g(z)f(x)g(y) + g(z)f(y)g(x) + g(z)F(x, y) + \gamma f(z) \int_K \int_G f(xtk \cdot y)dkd\mu(t) \\
 & \quad + \delta f(z) \int_K \int_K \int_G \int_G f(xtk \cdot ysk' \cdot a)dkdk'd\mu(s)d\mu(t) \\
 & \quad + \lambda f(z) \int_K \int_G F((xtk \cdot y), a)dkd\mu(t) + \int_K \int_G F((xtk \cdot y), z)dkd\mu(t).
 \end{aligned}$$

In view of Lemma 2.1, we get

$$\begin{aligned}
 & \int_K \int_K \int_G \int_G f((xtk \cdot y)sk' \cdot z)dkdk'd\mu(t)d\mu(s) \\
 &= g(z)f(x)g(y) + g(z)f(y)g(x) \\
 & \quad + g(z)F(x, y) + \gamma f(z) \int_K \int_G f(xtk \cdot y)dkd\mu(t) \\
 & \quad + \delta f(z) \int_K \int_K \int_G \int_G f(xtk \cdot ysk' \cdot a)dkdk'd\mu(s)d\mu(t) \\
 & \quad + \lambda f(z) \int_K \int_G F((xtk \cdot y), a)dkd\mu(t) + \int_K \int_G F((xtk \cdot y), z)dkd\mu(t)
 \end{aligned}$$



Hyers-Ulam stability of the Generalized Trigonometric Formulas

Ahmed Redouani,
Elhoucien Elqorachi and
Belaid Bouikhalene

Title Page

Contents



Go Back

Close

Quit

Page 15 of 29


$$\begin{aligned}
&= g(z)f(x)g(y) + g(z)f(y)g(x) + g(z)F(x, y) + \gamma f(z)f(x)g(y) \\
&\quad + \gamma f(z)f(y)g(x) + \gamma f(z)F(x, y) + \delta f(z)f(x) \int_K \int_G g(y sk' \cdot a) dk' d\mu(s) \\
&\quad + \delta f(z)g(x) \int_K \int_G f(y sk' \cdot a) dk' d\mu(s) + \delta f(z) \int_K \int_G F(x, y sk' \cdot a) dk' d\mu(s) \\
&\quad + \lambda f(z) \int_K \int_G F((x tk \cdot y), a) dk d\mu(t) + \int_K \int_G F((x tk \cdot y), z) dk d\mu(t).
\end{aligned}$$

By using again Lemma 2.1, we obtain

$$\begin{aligned}
&\int_K \int_K \int_G \int_G f((x tk \cdot y) sk' \cdot z) dk dk' d\mu(t) d\mu(s) \\
&\quad = \int_K \int_K \int_G \int_G f(x tk \cdot (y sk' \cdot z)) dk dk' d\mu(t) d\mu(s) \\
&\quad = f(x) \int_K \int_G g(y sk' \cdot z) dk' d\mu(s) + g(x) \int_K \int_G f(y sk' \cdot z) dk' d\mu(s) \\
&\quad\quad + \int_K \int_G F(x, (y sk' \cdot z)) dk' d\mu(s).
\end{aligned}$$

Hence, it follows that

$$\begin{aligned}
&f(x) \left[g(y)g(z) + \gamma g(y)f(z) + \delta f(z) \int_K \int_G g(y sk' \cdot a) dk' d\mu(s) \right. \\
&\quad \left. - \int_K \int_G g(y sk' \cdot z) dk' d\mu(s) \right] + g(x) \left[f(y)g(z) + \gamma f(y)f(z) \right. \\
&\quad \left. + \delta f(z) \int_K \int_G f(y sk' \cdot a) dk' d\mu(s) - \int_K \int_G f(y sk' \cdot z) dk' d\mu(s) \right]
\end{aligned}$$



Hyers-Ulam stability of the Generalized Trigonometric Formulas

Ahmed Redouani,
Elhoucien Elqorachi and
Belaid Bouikhalene

Title Page

Contents



Go Back

Close

Quit


Page 16 of 29

$$\begin{aligned}
&= \int_K \int_G F(x, (ysk' \cdot z)) dk' d\mu(s) - g(z)F(x, y) - \gamma f(z)F(x, y) \\
&\quad - \delta f(z) \int_K \int_G F(x, (ysk' \cdot a)) dk' d\mu(s) \\
&\quad - \lambda f(z) \int_K \int_G F((xtk \cdot y), a) dk d\mu(t) \\
&\quad - \int_K \int_G F((xtk \cdot y), z) dk' d\mu(s).
\end{aligned}$$

Since the right-hand side is bounded as a function of x for all fixed $y, z \in G$, then we get

$$\begin{aligned}
&g(z)F(x, y) + f(z) \left[\gamma F(x, y) + \delta \int_K \int_G F(x, ysk' \cdot a) dk' d\mu(s) \right. \\
&\quad \left. + \lambda \int_K \int_G F((xtk \cdot y), a) dk d\mu(t) \right] \\
&= \int_K \int_G F(x, (ysk' \cdot z)) dk' d\mu(s) - \int_K \int_G F((xtk \cdot y), z) dk d\mu(t).
\end{aligned}$$

Since the right-hand side is bounded as a function of z for all fixed $x, y \in G$, then we obtain $F(x, y) = 0$, for all $x, y \in G$. This is case (v) and the proof of Theorem 3.1 is completed. \square



Hyers-Ulam stability of the Generalized Trigonometric Formulas

Ahmed Redouani,
Elhoucien Elqorachi and
Belaid Bouikhalene

Title Page

Contents



Go Back

Close

Quit

Page 17 of 29

4. Stability of Equation (1.4)

In this section, we study the problem of the Hyers-Ulam stability of equation (1.4). It is a generalization of the stability of equation (1.2) proved by Székelyhidi in [30].

Theorem 4.1. *Let K be a compact subgroup of $\text{Mor}(G)$, let μ be a K -invariant measure with compact support. Let f, g be continuous complex-valued functions such that f satisfies $K(\mu)$ and the function*

$$(4.1) \quad (G, G) \ni (x, y) \longrightarrow \int_K \int_G f(xtk \cdot y) dk d\mu(t) - f(x)f(y) + g(x)g(y)$$


is bounded. Then,

- i) f, g are bounded or*
- ii) f is a solution of Badora's equation, g is bounded, or*
- iii) f, g are unbounded, $f + g$ or $f - g$ are bounded solutions of Badora's equation or*
- iv) There exists φ a solution of Badora's equation, there exists b a continuous bounded function on G and $\gamma \in \mathbb{C} \setminus \{\pm 1\}$ such that*

$$f = \frac{\gamma^2 \varphi - b}{\gamma^2 - 1}, \quad g = \frac{\gamma}{\gamma^2 - 1} (\varphi - b)$$

or

- v) f, g are solutions of (1.4).*



Hyers-Ulam stability of the Generalized Trigonometric Formulas

Ahmed Redouani,
Elhoucien Elqorachi and
Belaid Bouikhalene

Title Page

Contents



Go Back

Close

Quit

Page 18 of 29

Proof. If g is bounded, then we obtain that the function

$$G \times G \ni (x, y) \mapsto \int_K \int_G f(xtk \cdot y) dk d\mu(t) - f(x)f(y)$$

is bounded. So by [5, Theorem 3.1], we have either f is bounded or f is a solution of Badora's equation. This is cases (i) and (ii). If g is unbounded, then f is unbounded. As in the preceding proof, we distinguish two cases.

First case. Assume that there exist $\alpha, \beta \in \mathbb{C} \setminus \{0\}$ such that $\alpha f + \beta g$ is a bounded function on G , then there exists a constant $\gamma \in \mathbb{C} \setminus \{0\}$ such that $f = \gamma g + b$, where b is a bounded function on G . Hence the function

$$G \ni x \mapsto \int_K \int_G g(xtk \cdot y) dk d\mu(t) - \frac{(\gamma^2 - 1)g(y) + \gamma b(y)}{\gamma} g(x)$$


is bounded for all $y \in G$. It follows from [5, Theorem 3.1] that $\varphi = \frac{\gamma^2 - 1}{\gamma} g + b$ is a solution of Badora's equation. Hence, we obtain case (iii) for $\gamma^2 = 1$ and (iv) for $\gamma^2 \neq 1$.

Second case. For all $\alpha, \beta \in \mathbb{C} \setminus \{0\}$, $\alpha f + \beta g$ is an unbounded function on G . We put

$$H(x, y) = \int_K \int_G f(xtk \cdot y) dk d\mu(t) - f(x)f(y) + g(x)g(y), \quad x, y \in G$$

and follow some computation used by Székelyhidi in [30]. There exists $\gamma, \delta, \lambda \in \mathbb{C}$ and $a \in G$ such that

$$g(x) = \gamma f(x) + \delta \int_K \int_G f(xtk \cdot a) dk d\mu(t) + \lambda H(x, a), \quad x \in G.$$



Hyers-Ulam stability of the Generalized Trigonometric Formulas

Ahmed Redouani,
Elhoucien Elqorachi and
Belaid Bouikhalene

Title Page

Contents



Go Back

Close

Quit


Page 19 of 29

Now, for all $x, y, z \in G$, we get

$$\begin{aligned}
 & \int_K \int_K \int_G \int_G f((xsk \cdot y)tk' \cdot z)dkdk'd\mu(t)d\mu(s) \\
 &= f(z) \int_K \int_G f(xsk \cdot y)dkd\mu(s) - g(z) \int_K \int_G g(xsk \cdot y)dkd\mu(s) \\
 & \quad + \int_K \int_G H((xsk \cdot y), z)dkd\mu(s) \\
 &= f(x)f(y)f(z) - g(x)g(y)f(z) + f(z)H(x, y) - \gamma f(x)f(y)g(z) \\
 & \quad + \gamma g(x)g(y)g(z) - \gamma g(z)H(x, y) - \delta g(z)f(x) \int_K \int_G f(ytk' \cdot a)dk'd\mu(t) \\
 & \quad + \delta g(x)g(z) \int_K \int_G g(ytk' \cdot a)dk'd\mu(t) - \delta g(z) \int_K \int_G H(x, (ytk' \cdot a))dk'd\mu(t) \\
 & \quad - \lambda g(z) \int_K \int_G H((xsk \cdot y), a)dkd\mu(s) + \int_K \int_G H((xsk \cdot y), z)dkd\mu(s).
 \end{aligned}$$

On the other hand, we have

$$\begin{aligned}
 & \int_K \int_K \int_G \int_G f(xsk \cdot ytk' \cdot z)dkdk'd\mu(t)d\mu(s) \\
 &= \int_K \int_K \int_G \int_G f(xsk \cdot (ytk' \cdot z))dkdk'd\mu(t)d\mu(s) \\
 &= f(x) \int_K \int_G f(ytk' \cdot z)dk'd\mu(t) \\
 & \quad - g(x) \int_K \int_G g(ytk' \cdot z)dk'd\mu(t) + \int_K \int_G H(x, (ytk' \cdot z))dk'd\mu(t).
 \end{aligned}$$



Hyers-Ulam stability of the Generalized Trigonometric Formulas

Ahmed Redouani,
Elhoucien Elqorachi and
Belaid Bouikhalene

Title Page

Contents



Go Back

Close

Quit


Page 20 of 29

Consequently, we obtain

$$\begin{aligned}
 & f(x) \left[f(y)f(z) - \gamma f(y)g(z) - \delta g(z) \int_K \int_G f(ytk' \cdot a) dk' d\mu(t) \right. \\
 & \left. - \int_K \int_G f(ytk' \cdot a) dk' d\mu(t) \right] - g(x) \left[g(y)f(z) - \gamma g(y)g(z) \right. \\
 & \left. - \delta g(z) \int_K \int_G g(ytk' \cdot a) dk' d\mu(t) - \int_K \int_G g(ytk' \cdot z) dk' d\mu(t) \right] \\
 & = \int_K \int_G H(x, (ytk' \cdot z)) dk' d\mu(t) - f(z)H(x, y) + \gamma g(z)H(x, y) \\
 & \quad + \delta g(z) \int_K \int_G H(x, (ytk' \cdot a)) dk' d\mu(t) \\
 & \quad + \lambda g(z) \int_K \int_G H((xsk \cdot y), a) dk d\mu(s) \\
 & \quad - \int_K \int_G H((xsk \cdot y), z) dk d\mu(s).
 \end{aligned}$$

Since the right hand side is bounded as a function of x for all fixed $y, z \in G$, then we get

$$\begin{aligned}
 & f(z)[-H(x, y)] + g(z) \left[\gamma H(x, y) + \delta \int_K \int_G H(x, (ytk' \cdot a)) dk' d\mu(t) \right. \\
 & \quad \left. + \lambda \int_K \int_G H((xsk \cdot y), a) dk d\mu(s) \right] \\
 & = \int_K \int_G H(xsk \cdot y, z) dk d\mu(s) - \int_K \int_G H(x, (ytk' \cdot z)) dk' d\mu(t).
 \end{aligned}$$



Hyers-Ulam stability of the Generalized Trigonometric Formulas

Ahmed Redouani,
Elhoucien Elqorachi and
Belaid Bouikhalene

Title Page

Contents




Go Back

Close

Quit

Page 21 of 29

Since the right-hand side is bounded as a function of z for all fixed $x, y \in G$, we conclude that $H(x, y) = 0$, for all $x, y \in G$, which is case (v). This ends the proof of the theorem. \square



**Hyers-Ulam stability of the
Generalized Trigonometric
Formulas**

Ahmed Redouani,
Elhoucien Elqorachi and
Belaid Bouikhalene

Title Page

Contents



Go Back

Close

Quit

Page 22 of 29

5. Superstability of Equation (1.5)

In this subsection, we study a superstability of the functional equation

$$(5.1) \quad \int_K \int_G f(xtk \cdot y) dk d\mu(t) = f(x) + f(y) \quad x, y \in G.$$

Theorem 5.1. *Let μ be a K -invariant measure with compact support. Let $\delta : G \times G \mapsto \mathbb{R}^+$ be an arbitrary function and assume that there exists a sequence $(u_n) \in G$ such that*

$$\lim_{n \rightarrow +\infty} \delta(u_n x, y) = 0, \text{ for all } x, y \in G \text{ (uniform convergence).}$$

Let $f : G \mapsto \mathbb{C}$ be a continuous function, which satisfies the Kannappan type condition $K(\mu)$. If f satisfies the inequality


$$(5.2) \quad \left| \int_K \int_G f(xtk \cdot y) dk d\mu(t) - f(x) - f(y) \right| \leq \delta(x, y),$$

for all $x, y \in G$,

then f is a solution of equation (5.1).

Proof. Assume that $f \in C(G)$ is such that f satisfies $K(\mu)$ and inequality (5.2). It follows that there exists a sequence u_n such that $\lim_{n \rightarrow +\infty} L(u_n x, y) = 0$ (uniformly). Now, by Lemma 2.2, we get

$$(5.3) \quad \begin{aligned} L(u_n x, y) + \int_G \int_K L((u_n xtk \cdot y), z) dk d\mu(t) \\ = L(y, z) + \int_G \int_K L(u_n x, (yt \cdot z)) dk d\mu(t), \end{aligned}$$



Hyers-Ulam stability of the
Generalized Trigonometric
Formulas

Ahmed Redouani,
Elhoucien Elqorachi and
Belaid Bouikhalene

Title Page

Contents



Go Back


Close

Quit

Page 23 of 29

for all $x, y, z \in G$ and $n \in \mathbb{N}$. By letting $n \rightarrow +\infty$, we deduce the desired result and the proof of the theorem is complete. \square

Remark 1. *If K is a compact subgroup of $\text{Aut}(G)$, the condition $K(\mu)$ is not necessary.*



Hyers-Ulam stability of the Generalized Trigonometric Formulas

Ahmed Redouani,
Elhoucien Elqorachi and
Belaid Bouikhalene

Title Page

Contents



Go Back


Close

Quit

Page 24 of 29

References

- [1] R. BADORA, On Hyers-Ulam stability of Wilson's functional equation., *Aequationes Math.*, **60** (2000), 211–218.
- [2] R. BADORA, On a joint generalization of Cauchy's and d'Alembert's functional equations, *Aequationes Math.*, **43** (1992), 72–89.
- [3] D.G. BOURGIN, Classes of transformations and bordering transformations, *Bull. Amer. Math. Soc.*, **57** (1951), 223–237.
- [4] S. CZERWIK (Ed.), *Stability of Functional Equations of Ulam-Hyers-Rassias Type*, Hadronic Press, Inc., Palm Harbor, Florida, 2003.
- [5] E. EQORACHI, The stability of the generalized form for the Cauchy and d'Alembert functional equations, (submitted).
- [6] E. EQORACHI, M. AKKOUCHE, A. BAKALI AND B. BOUIKHALENE, Badora's Equation on non-Abelian locally compact groups, *Georgian Math. J. V.*, **11**(3) (2004), 449–466.
- [7] V.A. FAIZEV, Th.M. RASSIAS AND P.K. SAHOO, The space of (ψ, γ) -additive mappings on semigroups, *Transactions of the Amer. Math. Soc.*, **354**(11) (2002), 4455–4472.
- [8] D.H. HYERS, On the stability of the linear functional equation, *Proc. Nat. Acad. Sci. U.S.A.*, **27** (1941), 222–224.
- [9] D.H. HYERS, G. ISAC AND Th.M. RASSIAS, *Stability of Functional Equations in Several Variables*, (Birkhauser, Basel), 1998.



Hyers-Ulam stability of the
Generalized Trigonometric
Formulas

Ahmed Redouani,
Elhoucien Elqorachi and
Belaid Bouikhalene

Title Page

Contents




Go Back

Close

Quit

Page 25 of 29

- [10] D.H. HYERS AND Th.M. RASSIAS, Approximate homomorphisms, *Aequationes Math.*, **44** (1992), 125–153.
- [11] S.-M. JUNG, *Hyers-Ulam-Rassias Stability of Functional Equations in Mathematical Analysis*, Hardonic Press, Inc., Palm Harbor, Florida, 2001.
- [12] G. MAKSA AND Z. PÁLES, Hyperstability of a class of linear functional equations, *Acta Math. Acad. Paedagogicae Nyíregyháziensis*, **17** (2001), 107–112.
- [13] C.-G. PARK, On an approximate automorphism on a C^* -algebra, *Proc. Amer. Math. Soc.*, **132** (2004), 1739–1745.
- [14] C.-G. PARK, Homomorphisms between Lie JC^* -algebras and Cauchy-Rassias stability of Lie JC^* -algebras derivations, *J. Lie Theory*, **15** (2005), 393–414.
- [15] C.-G. PARK, Cauchy-Rassias stability of a generalized Trif's mapping in Banach modules and its applications, *Nonlinear Analysis- TMA*, **62**(2005), 595–613.
- [16] J.M. RASSIAS, On approximation of approximately linear mappings by linear mappings, *J. Funct. Anal.*, **46** (1982), 126–130.
- [17] J.M. RASSIAS, On approximation of approximately linear mappings by linear mappings, *Bull. Sci. Math.*, **108**(2) (1984), 445–446.
- [18] Th.M. RASSIAS, On the stability of mappings, *Rend. Sem. Mat. Fis. Milano*, **58** (1988), 91–99.



**Hyers-Ulam stability of the
Generalized Trigonometric
Formulas**

Ahmed Redouani,
Elhoucien Elqorachi and
Belaid Bouikhalene

Title Page

Contents




Go Back

Close

Quit

Page 26 of 29

- [19] Th.M. RASSIAS, On a modified Hyers-Ulam sequence, *J. Math. Anal. Appl.*, **158** (1991), 106–113.
- [20] Th.M. RASSIAS, On the stability of the linear mapping in Banach spaces, *Proc. Amer. Math. Soc.*, **72** (1978), 297–300.
- [21] Th.M. RASSIAS, The problem of S. M. Ulam for approximately multiplicative mappings, *J. Math. Anal. Appl.*, **246** (2000), 352–378.
- [22] Th.M. RASSIAS, On the stability of functional equations and a problem of Ulam, *Acta Applicandae Math.*, **62**(2000), 23–130.
- [23] Th.M. RASSIAS, On the stability of functional equations in Banach spaces, *J. Math. Anal. Appl.*, **251** (2000), 264–284.
- [24] Th.M. RASSIAS AND P. ŠEMREL, On the behaviour of mappings which do not satisfy Hyers stability, *Proc. Amer. Math. Soc.*, **114** (1992), 989–993.
- [25] Th.M. RASSIAS AND P. ŠEMREL, On the Hyers-Ulam stability of linear mappings, *J. Math. Anal. Appl.*, **173** (1993), 325–338.
- [26] J. SCHWAIGER, The functional equation of homogeneity and its stability properties, *Österreich. Akad. Wiss. Math.-Natur, Kl, Sitzungsber. Abt., II* (1996), 205, 3–12.
- [27] L. SZÉKELYHIDI, On a theorem of Baker Lawrence and Zorzitto, *Proc. Amer. Math. Soc.*, **84** (1982), 95–96.
- [28] L. SZÉKELYHIDI, On a stability theorem, *C.R. Math. Rep. Acad. Sci. Canada*, **3** (1981), 253–255.



**Hyers-Ulam stability of the
Generalized Trigonometric
Formulas**

Ahmed Redouani,
Elhoucien Elqorachi and
Belaid Bouikhalene

Title Page

Contents




Go Back

Close

Quit

Page 27 of 29

- [29] L. SZÉKELYHIDI, The stability of d'Alembert-type functional equations, *Acta Sci. Math. (Szeged)*, **44** (1982), 313–320.
- [30] L. SZÉKELYHIDI, The stability of the sine and cosine functional equations, *Proc. Amer. Math. Soc.*, **110** (1990), 109–115.
- [31] L. SZÉKELYHIDI, The stability of the of linear functional equations, *C.R. Math. Rep. Acad. Sci. Canada*, **3** (1981), 63–67.
- [32] L. SZÉKELYHIDI, Note on a stability theorem, *Canad. Math. Bull.*, **25** (1982), 500–501.
- [33] L. SZÉKELYHIDI, Remarks on Hyers's theorem, *Publ. Math. Debrecen*, **34** (1987), 131–135.
- [34] L. SZÉKELYHIDI, Fréchet's equation and Hyers theorem on noncommutative semigroups, *Ann. Polon. Math.*, **48** (1988), 183–189.
- [35] L. SZÉKELYHIDI, Stability of some functional equations in economics, *Rend. Sem. Mat. Fis. Milano*, **58** (1988), 169–176.
- [36] L. SZÉKELYHIDI, An abstract superstability theorem, *Abh. Math. Sem. Univ. Hamburg*, **59** (1989), 81–83.
- [37] L. SZÉKELYHIDI, Stability properties of functional equations in several variables. Manuscript.
- [38] J. TABOR, On function behaving like additive functions, *Aequationes. Math.*, **35** (1988), 164–185.



**Hyers-Ulam stability of the
Generalized Trigonometric
Formulas**

Ahmed Redouani,
Elhoucien Elqorachi and
Belaid Bouikhalene

Title Page

Contents




Go Back

Close

Quit

Page 28 of 29

- [39] J. TABOR, Quasi-additive functions, *Aequationes. Math.*, **39** (1990), 179–197.
- [40] J. TABOR, Approximate endomorphisms of the complex field, *J. Natur. Geom.Math.*, **I** (1992), 71–86.
- [41] S.M. ULAM, *A Collection of Mathematical Problems*, Interscience Publ. New York, 1961. Problems in Modern Mathematics, Wiley, New York, 1964.



**Hyers-Ulam stability of the
Generalized Trigonometric
Formulas**

Ahmed Redouani,
Elhoucien Elqorachi and
Belaid Bouikhalene

Title Page

Contents



Go Back

Close

Quit

Page 29 of 29