

S.Lahrech, A.Jaddar, A.Ouahab, and A.Mbarki

**SOME REMARKS ABOUT STRICTLY PSEUDOCONVEX
FUNCTIONS WITH RESPECT TO THE
CLARKE-ROCKAFELLAR SUBDIFFERENTIAL**

(submitted by A. V. Lapin)

ABSTRACT. Using the notion of radially Clarke-Rockafellar subdifferentiable functions (RCRS-functions), we characterize strictly pseudoconvex functions with respect to the Clarke-Rockafellar subdifferential in two different ways, and we study a maximization problem involving RCRS-strictly pseudoconvex functions over a convex set.

1. INTRODUCTION

Generalized convexity has proved to be a good tool in the study of some economic problems and in mathematical programming. Strict pseudoconvexity is a kind of generalized convexity that appeared recently as an important part of the class of pseudoconvex functions. The former class has been characterized by many authors (see for instance [1, 2, 4, 7, 10]). In this paper we will refine these results in section 2, using the Clarke Rockafellar subdifferential. While, in section 3 we give a necessary and sufficient condition for a point to be a maximum of a strictly pseudoconvex function over a convex set.

Let us recall some definitions and well known results in connection with what we shall do in the sequel. By X we mean a Banach space and

Key words and phrases. Strictly pseudoconvex functions, RCRS-functions, strongly RCRS-functions.

2000 Mathematical Subject Classification. 49J52, 49J50.

by X^* its topological dual, while $\langle \cdot, \cdot \rangle$ is the duality pairing between X and X^* . For x and y in X , the closed segment $[x, y]$ is the set $[x, y] = \{x + t(y - x); t \in [0, 1]\}$. By $[x, y)$ we denote the set $[x, y] \setminus \{y\}$. Given a lower semi-continuous (l.s.c.) function $f : X \rightarrow R \cup \{+\infty\}$ whose domain

$$\text{dom} f = \{x \in X; f(x) < +\infty\}$$

is nonempty. The Clarke-Rockafellar generalized directional derivative $f^\uparrow(x, v)$ of f at $x \in \text{dom} f$ along the direction v is defined by:

$$f^\uparrow(x, v) = \sup_{\varepsilon > 0} \limsup_{y \rightarrow_f x, t \searrow 0} \inf_{u \in B(v, \varepsilon)} t^{-1} [f(y + tu) - f(y)], \quad (1)$$

where by $y \rightarrow_f x$, we mean $y \rightarrow x$ and $f(y) \rightarrow f(x)$. Here, by $B(v, \varepsilon)$ we denote the open ball centered at v with radius ε . The Clarke-Rockafellar subdifferential of f at $x \in \text{dom} f$ is defined by

$$\partial f(x) = \{x^* \in X^* : \langle x^*, v \rangle \leq f^\uparrow(x, v) \quad \forall v \in X\}.$$

We adopt the convention $\partial f(x) = \emptyset$ when $x \notin \text{dom} f$.

A function f is said to be quasiconvex if for any $x, y \in X$ and $\lambda \in [0, 1]$ we have

$$f(\lambda x + (1 - \lambda)y) \leq \max\{f(x), f(y)\}. \quad (2)$$

f is said to be strictly quasiconvex if the inequality (2) is strict when $x \neq y$. f is said to be pseudoconvex (with respect to the Clarke-Rockafellar subdifferential) if for any x and y in X the following implication holds:

$$(\exists x^* \in \partial f(x) : \langle x^*, y - x \rangle \geq 0) \implies f(x) \leq f(y). \quad (3)$$

The relation between pseudoconvexity and quasiconvexity has been described in [2, 4, 7, 10] by the following result.

Theorem 1. *Let $f : X \mapsto R \cup \{+\infty\}$ be a l.s.c. function. Consider the propositions:*

- i)** f is pseudoconvex.
- ii)** f is quasiconvex and $(0 \in \partial f(x) \implies x \text{ is a global minimum of } f)$.

Then i) implies ii). If moreover, f is radially continuous, then ii) implies i).

Generally, in generalized convexity, there is a close link between the kind of convexity of a function and a corresponding kind of monotonicity of its subdifferential. Recall that a multifunction $T : X \rightarrow X^*$ is said to be pseudomonotone if for any $x, y \in X$, we have:

$$[\exists x^* \in T(x) : \langle x^*, y - x \rangle \geq 0] \implies \quad \forall y^* \in T(y) : \langle y^*, y - x \rangle \geq 0. \quad (4)$$

We have the following classical result:

Theorem 2. [2, 4, 7, 11] *Let $f : X \rightarrow R \cup \{+\infty\}$ be a l.s.c. function. Consider the propositions:*

- i) f is pseudoconvex.
- ii) ∂f is pseudomonotone.

Then i) implies ii). If moreover, f is radially continuous, then ii) implies i).

In this paper, we want to characterize strictly pseudoconvex functions with respect to the Clarke-Rockafellar subdifferential in two different ways. For this, we introduce the so what we call radially Clarke-Rockafellar subdifferentiable functions (RCRS-functions).

Let $f : X \mapsto R \cup \{+\infty\}$ be a l.s.c. function. We say that f is radially Clarke-Rockafellar subdifferentiable if for all $x, y \in X$ with $x \neq y$, there is $x_0 \in (x, y)$ such that $\partial f(x_0) \neq \emptyset$. Recall that an extended-real valued function $f : X \mapsto R \cup \{+\infty\}$ is said to be radially continuous if for all $x, y \in X$ f is continuous on $[x, y]$.

2. CHARACTERIZATION OF RCRS-STRICT PSEUDOCONVEX FUNCTIONS

In this section, we get analogous results to theorem 1 and theorem 2 for RCRS-strictly pseudoconvex functions.

An extended-real valued function $f : X \mapsto R \cup \{+\infty\}$ is said to be radially non constant if for all $x, y \in X$ with $x \neq y$, $f \not\equiv \text{constant}$ on $[x, y]$.

Definition 3. A function $f : X \mapsto R \cup \{+\infty\}$ is said to be strictly pseudoconvex (with respect to the Clarke-Rockafellar subdifferential) if for any different points $x, y \in X$, the following implication holds:

$$(\exists x^* \in \partial f(x) : \langle x^*, y - x \rangle \geq 0) \implies f(x) < f(y). \quad (5)$$

We can check immediately that a strict pseudoconvex function is pseudoconvex while the converse is not true in general as we can see for example for the function

$$f(x) = \begin{cases} \sqrt{|x| - 1} & \text{if } x \in]-\infty, -1] \cup [1, +\infty[, \\ 0 & \text{if } x \in [-1, 1]. \end{cases} \quad (6)$$

We can describe the relation between strict pseudoconvexity and strict quasiconvexity via the following result:

Theorem 4. *Let $f : X \mapsto R \cup \{+\infty\}$ be a l.s.c. function such that f is radially Clarke-Rockafellar subdifferentiable. Consider the following assertions:*

- i) f is strictly pseudoconvex.*
- ii) f is strictly quasiconvex and $(0 \in \partial f(x) \implies x \text{ is a strict global minimum of } f)$.*

Then i) implies ii). If moreover, f is radially continuous, then ii) implies i).

Proof. Let f be a strictly pseudoconvex function, then by theorem 1, f is quasiconvex. Let us prove now that f is strictly quasiconvex. Since f is quasiconvex, then according to Diewert [5], it suffices to prove that f is radially non constant. By the contrary, assume that there exists a closed segment $[x, y]$ ($x \neq y$) on which f is constant. Let $z \in (x, y)$. Then applying the strict pseudoconvexity property on x and z , we deduce

$$\forall z^* \in \partial f(z) \quad \langle z^*, x - z \rangle < 0.$$

Using the same argument for z and y , we obtain

$$\forall z^* \in \partial f(z) \quad \langle z^*, y - z \rangle < 0.$$

Therefore,

$$\forall z^* \in \partial f(z), \quad \langle z^*, x - y \rangle < 0 \quad \text{and} \quad \langle z^*, x - y \rangle > 0.$$

Consequently, for all $z \in (x, y)$ we have $\partial f(z) = \emptyset$. But this contradicts the fact that f is a RCRS-function. Thus, f is strictly quasiconvex. On the other hand, f is pseudoconvex. Therefore,

$$0 \in \partial f(x) \implies x \text{ is a strict global minimum of } f.$$

Conversely, assume that f satisfies the condition ii) and f is radially continuous. Then by theorem 1, f is pseudoconvex.

Let us prove now that f is strictly pseudoconvex. Assume by contradiction that there exist $x \neq y$ in X and $x^* \in \partial f(x)$ such that

$$\langle x^*, y - x \rangle \geq 0 \quad \text{and} \quad f(x) \geq f(y).$$

Then, It follows by pseudoconvexity property that

$$f(x) = f(y) \text{ and } \forall z \in [x, y], \quad f(z) \geq f(x) \geq f(y).$$

On the other hand, f is quasiconvex. Therefore,

$$f(z) = f(x) = f(y), \quad \forall z \in [x, y].$$

Consequently, f is not radially non constant on X . But this contradicts the fact that f is strictly quasiconvex. Thus, we achieve the proof.

Analogously to pseudomonotone multioperators, we define strictly pseudomonotone multioperators as follows:

Definition 5. A multioperator $T : X \rightarrow X^*$ is said to be strictly pseudomonotone if for any different points x and y in X , the following implication holds:

$$\exists x^* \in T(x) : \langle x^*, y - x \rangle \geq 0 \implies \forall y^* \in T(y) : \langle y^*, y - x \rangle > 0. \quad (7)$$

We have also a relation between strict pseudoconvexity of functions and strict monotonicity of their corresponding Clarke-Rockafellar subdifferentials.

Theorem 6. Let $f : X \mapsto R \cup \{+\infty\}$ be a l.s.c. function such that f is radially Clarke-Rockafellar subdifferentiable. Consider the following assertions

- i) f is strictly pseudoconvex.
- ii) ∂f is strictly pseudomonotone.

Then i) implies ii). if moreover, f is radially continuous, then ii) implies i).

Proof. The first implication can be easily proved, nevertheless we include it here for completeness. Assume that f is strictly pseudoconvex. Let us prove by the contrary that ∂f is strictly pseudomonotone. Suppose that there exist two different points $x, y \in X$, $x^* \in \partial f(x)$ and $y^* \in \partial f(y)$ such that

$$\langle x^*, y - x \rangle \geq 0 \quad \text{and} \quad \langle y^*, x - y \rangle \geq 0.$$

Since f is strictly pseudoconvex, then

$$f(x) < f(y) \quad \text{and} \quad f(x) > f(y).$$

Contradiction. Thus, ∂f is strictly pseudomonotone.

Conversely, assume that f satisfies the condition ii) and f is radially continuous. Let us prove that f is strictly pseudoconvex. By the contrary, assume that there exist two different points x and y in X , and $x^* \in \partial f(x)$ such that both inequalities

$$\langle x^*, y - x \rangle \geq 0 \quad \text{and} \quad f(x) \geq f(y)$$

hold. Then

$$\langle x^*, z - x \rangle \geq 0 \quad \forall z \in [x, y]. \quad (8)$$

By theorem 2, it follows that f is pseudoconvex. Therefore,

$$f(x) \leq f(z) \quad \forall z \in [x, y].$$

By theorem 1, f is quasiconvex. Consequently, we can easily see that f must be constant on $[x, y]$. On the other hand, by (8) and the strict pseudomonotonicity of ∂f , we have:

$$\langle z^*, z - x \rangle > 0, \quad \forall z \in (x, y) \quad \forall z^* \in \partial f(z). \quad (9)$$

Pick $z_0 \in (x, y)$ such that $\partial f(z_0) \neq \emptyset$ (such a z_0 exists since f is a RCRS-function). Choose any $z_0^* \in \partial f(z_0)$. Then, $\langle z_0^*, z_0 - x \rangle > 0$. Therefore, $\langle z_0^*, y - z_0 \rangle > 0$. Consequently, there is $\varepsilon > 0$ such that

$$\langle z_0^*, y' - z_0 \rangle > 0 \quad \forall y' \in B(y, \varepsilon).$$

By the pseudoconvexity of f , it follows that y is a global minimum of f . Hence, z_0 is also a global minimum of f . Thus, $0 \in \partial f(z_0)$ and this is in contradiction with (9).

3. MAXIMA OF STRONGLY RCRS-STRICT PSEUDOCONVEX FUNCTIONS

In this section, we study a maximization problem over a convex set involving a certain class of RCRS-strictly pseudoconvex functions called class of strongly RCRS-strictly pseudoconvex functions.

Let $f : X \mapsto R \cup \{+\infty\}$ be a l.s.c. function. We say that f is strongly radially Clarke-Rockafellar subdifferentiable if for all $x, y \in X$ with $x \neq y$ and for all $c : f(x) < c < f(y)$, there is $x_0 \in (x, y)$ such that $f(x_0) = c$ and $\partial f(x_0) \neq \emptyset$.

Let C be a nonempty convex set of X . Consider the following maximization problem:

$$(\mathcal{P}) \quad \max_{x \in C} f(x),$$

where the function f is assumed to be strictly pseudoconvex, l.s.c. and strongly radially Clarke-Rockafellar subdifferentiable.

Theorem 7. *Consider $\bar{x} \in C$ such that*

$$-\infty \leq \inf_C f < f(\bar{x}). \quad (10)$$

Then \bar{x} is a maximum of f over C if and only if for all $x \in C$ such that $f(x) = f(\bar{x})$ and all $x^ \in \partial f(x)$ we have:*

$$\langle x^*, y - x \rangle < 0 \quad \forall y \in C \setminus \{x\}. \quad (11)$$

Proof. Assume that \bar{x} is a solution of the problem (\mathcal{P}) . Let $x \in X$ such that $f(x) = f(\bar{x})$ and let $x^* \in \partial f(x)$. Then

$$f(y) \leq f(x), \quad \forall y \in C.$$

Since f is strictly pseudoconvex, then

$$\langle x^*, y - x \rangle < 0, \quad \forall y \in C \setminus \{x\}.$$

Conversely, suppose that there exists $z \in C$ such that $f(z) > f(\bar{x})$. By the hypotheses, there is $z_0 \in C$ such that $f(z_0) < f(\bar{x})$. Since f is strongly radially Clarke-Rockafellar subdifferentiable, then there is some $x_0 \in (z_0, z)$ such that $f(x_0) = f(\bar{x})$ and $\partial f(x_0) \neq \emptyset$. Pick any $x_0^* \in \partial f(x_0)$. Then

$$\langle x_0^*, z - x_0 \rangle < 0 \quad \text{and} \quad \langle x_0^*, z_0 - x_0 \rangle < 0.$$

Which is impossible. To prove that (11) holds when \bar{x} is a maximum, we use only the strict pseudoconvexity of f , the other conditions that appear in theorem 7 are needed only to prove that (11) implies that \bar{x} is a maximum. This result is a refinement of both theorem 2.1 of [8] where the function was supposed to be convex continuous and of theorem 4.1 of [7] where the function was assumed to be pseudoconvex and radially continuous.

REFERENCES

- [1] D. Aussel, Théorème de la valeur moyenne et convexité généralisée en analyse non régulière, université Blaise Pascal, Ph-D Thesis, (1994).
- [2] D. Aussel, Subdifferential properties of quasiconvex and pseudoconvex function: A unified Approach, J. Optim. Th. Appl. 97 (to appear).
- [3] F. H. Clarke, Yu. S. Ledyae, R. J. Stern and P. R. Wolenski, Nonsmooth analysis and control theory, Springer-Verlag, New York, (1998).
- [4] A. Daniilidis and N. Hadjisavvas, On subdifferentials of quasiconvex and pseudoconvex functions and cyclic monotonicity, Les presses de L'Université de Aegean, Greece, (1997).
- [5] W. E. Diewert, Alternative characterizations of six kinds of quasiconvexity in the nondifferentiable case with applications nonsmooth programming, Academic Press (1981).
- [6] A. Hassouni and A. Jaddar, On quasiconvex functions and applications to optimality conditions in nonlinear programming, Appl. Math. Lett. (to appear).
- [7] A. Hassouni and A. Jaddar, Sur les fonctions pseudoconvexes et applications l'optimisation globale, (to appear).
- [8] J.-B. Hiriart Urruty and Yu. S. Ledyae, A note on characterization of the global maxima of a (tangentially) convex functions over a convex set, J. Convex Anal., **3**, N. 1, 55–31 (1996).
- [9] O. L. Mangazarian, Pseudoconvex functions, SIAM J. Control, **3**, 281–290 (1965).
- [10] J.-P. Penot, Generalized convex functions in the light of non smooth analysis, Lecture notes in Economics and Math. Systems, **429**, Springer Verlag, 269–291 (1995).
- [11] J.-P. Penot, Are generalized derivatives useful for generalized convex functions, Proceeding of fifth symposium on generalized convexity, 3–59 (1996).

DEPT OF MATHEMATICS, FACULTY OF SCIENCE, MOHAMED FIRST UNIVERSITY, OUJDA, MOROCCO, (DYNAMICAL SYSTEM AND OPTIMIZATION GROUP, GAFO LABORATORY)

E-mail address: lahrech@sciences.univ-oujda.ac.ma

NATIONAL SCHOOL OF MANAGMENT, OUJDA, MOROCCO, (DYNAMICAL SYSTEM AND OPTIMIZATION GROUP, GAFO LABORATORY)

E-mail address: jaddar@sciences.univ-oujda.ac.ma

DEPT OF MATHEMATICS, FACULTY OF SCIENCE, MOHAMED FIRST UNIVERSITY, OUJDA, MOROCCO, (DYNAMICAL SYSTEM AND OPTIMIZATION GROUP, GAFO LABORATORY)

E-mail address: ouahab@sciences.univ-oujda.ac.ma

NATIONAL SCHOOL OF APPLIED SCIENCES, OUJDA, MOROCCO, (DYNAMICAL SYSTEM AND OPTIMIZATION GROUP, GAFO LABORATORY)

E-mail address: ambarki@ensa.univ-oujda.ac.ma

Received October 19, 2006