

Homogenization of Random Walk in Asymmetric Random Environment

Joseph G. Conlon

ABSTRACT. In this paper, the author investigates the scaling limit of a partial difference equation on the d dimensional integer lattice \mathbf{Z}^d , corresponding to a translation invariant random walk perturbed by a random vector field. In the case when the translation invariant walk scales to a Cauchy process he proves convergence to an effective equation on \mathbf{R}^d . The effective equation corresponds to a Cauchy process perturbed by a constant vector field. In the case when the translation invariant walk scales to Brownian motion he shows that the scaling limit, if it exists, depends on dimension. For $d = 1, 2$ he provides evidence that the scaling limit cannot be diffusion.

CONTENTS

1.	Introduction	31
2.	Proof of Theorem 1.1	35
3.	Proof of Theorem 1.2	46
4.	Proof of Theorem 1.3	51
5.	Proof of Theorem 1.4	54
	References	60

1. Introduction

In this paper we shall be concerned with homogenization of a nondivergence form elliptic equation. Let $(\Omega, \mathcal{F}, \mu)$ be a probability space and $\mathbf{b} : \Omega \rightarrow \mathbf{R}^d$ be a bounded measurable function $\mathbf{b}(\omega) = (b_1(\omega), \dots, b_d(\omega))$ with

$$(1.1) \quad |\mathbf{b}(\omega)|^2 = \sum_{i=1}^d |b_i(\omega)|^2 \leq 1, \quad \omega \in \Omega.$$

We assume that \mathbf{Z}^d acts on Ω by translation operators $\tau_x : \Omega \rightarrow \Omega$, $x \in \mathbf{Z}^d$, which are measure preserving ergodic and satisfy the properties $\tau_x \tau_y = \tau_{x+y}$, $x, y \in \mathbf{Z}^d$, $\tau_0 = \text{identity}$. Using these operators we can define a measurable function $\mathbf{b} : \mathbf{Z}^d \times \Omega \rightarrow \mathbf{R}^d$ by $\mathbf{b}(x, \omega) = \mathbf{b}(\tau_x \omega)$, $x \in \mathbf{Z}^d$, $\omega \in \Omega$.

Received October 4, 2001.

Mathematics Subject Classification. 35R60, 60H30, 60J60.

Key words and phrases. random walk, homogenization, random environment.

For $\varepsilon > 0$ let $\mathbf{Z}_\varepsilon^d = \varepsilon\mathbf{Z}^d$ be the ε scaled integer lattice. We define the integral of a function $g : \mathbf{Z}_\varepsilon^d \rightarrow \mathbf{R}$ by

$$\int_{\mathbf{Z}_\varepsilon^d} g(x) dx \stackrel{\text{def.}}{=} \varepsilon^d \sum_{x \in \mathbf{Z}_\varepsilon^d} g(x).$$

The space $L^2(\mathbf{Z}_\varepsilon^d)$ is then the space of square integrable functions $u : \mathbf{Z}_\varepsilon^d \rightarrow \mathbf{C}$ with norm, $\|u\|_\varepsilon$ satisfying

$$\|u\|_\varepsilon^2 = \int_{\mathbf{Z}_\varepsilon^d} |u(x)|^2 dx.$$

For $i = 1, \dots, d$, let $\mathbf{e}_i \in \mathbf{Z}^d$ be the element with entry 1 in the i th position and 0 in the other positions. Suppose $\gamma \in \mathbf{C}$ and $f : \mathbf{R}^d \rightarrow \mathbf{R}$ is a C^∞ function with compact support. Let $p : \mathbf{Z}^d \rightarrow \mathbf{R}$ be a probability density, whence

$$p(x) \geq 0, \quad x \in \mathbf{Z}^d, \quad \sum_{x \in \mathbf{Z}^d} p(x) = 1.$$

With any such p we can associate a translation invariant operator A_p on functions $u_\varepsilon : \mathbf{Z}_\varepsilon^d \rightarrow \mathbf{R}$ by

$$A_p u_\varepsilon(x) = u_\varepsilon(x) - \sum_{y \in \mathbf{Z}^d} p(y) u_\varepsilon(x + \varepsilon y).$$

For some probability densities p , the Markov chain with generator A_p scales in the large time limit to a Levy process [20]. If A_p is such a generator, we denote the index of the corresponding symmetric stable process by $|A_p|$. If p is given by

$$(1.2) \quad p(\pm \mathbf{e}_i) = 1/2d, \quad i = 1, \dots, d,$$

then $|A_p| = 2$ since the corresponding Levy process is Brownian motion. We also define a p which scales to a Cauchy process, whence $|A_p| = 1$. To do this consider the standard nearest neighbor random walk on \mathbf{Z}^{d+1} started at the point $(0, -1)$, $0 \in \mathbf{Z}^d$. Then

$$(1.3) \quad p(x) = \text{probability the walk first hits the hyperplane} \\ \{(y, 0) : y \in \mathbf{Z}^d\} \text{ in the point } (x, 0), \quad x \in \mathbf{Z}^d.$$

We shall be interested in solutions to the equation

$$(1.4) \quad A_p u_\varepsilon(x, \omega) - \gamma \sum_{i=1}^d b_i \left(\frac{x}{\varepsilon}, \omega \right) [u_\varepsilon(x + \varepsilon \mathbf{e}_i, \omega) - u_\varepsilon(x - \varepsilon \mathbf{e}_i, \omega)] + \varepsilon^{|A_p|} u_\varepsilon(x, \omega) \\ = \varepsilon^{|A_p|} f(x), \quad x \in \mathbf{Z}_\varepsilon^d, \quad \omega \in \Omega,$$

when p is given by either (1.2) or (1.3). For a function $g : \mathbf{X} \rightarrow \mathbf{C}$ where $\mathbf{X} = \mathbf{R}^d$ or $\mathbf{X} = \mathbf{Z}_\varepsilon^d$, let \hat{g} be its Fourier transform,

$$\hat{g}(\xi) = \int_{\mathbf{X}} g(x) e^{ix \cdot \xi} dx, \quad \xi \in \mathbf{R}^d.$$

Note that when $\mathbf{X} = \mathbf{Z}_\varepsilon^d$, then $\hat{g}(\xi)$ is periodic in ξ whence we can restrict ξ to the cube $[-\pi/\varepsilon, \pi/\varepsilon]^d$. Suppose now $\mathbf{b} \equiv 0$ in (1.4) and p is given by (1.2). Then

it is easy to see that (1.4) has a unique solution $u_\varepsilon(x)$ in $L^2(\mathbf{Z}_\varepsilon^d)$. Further, if $u(x)$, $x \in \mathbf{R}^d$, is the function which satisfies

$$[1 + |\xi|^2/2d]\hat{u}(\xi) = \hat{f}(\xi), \quad \xi \in \mathbf{R}^d,$$

then $\lim_{\varepsilon \rightarrow 0} \|u_\varepsilon - u\|_\varepsilon = 0$. Similarly if $\mathbf{b} \equiv 0$ and p is given by (1.3) then (1.4) has a unique solution $u_\varepsilon(x)$ in $L^2(\mathbf{Z}_\varepsilon^d)$. We show in §2 that if $u(x)$, $x \in \mathbf{R}^d$, is the function which satisfies

$$(1.5) \quad [1 + |\xi|]\hat{u}(\xi) = \hat{f}(\xi), \quad \xi \in \mathbf{R}^d,$$

then

$$(1.6) \quad \lim_{\varepsilon \rightarrow 0} \|u_\varepsilon - u\|_\varepsilon = 0.$$

We can also prove a corresponding theorem when $\mathbf{b} \neq 0$ in the case p is given by (1.3). In the following the expectation value of a random variable X on Ω is denoted by $\langle X \rangle$.

Theorem 1.1. *For $0 < \varepsilon \leq 1$, there exists a constant $\gamma_d > 0$, depending only on d , such that if $\gamma \in \mathbf{C}$ satisfies $|\gamma| < \gamma_d$ then (1.4) has a unique solution $u_\varepsilon(\cdot, \omega) \in L^2(\mathbf{Z}_\varepsilon^d)$, $\omega \in \Omega$. There exists a vector $q_\gamma \in \mathbf{C}^d$, $|q_\gamma| < 1$, analytic in γ , such that if $u(x)$, $x \in \mathbf{R}^d$, is the function which satisfies*

$$(1.7) \quad [1 + |\xi| + iq_\gamma \cdot \xi]\hat{u}(\xi) = \hat{f}(\xi), \quad \xi \in \mathbf{R}^d,$$

then there is the limit

$$(1.8) \quad \lim_{\varepsilon \rightarrow 0} \langle \|u_\varepsilon(\cdot, \cdot) - u\|_\varepsilon^2 \rangle = 0.$$

Theorem 1.1 is a homogenization result for p given by (1.3). It says that the scaling limit of the random walk, with transition probability (1.3), in a random environment described by $\mathbf{b}(x, \omega)$, is a Cauchy process with a constant drift q_γ . Corresponding results for uniformly elliptic partial differential equations in divergence form were first proved by Kozlov [11] and later by Papanicolaou and Varadhan [16]. In this case the scaling limit is Brownian motion. One can also prove a Brownian motion scaling limit for parabolic equations in divergence form when the coefficients are random in time as well as space [13]. It is possible to relax somewhat the uniform ellipticity assumption and still prove the scaling limit [5]. For a comprehensive survey of homogenization results in partial differential equations see the book of Oleinik et al [25]. Here we obtain the scaling limit of an asymmetric partial difference equation. The scaling limit of a divergence form partial difference equation has been obtained by Künnemann [12] and of a symmetric partial difference equation in non-divergence form by Lawler [14].

Evidently q_γ is in general nonzero even in the case $\langle \mathbf{b}(\cdot) \rangle = 0$. Nevertheless it is possible to impose some general conditions on $\mathbf{b}(\cdot)$ to ensure $q_\gamma = 0$. We say that the vector field $\mathbf{b}(\cdot)$ is reflection invariant if it has the property that

$$(1.9) \quad \left\langle \prod_{i=1}^n b_{k_i}(\tau_{x_i} \cdot) \right\rangle = (-1)^n \left\langle \prod_{i=1}^n b_{k_i}(\tau_{-x_i} \cdot) \right\rangle,$$

$$x_i \in \mathbf{Z}^d, \quad 1 \leq k_i \leq d, \quad i = 1, \dots, n, \quad n \geq 1.$$

Clearly (1.9) implies that $\langle \mathbf{b}(\cdot) \rangle = \mathbf{0}$. We show at the end of §2 that if $\mathbf{b}(\cdot)$ satisfies (1.9) then $q_\gamma = 0$ in (1.7).

Suppose now (1.9) holds. Let $u_\varepsilon(x) = \langle u_\varepsilon(x, \cdot) \rangle$, $x \in \mathbf{Z}_\varepsilon^d$, where $u_\varepsilon(x, \omega)$ is the unique solution to (1.4) given by Theorem 1.1. Suppose $\hat{p} : [-\pi, \pi]^d \rightarrow \mathbf{R}$ is the Fourier transform of the function p of (1.3) and \hat{f}_ε the Fourier transform of the function f restricted to the lattice \mathbf{Z}_ε^d . If $\hat{u}_\varepsilon(\xi)$ is the Fourier transform of $u_\varepsilon(x)$, then one can see that there is a $d \times d$ matrix $q_{\gamma, \varepsilon}(\zeta)$, $\zeta \in [-\pi, \pi]^d$, continuous in ζ and analytic in γ , such that

(1.10)

$$\hat{u}_\varepsilon(\xi)[1 + \varepsilon^{-1}\{1 - \hat{p}(\varepsilon\xi)\} - \varepsilon^{-1}e(\varepsilon\xi)q_{\gamma, \varepsilon}(\varepsilon\xi)e(-\varepsilon\xi)] = \hat{f}_\varepsilon(\xi), \quad \xi \in [-\pi/\varepsilon, \pi/\varepsilon],$$

where $e(\zeta)$, $\zeta \in \mathbf{R}^d$, is the vector $e(\zeta) = (e_1(\zeta), \dots, e_d(\zeta))$, with $e_k(\zeta) = 1 - e^{ie_k \cdot \zeta}$.

Theorem 1.2. *Suppose the variables $\mathbf{b}(\tau_x \cdot)$, $x \in \mathbf{Z}^d$, are independent with $\langle \mathbf{b}(\cdot) \rangle = 0$ and $q_{\gamma, \varepsilon}(\zeta)$ is defined by (1.10) for $|\gamma| < \gamma_d$, where γ_d is the constant of Theorem 1.1. Then there exists a positive constant $\gamma'_d < \gamma_d$, depending only on d , such that if $|\gamma| < \gamma'_d$ the matrix $q_{\gamma, \varepsilon}(\zeta)$ converges uniformly in ζ as $\varepsilon \rightarrow 0$ to a matrix $q_\gamma(\zeta)$, $\zeta \in [-\pi, \pi]^d$.*

Theorem 1.2 is a considerably deeper theorem than Theorem 1.1. One has to use the Calderon-Zygmund theorem [8, 22] that the Hilbert transform is bounded on L^p spaces for $p \neq 2$. In contrast, Theorem 1.1 can be proved by just using L^2 theory.

Next we consider the situation when p is given by (1.2) and $\mathbf{b} \neq 0$. We then have the following theorem:

Theorem 1.3. *For $0 < \varepsilon \leq 1$ and $\gamma \in \mathbf{C}$ satisfying $|\gamma| < \varepsilon/\sqrt{2d}$, equation (1.4) has a unique solution $u_\varepsilon(\cdot, \omega) \in L^2(\mathbf{Z}_\varepsilon^d)$, $\omega \in \Omega$. If (1.9) holds then there is a $d \times d$ matrix $q_{\gamma, \varepsilon}(\zeta)$, $\zeta \in [-\pi, \pi]^d$, continuous in ζ and analytic in γ , such that the Fourier transform $\hat{u}_\varepsilon(\xi)$ of $\langle u_\varepsilon(x, \cdot) \rangle$ satisfies*

(1.11)

$$\hat{u}_\varepsilon(\xi)[1 + \varepsilon^{-2}\{1 - \hat{p}(\varepsilon\xi)\} - \varepsilon^{-2}e(\varepsilon\xi)q_{\gamma, \varepsilon}(\varepsilon\xi)e(-\varepsilon\xi)] = \hat{f}_\varepsilon(\xi), \quad \xi \in [-\pi/\varepsilon, \pi/\varepsilon].$$

Note that (1.11) is simply the analogue for p given by (1.2) of (1.10) for p given by (1.3). Since $q_{\gamma, \varepsilon}(\zeta)$ is analytic in γ , we can write it in a series expansion,

$$(1.12) \quad q_{\gamma, \varepsilon}(\zeta) = \sum_{m=2}^{\infty} \gamma^m q_{m, \varepsilon}(\zeta),$$

which converges for $|\gamma| < \varepsilon/\sqrt{2d}$. In §5 we investigate the limiting behavior of $q_{m, \varepsilon}$ as $\varepsilon \rightarrow 0$. We restrict ourselves to the situation when the $\mathbf{b}(\tau_x \cdot)$, $x \in \mathbf{Z}^d$, are given by independent Bernoulli variables. Thus we assume that $b_j(\cdot) \equiv 0$, $j > 1$, and $b_1(\tau_x \cdot) = Y_x$, $x \in \mathbf{Z}^d$, where the variables Y_x , $x \in \mathbf{Z}^d$, are assumed to be i.i.d. Bernoulli, $Y_x = \pm 1$ with equal probability. In that case we have the following:

Theorem 1.4. *For $m = 2, 3, \dots$ the $d \times d$ matrix $q_{m, \varepsilon}(\zeta)$ has the properties:*

- (a) $q_{m, \varepsilon}(\zeta) \equiv 0$ if m is odd or $m = 2$.
- (b) The entries of the matrix $q_{m, \varepsilon}(\zeta) = [q_{m, \varepsilon, k, k'}(\zeta)]$, $1 \leq k, k' \leq d$, are zero if $k, k' > 1$.
- (c) The entries of the matrix $q_{m, \varepsilon}(0) = [q_{m, \varepsilon, k, k'}(0)]$ are zero if $k + k' > 2$.

- (d) Assume $d = 1$ and $\mathcal{K} \subset \mathbf{R}$ is a compact subset. Then $\varepsilon^{m/2} q_{m,\varepsilon}(\varepsilon\xi)$ converges uniformly for $\xi \in \mathcal{K}$ to a function $q_m(\xi)$, as $\varepsilon \rightarrow 0$. The function $q_4(\xi)$ is given by the formula,

$$(1.13) \quad q_4(\xi) = \frac{32}{18 + \xi^2}.$$

- (e) If $d = 2$ then $q_{4,\varepsilon,1,1}(0)$ converges to $+\infty$ as $\varepsilon \rightarrow 0$.
(f) If $d \geq 3$ then $q_{4,\varepsilon}(\zeta)$ converges uniformly for $\zeta \in [-\pi, \pi]^d$ to a matrix $q_4(\zeta) = [q_{4,k,k'}(\zeta)]$ satisfying $q_{4,1,1}(0) > 0$, as $\varepsilon \rightarrow 0$.

Unlike in the previous theorems, the results of Theorem 1.4 depend on the dimension d . For $d = 1$ the theorem suggests that if we set $\gamma = \gamma' \sqrt{\varepsilon}$ in (1.4), fix γ' and let $\varepsilon \rightarrow 0$, then we should obtain a scaling limit. Note however that one expects the resulting series obtained from (1.12) by letting $\varepsilon \rightarrow 0$ to be at most asymptotic in γ' , not analytic. The limit here is related to the limits obtained by Sinai [21] and Kesten [9] for one dimensional random walk in random environment. For $d = 2$ the theorem suggests that if we fix γ in (1.4) and let $\varepsilon \rightarrow 0$ no scaling limit exists. This appears to contradict a conjecture of Fisher [6] that two dimensional random walk in random environment has a diffusive scaling limit. In contrast Derrida and Luck [4] conjecture a nondiffusive scaling limit in two dimensions. A numerical study of the two dimensional equation (1.4) was made in [2] using a multigrid algorithm. It was observed that the algorithm gave considerable acceleration over pure iterative methods such as Gauss-Seidel. This suggests that there is some stability in the Fourier space behavior of the solution to (1.4) as $\varepsilon \rightarrow 0$. For $d \geq 3$ the theorem suggests a diffusive scaling limit for d dimensional random walk in random environment and that the effective diffusion constant is smaller than for the zero noise case.

There has been some recent work [17, 23, 24] on (1.4) with p given by (1.2) and under the assumption $\langle \mathbf{b}(\cdot) \rangle \neq 0$. This situation is very different to the situation studied in Theorem 1.4 since one expects now the drift to dominate diffusion. The methods used in [23, 24] are related to methods used to prove Anderson localisation for the random Schrodinger equation.

In this paper we shall adapt a method developed in [3] to prove the homogenization results for (1.4). The method consists of space translation in Ω followed by Fourier transformation in \mathbf{Z}^d . The space translation in Ω is similar in spirit to “viewing the medium from an observer sitting on a tagged particle”, as described in [10]. The main advantage here is that one obtains the coefficients of the effective homogenized equation from the zero Fourier mode equation in the new variables. The method also avoids use of Martingales, as for example occurs in the recent paper [15].

2. Proof of Theorem 1.1

We first prove (1.6). Thus we consider (1.4) with $\mathbf{b} \equiv 0$ and p given by (1.3). To obtain an expression for p , let us denote a point in \mathbf{Z}^{d+1} by (x, y) with $x \in \mathbf{Z}^d$, $y \in \mathbf{Z}$. For $n = 0, 1, 2, \dots$ let $G(x, y, n)$ be the probability density of the standard

random walk in \mathbf{Z}^{d+1} , started at the origin, after n steps. Thus

$$(2.1) \quad \sum_{x \in \mathbf{Z}^d} \sum_{y \in \mathbf{Z}} G(x, y, n) e^{i[x \cdot \xi + y \zeta]} = \left\{ \frac{1}{(d+1)} \left[\sum_{i=1}^d \cos(\mathbf{e}_i \cdot \xi) + \cos \zeta \right] \right\}^n.$$

By the reflection principle, the probability that the $d+1$ dimensional walk, started at $(0, -1)$ first hits the hyperplane $\{(y, 0) : y \in \mathbf{Z}^d\}$ after $n+1$ steps at the point $(x, 0)$ is given by

$$\frac{1}{2(d+1)} [G(x, 0, n) - G(x, 2, n)].$$

Hence we have

$$(2.2) \quad p(x) = \frac{1}{2(d+1)} \sum_{n=0}^{\infty} [G(x, 0, n) - G(x, 2, n)].$$

We can compute the Fourier transform of p by using (2.1). Thus

$$(2.3) \quad \hat{p}(\xi) = \sum_{x \in \mathbf{Z}^d} p(x) e^{ix \cdot \xi} = \frac{1}{4\pi(d+1)} \int_{-\pi}^{\pi} d\zeta \frac{1 - \cos 2\zeta}{1 - \left[\sum_{i=1}^d \cos(\mathbf{e}_i \cdot \xi) + \cos \zeta \right] / (d+1)}.$$

Lemma 2.1. *The limit (1.6) holds.*

Proof. We rewrite the formula of (2.3) for $\hat{p}(\xi)$ as

$$(2.4) \quad \hat{p}(\xi) = 1 - \frac{1}{2\pi} \int_{-\pi}^{\pi} d\zeta \frac{(1 + \cos \zeta) \sum_{i=1}^d [1 - \cos(\mathbf{e}_i \cdot \xi)]}{\left[(1 - \cos \zeta) + \sum_{i=1}^d [1 - \cos(\mathbf{e}_i \cdot \xi)] \right]},$$

whence, on performing the integration with respect to ζ , we obtain the formula,

$$(2.5) \quad 1 - \hat{p}(\xi) = \left\{ \left[1 + \frac{2}{\sum_{i=1}^d [1 - \cos(\mathbf{e}_i \cdot \xi)]} \right]^{1/2} - 1 \right\} \sum_{i=1}^d [1 - \cos(\mathbf{e}_i \cdot \xi)].$$

It follows from (2.3), (2.4) that $0 < \hat{p}(\xi) \leq 1$. We can also easily see from (2.5) that

$$(2.6) \quad \lim_{\varepsilon \rightarrow 0} \varepsilon^{-1} [1 - \hat{p}(\varepsilon \xi)] = |\xi|, \quad \xi \in \mathbf{R}^d.$$

Consider now (1.4) with $\mathbf{b} \equiv 0$. Taking the Fourier transform of (1.4), we see that (1.4) is equivalent to the equation,

$$(2.7) \quad [1 + \varepsilon - \hat{p}(\varepsilon \xi)] \hat{u}_\varepsilon(\xi) = \varepsilon \hat{f}_\varepsilon(\xi), \quad \xi \in \left[\frac{-\pi}{\varepsilon}, \frac{\pi}{\varepsilon} \right]^d,$$

where \hat{f}_ε is the Fourier transform of f regarded as a function on \mathbf{Z}_ε^d . Since f is C^∞ of compact support it is easy to see that (2.7) is uniquely solvable in $L^2 \left(\left[\frac{-\pi}{\varepsilon}, \frac{\pi}{\varepsilon} \right]^d \right)$ and

$$(2.8) \quad \hat{u}_\varepsilon(\xi) = \hat{f}_\varepsilon(\xi) / [1 + \varepsilon^{-1} \{1 - \hat{p}(\varepsilon \xi)\}], \quad \xi \in \left[\frac{-\pi}{\varepsilon}, \frac{\pi}{\varepsilon} \right]^d.$$

Let $\hat{u}(\xi)$ be the function given in (1.5). We decompose its Fourier inverse $u(x)$, $x \in \mathbf{R}^d$, as a sum, $u(x) = v_\varepsilon(x) + w_\varepsilon(x)$, where

$$v_\varepsilon(x) = \frac{1}{(2\pi)^d} \int_{[-\frac{\pi}{\varepsilon}, \frac{\pi}{\varepsilon}]^d} \hat{u}(\xi) e^{-ix \cdot \xi} d\xi.$$

Since $\hat{f}(\xi)$ is rapidly decreasing in ξ it follows that $\lim_{\varepsilon \rightarrow 0} \|w_\varepsilon\|_\varepsilon = 0$. We can see this from the identity

$$\|w_\varepsilon\|_\varepsilon^2 = \frac{1}{(2\pi)^d} \int_{[-\frac{\pi}{\varepsilon}, \frac{\pi}{\varepsilon}]^d} \left| \sum_{n \in \mathbf{Z}^d \setminus \{0\}} \hat{u}(\xi + \frac{2\pi n}{\varepsilon}) \right|^2 d\xi.$$

Thus we have

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \|u_\varepsilon - u\|_\varepsilon &= \lim_{\varepsilon \rightarrow 0} \|u_\varepsilon - v_\varepsilon\|_\varepsilon, \\ \|u_\varepsilon - v_\varepsilon\|_\varepsilon^2 &= \frac{1}{(2\pi)^d} \int_{[-\frac{\pi}{\varepsilon}, \frac{\pi}{\varepsilon}]^d} |\hat{u}_\varepsilon(\xi) - \hat{u}(\xi)|^2 d\xi, \end{aligned}$$

where $\hat{u}_\varepsilon(\xi)$ is given by (2.8) and $\hat{u}(\xi)$ by (1.5). The result follows from (2.6) and the dominated convergence theorem. \square

Next we wish to prove existence of a solution to (1.4) provided γ is sufficiently small, depending only on d . To do this we need to define the Green's function $G_{p,\eta}(x)$, $x \in \mathbf{Z}^d$, associated with p . Thus for $\eta > 0$, $G_{p,\eta}(x)$ is the solution to the equation

$$(2.9) \quad A_p G_{p,\eta}(x) + \eta G_{p,\eta}(x) = \delta(x), \quad x \in \mathbf{Z}^d,$$

where δ is the Kronecker δ function, $\delta(0) = 1$, $\delta(x) = 0$, $x \neq 0$. The Fourier transform $\hat{G}_{p,\eta}(\xi)$ of $G_{p,\eta}(x)$ is given by the formula

$$(2.10) \quad \hat{G}_{p,\eta}(\xi) = [1 + \eta - \hat{p}(\xi)]^{-1}, \quad \xi \in [-\pi, \pi]^d.$$

It is easy to see from (2.5) that

$$0 \leq \hat{G}_{p,\eta}(\xi) \leq C/[\eta + |\xi|],$$

for some universal constant C . We can also obtain a corresponding decay rate on the function $G_{p,\eta}(x)$.

Lemma 2.2. *Let $G_{p,\eta}(x)$, $x \in \mathbf{Z}^d$, be the solution to (2.9), and $0 < \eta \leq 1$. Then there is a constant C_d , depending only on d such that if $d \geq 2$,*

$$(2.11) \quad 0 \leq G_{p,\eta}(x) \leq C_d / [1 + \eta^2 |x|^2] [1 + |x|^{d-1}], \quad x \in \mathbf{Z}^d.$$

Proof. Consider the random walk on \mathbf{Z}^d with transition probability p given by (2.2). Let $p_n(x)$, $n = 0, 1, 2, \dots$ be the probability that the walk, started at the origin hits x after n steps. Thus $p_0(x) = \delta(x)$ and $p_1(x) = p(x)$, $x \in \mathbf{Z}^d$. It is easy to see that the solution of (2.9) is given by the formula,

$$(2.12) \quad G_{p,\eta}(x) = \sum_{m=0}^{\infty} (1 + \eta)^{-m-1} p_m(x), \quad x \in \mathbf{Z}^d.$$

We can derive a formula for $p_m(x)$, $m \geq 2$, similar to the formula (2.2) for $p(x)$. In fact we have

$$(2.13) \quad p_m(x) = \frac{1}{2(d+1)} \sum_{n=0}^{\infty} [G(x, m-1, n) - G(x, m+1, n)].$$

It is easy to see from this and (2.12) that

$$(2.14) \quad G_{p,0}(x) = \delta(x) + \frac{1}{2(d+1)} \sum_{n=0}^{\infty} [G(x, 0, n) + G(x, 1, n)].$$

If we use now the bound

$$0 \leq G(x, y, n) \leq \frac{C_d}{1 + n^{(d+1)/2}} \exp \left[-\min\{|x| + |y|, (|x|^2 + |y|^2)/(n+1)\} / C_d \right],$$

for the standard random walk on \mathbf{Z}^{d+1} , then the inequality (2.11) for $\eta = 0$ follows from (2.14). To obtain (2.11) for $\eta > 0$ we need to use the inequality,

$$0 \leq G(x, y-1, n) - G(x, y+1, n) \leq \frac{C_d y}{1 + n^{(d+3)/2}} \exp \left[-\min\{|x| + |y|, (|x|^2 + |y|^2)/(n+1)\} / C_d \right],$$

$$x \in \mathbf{Z}^d, y \in \mathbf{Z} \quad y \geq 0.$$

It follows from this and (2.13) that

$$0 \leq p_m(x) \leq \frac{C_d m}{1 + [|x|^2 + m^2]^{(d+1)/2}}, \quad m \geq 1.$$

The inequality (2.11) follows now from this last inequality by estimating the sum (2.12). \square

Lemma 2.3. *Suppose $d \geq 2$ and $f : \mathbf{R}^d \rightarrow \mathbf{R}$ is a C^∞ function with compact support. Then there exists a constant γ_d , depending only on d such that if $|\gamma| < \gamma_d$ and $0 < \varepsilon \leq 1$, then (1.4) has a unique solution $u_\varepsilon(x, \omega)$ in $L^2(\mathbf{Z}_\varepsilon^d)$. Further, there is a constant C_d , depending only on d , such that $\|u_\varepsilon(\cdot, \omega)\|_\varepsilon \leq C_d \|f\|_\varepsilon$.*

Proof. Suppose $u_\varepsilon(x, \omega)$ is in $L^2(\mathbf{Z}_\varepsilon^d)$ and satisfies (1.4). Let $v_\varepsilon(x, \omega)$ be defined by

$$v_\varepsilon(x, \omega) = \varepsilon^{-1} [A_p u_\varepsilon(x, \omega) + \varepsilon u_\varepsilon(x, \omega)].$$

It is easy to see that $v_\varepsilon(x, \omega) \in L^2(\mathbf{Z}_\varepsilon^d)$. In view of (2.9) it follows that

$$(2.15) \quad u_\varepsilon(x, \omega) = \varepsilon^{1-d} \int_{\mathbf{Z}_\varepsilon^d} dy G_{p,\varepsilon} \left(\frac{x-y}{\varepsilon} \right) v_\varepsilon(y, \omega).$$

Now for $i = 1, \dots, d$ let us define operators T_i on $L^2(\mathbf{Z}_\varepsilon^d)$ by

$$T_i w(x) = \varepsilon^{-d} \int_{\mathbf{Z}_\varepsilon^d} dy \left[G_{p,\varepsilon} \left(\frac{x-y}{\varepsilon} + \mathbf{e}_i \right) - G_{p,\varepsilon} \left(\frac{x-y}{\varepsilon} - \mathbf{e}_i \right) \right] w(y).$$

It is easy to see that T_i is a bounded operator on $L^2(\mathbf{Z}_\varepsilon^d)$. It follows from (2.10) that the norm of T_i is bounded by

$$(2.16) \quad \|T_i\| \leq \sup_{\xi \in [-\pi, \pi]^d} \left[\frac{2|\sin(\mathbf{e}_i \cdot \xi)|}{1 + \varepsilon - \hat{p}(\xi)} \right].$$

We can see from (2.5) that there is a constant c_d , $0 < c_d < 1$, depending only on d such that

$$(2.17) \quad 1 - \hat{p}(\xi) \geq c_d |\xi|, \quad \xi \in [-\pi, \pi]^d.$$

It follows from this and (2.16) that $\|T_i\| \leq 2/c_d$. Observe now that (1.4) implies that $v_\varepsilon(x, \omega) \in L^2(\mathbf{Z}_\varepsilon^d)$ satisfies the equation

$$(2.18) \quad v_\varepsilon(x, \omega) - \gamma \sum_{i=1}^d b_i(x/\varepsilon, \omega) T_i v_\varepsilon(x, \omega) = f(x).$$

In view of (1.1), if we take $\gamma_d = c_d/2$ it follows that for $|\gamma| < \gamma_d$ the function $v_\varepsilon(x, \omega)$ is uniquely determined by (2.18). In view of (2.15) it follows that $u_\varepsilon(x, \omega) \in L^2(\mathbf{Z}_\varepsilon^d)$ is the unique solution to (1.4). To prove existence note that (2.18) has a solution $v_\varepsilon(x, \omega) \in L^2(\mathbf{Z}_\varepsilon^d)$ provided $|\gamma| < \gamma_d$. If we then define $u_\varepsilon(x, \omega)$ by (2.15) one can easily see that $u_\varepsilon(x, \omega)$ satisfies (1.4). \square

We have proven existence of a unique solution $u_\varepsilon(x, \omega)$ to (1.4) provided $|\gamma| < \gamma_d$. Next we follow a development similar to the one in [3]. Thus we put $v_\varepsilon(x, \omega) = u_\varepsilon(x, \tau_{-x/\varepsilon}\omega)$, $x \in \mathbf{Z}_\varepsilon^d$, $\omega \in \Omega$, whence $u_\varepsilon(x, \omega) = v_\varepsilon(x, \tau_{x/\varepsilon}\omega)$. Rewriting (1.4) in terms of v_ε , we obtain the equation,

$$\begin{aligned} v_\varepsilon(x, \tau_{x/\varepsilon}\omega) - \sum_{y \in \mathbf{Z}^d} p(y) v_\varepsilon(x + \varepsilon y, \tau_y \tau_{x/\varepsilon}\omega) \\ - \gamma \sum_{i=1}^d b_i(\tau_{x/\varepsilon}\omega) [v_\varepsilon(x + \varepsilon \mathbf{e}_i, \tau_{\mathbf{e}_i} \tau_{x/\varepsilon}\omega) \\ - v_\varepsilon(x - \varepsilon \mathbf{e}_i, \tau_{-\mathbf{e}_i} \tau_{x/\varepsilon}\omega)] + \varepsilon v_\varepsilon(x, \tau_{x/\varepsilon}\omega) = \varepsilon f(x), \quad x \in \mathbf{Z}_\varepsilon^d, \quad \omega \in \Omega. \end{aligned}$$

This last equation is evidently the same as

$$(2.19) \quad \begin{aligned} v_\varepsilon(x, \omega) - \sum_{y \in \mathbf{Z}^d} p(y) v_\varepsilon(x + \varepsilon y, \tau_y \omega) \\ - \gamma \sum_{i=1}^d b_i(\omega) [v_\varepsilon(x + \varepsilon \mathbf{e}_i, \tau_{\mathbf{e}_i} \omega) - v_\varepsilon(x - \varepsilon \mathbf{e}_i, \tau_{-\mathbf{e}_i} \omega)] \\ + \varepsilon v_\varepsilon(x, \omega) = \varepsilon f(x), \quad x \in \mathbf{Z}_\varepsilon^d, \quad \omega \in \Omega. \end{aligned}$$

We write now $v_\varepsilon(x, \omega) = u_\varepsilon(x) + \psi_\varepsilon(x, \omega)$, $x \in \mathbf{Z}_\varepsilon^d$, $\omega \in \Omega$, where $\langle \psi_\varepsilon(x, \cdot) \rangle = 0$, $x \in \mathbf{Z}_\varepsilon^d$. It follows that

$$u_\varepsilon(x) = \langle v_\varepsilon(x, \cdot) \rangle = \langle u_\varepsilon(x, \cdot) \rangle, \quad x \in \mathbf{Z}_\varepsilon^d.$$

Substituting into (2.19) we obtain the equation

$$\begin{aligned}
(2.20) \quad u_\varepsilon(x) &- \sum_{y \in \mathbf{Z}^d} p(y) u_\varepsilon(x + \varepsilon y) \\
&- \gamma \sum_{i=1}^d b_i(\omega) [u_\varepsilon(x + \varepsilon \mathbf{e}_i) - u_\varepsilon(x - \varepsilon \mathbf{e}_i)] + \varepsilon u_\varepsilon(x) \\
&+ \psi_\varepsilon(x, \omega) - \sum_{y \in \mathbf{Z}^d} p(y) \psi_\varepsilon(x + \varepsilon y, \tau_y \omega) \\
&- \gamma \sum_{i=1}^d b_i(\omega) [\psi_\varepsilon(x + \varepsilon \mathbf{e}_i, \tau_{\mathbf{e}_i} \omega) - \psi_\varepsilon(x - \varepsilon \mathbf{e}_i, \tau_{-\mathbf{e}_i} \omega)] \\
&+ \varepsilon \psi_\varepsilon(x, \omega) = \varepsilon f(x), \quad x \in \mathbf{Z}_\varepsilon^d, \quad \omega \in \Omega.
\end{aligned}$$

Taking the expected value of this last equation, we obtain the equation,

$$\begin{aligned}
(2.21) \quad A_p u_\varepsilon(x) &- \gamma \sum_{i=1}^d \langle b_i(\cdot) \rangle [u_\varepsilon(x + \varepsilon \mathbf{e}_i) - u_\varepsilon(x - \varepsilon \mathbf{e}_i)] + \varepsilon u_\varepsilon(x) \\
&- \gamma \left\langle \sum_{i=1}^d b_i(\cdot) [\psi_\varepsilon(x + \varepsilon \mathbf{e}_i, \tau_{\mathbf{e}_i} \cdot) - \psi_\varepsilon(x - \varepsilon \mathbf{e}_i, \tau_{-\mathbf{e}_i} \cdot)] \right\rangle \\
&= \varepsilon f(x), \quad x \in \mathbf{Z}_\varepsilon^d.
\end{aligned}$$

If we subtract now (2.21) from (2.20) we obtain

$$\begin{aligned}
(2.22) \quad &\psi_\varepsilon(x, \omega) - \sum_{y \in \mathbf{Z}^d} p(y) \psi_\varepsilon(x + \varepsilon y, \tau_y \omega) \\
&- \gamma P \sum_{i=1}^d b_i(\omega) [\psi_\varepsilon(x + \varepsilon \mathbf{e}_i, \tau_{\mathbf{e}_i} \omega) - \psi_\varepsilon(x - \varepsilon \mathbf{e}_i, \tau_{-\mathbf{e}_i} \omega)] + \varepsilon \psi_\varepsilon(x, \omega) \\
&= \gamma \sum_{i=1}^d [b_i(\omega) - \langle b_i(\cdot) \rangle] [u_\varepsilon(x + \varepsilon \mathbf{e}_i) - u_\varepsilon(x - \varepsilon \mathbf{e}_i)], \quad x \in \mathbf{Z}_\varepsilon^d, \quad \omega \in \Omega,
\end{aligned}$$

where $P : L^2(\Omega) \rightarrow L^2(\Omega)$ is the projection orthogonal to the constant function. Thus a solution $u_\varepsilon(x, \omega)$ of (1.4) which is in $L^2(\mathbf{Z}_\varepsilon^d \times \Omega)$ yields functions $u_\varepsilon(x)$ in $L^2(\mathbf{Z}_\varepsilon^d)$ and $\psi_\varepsilon(x, \omega)$ in $L^2(\mathbf{Z}_\varepsilon^d \times \Omega)$ which satisfy $\langle \psi_\varepsilon(x, \cdot) \rangle = 0$, $x \in \mathbf{Z}_\varepsilon^d$, and the equations (2.21), (2.22). Conversely, if we can find functions $u_\varepsilon(x)$ in $L^2(\mathbf{Z}_\varepsilon^d)$ and $\psi_\varepsilon(x, \omega)$ in $L^2(\mathbf{Z}_\varepsilon^d \times \Omega)$ which have the property that $\langle \psi_\varepsilon(x, \cdot) \rangle = 0$, $x \in \mathbf{Z}_\varepsilon^d$, and satisfy (2.21), (2.22) then we can construct from them the solution $u_\varepsilon(x, \omega)$ of (1.4) and $u_\varepsilon(x, \omega)$ is in $L^2(\mathbf{Z}_\varepsilon^d \times \Omega)$.

We concentrate now on finding solutions to the system (2.21), (2.22) of equations. To do this we Fourier transform the equations. The Fourier transform of (2.21) is

given by

$$\begin{aligned}
 (2.23) \quad \hat{u}_\varepsilon(\xi) & \left\{ 1 + \varepsilon^{-1} [1 - \hat{p}(\varepsilon\xi)] + 2\gamma i \sum_{j=1}^d \langle b_j(\cdot) \rangle \frac{\sin(\mathbf{e}_j \cdot \varepsilon\xi)}{\varepsilon} \right\} \\
 & + \gamma \varepsilon^{-1} \left\langle \sum_{j=1}^d b_j(\cdot) \left[e^{i\varepsilon\mathbf{e}_j \cdot \xi} \hat{\psi}_\varepsilon(\xi, \tau_{-\mathbf{e}_j} \cdot) - e^{-i\varepsilon\mathbf{e}_j \cdot \xi} \hat{\psi}_\varepsilon(\xi, \tau_{\mathbf{e}_j} \cdot) \right] \right\rangle \\
 & = \hat{f}_\varepsilon(\xi), \quad \xi \in \left[\frac{-\pi}{\varepsilon}, \frac{\pi}{\varepsilon} \right]^d.
 \end{aligned}$$

The Fourier transform of (2.22) is given by

$$\begin{aligned}
 (2.24) \quad \hat{\psi}_\varepsilon(\xi, \omega) & - \sum_{y \in \mathbf{Z}^d} p(y) e^{-i\varepsilon y \cdot \xi} \hat{\psi}_\varepsilon(\xi, \tau_y \omega) \\
 & + \gamma P \sum_{j=1}^d b_j(\omega) \left[e^{i\varepsilon\mathbf{e}_j \cdot \xi} \hat{\psi}_\varepsilon(\xi, \tau_{-\mathbf{e}_j} \omega) - e^{-i\varepsilon\mathbf{e}_j \cdot \xi} \hat{\psi}_\varepsilon(\xi, \tau_{\mathbf{e}_j} \omega) \right] + \varepsilon \hat{\psi}_\varepsilon(\xi, \omega) \\
 & = \gamma \sum_{j=1}^d [b_j(\omega) - \langle b_j(\cdot) \rangle] [e^{-i\varepsilon\mathbf{e}_j \cdot \xi} - e^{i\varepsilon\mathbf{e}_j \cdot \xi}] \hat{u}_\varepsilon(\xi), \quad \xi \in \left[\frac{-\pi}{\varepsilon}, \frac{\pi}{\varepsilon} \right]^d, \quad \omega \in \Omega.
 \end{aligned}$$

Suppose now that $\Psi_\varepsilon^k(\zeta, \omega)$, $1 \leq k \leq d$, $\zeta \in [-\pi, \pi]^d$, $\omega \in \Omega$, are functions in $L^2([-\pi, \pi]^d \times \Omega)$ which satisfy the equation,

$$\begin{aligned}
 (2.25) \quad \Psi_\varepsilon^k(\zeta, \omega) & - \sum_{y \in \mathbf{Z}^d} p(y) e^{-iy \cdot \zeta} \Psi_\varepsilon^k(\zeta, \tau_y \omega) \\
 & + \gamma P \sum_{j=1}^d b_j(\omega) \left[e^{i\mathbf{e}_j \cdot \zeta} \Psi_\varepsilon^k(\zeta, \tau_{-\mathbf{e}_j} \omega) - e^{-i\mathbf{e}_j \cdot \zeta} \Psi_\varepsilon^k(\zeta, \tau_{\mathbf{e}_j} \omega) \right] + \varepsilon \Psi_\varepsilon^k(\zeta, \omega) \\
 & = b_k(\omega) - \langle b_k(\cdot) \rangle, \quad \zeta \in [-\pi, \pi]^d, \quad \omega \in \Omega, \quad 1 \leq k \leq d.
 \end{aligned}$$

Then it is clear that the function

$$(2.26) \quad \hat{\psi}_\varepsilon(\xi, \omega) = -2i\gamma \hat{u}_\varepsilon(\xi) \sum_{k=1}^d \sin(\varepsilon \mathbf{e}_k \cdot \xi) \Psi_\varepsilon^k(\varepsilon\xi, \omega)$$

is in $L^2\left(\left[\frac{-\pi}{\varepsilon}, \frac{\pi}{\varepsilon}\right]^d \times \Omega\right)$ provided $\hat{u}_\varepsilon(\xi)$ is bounded. Furthermore $\hat{\psi}_\varepsilon(\xi, \omega)$ given by (2.25), (2.26) satisfies (2.24). If we substitute for $\hat{\psi}_\varepsilon$ from (2.26) into (2.23) then

we obtain the equation,

$$(2.27) \quad \hat{u}_\varepsilon(\xi) \left\{ 1 + \varepsilon^{-1} [1 - \hat{p}(\varepsilon\xi)] + 2\gamma i \sum_{j=1}^d \langle b_j(\cdot) \rangle \frac{\sin(\varepsilon \mathbf{e}_j \cdot \xi)}{\varepsilon} \right. \\ \left. - 2i\gamma^2 \left\langle \sum_{j,k=1}^d b_j(\cdot) \frac{\sin(\varepsilon \mathbf{e}_k \cdot \xi)}{\varepsilon} \left[e^{i\varepsilon \mathbf{e}_j \cdot \xi} \Psi_\varepsilon^k(\varepsilon\xi, \tau_{-\mathbf{e}_j} \cdot) - e^{-i\varepsilon \mathbf{e}_j \cdot \xi} \Psi_\varepsilon^k(\varepsilon\xi, \tau_{\mathbf{e}_j} \cdot) \right] \right\rangle \right\} \\ = \hat{f}_\varepsilon(\xi), \quad \xi \in \left[\frac{-\pi}{\varepsilon}, \frac{\pi}{\varepsilon} \right]^d.$$

Lemma 2.4. *Suppose $d \geq 2$ and $\gamma_d = c_d/2$, where c_d satisfies (2.17). Then if $|\gamma| < \gamma_d$ and $0 < \varepsilon \leq 1$, Equation (2.25) has a unique solution $\Psi_\varepsilon^k(\zeta, \omega) \in L^2(\Omega)$. Furthermore, $\Psi_\varepsilon^k(\zeta, \omega)$ viewed as a function from $[-\pi, \pi]^d$ to $L^2(\Omega)$ is continuous and satisfies an inequality,*

$$(2.28) \quad \varepsilon^2 \sup_{\zeta \in [-\pi, \pi]^d} \langle |\Psi_\varepsilon^k(\zeta, \cdot)|^2 \rangle \leq C_d,$$

where C_d depends only on d .

Proof. We follow an argument similar to that in Lemma 2.3. Suppose $\Psi_\varepsilon^k(\zeta, \omega) \in L^2(\Omega)$ and satisfies (2.25). Let $\Phi_\varepsilon^k(\zeta, \omega)$ be defined by

$$(2.29) \quad \Phi_\varepsilon^k(\zeta, \omega) = \Psi_\varepsilon^k(\zeta, \omega) - \sum_{y \in \mathbf{Z}^d} p(y) e^{-iy \cdot \zeta} \Psi_\varepsilon^k(\zeta, \tau_y \omega) + \varepsilon \Psi_\varepsilon^k(\zeta, \omega).$$

It is easy to see that $\Phi_\varepsilon^k(\zeta, \omega) \in L^2(\Omega)$. Further, it follows from (2.9) that

$$(2.30) \quad \Psi_\varepsilon^k(\zeta, \omega) = \sum_{x \in \mathbf{Z}^d} G_{p,\varepsilon}(x) e^{-ix \cdot \zeta} \Phi_\varepsilon^k(\zeta, \tau_x \omega).$$

Now for $\eta > 0$, $\zeta \in [-\pi, \pi]^d$, $j = 1, \dots, d$ we define operators $T_{j,\eta,\zeta}$ on $L^2(\Omega)$ by

$$(2.31) \quad T_{j,\eta,\zeta} \varphi(\omega) = \sum_{x \in \mathbf{Z}^d} [G_{p,\eta}(x - \mathbf{e}_j) - G_{p,\eta}(x + \mathbf{e}_j)] e^{-ix \cdot \zeta} \varphi(\tau_x \omega).$$

We can see from Bochner's theorem [19] that $T_{j,\eta,\zeta}$ is bounded on $L^2(\Omega)$ and

$$(2.32) \quad \|T_{j,\eta,\zeta}\| \leq \sup_{\zeta' \in [-\pi, \pi]^d} \frac{2|\sin(\mathbf{e}_j \cdot \zeta')|}{1 + \eta - \hat{p}(\zeta')} \leq \frac{2}{c_d},$$

where the constant c_d is from (2.17). Further, (2.25), (2.29), (2.30) imply that $\Phi_\varepsilon^k(\zeta, \omega)$ satisfies the equation,

$$(2.33) \quad \Phi_\varepsilon^k(\zeta, \omega) - \gamma P \sum_{j=1}^d b_j(\omega) T_{j,\varepsilon,\zeta} \Phi_\varepsilon^k(\zeta, \omega) = b_k(\omega) - \langle b_k(\cdot) \rangle.$$

Hence if we take $\gamma_d = c_d/2$, it follows that for $|\gamma| < \gamma_d$ the function $\Phi_\varepsilon^k(\zeta, \omega)$ is uniquely determined by Equation (2.33). In view of (2.30) it follows that $\Psi_\varepsilon^k(\zeta, \omega)$ is the unique solution of (2.25) in $L^2(\Omega)$. We can similarly prove existence for $|\gamma| < \gamma_d$. The inequality of (2.28) follows from (2.30) on using Bochner's theorem and the fact that $\Phi_\varepsilon^k(\zeta, \cdot) \in L^2(\Omega)$ is bounded independently of ζ, ε . \square

It is clear now that we can construct the solutions to (2.21), (2.22), by using Lemma 2.4 and Equations (2.26) and (2.27). Next we wish to show that the limit of the coefficient of $\hat{u}_\varepsilon(\xi)$ in (2.27) as $\varepsilon \rightarrow 0$ exists. Evidently

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \left\{ 1 + \varepsilon^{-1}[1 - \hat{p}(\varepsilon\xi)] + 2\gamma i \sum_{j=1}^d \langle b_j(\cdot) \rangle \frac{\sin(\varepsilon \mathbf{e}_j \cdot \xi)}{\varepsilon} \right\} \\ = 1 + |\xi| + 2\gamma i \sum_{j=1}^d \langle b_j(\cdot) \rangle (\mathbf{e}_j \cdot \xi), \end{aligned}$$

so we are left to show existence of the term involving the function Ψ_ε^k . To prove this first observe from (2.30), (2.31) that

$$e^{-i\mathbf{e}_j \cdot \zeta} \Psi_\varepsilon^k(\zeta, \tau_{\mathbf{e}_j} \omega) - e^{i\mathbf{e}_j \cdot \zeta} \Psi_\varepsilon^k(\zeta, \tau_{-\mathbf{e}_j} \omega) = T_{j,\varepsilon,\zeta} \Phi_\varepsilon^k(\zeta, \omega).$$

Next we prove a version of the von Neumann ergodic theorem [19].

Lemma 2.5. *Let $\varphi \in L^2(\Omega)$ and $\xi \in \mathbf{R}^d$. Then if $\langle \varphi \rangle = 0$ the function $T_{j,\varepsilon,\xi} \varphi$ converges in $L^2(\Omega)$ to the function $T_{j,0,0} \varphi$.*

Proof. First note that we can define the operator $T_{j,0,\zeta}$, $\zeta \in [-\pi, \pi]^d$, by (2.31) and from (2.32) it is a bounded operator on $L^2(\Omega)$. To be more precise, observe that for $\eta > 0$, $T_{j,\eta,\zeta} \varphi = 0$ if $\varphi \in L^2(\Omega)$ satisfies $\partial_{j,\zeta} \varphi = 0$, where $\partial_{j,\zeta}$ is the operator on $L^2(\Omega)$ given by

$$\partial_{j,\zeta} \varphi(\omega) = e^{-i\mathbf{e}_j \cdot \zeta} \varphi(\tau_{\mathbf{e}_j} \omega) - e^{i\mathbf{e}_j \cdot \zeta} \varphi(\tau_{-\mathbf{e}_j} \omega).$$

Observe that the adjoint $\partial_{j,\zeta}^*$ of $\partial_{j,\zeta}$ is given by $\partial_{j,\zeta}^* = -\partial_{j,\zeta}$. Suppose now $\varphi = \partial_{j,\zeta} \psi$ where $\psi \in L^2(\Omega)$. Then from (2.31) we have that

(2.34)

$$T_{j,\eta,\zeta} \varphi(\omega) = \sum_{x \in \mathbf{Z}^d} [G_{p,\eta}(x + 2\mathbf{e}_j) + G_{p,\eta}(x - 2\mathbf{e}_j) - 2G_{p,\eta}(x)] e^{-ix \cdot \zeta} \psi(\tau_x \omega).$$

We can readily see, using the argument of Lemma 2.2 that there is a constant C_d , depending only on d such that

$$\begin{aligned} |G_{p,\eta}(x + 2\mathbf{e}_j) + G_{p,\eta}(x - 2\mathbf{e}_j) - 2G_{p,\eta}(x)| \\ \leq C_d/[1 + \eta^2|x|^2][1 + |x|^{d+1}], \quad x \in \mathbf{Z}^d. \end{aligned}$$

Hence if $\varphi = \partial_{j,\zeta} \psi$, $\psi \in L^2(\Omega)$, we can define $T_{j,0,\zeta} \varphi$ by setting $\eta = 0$ in the formula (2.34). If $\partial_{j,\zeta} \varphi = 0$ we set $T_{j,0,\zeta} \varphi = 0$. In view of (2.32) this defines $T_{j,0,\zeta}$ as a bounded operator on $L^2(\Omega)$ which has the property that, for any $\varphi \in L^2(\Omega)$, $T_{j,\eta,\zeta} \varphi$ converges in $L^2(\Omega)$ to $T_{j,0,\zeta} \varphi$ as $\eta \rightarrow 0$.

So far we have not used the ergodicity of the translation operators τ_x , $x \in \mathbf{Z}^d$. Suppose now $\varphi \in L^2(\Omega)$ and $\langle \varphi \rangle = 0$. Then ergodicity of the translation operators implies that for any $\delta > 0$ there exists $\psi_\delta \in L^2(\Omega)$ such that $\|\varphi - \partial_{j,0} \psi_\delta\| < \delta$. Now using the fact that

$$\partial_{j,0} \psi_\delta = \partial_{j,\varepsilon\xi} \psi_\delta + \psi'_{\delta,\varepsilon}$$

where $\psi'_{\delta,\varepsilon} \in L^2(\Omega)$ satisfies $\|\psi'_{\delta,\varepsilon}\| \leq 2\varepsilon|\mathbf{e}_j \cdot \xi| \|\psi_\delta\|$, it follows from (2.32), (2.34) that $T_{j,\varepsilon,\xi}[\partial_{j,0} \psi_\delta]$ converges in $L^2(\Omega)$ as $\varepsilon \rightarrow 0$ to the function $T_{j,0,0}[\partial_{j,0} \psi_\delta]$. The result then follows from (2.32) on letting $\delta \rightarrow 0$. \square

Lemma 2.5 enables us to define the vector $q_\gamma \in \mathbf{R}^d$ of (1.7).

Lemma 2.6. *There exists a constant $\gamma_d > 0$, depending only on d , such that if $|\gamma| < \gamma_d$ and $0 < \varepsilon \leq 1$, Equation (2.27) has a unique solution $\hat{u}_\varepsilon \in L^2\left(\left[-\frac{\pi}{\varepsilon}, \frac{\pi}{\varepsilon}\right]^d\right)$. There is a vector $q_\gamma \in \mathbf{C}^d$, analytic in γ and satisfying $|q_\gamma| < 1$, such that if $u(x)$, $x \in \mathbf{R}^d$, has Fourier transform satisfying (1.7) and $u_\varepsilon(x)$, $x \in \mathbf{Z}_\varepsilon^d$, is the Fourier inverse of \hat{u}_ε , then*

$$(2.35) \quad \lim_{\varepsilon \rightarrow 0} \|u_\varepsilon - u\|_\varepsilon = 0.$$

Proof. First observe that we can rewrite (2.27) as

$$(2.36) \quad \hat{u}_\varepsilon(\xi) \left\{ 1 + \varepsilon^{-1} [1 - \hat{p}(\varepsilon\xi)] + 2\gamma i \sum_{j=1}^d \langle b_j(\cdot) \rangle \frac{\sin(\varepsilon \mathbf{e}_j \cdot \xi)}{\varepsilon} \right. \\ \left. + 2i\gamma^2 \left\langle \sum_{j,k=1}^d b_j(\cdot) \frac{\sin(\varepsilon \mathbf{e}_k \cdot \xi)}{\varepsilon} T_{j,\varepsilon,\varepsilon\xi} \Phi_\varepsilon^k(\varepsilon\xi, \cdot) \right\rangle \right\} \\ = \hat{f}_\varepsilon(\xi), \quad \xi \in \left[\frac{-\pi}{\varepsilon}, \frac{\pi}{\varepsilon} \right]^d,$$

where $\Phi_\varepsilon^k(\zeta, \omega)$ is the solution of (2.33). Note that from (2.36), (2.17) and (2.32) one has that $\hat{u}_\varepsilon \in L^2\left(\left[-\frac{\pi}{\varepsilon}, \frac{\pi}{\varepsilon}\right]^d\right)$ provided $0 < \varepsilon \leq 1$, $|\gamma| < \gamma_d$ and γ_d is sufficiently small, depending only on d . It is also clear from Lemma 2.5 and (2.32) that for any $\varphi \in L^2(\Omega)$ with $\langle \varphi \rangle = 0$ and nonnegative integer m ,

$$\lim_{\varepsilon \rightarrow 0} T_{j,\varepsilon,\varepsilon\xi}^m \varphi = T_{j,0,0}^m \varphi,$$

where the convergence is in the L^2 norm. It follows then from (2.33) that

$$\lim_{\varepsilon \rightarrow 0} T_{j,\varepsilon,\varepsilon\xi} \Phi_\varepsilon^k(\varepsilon\xi, \cdot) = T_{j,0,0} \Phi_0^k(0, \cdot).$$

Hence from (2.36) we define the vector $q_\gamma = (q_1, \dots, q_d)$ by

$$(2.37) \quad q_k = 2\gamma \langle b_k(\cdot) \rangle + 2\gamma^2 \sum_{j=1}^d \langle b_j(\cdot) T_{j,0,0} \Phi_0^k(0, \cdot) \rangle.$$

Evidently we may choose γ_d sufficiently small, depending only on d , such that $|q_\gamma| < 1$. The limit (2.35) now follows from the argument of Lemma 2.1. \square

Lemma 2.7. *Let $\Psi_\varepsilon^k(\zeta, \omega)$ be the function defined in Lemma 2.4 for $|\gamma| < \gamma_d$, $0 < \varepsilon \leq 1$. Then for any $\xi \in \mathbf{R}^d$, there is the limit,*

$$(2.38) \quad \lim_{\varepsilon \rightarrow 0} \varepsilon^2 \langle |\Psi_\varepsilon^k(\varepsilon\xi, \cdot)|^2 \rangle = 0.$$

Proof. We first consider the zeroth order contribution in γ to Ψ_ε^k . Thus in (2.33) we make the approximation $\Phi_\varepsilon^k(\zeta, \omega) \simeq b_k(\omega) - \langle b_k(\cdot) \rangle$. In this approximation we have from (2.30) and Bochner's theorem the identity,

$$(2.39) \quad \varepsilon^2 \langle |\Psi_\varepsilon^k(\varepsilon\xi, \cdot)|^2 \rangle = \int_{[-\pi, \pi]^d} d\mu_\varphi(\zeta) \left[\frac{\varepsilon}{1 + \varepsilon - \hat{p}(\zeta - \varepsilon\xi)} \right]^2,$$

where $d\mu_\varphi$ is the finite positive Borel measure on $[-\pi, \pi]^d$, satisfying

$$\langle \varphi(\tau_x \cdot) \overline{\varphi(\cdot)} \rangle = \int_{[-\pi, \pi]^d} e^{ix \cdot \zeta} d\mu_\varphi(\zeta), \quad x \in \mathbf{Z}^d,$$

and $\varphi(\omega) = b_k(\omega) - \langle b_k(\cdot) \rangle$. Observe now that

$$(2.40) \quad \left[\frac{\varepsilon}{1 + \varepsilon - \hat{p}(\zeta - \varepsilon\xi)} \right]^2 \leq 1, \quad \zeta \in [-\pi, \pi]^d,$$

$$(2.41) \quad \lim_{\varepsilon \rightarrow 0} \left[\frac{\varepsilon}{1 + \varepsilon - \hat{p}(\zeta - \varepsilon\xi)} \right]^2 = 0, \quad \zeta \in [-\pi, \pi]^d \setminus \{0\}.$$

Since $\langle \varphi \rangle = 0$, one has by ergodicity of the translation operators τ_x , $x \in \mathbf{Z}^d$, that $\mu_\varphi(\{0\}) = 0$. It follows now by dominated convergence that (2.38) holds in this approximation.

To deal with the general case note that (2.39) holds with $\varphi(\omega) = \varphi_\varepsilon(\omega) = \Phi_\varepsilon^k(\varepsilon\xi, \omega)$. The result follows then from the above argument if we use the fact that $d\mu_{\phi_\varepsilon} \leq 2d\mu_{\phi_0} + 2d\mu_{\phi_\varepsilon - \phi_0}$ and the fact that $\lim_{\varepsilon \rightarrow 0} \mu_{\phi_\varepsilon - \phi_0}([-\pi, \pi]^d) = \|\phi_\varepsilon - \phi_0\|^2 = 0$. \square

Proof of Theorem 1.1. We just need to prove (1.8). In view of Lemma 2.6 and (2.26) it will be sufficient to show that

$$\lim_{\varepsilon \rightarrow 0} \int_{[-\frac{\pi}{\varepsilon}, \frac{\pi}{\varepsilon}]^d} |\hat{f}_\varepsilon(\xi)|^2 |\xi|^2 \varepsilon^2 \langle |\Psi_\varepsilon^k(\varepsilon\xi, \cdot)|^2 \rangle d\xi = 0.$$

This follows by dominated convergence from (2.28) and Lemma 2.7. \square

Finally we show that if (1.9) holds then the vector q_γ defined by (2.37) is zero.

Lemma 2.8. *Suppose (1.9) holds and the conditions of Theorem 1.1. Then $q_\gamma = 0$ in (1.7).*

Proof. In view of (2.37) we need to show that

$$(2.42) \quad \langle b_j(\cdot) T_{j,0,0} \Phi_0^k(0, \cdot) \rangle = 0, \quad 1 \leq j, k \leq d,$$

where Φ_0^k satisfies (2.33). To do this let $L_R^2(\Omega)$ be the subset of functions $\Phi \in L^2(\Omega)$ which have the property that

$$\left\langle \Phi(\tau_x \cdot) \prod_{i=1}^n b_{k_i}(\tau_{x_i} \cdot) \right\rangle = (-1)^{n+1} \left\langle \Phi(\tau_{-x} \cdot) \prod_{i=1}^n b_{k_i}(\tau_{-x_i} \cdot) \right\rangle, \\ x, x_i \in \mathbf{Z}^d, \quad 1 \leq k_i \leq d, \quad i = 1, \dots, n, \quad n \geq 0.$$

Evidently $\Phi \in L_R^2(\Omega)$ implies $\langle \Phi \rangle = 0$. In view of (1.9) $b_i(\cdot) \in L_R^2(\Omega)$, $1 \leq i \leq d$. Observe now that for any $\eta > 0$ the operator $b_j(\omega) T_{j,\eta,0}$ takes $L_R^2(\Omega)$ into $L_R^2(\Omega)$. This follows from the fact that $G_{p,\eta}(x) = G_{p,\eta}(-x)$, $x \in \mathbf{Z}^d$. Now let $\Phi_\eta^k(\omega)$ be the solution in $L_R^2(\Omega)$ to the equation

$$(2.43) \quad \Phi_\eta^k(\omega) - \gamma \sum_{j=1}^d b_j(\omega) T_{j,\eta,0} \Phi_\eta^k(\omega) = b_k(\omega).$$

Clearly a unique solution Φ_η^k exists if $|\gamma| < \gamma_d$. Comparing (2.43) and (2.33) one sees that $\Phi_\varepsilon^k(0, \omega) = \Phi_\varepsilon^k(\omega) \in L_R^2(\Omega)$. Taking the limit as $\varepsilon \rightarrow 0$ we conclude that $\Phi_0^k(0, \omega) \in L_R^2(\Omega)$, whence (2.42) follows. \square

3. Proof of Theorem 1.2

Since $\langle \mathbf{b}(\cdot) \rangle = 0$ we have from (2.33), (2.36) that

$$e(\varepsilon\xi)q_{\gamma,\varepsilon}(\varepsilon\xi)e(-\varepsilon\xi) = -2i \sum_{k=1}^d \sin(\varepsilon \mathbf{e}_k \cdot \xi) \sum_{n=1}^{\infty} \gamma^{n+1} \left\langle \left[\sum_{j=1}^d b_j(\cdot) T_{j,\varepsilon,\varepsilon\xi} P \right]^n b_k(\cdot) \right\rangle.$$

It is easy to see from the argument of Lemma 2.8 that if (1.9) holds, then

$$(3.1) \quad \left\langle \left[\sum_{j=1}^d b_j(\cdot) T_{j,\varepsilon,0} P \right]^n b_k(\cdot) \right\rangle = 0, \quad 1 \leq k \leq d, \quad n \geq 1.$$

Hence we may write

$$(3.2) \quad e(\varepsilon\xi)q_{\gamma,\varepsilon}(\varepsilon\xi)e(-\varepsilon\xi) = -2i \sum_{k=1}^d \sin(\varepsilon \mathbf{e}_k \cdot \xi) \sum_{n=0}^{\infty} \gamma^{n+2} \sum_{m=0}^n \left\langle \left[\sum_{j=1}^d b_j(\cdot) T_{j,\varepsilon,0} P \right]^m \left[\sum_{j=1}^d b_j(\cdot) \{T_{j,\varepsilon,\varepsilon\xi} - T_{j,\varepsilon,0}\} P \right] \left[\sum_{j=1}^d b_j(\cdot) T_{j,\varepsilon,\varepsilon\xi} P \right]^{n-m} b_k(\cdot) \right\rangle.$$

If $x \in \mathbf{Z}^d$ with $x \cdot \mathbf{e}_1 > 0$, we denote by $L_1(x)$ all the points of \mathbf{Z}^d which lie on the line segment joining the point $x - (x \cdot \mathbf{e}_1)\mathbf{e}_1$ and x , excluding the point x . If $x \cdot \mathbf{e}_1 = 0$, then $L_1(x)$ is the empty set. If $x \cdot \mathbf{e}_1 < 0$, then $L_1(x)$ is all the points of \mathbf{Z}^d which lie on the line segment joining the point $x - (x \cdot \mathbf{e}_1)\mathbf{e}_1$ and x , excluding the point $x - (x \cdot \mathbf{e}_1)\mathbf{e}_1$. Similarly if $x \cdot \mathbf{e}_2 > 0$ we denote by $L_2(x)$ all the points of \mathbf{Z}^d which lie on the line segment joining the point $x - (x \cdot \mathbf{e}_1)\mathbf{e}_1 - (x \cdot \mathbf{e}_2)\mathbf{e}_2$ and $x - (x \cdot \mathbf{e}_1)\mathbf{e}_1$, excluding the point $x - (x \cdot \mathbf{e}_1)\mathbf{e}_1$ etc. For $\eta > 0$, $\zeta \in [-\pi, \pi]^d$, $k = 1, \dots, d$, we define operators $S_{k,\eta,\zeta}$ on $L^2(\Omega)$ by

$$(3.3) \quad S_{k,\eta,\zeta} \varphi(\omega) = \sum_{j=1}^d b_j(\omega) \sum_{x \in \mathbf{Z}^d} [G_{p,\eta}(x - \mathbf{e}_j) - G_{p,\eta}(x + \mathbf{e}_j)] \operatorname{sgn}(x \cdot \mathbf{e}_k) \sum_{y \in L_k(x)} e^{-iy \cdot \zeta} \varphi(\tau_x \omega).$$

We can see now from (3.2) and (2.31) that the matrix $q_{\gamma,\varepsilon}(\zeta) = [q_{\gamma,\varepsilon,k,k'}(\zeta)]$, $1 \leq k, k' \leq d$, is given by the formula

(3.4)

$$\begin{aligned} q_{\gamma,\varepsilon,k,k'}(\zeta) = & \\ & - \sum_{n=0}^{\infty} \gamma^{n+2} \sum_{m=0}^n \left\langle \left[\sum_{j=1}^d b_j(\cdot) T_{j,\varepsilon,0} P \right]^m S_{k',\varepsilon,\zeta} P \left[\sum_{j=1}^d b_j(\cdot) T_{j,\varepsilon,\zeta} P \right]^{n-m} b_{k'}(\cdot) \right\rangle \\ & - \sum_{n=0}^{\infty} \gamma^{n+2} \sum_{m=0}^n \left\langle \left[\sum_{j=1}^d b_j(\cdot) T_{j,\varepsilon,0} P \right]^m e^{-i\mathbf{e}_k \cdot \zeta} S_{k,\varepsilon,\zeta} P \left[\sum_{j=1}^d b_j(\cdot) T_{j,\varepsilon,\zeta} P \right]^{n-m} b_{k'}(\cdot) \right\rangle. \end{aligned}$$

We proceed as in [3] to consider the case when the $\mathbf{b}(\tau_x \cdot)$, $x \in \mathbf{Z}^d$, are given by independent Bernoulli variables. Thus we assume that $b_j(\cdot) \equiv 0$, $j > 1$, and $b_1(\tau_x \cdot) = Y_x$, $x \in \mathbf{Z}^d$, where the variables Y_x , $x \in \mathbf{Z}^d$, are assumed to be i.i.d. Bernoulli, $Y_x = \pm 1$ with equal probability. Using the notation of §4 of [3] we introduce the Fock spaces $\mathcal{F}^r(\mathbf{Z}^d)$, $1 < r < \infty$, with $\mathcal{F}^2(\mathbf{Z}^d)$ isomorphic to $L^2(\Omega)$. Now we have seen from (2.32) that $T_{1,\varepsilon,\zeta}$ is a bounded operator on $L^2(\Omega)$ with norm $\|T_{1,\varepsilon,\zeta}\|_2 \leq 2/c_d$. We have also seen in Lemma 2.5 that

$$(3.5) \quad \lim_{\varepsilon \rightarrow 0} \|[T_{1,\varepsilon,\zeta} - T_{1,0,\zeta}]\varphi\|_2 = 0, \quad \varphi \in L^2(\Omega).$$

We note now as in [3] that the Calderon-Zygmund theorem implies that a similar result holds for all Fock spaces $\mathcal{F}^r(\mathbf{Z}^d)$, $1 < r < \infty$. We shall show that the limit in (3.5) is uniform for $\zeta \in [-\pi, \pi]^d$, provided one projects orthogonal to the constant function.

Lemma 3.1. *Suppose $1 < r < \infty$ and regard $T_{1,\varepsilon,\zeta}$ as an operator on $\mathcal{F}^r(\mathbf{Z}^d)$. Then there is a constant $C_{r,d}$ depending only on r, d such that $\|T_{1,\varepsilon,\zeta}\|_r \leq C_{r,d}$. There is also the limit,*

$$(3.6) \quad \lim_{\varepsilon \rightarrow 0} \left\{ \sup_{\zeta \in [-\pi, \pi]^d} \|[T_{1,\varepsilon,\zeta} P - T_{1,0,\zeta} P]\varphi\|_r \right\} = 0, \quad \varphi \in \mathcal{F}^r(\mathbf{Z}^d).$$

Proof. Let $T_{\varepsilon,\zeta}$ be the integral operator on $L^r(\mathbf{Z}^d)$ defined by

$$T_{\varepsilon,\zeta} f(x) = \sum_{y \in \mathbf{Z}^d} K_{\varepsilon}(x-y) e^{-i(x-y) \cdot \zeta} f(y), \quad x \in \mathbf{Z}^d,$$

where $K_{\varepsilon}(x) = G_{p,\varepsilon}(x - \mathbf{e}_1) - G_{p,\varepsilon}(x + \mathbf{e}_1)$. It follows from (2.17) that

$$(3.7) \quad |\hat{K}_{\varepsilon}(\xi)| \leq 2/c_d, \quad \xi \in [-\pi, \pi]^d.$$

We may also apply the argument of Lemma 2.2 to see that there is a constant $C_d > 0$, depending only on d , such that

$$(3.8) \quad \begin{aligned} |K_{\varepsilon}(x)| &\leq C_d/[1 + \varepsilon^2|x|^2][1 + |x|^d], \quad x \in \mathbf{Z}^d, \\ |K_{\varepsilon}(x + \mathbf{e}_i) - K_{\varepsilon}(x)| &\leq C_d/[1 + \varepsilon^2|x|^2][1 + |x|^{d+1}], \quad x \in \mathbf{Z}^d. \end{aligned}$$

It follows now from (3.7), (3.8) that $T_{\varepsilon,\zeta}$ is a bounded operator on $L^r(\mathbf{Z}^d)$ and there is a constant $C_{r,d}$ depending only on r, d such that $\|T_{\varepsilon,\zeta}\|_r \leq C_{r,d}$. This is

a discrete version of the Calderon-Zygmund Theorem [22]. Now, arguing as in [3], we see that this also implies $\|T_{1,\varepsilon,\zeta}\|_r \leq C_{r,d}$.

Next we show that

$$(3.9) \quad \lim_{\varepsilon \rightarrow 0} \left\{ \sup_{\zeta \in [-\pi, \pi]^d} \|[T_{\varepsilon,\zeta} - T_{0,\zeta}]f\|_r \right\} = 0, \quad f \in L^r(\mathbf{Z}^d).$$

We first show that (3.9) holds for $r = 2$. We can do this by going to the Fourier representation of $T_{\varepsilon,\zeta}$, which we denote $\hat{T}_{\varepsilon,\zeta}$. Thus,

$$(3.10) \quad \hat{T}_{\varepsilon,\zeta} \hat{f}(\xi) = \hat{K}_\varepsilon(\xi - \zeta) \hat{f}(\xi), \quad \xi \in [-\pi, \pi]^d.$$

Observe now that since $\hat{f} \in L^2([-\pi, \pi]^d)$, for any $\delta > 0$ there exists $\nu > 0$ such that

$$(3.11) \quad \int_{|\xi - \zeta| < \nu} |\hat{f}(\xi)|^2 d\xi < \delta, \quad \zeta \in [-\pi, \pi]^d.$$

It is easy to see that for any $\nu > 0$, $\hat{K}_\varepsilon(\xi)$ converges uniformly to $\hat{K}_0(\xi)$ in the region $\xi \in [-\pi, \pi]^d$, $|\xi| > \nu$. It follows from this and (3.7), (3.10), (3.11) that (3.9) holds for $r = 2$.

To prove (3.9) for general r , $1 < r < \infty$, we define for $n = 1, 2, \dots$, $\zeta \in [-\pi, \pi]^d$ an operator $A_{n,\zeta}$ on functions $f: \mathbf{Z}^d \rightarrow \mathbf{C}$ by

$$A_{n,\zeta} f(x) = \sum_{z \in \mathbf{Z}^d} \varphi_n(z) e^{iz \cdot \zeta} f(x+z), \quad x \in \mathbf{Z}^d.$$

Here φ_n is the probability density for the standard random walk in \mathbf{Z}^d after n steps of the walk. Thus $\varphi_n(z) \geq 0$, $z \in \mathbf{Z}^d$, and

$$\sum_{z \in \mathbf{Z}^d} \varphi_n(z) = 1.$$

It follows that $A_{n,\zeta}$ is a bounded operator on $L^r(\mathbf{Z}^d)$ with $\|A_{n,\zeta}\|_r \leq 1$. Observe now that

$$T_{\varepsilon,\zeta}[f - A_{n,\zeta}f](x) = \sum_{y \in \mathbf{Z}^d} K_{\varepsilon,n}(x-y) e^{-i(x-y) \cdot \zeta} f(y), \quad x \in \mathbf{Z}^d,$$

where the function $K_{\varepsilon,n}$ is given by

$$K_{\varepsilon,n}(x) = \sum_{z \in \mathbf{Z}^d} [K_\varepsilon(x) - K_\varepsilon(x+z)] \varphi_n(z).$$

It is easy to see from (3.8) that $K_{\varepsilon,n} \in L^1(\mathbf{Z}^d)$ with norm $\|K_{\varepsilon,n}\|_1$ satisfying $\|K_{\varepsilon,n}\|_1 \leq C_{d,n}$, where the constant $C_{d,n}$ depends only on d and n . Further, for any $x \in \mathbf{Z}^d$, $K_{\varepsilon,n}(x)$ converges to $K_{0,n}(x)$. We conclude that for any function $f \in L^r(\mathbf{Z}^d)$ one has

$$\lim_{\varepsilon \rightarrow 0} \left\{ \sup_{\zeta \in [-\pi, \pi]^d} \|[T_{\varepsilon,\zeta} - T_{0,\zeta}][f - A_{n,\zeta}f]\|_r \right\} = 0,$$

for any $n = 1, 2, \dots$. The result (3.9) follows then if we can show that for any $f \in L^r(\mathbf{Z}^d)$, $1 < r < \infty$, and $\delta > 0$, there exists a positive integer N such that

$$\|A_{n,\zeta}f\|_r < \delta, \quad n \geq N, \quad \zeta \in [-\pi, \pi]^d.$$

This last inequality follows by approximating $f \in L^r(\mathbf{Z}^d)$ with a function of finite support. A similar argument implies (3.6). \square

Lemma 3.2. *Suppose $d \geq 2$, $1 < r < d$ and $q < \infty$ satisfies $1/r > 1/d + 1/q$. Then the operator $S_{k,\eta,\zeta}$ of (3.3) is a bounded operator from the space $\mathcal{F}^r(\mathbf{Z}^d)$ to $\mathcal{F}^q(\mathbf{Z}^d)$. There is a constant $C_{r,q,d}$, depending only on r, q, d , such that the norm $\|S_{k,\eta,\zeta}\|_{r,q}$ of the operator $S_{k,\eta,\zeta}$ satisfies the inequality $\|S_{k,\eta,\zeta}\|_{r,q} \leq C_{r,q,d}$. There is also the limit*

$$(3.12) \quad \lim_{\varepsilon \rightarrow 0} \left\{ \sup_{\zeta \in [-\pi, \pi]^d} \|[S_{k,\varepsilon,\zeta}P - S_{k,0,\zeta}P]\varphi\|_{r,q} \right\} = 0, \quad \varphi \in \mathcal{F}^r(\mathbf{Z}^d).$$

Proof. We proceed as in Lemma 3.1. Let $S_{\varepsilon,\zeta}$ be the operator on functions $f \in L^r(\mathbf{Z}^d)$ defined by

$$S_{\varepsilon,\zeta}f(x) = \sum_{y \in \mathbf{Z}^d} K_{\varepsilon,\zeta}(x-y)f(y), \quad x \in \mathbf{Z}^d,$$

where

$$K_{\varepsilon,\zeta}(x) = [G_{p,\varepsilon}(x - \mathbf{e}_1) - G_{p,\varepsilon}(x + \mathbf{e}_1)] \operatorname{sgn}(x \cdot \mathbf{e}_k) \sum_{z \in L_1(x)} e^{-iz \cdot \zeta}.$$

It follows from (3.8) that there is a constant $C_d > 0$, depending only on d , such that

$$|K_{\varepsilon,\zeta}(x)| \leq C_d/[1 + |x|^{d-1}], \quad x \in \mathbf{Z}^d.$$

Hence $K_{\varepsilon,\zeta} \in L^s(\mathbf{Z}^d)$ for any $s > d/(d-1)$. It follows now by Young's inequality that $S_{\varepsilon,\zeta}f \in L^q(\mathbf{Z}^d)$ for any $q > 0$ satisfying $1/r > 1/d + 1/q$. Further, there is a constant $C_{r,q,d}$, depending only on r, q, d , such that the norm $\|S_{\varepsilon,\zeta}\|_{r,q}$ of $S_{\varepsilon,\zeta}$ as an operator from $L^r(\mathbf{Z}^d)$ to $L^q(\mathbf{Z}^d)$ satisfies $\|S_{\varepsilon,\zeta}\|_{r,q} \leq C_{r,q,d}$. It is easy to see that for any $x \in \mathbf{Z}^d$,

$$\lim_{\varepsilon \rightarrow 0} \left\{ \sup_{\zeta \in [-\pi, \pi]^d} |K_{\varepsilon,\zeta}(x) - K_{0,\zeta}(x)| \right\} = 0.$$

Hence, if $S_{0,\zeta}$ is the operator with kernel $K_{0,\zeta}$, one has

$$\lim_{\varepsilon \rightarrow 0} \left\{ \sup_{\zeta \in [-\pi, \pi]^d} \|[S_{\varepsilon,\zeta} - S_{0,\zeta}]f\|_q \right\} = 0, \quad f \in L^r(\mathbf{Z}^d),$$

provided $q < \infty$ satisfies $1/r > 1/d + 1/q$. Now, arguing as in [3] we obtain the result. \square

Corollary 3.1. *Suppose $b_j(\cdot) \equiv 0$, $j > 1$ and $b_1(\tau_x \cdot) = Y_x$, $x \in \mathbf{Z}^d$, where the Y_x , $x \in \mathbf{Z}^d$, are independent Bernoulli variables with zero mean. Then Theorem 1.2 holds.*

Proof. Observe that $b_k(\cdot) \in \mathcal{F}^r(\mathbf{Z}^d)$ for all $r \geq 1$. The result follows now from Lemmas 3.1, 3.2 and the representation (3.4) for $q_{\gamma,\varepsilon}(\zeta)$. \square

Next we wish to generalize our method to all $\mathbf{b}(\cdot)$ such that $\mathbf{b}(\tau_x \cdot)$, $x \in \mathbf{Z}^d$, are independent random variables. To do this we pursue the method developed in §5 of [3]. Thus we define a spin space S as all $s = (m_1, \dots, m_d)$ where the m_i are

nonnegative integers satisfying $|s| = m_1 + \cdots + m_d \geq 1$. For $s \in S$ we define a random variable b_s by

$$b_s(\cdot) = \prod_{i=1}^d b_i(\cdot)^{m_i}, \quad s = (m_1, \dots, m_d),$$

and a random variable $Y_{0,s} = b_s - \langle b_s \rangle$. Then for $s \in S$, $n \in \mathbf{Z}^d$ we define the variable $Y_{n,s}$ to be the translate of $Y_{0,s}$, i.e., $Y_{n,s}(\cdot) = Y_{0,s}(\tau_n \cdot)$. For $1 < r < \infty$ we may define Fock spaces $\mathcal{F}_S^r(\mathbf{Z}^d)$ of many particle wave functions where the particles move in \mathbf{Z}^d and have spin in S . We can also, for any $\delta > 0$, define a mapping U_δ from $\mathcal{F}_S^2(\mathbf{Z}^d)$ to functions on Ω by

$$(3.13) \quad U_\delta \psi = \psi_0 + \sum_{N=1}^{\infty} \sum_{\substack{\{n_1, \dots, n_N\} \in \mathbf{Z}^{d,N} \\ s_i \in S, 1 \leq i \leq N}} \psi_N(n_1, s_1, n_2, s_2, \dots, n_N, s_N) \\ \times \delta^{|s_1| + \cdots + |s_N|} Y_{n_1, s_1} Y_{n_2, s_2} \cdots Y_{n_N, s_N}.$$

Following the argument of Lemma 5.1 of [3] we have:

Lemma 3.3. *There exists $\delta > 0$, depending only on d , such that the mapping U_δ of (3.13) is a bounded operator from $\mathcal{F}_S^2(\mathbf{Z}^d)$ to $L^2(\Omega)$ satisfying $\|U_\delta\| \leq 1$.*

We shall use the operator $U_\delta = U$ of Lemma 3.3 to transfer the equation of (2.33) from $L^2(\Omega)$ to $\mathcal{F}_S^2(\mathbf{Z}^d)$. Thus we shall need to construct operators $P_{\mathcal{F}}$, $B_{j,\mathcal{F}}$, $T_{j,\varepsilon,\zeta,\mathcal{F}}$, $1 \leq j \leq d$, with the property that they are bounded operators on $\mathcal{F}_S^2(\mathbf{Z}^d)$ and satisfy

$$(3.14) \quad UP_{\mathcal{F}} = PU, \quad UB_{j,\mathcal{F}} = b_j(\cdot)U, \quad UT_{j,\varepsilon,\zeta,\mathcal{F}} = T_{j,\varepsilon,\zeta}U.$$

The simplest of these operators to construct is $P_{\mathcal{F}}$. Thus if $\psi = \{\psi_N : N = 0, 1, 2, \dots\} \in \mathcal{F}_S^2(\mathbf{Z}^d)$ then $P_{\mathcal{F}}\psi = \{(P_{\mathcal{F}}\psi)_N : N = 0, 1, 2, \dots\}$, where $(P_{\mathcal{F}}\psi)_0 = 0$, $(P_{\mathcal{F}}\psi)_N = \psi_N$, $N \geq 1$. Similarly we have $T_{j,\varepsilon,\zeta,\mathcal{F}}\psi = \{(T_{j,\varepsilon,\zeta,\mathcal{F}}\psi)_N : N = 0, 1, 2, \dots\}$, where

$$(T_{j,\varepsilon,\zeta,\mathcal{F}}\psi)_0 = [e^{-i\zeta \cdot \mathbf{e}_j} - e^{i\zeta \cdot \mathbf{e}_j}] \hat{G}_{p,\varepsilon}(-\zeta) \psi_0, \\ (T_{j,\varepsilon,\zeta,\mathcal{F}}\psi)_N(n_1, s_1, \dots, n_N, s_N) = \\ \sum_{x \in \mathbf{Z}^d} [G_{p,\varepsilon}(x - \mathbf{e}_j) - G_{p,\varepsilon}(x + \mathbf{e}_j)] e^{-ix \cdot \zeta} \psi_N(n_1 - x, s_1, \dots, n_N - x, s_N), \quad N \geq 1.$$

The most complicated of the operators to define is $B_{j,\mathcal{F}}$. We have

$$(B_{j,\mathcal{F}}\psi)_N(n_1, s_1, \dots, n_N, s_N) = \langle b_j(\cdot) \rangle \psi_N(n_1, s_1, \dots, n_N, s_N) \\ + \sum_{s \in S} [\langle b_j b_s \rangle - \langle b_j \rangle \langle b_s \rangle] \delta^{|s|} \psi_{N+1}(0, s, n_1, s_1, \dots, n_N, s_N),$$

if $n_k \neq 0$, $1 \leq k \leq N$, $N \geq 0$;

$$\begin{aligned}
(B_{j,\mathcal{F}}\psi)_N(0, s_1, n_2, s_2, \dots, n_N, s_N) &= 0, \\
&\text{if } n_k \neq 0, \quad 2 \leq k \leq N, \quad s_1 = (m_1, \dots, m_d), \quad m_j = 0; \\
(B_{j,\mathcal{F}}\psi)_N(0, s_1, n_2, s_2, \dots, n_N, s_N) &= \delta^{-1}\psi_N(0, s'_1, n_2, s_2, \dots, n_N, s_N), \\
&\text{if } n_k \neq 0, \quad 2 \leq k \leq N, \quad s_1 = (m_1, \dots, m_d), \quad m_j > 0, \quad |s_1| > 1, \\
&\quad s'_1 = (m_1, \dots, m_{j-1}, m_j - 1, m_{j+1}, \dots, m_d); \\
(B_{j,\mathcal{F}}\psi)_N(0, s_1, n_2, s_2, \dots, n_N, s_N) &= \delta^{-1}\psi_{N-1}(n_2, s_2, \dots, n_N, s_N) \\
&\quad - \sum_{s \in \mathcal{S}} \langle b_s \rangle \delta^{|s|-1} \psi_N(0, s, n_2, s_2, \dots, n_N, s_N),
\end{aligned}$$

if $n_k \neq 0$, $2 \leq k \leq N$ and $s_1 = (0, 0, \dots, 1, 0, \dots, 0)$, with 1 in the i th position.

Lemma 3.4. For $1 \leq k \leq d$ let $\varphi_k \in \mathcal{F}_S^2(\mathbf{Z}^d)$ be the wave function $\varphi_k = \{\varphi_{k,N} : N = 0, 1, 2, \dots\}$ where $\varphi_{k,N} \equiv 0$ if $N \neq 1$ and $\varphi_{k,1}$ is defined by

$$\varphi_{k,1}(n, s) = 1 \quad \text{if } n = 0, \quad s = (0, \dots, 0, 1, 0, \dots, 0),$$

with 1 in the k th position, $\varphi_{k,1}(n, s) = 0$, otherwise. Then there exists a constant γ_d depending only on d such that if $|\gamma| < \gamma_d$, the equation

$$(3.15) \quad \Phi_{\varepsilon,\mathcal{F}}^k(\zeta) - \gamma P_{\mathcal{F}} \sum_{j=1}^d B_{j,\mathcal{F}} T_{j,\varepsilon,\zeta,\mathcal{F}} \Phi_{\varepsilon,\mathcal{F}}^k(\zeta) = \varphi_k,$$

has a unique solution $\Phi_{\varepsilon,\mathcal{F}}^k(\zeta) \in \mathcal{F}_S^2(\mathbf{Z}^d)$. Further the solution to (2.33) is given by the function $\Phi_{\varepsilon}^k(\zeta, \omega) = U \Phi_{\varepsilon,\mathcal{F}}^k(\zeta) \in L^2(\Omega)$.

Proof. Evidently $P_{\mathcal{F}}$ is a bounded operator on $\mathcal{F}_S^2(\mathbf{Z}^d)$ with norm 1. We can also easily see that $T_{j,\varepsilon,\zeta,\mathcal{F}}$ is a bounded operator on $\mathcal{F}_S^2(\mathbf{Z}^d)$ with norm bounded by the RHS of (2.32). Also $B_{j,\mathcal{F}}$ is bounded with norm depending only on d . Hence we may choose γ_d small enough, depending only on d , such that (3.15) is uniquely solvable. The fact that $U \Phi_{\varepsilon,\mathcal{F}}^k(\zeta)$ solves (2.33) comes from (3.14) and $U \varphi_k = b_k(\cdot) - \langle b_k \rangle$. \square

Proof of Theorem 1.2. Observe from (2.36) that $q_{\varepsilon,\gamma}(\zeta)$ is defined in terms of $\Phi_{\varepsilon}^k(\zeta, \omega)$. In view of Lemma 3.4 we get a representation of $q_{\varepsilon,\gamma}(\zeta)$ similar to (3.4) but with the operators $T_{j,\varepsilon,\zeta}$, $S_{k,\varepsilon,\zeta}$, $b_j(\cdot)$ on $L^2(\Omega)$ replaced by the corresponding operators $T_{j,\varepsilon,\zeta,\mathcal{F}}$, $S_{k,\varepsilon,\zeta,\mathcal{F}}$, $B_{j,\mathcal{F}}$ on Fock space. Here the operator $S_{k,\varepsilon,\zeta,\mathcal{F}}$ satisfies $U S_{k,\varepsilon,\zeta,\mathcal{F}} = S_{k,\varepsilon,\zeta} U$. We now argue exactly as in Corollary 3.1. \square

4. Proof of Theorem 1.3

We assume in this section that the function p is given by (1.2). Let $G_{p,\eta}(x)$ be the function defined by (2.9). Then $\hat{G}_{p,\eta}(\xi)$ satisfies the inequality,

$$0 \leq \hat{G}_{p,\eta}(\xi) \leq C/[\eta + |\xi|^2],$$

for some universal constant C . One can also see that if $d \geq 2$ and $0 < \eta < 1$, there is a constant C_d , depending only on d , such that

$$(4.1) \quad |G_{p,\eta}(x - \mathbf{e}_k) - G_{p,\eta}(x + \mathbf{e}_k)| \leq C_d \exp[-\sqrt{\eta}|x|/C_d] / [1 + |x|^{d-1}].$$

Next we prove the analogue of Lemma 2.3.

Lemma 4.1. *Suppose $d \geq 2$ and $f : \mathbf{R}^d \rightarrow \mathbf{R}$ is a C^∞ function with compact support. Then if $\gamma \in \mathbf{C}$ satisfies $|\gamma| < \varepsilon/\sqrt{2d}$ and $0 < \varepsilon \leq 1$, Equation (1.4) has a unique solution $u_\varepsilon(x, \omega)$ in $L^2(\mathbf{Z}_\varepsilon^d)$. Further, there is a constant C_d , depending only on d , such that $\|u_\varepsilon(\cdot, \omega)\|_\varepsilon \leq C_d \|f\|_\varepsilon$.*

Proof. Suppose $u_\varepsilon(x, \omega)$ is in $L^2(\mathbf{Z}_\varepsilon^d)$ and satisfies (1.4). Let $v_\varepsilon(x, \omega)$ be defined by

$$v_\varepsilon(x, \omega) = \varepsilon^{-2}[A_p u_\varepsilon(x, \omega) + \varepsilon^2 u_\varepsilon(x, \omega)].$$

It is easy to see that $v_\varepsilon(x, \omega) \in L^2(\mathbf{Z}_\varepsilon^d)$. In view of (2.9) it follows that

$$(4.2) \quad u_\varepsilon(x, \omega) = \varepsilon^{2-d} \int_{\mathbf{Z}_\varepsilon^d} dy G_{p, \varepsilon^2} \left(\frac{x-y}{\varepsilon} \right) v_\varepsilon(y, \omega).$$

Now for $i = 1, \dots, d$ let us define operators T_i on $L^2(\mathbf{Z}_\varepsilon^d)$ by

$$T_i w(x) = \varepsilon^{1-d} \int_{\mathbf{Z}_\varepsilon^d} dy \left[G_{p, \varepsilon^2} \left(\frac{x-y}{\varepsilon} + \mathbf{e}_i \right) - G_{p, \varepsilon^2} \left(\frac{x-y}{\varepsilon} - \mathbf{e}_i \right) \right] w(y).$$

It is easy to see that T_i is a bounded operator on $L^2(\mathbf{Z}_\varepsilon^d)$. It follows from (2.10) that the norm of T_i is bounded by

$$(4.3) \quad \|T_i\| \leq \sup_{\xi \in [-\pi, \pi]^d} \left[\frac{2\varepsilon |\sin(\mathbf{e}_i \cdot \xi)|}{1 + \varepsilon^2 - \hat{p}(\xi)} \right].$$

Now (4.3) and (1.1) imply that the operator $\sum_{i=1}^d b_i(\cdot, \omega) T_i$ is bounded on $L^2(\mathbf{Z}_\varepsilon^d)$ and has norm satisfying the inequality

$$\begin{aligned} \left\| \sum_{i=1}^d b_i(\cdot, \omega) T_i \right\| &\leq \sup_{\xi \in [-\pi, \pi]^d} \left[\frac{2\varepsilon \left\{ \sum_{i=1}^d \sin^2(\mathbf{e}_i \cdot \xi) \right\}^{1/2}}{1 + \varepsilon^2 - \hat{p}(\xi)} \right] \\ &\leq 4 \sup_{z > 0} \left[\frac{\varepsilon z}{\varepsilon^2 + 2z^2/d} \right] = \sqrt{2d}. \end{aligned}$$

Observe now that (1.4) implies that $v_\varepsilon(x, \omega) \in L^2(\mathbf{Z}_\varepsilon^d)$ satisfies the equation

$$(4.4) \quad v_\varepsilon(x, \omega) - \gamma \varepsilon^{-1} \sum_{i=1}^d b_i(x/\varepsilon, \omega) T_i v_\varepsilon(x, \omega) = f(x).$$

Hence if $|\gamma| < \varepsilon/\sqrt{2d}$, the function $v_\varepsilon(x, \omega)$ is uniquely determined by (4.4). In view of (4.2) it follows that $u_\varepsilon(x, \omega) \in L^2(\mathbf{Z}_\varepsilon^d)$ is the unique solution to (1.4). To prove existence note that (4.4) has a solution $v_\varepsilon(x, \omega) \in L^2(\mathbf{Z}_\varepsilon^d)$ provided $|\gamma| < \varepsilon/\sqrt{2d}$. If we then define $u_\varepsilon(x, \omega)$ by (4.2) one can easily see that $u_\varepsilon(x, \omega)$ satisfies (1.4). \square

In order to obtain Equation (1.11) we follow a development similar to the one following Lemma 2.3. Thus we put $v_\varepsilon(x, \omega) = u_\varepsilon(x, \tau_{-x/\varepsilon} \omega)$, $x \in \mathbf{Z}_\varepsilon^d, \omega \in \Omega$, and write $v_\varepsilon(x, \omega) = u_\varepsilon(x) + \psi_\varepsilon(x, \omega)$, $x \in \mathbf{Z}_\varepsilon^d, \omega \in \Omega$, where $\langle \psi_\varepsilon(x, \cdot) \rangle = 0$, $x \in \mathbf{Z}_\varepsilon^d$. It follows that

$$u_\varepsilon(x) = \langle v_\varepsilon(x, \cdot) \rangle = \langle u_\varepsilon(x, \cdot) \rangle, \quad x \in \mathbf{Z}_\varepsilon^d.$$

Furthermore, the Fourier transform $\hat{u}_\varepsilon(\xi)$ of $u_\varepsilon(x)$ satisfies the equation

$$(4.5) \quad \hat{u}_\varepsilon(\xi) \left\{ 1 + \varepsilon^{-2} [1 - \hat{p}(\varepsilon\xi)] + 2\gamma i \sum_{j=1}^d \langle b_j(\cdot) \rangle \frac{\sin(\mathbf{e}_j \cdot \varepsilon\xi)}{\varepsilon^2} \right\} \\ + \gamma \varepsilon^{-2} \left\langle \sum_{j=1}^d b_j(\cdot) \left[e^{i\varepsilon\mathbf{e}_j \cdot \xi} \hat{\psi}_\varepsilon(\xi, \tau_{-\mathbf{e}_j \cdot \cdot}) - e^{-i\varepsilon\mathbf{e}_j \cdot \xi} \hat{\psi}_\varepsilon(\xi, \tau_{\mathbf{e}_j \cdot \cdot}) \right] \right\rangle \\ = \hat{f}_\varepsilon(\xi), \quad \xi \in \left[\frac{-\pi}{\varepsilon}, \frac{\pi}{\varepsilon} \right]^d.$$

The Fourier transform of $\psi_\varepsilon(x, \omega)$ is given in terms of $\hat{u}_\varepsilon(\xi)$ by

$$(4.6) \quad \hat{\psi}_\varepsilon(\xi, \omega) = -2i\gamma \hat{u}_\varepsilon(\xi) \sum_{k=1}^d \sin(\varepsilon\mathbf{e}_k \cdot \xi) \Psi_\varepsilon^k(\varepsilon\xi, \omega)$$

Here $\Psi_\varepsilon^k(\zeta, \omega)$, $1 \leq k \leq d$, $\zeta \in [-\pi, \pi]^d$, $\omega \in \Omega$, are functions in $L^2([-\pi, \pi]^d \times \Omega)$ which satisfy the equation

$$(4.7) \quad \Psi_\varepsilon^k(\zeta, \omega) - \sum_{y \in \mathbf{Z}^d} p(y) e^{-iy \cdot \zeta} \Psi_\varepsilon^k(\zeta, \tau_y \omega) \\ + \gamma P \sum_{j=1}^d b_j(\omega) \left[e^{i\mathbf{e}_j \cdot \zeta} \Psi_\varepsilon^k(\zeta, \tau_{-\mathbf{e}_j} \omega) - e^{-i\mathbf{e}_j \cdot \zeta} \Psi_\varepsilon^k(\zeta, \tau_{\mathbf{e}_j} \omega) \right] + \varepsilon^2 \Psi_\varepsilon^k(\zeta, \omega) \\ = b_k(\omega) - \langle b_k(\cdot) \rangle, \quad \zeta \in [-\pi, \pi]^d, \quad \omega \in \Omega, \quad 1 \leq k \leq d.$$

If we substitute for $\hat{\psi}_\varepsilon$ from (4.6) into (4.5) then we obtain the equation,

$$(4.8) \quad \hat{u}_\varepsilon(\xi) \left\{ 1 + \varepsilon^{-2} [1 - \hat{p}(\varepsilon\xi)] + 2\gamma i \sum_{j=1}^d \langle b_j(\cdot) \rangle \frac{\sin(\varepsilon\mathbf{e}_j \cdot \xi)}{\varepsilon^2} \right. \\ \left. - 2i\gamma^2 \left\langle \sum_{j,k=1}^d b_j(\cdot) \frac{\sin(\varepsilon\mathbf{e}_k \cdot \xi)}{\varepsilon^2} \left[e^{i\varepsilon\mathbf{e}_j \cdot \xi} \Psi_\varepsilon^k(\varepsilon\xi, \tau_{-\mathbf{e}_j \cdot \cdot}) \right. \right. \right. \\ \left. \left. \left. - e^{-i\varepsilon\mathbf{e}_j \cdot \xi} \Psi_\varepsilon^k(\varepsilon\xi, \tau_{\mathbf{e}_j \cdot \cdot}) \right] \right\rangle \right\} = \hat{f}_\varepsilon(\xi), \quad \xi \in \left[\frac{-\pi}{\varepsilon}, \frac{\pi}{\varepsilon} \right]^d.$$

Let us assume now that (1.9) holds. Then, following the development at the beginning of §3, we have that the matrix $q_{\gamma, \varepsilon}(\zeta) = [q_{\gamma, \varepsilon, k, k'}(\zeta)]$, $1 \leq k, k' \leq d$ of (1.11) is given by the formula

$$(4.9) \quad q_{\gamma, \varepsilon, k, k'}(\zeta) = \\ - \sum_{n=0}^{\infty} \gamma^{n+2} \sum_{m=0}^n \left\langle \left[\sum_{j=1}^d b_j(\cdot) T_{j, \varepsilon^2, 0} P \right]^m S_{k', \varepsilon^2, \zeta} P \left[\sum_{j=1}^d b_j(\cdot) T_{j, \varepsilon^2, \zeta} P \right]^{n-m} b_k(\cdot) \right\rangle \\ - \sum_{n=0}^{\infty} \gamma^{n+2} \sum_{m=0}^n \left\langle \left[\sum_{j=1}^d b_j(\cdot) T_{j, \varepsilon^2, 0} P \right]^m e^{-i\mathbf{e}_k \cdot \zeta} S_{k, \varepsilon^2, \zeta} P \left[\sum_{j=1}^d b_j(\cdot) T_{j, \varepsilon^2, \zeta} P \right]^{n-m} b_{k'}(\cdot) \right\rangle,$$

where the operator $T_{j,\eta,\zeta}$ is defined by (2.31) and $S_{k,\eta,\zeta}$ by (3.3), with p given by (1.2). One can see from the proof of Lemma 4.1 that $q_{\gamma,\varepsilon}(\zeta)$ is continuous in ζ and analytic in γ provided $|\gamma| < \varepsilon/\sqrt{2d}$.

5. Proof of Theorem 1.4

The matrix $q_{m,\varepsilon}(\zeta)$ is given by the coefficient of γ^m in the expansion (4.9). Since the $\mathbf{b}(\tau_x \cdot)$ are given by Bernoulli variables, (a) and (b) of Theorem 1.4 follow easily from (4.9). Next we obtain a formula for $q_{4,\varepsilon}(\zeta)$. Let $K_\varepsilon(x)$ be defined by

$$(5.1) \quad K_\varepsilon(x) = G_{p,\varepsilon^2}(x - \mathbf{e}_1) - G_{p,\varepsilon^2}(x + \mathbf{e}_1), \quad x \in \mathbf{Z}^d.$$

For $x \in \mathbf{Z}^d$, let x^R be the reflection of x in the plane through the origin with normal \mathbf{e}_1 . It is clear that

$$(5.2) \quad K_\varepsilon(x^R) = -K_\varepsilon(x), \quad x \in \mathbf{Z}^d.$$

For $1 \leq k \leq d$, $\zeta \in [-\pi, \pi]^d$, define $a_k(\zeta)$, $b_k(\zeta)$ by

$$(5.3) \quad \begin{aligned} a_{k,\varepsilon}(\zeta) &= \sum_{x \in \mathbf{Z}^d} [K_\varepsilon(x)]^3 \operatorname{sgn}(x \cdot \mathbf{e}_k) \sum_{y \in L_k(x)} e^{-iy \cdot \zeta}, \\ b_{k,\varepsilon}(\zeta) &= \sum_{x \in \mathbf{Z}^d} [K_\varepsilon(x)]^3 e^{ix \cdot \zeta} \operatorname{sgn}(x \cdot \mathbf{e}_k) \sum_{y \in L_k(x)} e^{-iy \cdot \zeta}. \end{aligned}$$

Note that

$$(5.4) \quad a_{k,\varepsilon}(0) = b_{k,\varepsilon}(0) = \sum_{x \in \mathbf{Z}^d} [K_\varepsilon(x)]^3 (x \cdot \mathbf{e}_k).$$

Then one can see from (4.9) that

$$(5.5) \quad q_{4,\varepsilon,1,1}(\zeta) = [1 + e^{-i\mathbf{e}_1 \cdot \zeta}] [2a_{1,\varepsilon}(\zeta) - b_{1,\varepsilon}(\zeta)].$$

One also has for $1 < k \leq d$,

$$(5.6) \quad \begin{aligned} q_{4,\varepsilon,1,k}(\zeta) &= 2a_{k,\varepsilon}(\zeta) - b_{k,\varepsilon}(\zeta), \\ q_{4,\varepsilon,1,k}(\zeta) &= e^{-i\mathbf{e}_k \cdot \zeta} [2a_{k,\varepsilon}(\zeta) - b_{k,\varepsilon}(\zeta)]. \end{aligned}$$

Since $a_{k,\varepsilon}(0) = b_{k,\varepsilon}(0)$ it follows that $q_{4,\varepsilon,1,1}(0) = 2a_{1,\varepsilon}(0)$. From (5.2) and (5.4) we see that $a_{k,\varepsilon}(0) = 0$ if $k > 1$. Hence $q_{4,\varepsilon,k,k'}(0) = 0$ for $k + k' > 2$.

Lemma 5.1. (a) *Let $d = 1$ and $\mathcal{K} \subset \mathbf{R}$ be a compact set. Then $\varepsilon^2 a_{1,\varepsilon}(\varepsilon\xi)$ converges uniformly for $\xi \in \mathcal{K}$ to a function $a_1(\xi)$, as $\varepsilon \rightarrow 0$. Similarly $\varepsilon^2 b_{1,\varepsilon}(\varepsilon\xi)$ converges to a function $b_1(\xi)$.*

(b) *If $d \geq 3$ and $1 \leq k \leq d$, then $a_{k,\varepsilon}(\zeta)$ converges uniformly for $\zeta \in [-\pi, \pi]$ to a function $a_k(\zeta)$, as $\varepsilon \rightarrow 0$. Similarly $b_{k,\varepsilon}(\zeta)$ converges to a function $b_k(\zeta)$.*

Proof. (a) It is easy to see that for $d = 1$, $\eta > 0$, one has

$$G_{p,\eta}(x) = \frac{[1 + \eta - \sqrt{2\eta + \eta^2}]^{|x|}}{\sqrt{2\eta + \eta^2}}, \quad x \in \mathbf{Z}.$$

Hence if for $\varepsilon > 0$, we define $\nu(\varepsilon)$ by

$$(5.7) \quad \nu(\varepsilon) = 1 + \varepsilon^2 - \varepsilon\sqrt{2 + \varepsilon^2},$$

then the function $K_\varepsilon(x)$ of (5.1) is given by

$$(5.8) \quad K_\varepsilon(x) = 2\nu(\varepsilon)^{|x|}, \quad x \in \mathbf{Z}, \quad x > 0.$$

Substituting from (5.8) into (5.3) we obtain the identity

$$a_{1,\varepsilon}(\zeta) = \frac{8\nu(\varepsilon)^3}{1 - \nu(\varepsilon)^3} \left[\frac{1}{1 - \nu(\varepsilon)^3 e^{-i\zeta}} + \frac{e^{i\zeta}}{1 - \nu(\varepsilon)^3 e^{i\zeta}} \right].$$

One can see that

$$\lim_{\varepsilon \rightarrow 0} \frac{\varepsilon}{1 - \nu(\varepsilon)^3 e^{-i\varepsilon\xi}} = \frac{1}{3\sqrt{2} + i\xi}.$$

We conclude that

$$\lim_{\varepsilon \rightarrow 0} \varepsilon^2 a_{1,\varepsilon}(\varepsilon\xi) = \frac{16}{18 + \xi^2}.$$

Since for $d = 1$ one has $a_{1,\varepsilon}(\zeta) = b_{1,\varepsilon}(-\zeta)$, the formula (1.13) follows now from the last equation and (5.5).

(b) This follows from the fact that for $d \geq 3$, the limit $\lim_{\varepsilon \rightarrow 0} K_\varepsilon(x) = K_0(x)$ exists for all $x \in \mathbf{Z}^d$, and the bound (4.1). \square

Lemma 5.2. (a) For $d \geq 3$, one has $a_1(0) > 0$.

(b) For $d = 2$ one has $\lim_{\varepsilon \rightarrow 0} a_{1,\varepsilon}(0) = +\infty$.

Proof. (a) In view of (5.2) and (5.4) it is sufficient to show that $K_0(x) > 0$ if $x \in \mathbf{Z}^d$ satisfies $x \cdot \mathbf{e}_1 > 0$. To see this observe from (2.9) and (5.2) that $K_\varepsilon(x)$ satisfies the equation

$$A_p K_\varepsilon(x) + \varepsilon^2 K_\varepsilon(x) = g(x), \quad x \in \mathbf{Z}^d, \quad x \cdot \mathbf{e}_1 > 0,$$

with Dirichlet boundary condition $K_\varepsilon(x) = 0$, $x \cdot \mathbf{e}_1 = 0$. Here g is the function, $g(x) = 1$, $x = \mathbf{e}_1$, $g(x) = 0$ otherwise. The positivity of $K_0(x)$ follows now from the maximum principle.

(b) We shall show that there exists $\alpha > 0$ and a positive constant c_α , depending only on α such that for $d = 2$,

$$(5.9) \quad \lim_{\varepsilon \rightarrow 0} K_\varepsilon(n\mathbf{e}_1 + m\mathbf{e}_2) \geq \frac{c_\alpha}{n}, \quad |m| < \alpha n, \quad n = 1, 2, \dots$$

To prove (5.9) we use the fact that for $d = 1$ one has

$$K_\varepsilon(n) = 2\nu(\varepsilon)^n = \frac{2}{\pi} \int_0^\pi \frac{\sin(n\zeta) \sin(\zeta)}{2 \sin^2(\zeta/2) + \varepsilon^2} d\zeta \quad n \geq 1.$$

Since one also has for $d = 2$ the representation

$$K_\varepsilon(n\mathbf{e}_1 + m\mathbf{e}_2) = \frac{2}{\pi^2} \int_0^\pi \int_0^\pi \frac{\cos(m\xi) \sin(n\zeta) \sin(\zeta)}{\sin^2(\zeta/2) + \sin^2(\xi/2) + \varepsilon^2} d\zeta d\xi,$$

we obtain for $d = 2$ the formula

$$\lim_{\varepsilon \rightarrow 0} K_\varepsilon(n\mathbf{e}_1 + m\mathbf{e}_2) = \frac{4}{\pi} \int_0^\pi \nu(\sqrt{2} \sin(\xi/2))^n \cos(m\xi) d\xi, \quad m, n \in \mathbf{Z}, \quad n \geq 1.$$

Now there exist universal positive constants c, C such that

$$\exp[-Cn\xi] \leq \nu(\sqrt{2} \sin(\xi/2))^n \leq \exp[-cn\xi], \quad n \geq 1, \quad 0 \leq \xi \leq \pi.$$

We conclude that the LHS of (5.9) is bounded below by

$$\frac{4}{\pi} \int_0^\pi \exp[-Cn\xi] d\xi = \frac{4}{C\pi n} \{1 - \exp[-C\pi n]\},$$

if $m = 0$, and by

$$\begin{aligned} \frac{2\sqrt{2}}{\pi} \int_0^{\pi/4|m|} \exp[-Cn\xi] d\xi - \frac{4}{\pi} \int_{\pi/4|m|}^\infty \exp[-c\xi] d\xi \\ = \frac{2\sqrt{2}}{C\pi n} \left\{ 1 - \exp\left[-\frac{C\pi n}{4|m|}\right] \right\} - \frac{4}{c\pi n} \exp\left[-\frac{c\pi n}{4|m|}\right], \end{aligned}$$

if $m \neq 0$. The inequality (5.9) clearly follows from the last two identities. The result follows now from (5.4) and (5.9) since the argument of Part (a) implies $K_\varepsilon(n\mathbf{e}_1 + m\mathbf{e}_2) > 0$, $m, n \in \mathbf{Z}$, $n > 0$. \square

We are left now to prove (c) and (d) of Theorem 1.4 for $m > 4$. In order to do this we need to obtain a suitable representation of $q_{m,\varepsilon}(\zeta)$ for general m . We first consider the situation $\zeta = 0$. Recall that we are assuming the $b_j(\cdot) \equiv 0$, $j > 1$, and $b_1(\tau_x \cdot) = Y_x$, $x \in \mathbf{Z}^d$, where the variables Y_x , $x \in \mathbf{Z}^d$, are assumed to be i.i.d. Bernoulli, $Y_x = \pm 1$ with equal probability.

Lemma 5.3. *Suppose $m \geq 4$ is even. Then $q_{m,\varepsilon,k,k'}(0) = 0$ if $k + k' > 2$ and $q_{m,\varepsilon,1,1}(0)$ has the representation*

$$(5.10) \quad q_{m,\varepsilon,1,1}(0) = -2 \sum_{x_1, \dots, x_{m-1} \in \mathbf{Z}^d} \left(\sum_{\alpha=1}^{m-1} x_\alpha \cdot \mathbf{e}_1 \right) K_\varepsilon(x_1) \cdots K_\varepsilon(x_{m-1}) \\ \langle Y_0 Y_{x_1} Y_{x_1+x_2} \cdots Y_{x_1+\dots+x_{m-1}} \rangle.$$

Proof. First note that by the argument of Lemma 2.8 the representation (4.9) for $q_{\gamma,\varepsilon}(0)$ continues to hold when we delete the projection operator P from the RHS of the equation. It is easy to see from this that (5.10) holds. If $k > 1$ then one has a similar representation to (5.10) for $q_{m,\varepsilon,1,k}(0)$ and $q_{m,\varepsilon,k,1}(0)$ but with \mathbf{e}_1 in (5.10) replaced by \mathbf{e}_k . In that case (5.2) implies $q_{m,\varepsilon,k,1}(0) = q_{m,\varepsilon,1,k}(0) = 0$. \square

In order to prove (d) of Theorem 1.4 for general m we use (5.10) to obtain a representation of $q_{m,\varepsilon,1,1}(0)$ as a sum indexed by certain types of graph. We shall use the terminology of [1]. For $q = 2, 3, \dots$ let \mathcal{F}_q be the set of unlabeled, connected, directed multigraphs on q vertices with the properties:

- (A) The graph has no loops.
- (B) Each vertex has equal indegree and outdegree.
- (C) The degree of each vertex is a multiple of 4.
- (D) The number of edges in the graph is $4q$.

For a graph $G \in \mathcal{F}_q$ let $V[G]$ denote the vertex set of G and $E[G]$ be the set of directed edges of G . Each directed edge e has two vertices e_+ and e_- , with the direction of the edge being from e_- to e_+ . We associate with G and a directed edge $e \in E[G]$, a number $K_\varepsilon(G, e)$ defined by

$$(5.11) \quad K_\varepsilon(G, e) = \sum_{\{y_v \in \mathbf{Z}^d : v \in V[G]\}} (y_{e_-} \cdot \mathbf{e}_1) \delta(y_{e_+}) \prod_{e' \in E[G], e' \neq e} K_\varepsilon(y_{e'_+} - y_{e'_-}).$$

In (5.11) $\delta(x)$, $x \in \mathbf{Z}^d$ is the Kronecker delta, $\delta(0) = 1$, $\delta(x) = 0$, $x \neq 0$.

Lemma 5.4. *Let $m \geq 4$ be an even integer and $q = m/2$. Then*

$$(5.12) \quad q_{m,\varepsilon,1,1}(0) = \sum_{G \in \mathcal{F}_q, e \in E[G]} c(G, e) K_\varepsilon(G, e),$$

for some integers $c(G, e)$ depending only on the graph G and the directed edge $e \in E[G]$.

Proof. Observe that

$$K_\varepsilon(x_1) \dots K_\varepsilon(x_{m-1}) \langle Y_0 Y_{x_1} Y_{x_1+x_2} \dots Y_{x_1+\dots+x_{m-1}} \rangle \neq 0,$$

only if the sequence $0, x_1, x_1 + x_2, \dots, x_1 + \dots + x_{m-1}$ of points in \mathbf{Z}^d has the property that it consists of q pairs of points with the property that no two adjacent points of the sequence are the same. The q pairs of points do not have to be distinct. Hence we may write

$$(5.13) \quad q_{m,\varepsilon,1,1}(0) = -2 \sum_{i=1}^q a_i,$$

where a_i corresponds to the term on the RHS of (5.10) where i of the q pairs are distinct. For a graph $G \in \mathcal{F}_q$ let $K'_\varepsilon(G, e)$ be defined in the same way as $K_\varepsilon(G, e)$ except that the summation on the RHS of (5.11) is over all $y_v \in \mathbf{Z}^d : v \in V[G]$ such that the q points $y_v \in \mathbf{Z}^d$ are distinct. Define \mathcal{F}'_q as the subset of \mathcal{F}_q consisting of regular graphs of degree 4 i.e., the degree of each vertex is exactly 4. Now by (B) of the definition of \mathcal{F}_q and Euler's theorem [1] a graph $G \in \mathcal{F}_q$ has an Eulerian path. Let $c(G)$ be the number of Eulerian paths. For a graph $G \in \mathcal{F}_q$, one can group directed edges into equivalence classes where two directed edges e, e' are equivalent if there is a graph isomorphism of G taking e to e' . Let $E'[G]$ be this set of equivalence classes. Then we may define $K'_\varepsilon(G, \hat{e})$ for $\hat{e} \in E'[G]$ by $K'_\varepsilon(G, \hat{e}) = K'_\varepsilon(G, e)$ for any $e \in \hat{e}$. Then one can see that a_q of (5.13) is given by

$$a_q = \sum_{G \in \mathcal{F}'_q, \hat{e} \in E'[G]} c(G) K'_\varepsilon(G, \hat{e}).$$

Since we can obtain a similar representation for the a_i , $1 \leq i < q$, we have the formula (5.12) up to replacing $K'_\varepsilon(G, e)$ by $K_\varepsilon(G, e)$. The result therefore holds if we can show that for any graph $G \in \mathcal{F}_q$ and $e \in E[G]$ there exist integers $c_{G,e}(G', e')$, $G' \in \mathcal{F}_q$, $e' \in E[G']$ such that

$$(5.14) \quad K'_\varepsilon(G, e) = \sum_{G' \in \mathcal{F}_q, e' \in E[G']} c_{G,e}(G', e') K_\varepsilon(G', e').$$

The previous identity follows from Mayer's trick to obtain an expansion for an non-ideal gas [7]. That is for a function $\varphi : \mathbf{Z}^d \rightarrow \mathbf{R}$, one writes

$$(5.15) \quad \exp \left[- \sum_{1 \leq i < j \leq N} \varphi(y_i - y_j) \right] = \prod_{1 \leq i < j \leq N} [1 + f_{i,j}],$$

where $f_{i,j} = \exp[-\varphi(y_i - y_j)] - 1$, and expands the RHS of (5.15) out. Now (5.14) follows from (5.15) on taking φ to be the function, $\varphi(y) = 0$, $y \neq 0$, $\varphi(0) = +\infty$. \square

Lemma 5.5. *Suppose $d = 1$, $q \geq 2$ and $G \in \mathcal{F}_q$, $e \in G$. If $|G|$ is the number of vertices of G , then the limit*

$$(5.16) \quad \lim_{\varepsilon \rightarrow 0} \varepsilon^{|G|} K_\varepsilon(G, e) = K(G, e)$$

exists.

Proof. Evidently it is sufficient to show that (5.16) holds with $K_\varepsilon(G, e)$ replaced by $K'_\varepsilon(G, e)$. Suppose $|G| = k \leq q$ and label the vertices of G as $1, 2, \dots, k$ with $e_+ = 1$, $e_- = k$. Then

$$K'_\varepsilon(G, e) = \sum_{\pi \in \mathcal{S}_k} K_{\varepsilon, \pi}(G, e),$$

where \mathcal{S}_k is the group of permutations on $1, 2, \dots, k$ and

$$K_{\varepsilon, \pi}(G, e) = \sum_{y_{\pi_1} < y_{\pi_2} < \dots < y_{\pi_k}} \delta(y_1) y_k \prod_{e' \in E[G], e' \neq e} K_\varepsilon(y_{e'_+} - y_{e'_-}).$$

In view of (5.8) there are positive integers N_j , $1 \leq j \leq k-1$ such that

$$K_{\varepsilon, \pi}(G, e) = \pm \sum_{y_{\pi_1} < y_{\pi_2} < \dots < y_{\pi_k}} \delta(y_1) y_k \prod_{j=1}^{k-1} 2^{N_j} \nu(\varepsilon)^{N_j [y_{\pi(j+1)} - y_{\pi(j)}]}.$$

Suppose now that $\pi k_1 = 1$, $\pi k_2 = k$ and $k_1 < k_2$. Then

$$(5.17) \quad K_{\varepsilon, \pi}(G, e) = \sum_{i=k_1}^{k_2-1} K_{\varepsilon, \pi, i}(G, e),$$

where

$$K_{\varepsilon, \pi, i}(G, e) = \pm \sum_{y_{\pi_1} < y_{\pi_2} < \dots < y_{\pi_k}} \delta(y_1) [y_{\pi(i+1)} - y_{\pi(i)}] \prod_{j=1}^{k-1} 2^{N_j} \nu(\varepsilon)^{N_j [y_{\pi(j+1)} - y_{\pi(j)}]}.$$

It is easy to see now just as in Lemma 5.1 that the limit

$$\lim_{\varepsilon \rightarrow 0} \varepsilon^k K_{\varepsilon, \pi, i}(G, e)$$

exists. Hence from (5.17) the limit,

$$(5.18) \quad \lim_{\varepsilon \rightarrow 0} \varepsilon^k K_{\varepsilon, \pi}(G, e)$$

exists if $k_1 < k_2$. A similar argument shows that the limit (5.18) also exists if $k_1 > k_2$. It follows then from the previous identities that the limit (5.16) exists. \square

Remark 1. Lemma 5.5 proves that (d) of Theorem 1.4 holds with $\xi = 0$. Observe that the number of graphs in \mathcal{F}_q is of order $q!$. One can see this from an asymptotic formula for the number A_q of labeled 4-regular simple graphs on q vertices [18],

$$A_q \sim [(4q)! e^{-15/4}] / [(2q)! 96^q].$$

Next we wish to obtain the analogue of Lemma 5.4 for $q_{m, \varepsilon, k, k'}(\zeta)$ with $\zeta \in [-\pi, \pi]^d$ and at least one of k, k' being 1. To do this we define for any $G \in \mathcal{F}_q$, $e \in E[G]$, a function $K_\varepsilon(G, e, \zeta)$ by

$$(5.19) \quad K_\varepsilon(G, e, \zeta) = \sum_{\{y_v \in \mathbf{Z}^d : v \in V[G]\}} \exp[-iy_{e_-} \cdot \zeta] \delta(y_{e_+}) \prod_{e' \in E[G], e' \neq e} K_\varepsilon(y_{e'_+} - y_{e'_-}).$$

Note that $K_\varepsilon(G, e, \zeta)$ is defined similarly to $K_\varepsilon(G, e)$ of (5.11) and has the property $K_\varepsilon(G, e, 0) = 0$. For $1 \leq k \leq d$ we define $K_{\varepsilon,k}(G, e, \zeta)$ by

$$(5.20) \quad K_{\varepsilon,k}(G, e, \zeta) = \sum_{\{y_v \in \mathbf{Z}^d : v \in V[G]\}} \delta(y_{e_+}) \left[\prod_{e' \in E[G], e' \neq e} K_\varepsilon(y_{e'_+} - y_{e'_-}) \right] \\ \left[\sum_{e' \in E[G], e' \neq e} \exp[i(y_{e'_+} - y_{e'_-}) \cdot \zeta] \operatorname{sgn}[(y_{e'_+} - y_{e'_-}) \cdot \mathbf{e}_k] \sum_{y \in L_k(y_{e'_+} - y_{e'_-})} e^{-iy \cdot \zeta} \right].$$

Note that $K_\varepsilon(G, e)$ of (5.11) is given by $K_\varepsilon(G, e) = K_{\varepsilon,1}(G, e, 0)$. In the following lemma we need to define \mathcal{F}_q for $q = 0, 1$. The set \mathcal{F}_1 is the single point vertex graph G with $K_\varepsilon(G, e, \zeta) = 0$. The set \mathcal{F}_0 is also the single point vertex graph G with $K_\varepsilon(G, e, \zeta) = 1$.

Lemma 5.6. *Let $m \geq 4$ be an even integer and $q = m/2$. Define $A_{m,\varepsilon,k}(\zeta)$ by*

$$(5.21) \quad A_{m,\varepsilon,k}(\zeta) = \sum_{q'=2}^q \sum_{G \in \mathcal{F}_{q'}, G' \in \mathcal{F}_{q-q'}} \sum_{e \in E[G], e' \in E[G']} c(G, e, G', e') K_{\varepsilon,k}(G, e, \zeta) K_\varepsilon(G', e', \zeta),$$

for suitable integers $c(G, e, G', e')$ depending only on the graphs G, G' and the directed edges $e \in E[G], e' \in E[G']$. Then

$$\begin{aligned} q_{m,\varepsilon,1,1}(\zeta) &= [1 + e^{-i\mathbf{e}_1 \cdot \zeta}] A_{m,\varepsilon,1}(\zeta), \\ q_{m,\varepsilon,1,k}(\zeta) &= A_{m,\varepsilon,k}(\zeta), \quad 1 < k \leq d, \\ q_{m,\varepsilon,k,1}(\zeta) &= e^{-i\mathbf{e}_k \cdot \zeta} A_{m,\varepsilon,k}(\zeta), \quad 1 < k \leq d. \end{aligned}$$

Proof. We define $A_{m,\varepsilon,k'}(\zeta)$ as the coefficient of γ^m in the first term on the RHS of (4.9). Evidently the formulas for $q_{m,\varepsilon,k,k'}(\zeta)$ in the statement of the lemma follow from this. We need then to establish (5.21). This follows by the same argument as in Lemma 5.4. Note now that, unlike in Lemmas 5.3 and 5.4, the projection operators P in (4.9) make a contribution. Thus we obtain the factorisations $K_{\varepsilon,k}(G, e, \zeta) K_\varepsilon(G', e', \zeta)$ in (5.21) with $G \in \mathcal{F}_{q'}, G' \in \mathcal{F}_{q''}$, where $q' + q'' = q$. \square

Lemma 5.7. *Suppose $d = 1$ and $\mathcal{K} \subset \mathbf{R}$ is a compact set. Let $q \geq 2$ and $G \in \mathcal{F}_q$, $e \in G$. If $|G|$ is the number of vertices of G , then the limits*

$$(5.22) \quad \lim_{\varepsilon \rightarrow 0} \varepsilon^{|G|-1} K_\varepsilon(G, e, \varepsilon \xi) = K(G, e, \xi)$$

$$(5.23) \quad \lim_{\varepsilon \rightarrow 0} \varepsilon^{|G|} K_{\varepsilon,1}(G, e, \varepsilon \xi) = K_1(G, e, \xi)$$

exist, uniformly for $\xi \in \mathcal{K}$.

Proof. The proof of (5.22) follows exactly the proof of Lemma 5.5. To prove (5.23) we proceed similarly. Thus we focus on one term in the summation over $e' \in E[G], e' \neq e$ in (5.20). As in Lemma 5.5 we fix a permutation π and consider the sum over $y_{\pi 1} < y_{\pi 2} < \dots < y_{\pi k}$. Suppose that $\pi k_1 = e'_-$, $\pi k_2 = e'_+$ and

$k_1 < k_2$. Then

$$\operatorname{sgn}(y_{e'_+} - y_{e'_-}) \sum_{y \in L_1(y_{e'_+} - y_{e'_-})} e^{-iy\zeta} = \sum_{i=k_1}^{k_2-1} \left(\exp[-i(y_{\pi(i)} - y_{\pi(k_1)})\zeta] \sum_{z=0}^{y_{\pi(i+1)} - y_{\pi(i)} - 1} e^{-iz\zeta} \right).$$

If we consider a fixed i on the RHS of the previous equation and sum over $y_{\pi_1} < y_{\pi_2} < \dots < y_{\pi k}$, then we see as in Lemma 5.5 that the contribution of this summation to (5.23) converges uniformly for $\xi \in \mathcal{K}$ as $\varepsilon \rightarrow 0$. Since we can similarly argue for all the other contributions we conclude that the uniform limit (5.23) exists. \square

It is clear that (d) of Theorem 1.4 follows from Lemmas 5.6 and 5.7.

References

- [1] B. Bollobas, *Modern Graph Theory*, Graduate Texts in Mathematics, 184, Springer, New York, 1998, MR 99h:05001, Zbl 0902.05016.
- [2] J. Conlon and B. von Dohlen, *Numerical simulations of random walk in random environment*, J. Statist. Phys. **92** (1998), 571–586, MR 99j:82039, Zbl 0936.60089.
- [3] J. Conlon and A. Naddaf, *On homogenisation of elliptic equations with random coefficients*, Electronic Journal of Probability **5** (2000), paper 9, 1–58, CMP 1 768 843, Zbl 0956.35013.
- [4] B. Derrida and J. Luck, *Diffusion on a random lattice: Weak-disorder expansion in arbitrary dimension*, Phys. Rev. B **28** (1983), 7183–7190.
- [5] A. Fannjiang and G. Papanicolaou, *Diffusion in turbulence*, Prob. Theory Relat. Fields **105** (1996), 279–334, MR 98d:60156, Zbl 0847.60062.
- [6] D. Fisher, *Random walks in random environments*, Phys. Rev. A **30** (1984), 960–964, MR 85h:82028.
- [7] G. Ford and G. Uhlenbeck, *Lectures in Statistical Mechanics*, Lectures in Applied Mathematics, 1, American Math Society, Providence 1963, MR 27 #1241, Zbl 0111.43802.
- [8] D. Gilbarg and A. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Grundlehren der Mathematischen Wissenschaften, 224, 2nd edition, Springer, New York 1983, MR 86c:35035, Zbl 0562.35001.
- [9] H. Kesten, *The limit distribution of Sinai’s random walk in random environment*, Phys. A. **138** (1986), 299–309, MR 88b:60165, Zbl 0666.60065.
- [10] C. Kipnis and S.R.S. Varadhan, *Central limit theorem for additive functionals of reversible Markov processes and applications to simple exclusions*, Commun. Math. Phys. **104** (1986), 1–19, MR 87i:60038, Zbl 0588.60058.
- [11] S. Kozlov, *Averaging of random structures*, Dokl. Akad. Nauk. SSSR **241** (1978), 1016–1019, MR 80e:60078.
- [12] R. Künnemann, *The diffusion limit for reversible jump processes on \mathbf{Z} with ergodic random bond conductivities*, Commun. Math. Phys. **90** (1983), 27–68, MR 84m:60102, Zbl 0523.60097.
- [13] C. Landim, S. Olla, and H.-T. Yau, *Convection-diffusion equation with space-time ergodic random flow*, Probab. Theory Relat. Fields **112** (1998), 203–220, MR 99j:35084, Zbl 0914.60070.
- [14] G. Lawler, *Weak convergence of a random walk in a random environment*, Commun. Math. Phys. **87** (1982), 81–87, MR 84b:60093, Zbl 0502.60056.
- [15] A. Lejay, *Homogenization of divergence-form operators with lower-order terms in random media*, Probab. Theory Relat. Fields **120** (2001), 255–276, CMP 1 869 029.
- [16] G. Papanicolaou and S.R.S. Varadhan, *Boundary value problems with rapidly oscillating random coefficients*, Volume 2 of Coll. Math. Soc. Janos Bolya **27**, Random Fields, Amsterdam, North Holland Publ. Co. 1981, pp. 835–873, MR 84k:58233, Zbl 0499.60059.
- [17] A. Pisztora, T. Povel and O. Zeitouni, *Precise large deviation estimates for a one-dimensional walk in a random environment*, Probab. Theory Related Fields **113** (1999), 191–219, MR 99m:60048, Zbl 0922.60059.

- [18] R. Read and N. Wormald, *Number of labelled 4-regular graphs*, J. Graph Theory **4** (1980), 203–212, MR 81e:05085, Zbl 0427.05040.
- [19] M. Reed and B. Simon, *Methods of Mathematical Physics I: Functional Analysis*, Academic Press, 1972, MR 85e:46002, Zbl 0242.46001.
- [20] K. Sato, *Lévy Processes and Infinitely Divisible Distributions*, Cambridge Studies in Advanced Mathematics, 68, Cambridge University Press, 1999, CMP 1 739 520, Zbl 0973.60001.
- [21] Y. Sinai, *Limiting behavior of a one-dimensional random walk in a random medium*, Theory Prob. Appl. **27** (1982), 256–268, MR 83k:60078.
- [22] E. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton University Press, 1970, MR 58 #2467, Zbl 0207.13501.
- [23] A. Sznitman, *Slowdown and neutral pockets for a random walk in random environment*, Probab. Theory Related Fields **115** (1999), 287–323, MR 2001a:60035, Zbl 0947.60095.
- [24] A. Sznitman, *Slowdown estimates and central limit theorem for random walks in random environment*, J. Eur. Math. Soc. **2** (2000), 93–143, CMP 1 763 302.
- [25] V. Zhikov, S. Kozlov, and O. Oleinik, *Homogenization of Differential Operators and Integral Functionals*, Springer Verlag, Berlin, 1994, MR 96h:35003b.

UNIVERSITY OF MICHIGAN, DEPARTMENT OF MATHEMATICS, ANN ARBOR, MI 48109-1109
conlon@math.lsa.umich.edu <http://www.math.lsa.umich.edu/~conlon/>

This paper is available via <http://nyjm.albany.edu:8000/j/2002/8-3.html>.