

C^* -algebras associated with textile dynamical systems

Kengo Matsumoto

ABSTRACT. A C^* -symbolic dynamical system $(\mathcal{A}, \rho, \Sigma)$ is a finite family $\{\rho_\alpha\}_{\alpha \in \Sigma}$ of endomorphisms of a C^* -algebra \mathcal{A} with some conditions. It yields a C^* -algebra \mathcal{O}_ρ from an associated Hilbert C^* -bimodule. In this paper, we will extend the notion of C^* -symbolic dynamical system to C^* -textile dynamical system $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ which consists of two C^* -symbolic dynamical systems $(\mathcal{A}, \rho, \Sigma^\rho)$ and $(\mathcal{A}, \eta, \Sigma^\eta)$ with certain commutation relations κ between their endomorphisms $\{\rho_\alpha\}_{\alpha \in \Sigma^\rho}$ and $\{\eta_a\}_{a \in \Sigma^\eta}$. C^* -textile dynamical systems yield two-dimensional subshifts and C^* -algebras $\mathcal{O}_{\rho, \eta}^\kappa$. We will study their structure of the algebras $\mathcal{O}_{\rho, \eta}^\kappa$ and present its K-theory formulae.

CONTENTS

1. Introduction	1180
2. λ -graph systems, C^* -symbolic dynamical systems and their C^* -algebras	1183
3. C^* -textile dynamical systems and two-dimensional subshifts	1186
4. C^* -textile dynamical systems and their C^* -algebras	1190
5. Condition (I) for C^* -textile dynamical systems	1195
6. Concrete realization	1204
7. K-Theory machinery	1208
8. K-Theory formulae	1217
9. Examples	1234
9.1. LR-textile λ -graph systems	1234
9.2. Tensor products	1240
10. Concluding remark	1242
References	1242

Received April, 4, 2012.

2010 *Mathematics Subject Classification*. Primary 46L35, 37B10; Secondary 46L55, 37B10.

Key words and phrases. C^* -algebra, symbolic dynamics, subshift, K-group, tiling.

This work was supported by JSPS KAKENHI Grant Numbers 20540215, 23540237 and 15K04896.

1. Introduction

In [24], the author has introduced a notion of λ -graph system as presentations of subshifts. The λ -graph systems are labeled Bratteli diagram with shift transformation. They yield C^* -algebras so that its K-theory groups are related to topological conjugacy invariants of the underlying symbolic dynamical systems. The class of these C^* -algebras include the Cuntz–Krieger algebras. He has extended the notion of λ -graph system to C^* -symbolic dynamical system, which is a generalization of both a λ -graph system and an automorphism of a unital C^* -algebra. It is a finite family $\{\rho_\alpha\}_{\alpha \in \Sigma}$ of endomorphisms of a unital C^* -algebra \mathcal{A} such that $\rho_\alpha(Z_{\mathcal{A}}) \subset Z_{\mathcal{A}}$, $\alpha \in \Sigma$ and $\sum_{\alpha \in \Sigma} \rho_\alpha(1) \geq 1$ where $Z_{\mathcal{A}}$ denotes the center of \mathcal{A} . A finite labeled graph \mathcal{G} gives rise to a C^* -symbolic dynamical system $(\mathcal{A}_{\mathcal{G}}, \rho^{\mathcal{G}}, \Sigma)$ such that $\mathcal{A} = \mathbb{C}^N$ for some $N \in \mathbb{N}$. A λ -graph system \mathfrak{L} is a generalization of a finite labeled graph and yields a C^* -symbolic dynamical system $(\mathcal{A}_{\mathfrak{L}}, \rho^{\mathfrak{L}}, \Sigma)$ such that $\mathcal{A}_{\mathfrak{L}}$ is $C(\Omega_{\mathfrak{L}})$ for some compact Hausdorff space $\Omega_{\mathfrak{L}}$ with $\dim \Omega_{\mathfrak{L}} = 0$. It also yields a C^* -algebra $\mathcal{O}_{\mathfrak{L}}$. A C^* -symbolic dynamical system $(\mathcal{A}, \rho, \Sigma)$ provides a subshift Λ_ρ over Σ and a Hilbert C^* -bimodule $\mathcal{H}_{\mathcal{A}}^\rho$ over \mathcal{A} . The C^* -algebra \mathcal{O}_ρ for $(\mathcal{A}, \rho, \Sigma)$ may be realized as a Cuntz–Pimsner algebra from the Hilbert C^* -bimodule $\mathcal{H}_{\mathcal{A}}^\rho$ ([27], cf. [15], [39]). We call the algebra \mathcal{O}_ρ the C^* -symbolic crossed product of \mathcal{A} by the subshift Λ_ρ . If $\mathcal{A} = C(X)$ with $\dim X = 0$, there exists a λ -graph system \mathfrak{L} such that the subshift Λ_ρ is the subshift $\Lambda_{\mathfrak{L}}$ presented by \mathfrak{L} and the C^* -algebra \mathcal{O}_ρ is the C^* -algebra $\mathcal{O}_{\mathfrak{L}}$ associated with \mathfrak{L} . In particular, $\mathcal{A} = \mathbb{C}^N$, the subshift Λ_ρ is a sofic shift and \mathcal{O}_ρ is a Cuntz–Krieger algebra. If $\Sigma = \{\alpha\}$ an automorphism α of a unital C^* -algebra \mathcal{A} , the C^* -algebra \mathcal{O}_ρ is the ordinary crossed product $\mathcal{A} \rtimes_{\alpha} \mathbb{Z}$.

G. Robertson–T. Steger [43] have initiated a certain study of higher dimensional analogue of Cuntz–Krieger algebras from the view point of tiling systems of 2-dimensional plane. After their work, A. Kumjian–D. Pask [19] have generalized their construction to introduce the notion of higher rank graphs and its C^* -algebras. The C^* -algebras constructed from higher rank graphs are called the higher rank graph C^* -algebras. Since then, there have been many studies on these C^* -algebras by many authors (cf. [1], [9], [10], [11], [13], [16], [19], [36], [42], [43], etc.).

M. Nasu in [34] has introduced the notion of textile system which is useful in analyzing automorphisms and endomorphisms of topological Markov shifts. A textile system also gives rise to a two-dimensional tiling called Wang tiling. Among textile systems, LR textile systems have specific properties that consist of two commuting symbolic matrices. In [28], the author has extended the notion of textile systems to λ -graph systems and has defined a notion of textile systems on λ -graph systems, which are called textile λ -graph systems for short. C^* -algebras associated to textile systems have been initiated by V. Deaconu ([9]).

In this paper, we will extend the notion of C^* -symbolic dynamical system to C^* -textile dynamical system which is a higher dimensional analogue of C^* -symbolic dynamical system. The C^* -textile dynamical system $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ consists of two C^* -symbolic dynamical systems $(\mathcal{A}, \rho, \Sigma^\rho)$ and $(\mathcal{A}, \eta, \Sigma^\eta)$ with the following commutation relations between ρ and η through κ . Set

$$\begin{aligned} \Sigma^{\rho\eta} &= \{(\alpha, b) \in \Sigma^\rho \times \Sigma^\eta \mid \eta_b \circ \rho_\alpha \neq 0\}, \\ \Sigma^{\eta\rho} &= \{(a, \beta) \in \Sigma^\eta \times \Sigma^\rho \mid \rho_\beta \circ \eta_a \neq 0\}. \end{aligned}$$

We require that there exists a bijection $\kappa : \Sigma^{\rho\eta} \longrightarrow \Sigma^{\eta\rho}$, which we fix and call a specification. Then the required commutation relations are

$$(1.1) \quad \eta_b \circ \rho_\alpha = \rho_\beta \circ \eta_a \quad \text{if } \kappa(\alpha, b) = (a, \beta).$$

A C^* -textile dynamical system provides a two-dimensional subshifts and a C^* -algebra $\mathcal{O}_{\rho, \eta}^\kappa$. The C^* -algebra $\mathcal{O}_{\rho, \eta}^\kappa$ is defined to be the universal C^* -algebra $C^*(x, S_\alpha, T_a; x \in \mathcal{A}, \alpha \in \Sigma^\rho, a \in \Sigma^\eta)$ generated by $x \in \mathcal{A}$ and two families of partial isometries $S_\alpha, \alpha \in \Sigma^\rho, T_a, a \in \Sigma^\eta$ subject to the following relations called $(\rho, \eta; \kappa)$:

$$(1.2) \quad \sum_{\beta \in \Sigma^\rho} S_\beta S_\beta^* = 1, \quad x S_\alpha S_\alpha^* = S_\alpha S_\alpha^* x, \quad S_\alpha^* x S_\alpha = \rho_\alpha(x),$$

$$(1.3) \quad \sum_{b \in \Sigma^\eta} T_b T_b^* = 1, \quad x T_a T_a^* = T_a T_a^* x, \quad T_a^* x T_a = \eta_a(x),$$

$$(1.4) \quad S_\alpha T_b = T_a S_\beta \quad \text{if } \kappa(\alpha, b) = (a, \beta)$$

for all $x \in \mathcal{A}$ and $\alpha \in \Sigma^\rho, a \in \Sigma^\eta$.

In Section 3, we will construct a tiling system in the plane from a C^* -textile dynamical system. The resulting tiling system is a two-dimensional subshift. In Section 4, we will study some basic properties of the C^* -algebra $\mathcal{O}_{\rho, \eta}^\kappa$. In Section 5, we will introduce a condition called (I) on $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ which will be studied as a generalization of the condition (I) on C^* -symbolic dynamical system [26] (cf. [8], [25]). In Section 6, we will realize the C^* -algebra $\mathcal{O}_{\rho, \eta}^\kappa$ as a Cuntz–Pimsner algebra associated with a certain Hilbert C^* -bimodule in a concrete way. We will have the following theorem.

Theorem 1.1. *Let $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ be a C^* -textile dynamical system satisfying condition (I). Then the C^* -algebra $\mathcal{O}_{\rho, \eta}^\kappa$ is a unique concrete C^* -algebra subject to the relations $(\rho, \eta; \kappa)$. If $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ is irreducible, $\mathcal{O}_{\rho, \eta}^\kappa$ is simple.*

A C^* -textile dynamical system $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ is said to form square if the C^* -subalgebra of \mathcal{A} generated by the projections $\rho_\alpha(1), \alpha \in \Sigma^\rho$ and the C^* -subalgebra of \mathcal{A} generated by the projections $\eta_a(1), a \in \Sigma^\eta$ coincide. It is said to have trivial K_1 if $K_1(\mathcal{A}) = \{0\}$. In Section 7 and Section 8, we

will restrict our interest to the C^* -textile dynamical systems forming square to prove the following K-theory formulae:

Theorem 1.2. *Suppose that $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ forms square and has trivial K_1 . Then there exist short exact sequences for $K_0(\mathcal{O}_{\rho, \eta}^\kappa)$ and $K_1(\mathcal{O}_{\rho, \eta}^\kappa)$ such that*

$$\begin{aligned} 0 &\longrightarrow K_0(\mathcal{A})/((\text{id} - \lambda_\eta)K_0(\mathcal{A}) + (\text{id} - \lambda_\rho)K_0(\mathcal{A})) \\ &\longrightarrow K_0(\mathcal{O}_{\rho, \eta}^\kappa) \\ &\longrightarrow \text{Ker}(\text{id} - \lambda_\eta) \cap \text{Ker}(\text{id} - \lambda_\rho) \text{ in } K_0(\mathcal{A}) \longrightarrow 0 \end{aligned}$$

and

$$\begin{aligned} 0 &\longrightarrow (\text{Ker}(\text{id} - \lambda_\eta) \text{ in } K_0(\mathcal{A}))/(\text{id} - \lambda_\rho)(\text{Ker}(\text{id} - \lambda_\eta) \text{ in } K_0(\mathcal{A})) \\ &\longrightarrow K_1(\mathcal{O}_{\rho, \eta}^\kappa) \\ &\longrightarrow \text{Ker}(\text{id} - \bar{\lambda}_\rho) \text{ in } (K_0(\mathcal{A})/(\text{id} - \lambda_\eta)K_0(\mathcal{A})) \longrightarrow 0 \end{aligned}$$

where the endomorphisms $\lambda_\rho, \lambda_\eta : K_0(\mathcal{A}) \longrightarrow K_0(\mathcal{A})$ are defined by

$$\begin{aligned} \lambda_\rho([p]) &= \sum_{\alpha \in \Sigma^\rho} [\rho_\alpha(p)] \in K_0(\mathcal{A}) \text{ for } [p] \in K_0(\mathcal{A}), \\ \lambda_\eta([p]) &= \sum_{a \in \Sigma^\eta} [\eta_a(p)] \in K_0(\mathcal{A}) \text{ for } [p] \in K_0(\mathcal{A}) \end{aligned}$$

and $\bar{\lambda}_\rho$ denotes an endomorphism on $K_0(\mathcal{A})/(\text{id} - \lambda_\eta)K_0(\mathcal{A})$ induced by λ_ρ .

Let A, B be mutually commuting $N \times N$ matrices with entries in non-negative integers. Let $G_A = (V_A, E_A), G_B = (V_B, E_B)$ be directed graphs with common vertex set $V_A = V_B$, whose transition matrices are A, B respectively. Let $\mathcal{M}_A, \mathcal{M}_B$ denote symbolic matrices for G_A, G_B whose components consist of formal sums of the directed edges of G_A, G_B respectively. Let Σ^{AB}, Σ^{BA} be the sets of the pairs of the concatenated directed edges in $E_A \times E_B, E_B \times E_A$ respectively. By the condition $AB = BA$, one may take a bijection $\kappa : \Sigma^{AB} \longrightarrow \Sigma^{BA}$ which gives rise to a specified equivalence $\mathcal{M}_A \mathcal{M}_B \stackrel{\kappa}{\cong} \mathcal{M}_B \mathcal{M}_A$. We then have a C^* -textile dynamical system written as $(\mathcal{A}, \rho^A, \rho^B, \Sigma^A, \Sigma^B, \kappa)$. The associated C^* -algebra is denoted by $\mathcal{O}_{A, B}^\kappa$. The C^* -algebra $\mathcal{O}_{A, B}^\kappa$ is realized as a 2-graph C^* -algebra constructed by Kumjian–Pask ([19]). It is also seen in Deaconu’s paper [9]. We will see the following proposition in Section 9.

Proposition 1.3. *Keep the above situations. There exist short exact sequences for $K_0(\mathcal{O}_{A, B}^\kappa)$ and $K_1(\mathcal{O}_{A, B}^\kappa)$ such that*

$$\begin{aligned} 0 &\longrightarrow \mathbb{Z}^N/((1 - A)\mathbb{Z}^N + (1 - B)\mathbb{Z}^N) \\ &\longrightarrow K_0(\mathcal{O}_{A, B}^\kappa) \\ &\longrightarrow \text{Ker}(1 - A) \cap \text{Ker}(1 - B) \text{ in } \mathbb{Z}^N \longrightarrow 0 \end{aligned}$$

and

$$\begin{aligned} 0 &\longrightarrow (\text{Ker}(1 - B) \text{ in } \mathbb{Z}^N)/(1 - A)(\text{Ker}(1 - B) \text{ in } \mathbb{Z}^N) \\ &\longrightarrow K_1(\mathcal{O}_{A,B}^k) \\ &\longrightarrow \text{Ker}(1 - \bar{A}) \text{ in } (\mathbb{Z}^N/(1 - B)\mathbb{Z}^N) \longrightarrow 0, \end{aligned}$$

where \bar{A} is an endomorphism on the abelian group $\mathbb{Z}^N/(1 - B)\mathbb{Z}^N$ induced by the matrix A .

Throughout the paper, we will denote by \mathbb{Z}_+ the set of nonnegative integers and by \mathbb{N} the set of positive integers.

This paper is a revised version of the paper: arXiv:1106.5092v1.

2. λ -graph systems, C*-symbolic dynamical systems and their C*-algebras

In this section, we will briefly review λ -graph systems and C*-symbolic dynamical systems. Throughout the section, Σ denotes a finite set with its discrete topology, that is called an alphabet. Each element of Σ is called a symbol. Let $\Sigma^{\mathbb{Z}}$ be the infinite product space $\prod_{i \in \mathbb{Z}} \Sigma_i$, where $\Sigma_i = \Sigma$, endowed with the product topology. The transformation σ on $\Sigma^{\mathbb{Z}}$ given by $\sigma((x_i)_{i \in \mathbb{Z}}) = (x_{i+1})_{i \in \mathbb{Z}}$ is called the full shift over Σ . Let Λ be a shift invariant closed subset of $\Sigma^{\mathbb{Z}}$ i.e. $\sigma(\Lambda) = \Lambda$. The topological dynamical system $(\Lambda, \sigma|_{\Lambda})$ is called a two-sided subshift, written as Λ for brevity. A word $\mu = (\mu_1, \dots, \mu_k)$ of Σ is said to be admissible for Λ if there exists $(x_i)_{i \in \mathbb{Z}} \in \Lambda$ such that $\mu_1 = x_1, \dots, \mu_k = x_k$. Let us denote by $|\mu|$ the length k of μ . Let $B_k(\Lambda)$ be the set of admissible words of Λ with length k . The union $\cup_{k=0}^{\infty} B_k(\Lambda)$ is denoted by $B_*(\Lambda)$ where $B_0(\Lambda)$ denotes the empty word. For two words $\mu = (\mu_1, \dots, \mu_k), \nu = (\nu_1, \dots, \nu_n)$, we write a new word $\mu\nu = (\mu_1, \dots, \mu_k, \nu_1, \dots, \nu_n)$.

There is a class of subshifts called sofic shifts, that are presented by finite labeled graphs ([14], [17], [18]). λ -graph systems are generalization of finite labeled graphs. Any subshift is presented by a λ -graph system. Let

$$\mathfrak{L} = (V, E, \lambda, \iota)$$

be a λ -graph system over Σ with vertex set $V = \cup_{l \in \mathbb{Z}_+} V_l$ and edge set $E = \cup_{l \in \mathbb{Z}_+} E_{l,l+1}$ that is labeled with symbols in Σ by a map $\lambda : E \rightarrow \Sigma$, and that is supplied with surjective maps $\iota (= \iota_{l,l+1}) : V_{l+1} \rightarrow V_l$ for $l \in \mathbb{Z}_+$. Here the vertex sets $V_l, l \in \mathbb{Z}_+$ and the edge sets $E_{l,l+1}, l \in \mathbb{Z}_+$ are finite disjoint sets for each $l \in \mathbb{Z}_+$. An edge e in $E_{l,l+1}$ has its source vertex $s(e)$ in V_l and its terminal vertex $t(e)$ in V_{l+1} respectively. Every vertex in V has a successor and every vertex in V_l for $l \in \mathbb{N}$ has a predecessor. It is then required that for vertices $u \in V_{l-1}$ and $v \in V_{l+1}$, there exists a bijective correspondence between the set of edges $e \in E_{l,l+1}$ such that $t(e) = v, \iota(s(e)) = u$ and the set of edges $f \in E_{l-1,l}$ such that $s(f) = u, t(f) = \iota(v)$, preserving their labels ([24]). We assume that \mathfrak{L} is left-resolving, which means that $t(e) \neq t(f)$

whenever $\lambda(e) = \lambda(f)$ for $e, f \in E_{l,l+1}$. Let us denote by $\{v_1^l, \dots, v_{m(l)}^l\}$ the vertex set V_l at level l . For $i = 1, 2, \dots, m(l)$, $j = 1, 2, \dots, m(l+1)$, $\alpha \in \Sigma$ we put

$$A_{l,l+1}(i, \alpha, j) = \begin{cases} 1 & \text{if } s(e) = v_i^l, \lambda(e) = \alpha, t(e) = v_j^{l+1} \text{ for some } e \in E_{l,l+1}, \\ 0 & \text{otherwise,} \end{cases}$$

$$I_{l,l+1}(i, j) = \begin{cases} 1 & \text{if } \iota_{l,l+1}(v_j^{l+1}) = v_i^l, \\ 0 & \text{otherwise.} \end{cases}$$

The C^* -algebra $\mathcal{O}_{\mathfrak{L}}$ associated with \mathfrak{L} is the universal C^* -algebra generated by partial isometries $S_\alpha, \alpha \in \Sigma$ and projections $E_i^l, i = 1, 2, \dots, m(l), l \in \mathbb{Z}_+$ subject to the following operator relations called (\mathfrak{L}) :

$$(2.1) \quad \sum_{\beta \in \Sigma} S_\beta S_\beta^* = 1,$$

$$(2.2) \quad \sum_{i=1}^{m(l)} E_i^l = 1, \quad E_i^l = \sum_{j=1}^{m(l+1)} I_{l,l+1}(i, j) E_j^{l+1},$$

$$(2.3) \quad S_\alpha S_\alpha^* E_i^l = E_i^l S_\alpha S_\alpha^*,$$

$$(2.4) \quad S_\alpha^* E_i^l S_\alpha = \sum_{j=1}^{m(l+1)} A_{l,l+1}(i, \alpha, j) E_j^{l+1},$$

for $i = 1, 2, \dots, m(l), l \in \mathbb{Z}_+, \alpha \in \Sigma$. If \mathfrak{L} satisfies λ -condition (I) and is λ -irreducible, the C^* -algebra $\mathcal{O}_{\mathfrak{L}}$ is simple and purely infinite ([25], [26]).

Let $\mathcal{A}_{\mathfrak{L},l}$ be the C^* -subalgebra of $\mathcal{O}_{\mathfrak{L}}$ generated by the projections $E_i^l, i = 1, \dots, m(l)$. We denote by $\mathcal{A}_{\mathfrak{L}}$ the C^* -subalgebra of $\mathcal{O}_{\mathfrak{L}}$ generated by all the projections $E_i^l, i = 1, \dots, m(l), l \in \mathbb{Z}_+$. As $\mathcal{A}_{\mathfrak{L},l} \subset \mathcal{A}_{\mathfrak{L},l+1}$ and $\cup_{l \in \mathbb{Z}_+} \mathcal{A}_{\mathfrak{L},l}$ is dense in \mathcal{A} , the algebra $\mathcal{A}_{\mathfrak{L}}$ is a commutative AF-algebra. For $\alpha \in \Sigma$, put

$$\rho_\alpha^{\mathfrak{L}}(X) = S_\alpha^* X S_\alpha \quad \text{for } X \in \mathcal{A}_{\mathfrak{L}}.$$

Then $\{\rho_\alpha^{\mathfrak{L}}\}_{\alpha \in \Sigma}$ yields a family of $*$ -endomorphisms of $\mathcal{A}_{\mathfrak{L}}$ such that $\rho_\alpha^{\mathfrak{L}}(1) \neq 0, \sum_{\alpha \in \Sigma} \rho_\alpha^{\mathfrak{L}}(1) \geq 1$ and for any nonzero $x \in \mathcal{A}_{\mathfrak{L}}, \rho_\alpha^{\mathfrak{L}}(x) \neq 0$ for some $\alpha \in \Sigma$.

The situations above are generalized to C^* -symbolic dynamical systems as follows. Let \mathcal{A} be a unital C^* -algebra. In what follows, an endomorphism of \mathcal{A} means a $*$ -endomorphism of \mathcal{A} that does not necessarily preserve the unit $1_{\mathcal{A}}$ of \mathcal{A} . The unit $1_{\mathcal{A}}$ is denoted by 1 unless we specify. Denote by $Z_{\mathcal{A}}$ the center of \mathcal{A} . Let $\rho_\alpha, \alpha \in \Sigma$ be a finite family of endomorphisms of \mathcal{A} indexed by symbols of a finite set Σ . We assume that $\rho_\alpha(Z_{\mathcal{A}}) \subset Z_{\mathcal{A}}, \alpha \in \Sigma$. The family $\rho_\alpha, \alpha \in \Sigma$ of endomorphisms of \mathcal{A} is said to be *essential* if $\rho_\alpha(1) \neq 0$ for all $\alpha \in \Sigma$ and $\sum_{\alpha} \rho_\alpha(1) \geq 1$. It is said to be *faithful* if for any nonzero $x \in \mathcal{A}$ there exists a symbol $\alpha \in \Sigma$ such that $\rho_\alpha(x) \neq 0$.

Definition 2.1 (cf. [27]). A *C*-symbolic dynamical system* is a triplet $(\mathcal{A}, \rho, \Sigma)$ consisting of a unital C*-algebra \mathcal{A} and an essential and faithful finite family $\{\rho_\alpha\}_{\alpha \in \Sigma}$ of endomorphisms of \mathcal{A} .

As in the above discussion, we have a C*-symbolic dynamical system $(\mathcal{A}_\mathfrak{L}, \rho^\mathfrak{L}, \Sigma)$ from a λ -graph system \mathfrak{L} . In [27], [29], [30], we have defined a C*-symbolic dynamical system in a less restrictive way than the above definition. Instead of the above condition $\sum_{\alpha \in \Sigma} \rho_\alpha(1) \geq 1$ with $\rho_\alpha(Z_\mathcal{A}) \subset Z_\mathcal{A}, \alpha \in \Sigma$, we have used the condition in the papers that the closed ideal generated by $\rho_\alpha(1), \alpha \in \Sigma$ coincides with \mathcal{A} . All of the examples appeared in the papers [27], [29], [30] satisfy the condition $\sum_{\alpha \in \Sigma} \rho_\alpha(1) \geq 1$ with $\rho_\alpha(Z_\mathcal{A}) \subset Z_\mathcal{A}, \alpha \in \Sigma$, and all discussions in the papers well work under the above new definition.

A C*-symbolic dynamical system $(\mathcal{A}, \rho, \Sigma)$ yields a subshift Λ_ρ over Σ such that a word $(\alpha_1, \dots, \alpha_k)$ of Σ is admissible for Λ_ρ if and only if

$$(\rho_{\alpha_k} \circ \dots \circ \rho_{\alpha_1})(1) \neq 0$$

([27, Proposition 2.1]). We say that a subshift Λ acts on a C*-algebra \mathcal{A} if there exists a C*-symbolic dynamical system $(\mathcal{A}, \rho, \Sigma)$ such that the associated subshift Λ_ρ is Λ .

The C*-algebra \mathcal{O}_ρ associated with a C*-symbolic dynamical system

$$(\mathcal{A}, \rho, \Sigma)$$

has been originally constructed in [27] as a C*-algebra by using the Pimsner’s general construction of C*-algebras from Hilbert C*-bimodules [39] (cf. [15] etc.). It is realized as the universal C*-algebra $C^*(x, S_\alpha; x \in \mathcal{A}, \alpha \in \Sigma)$ generated by $x \in \mathcal{A}$ and partial isometries $S_\alpha, \alpha \in \Sigma$ subject to the following relations called (ρ) :

$$\sum_{\beta \in \Sigma} S_\beta S_\beta^* = 1, \quad x S_\alpha S_\alpha^* = S_\alpha S_\alpha^* x, \quad S_\alpha^* x S_\alpha = \rho_\alpha(x)$$

for all $x \in \mathcal{A}$ and $\alpha \in \Sigma$. The C*-algebra \mathcal{O}_ρ is a generalization of the C*-algebra $\mathcal{O}_\mathfrak{L}$ associated with the λ -graph system \mathfrak{L} .

A C*-symbolic dynamical system $(\mathcal{A}, \rho, \Sigma)$ is said to be *free* if there exists a unital increasing sequence $\mathcal{A}_0 \subset \mathcal{A}_1 \subset \dots \subset \mathcal{A}$ of C*-subalgebras of \mathcal{A} such that:

- (1) $\rho_\alpha(\mathcal{A}_l) \subset \mathcal{A}_{l+1}$ for all $l \in \mathbb{Z}_+$ and $\alpha \in \Sigma$.
- (2) $\cup_{l \in \mathbb{Z}_+} \mathcal{A}_l$ is dense in \mathcal{A} .
- (3) For $j \leq l$ there exists a projection $q \in \mathcal{D}_\rho \cap \mathcal{A}_l'$ such that:
 - (i) $qx \neq 0$ for $0 \neq x \in \mathcal{A}_l$,
 - (ii) $\phi_\rho^n(q)q = 0$ for all $n = 1, 2, \dots, j$,
 where \mathcal{D}_ρ is the C*-subalgebra of \mathcal{O}_ρ generated by elements

$$S_{\mu_1} \cdots S_{\mu_k} x S_{\mu_k}^* \cdots S_{\mu_1}^*$$

for $(\mu_1, \dots, \mu_k) \in B_*(\Lambda_\rho)$ and $x \in \mathcal{A}$, and

$$\phi_\rho(X) = \sum_{\alpha \in \Sigma} S_\alpha X S_\alpha^*, \quad X \in \mathcal{D}_\rho.$$

The freeness has been called condition (I) in [30]. If in particular, one may take the above subalgebras $\mathcal{A}_l \subset \mathcal{A}, l = 0, 1, 2, \dots$ to be of finite dimensional, then $(\mathcal{A}, \rho, \Sigma)$ is said to be *AF-free*. $(\mathcal{A}, \rho, \Sigma)$ is said to be *irreducible* if there is no nontrivial ideal of \mathcal{A} invariant under the positive operator λ_ρ on \mathcal{A} defined by $\lambda_\rho(x) = \sum_{\alpha \in \Sigma} \rho_\alpha(x)$, $x \in \mathcal{A}$. It has been proved that if $(\mathcal{A}, \rho, \Sigma)$ is free and irreducible, then the C^* -algebra \mathcal{O}_ρ is simple ([30]).

3. C^* -textile dynamical systems and two-dimensional subshifts

Let Σ be a finite set. The two-dimensional full shift over Σ is defined to be

$$\Sigma^{\mathbb{Z}^2} = \{(x_{i,j})_{(i,j) \in \mathbb{Z}^2} \mid x_{i,j} \in \Sigma\}.$$

An element $x \in \Sigma^{\mathbb{Z}^2}$ is regarded as a function $x : \mathbb{Z}^2 \rightarrow \Sigma$ which is called a configuration on \mathbb{Z}^2 . For $x \in \Sigma^{\mathbb{Z}^2}$ and $F \subset \mathbb{Z}^2$, let x_F denote the restriction of x to F . For a vector $m = (m_1, m_2) \in \mathbb{Z}^2$, let $\sigma^m : \Sigma^{\mathbb{Z}^2} \rightarrow \Sigma^{\mathbb{Z}^2}$ be the translation along vector m defined by

$$\sigma^m((x_{i,j})_{(i,j) \in \mathbb{Z}^2}) = (x_{i+m_1, j+m_2})_{(i,j) \in \mathbb{Z}^2}.$$

A subset $X \subset \Sigma^{\mathbb{Z}^2}$ is said to be translation invariant if $\sigma^m(X) = X$ for all $m \in \mathbb{Z}^2$. It is obvious to see that a subset $X \subset \Sigma^{\mathbb{Z}^2}$ is translation invariant if and only if X is invariant only both horizontally and vertically, that is, $\sigma^{(1,0)}(X) = X$ and $\sigma^{(0,1)}(X) = X$. For $k \in \mathbb{Z}_+$, put

$$[-k, k]^2 = \{(i, j) \in \mathbb{Z}^2 \mid -k \leq i, j \leq k\} = [-k, k] \times [-k, k].$$

A metric d on $\Sigma^{\mathbb{Z}^2}$ is defined by for $x, y \in \Sigma^{\mathbb{Z}^2}$ with $x \neq y$

$$d(x, y) = \frac{1}{2^k} \quad \text{if} \quad x_{(0,0)} = y_{(0,0)},$$

where $k = \max\{k \in \mathbb{Z}_+ \mid x_{[-k,k]^2} = y_{[-k,k]^2}\}$. If $x_{(0,0)} \neq y_{(0,0)}$, put $k = -1$ on the above definition. If $x = y$, we set $d(x, y) = 0$. A two-dimensional subshift X is defined to be a closed, translation invariant subset of $\Sigma^{\mathbb{Z}^2}$ (cf. [21, p.467]). A finite subset $F \subset \mathbb{Z}^2$ is said to be a shape. A pattern f on a shape F is a function $f : F \rightarrow \Sigma$. For a list \mathfrak{F} of patterns, put

$$X_{\mathfrak{F}} = \{(x_{i,j})_{(i,j) \in \mathbb{Z}^2} \mid \sigma^m(x)|_F \notin \mathfrak{F} \text{ for all } m \in \mathbb{Z}^2 \text{ and } F \subset \mathbb{Z}^2\}.$$

It is well-known that a subset $X \subset \Sigma^{\mathbb{Z}^2}$ is a two-dimensional subshift if and only if there exists a list \mathfrak{F} of patterns such that $X = X_{\mathfrak{F}}$.

We will define a certain property of two-dimensional subshift as follows:

Definition 3.1. A two-dimensional subshift X is said to have the *diagonal property* if for $(x_{i,j})_{(i,j) \in \mathbb{Z}^2}, (y_{i,j})_{(i,j) \in \mathbb{Z}^2} \in X$, the conditions

$$x_{i,j} = y_{i,j}, \quad x_{i+1,j-1} = y_{i+1,j-1}$$

imply

$$x_{i,j-1} = y_{i,j-1}, \quad x_{i+1,j} = y_{i+1,j}.$$

A two-dimensional subshift having the diagonal property is called a *textile dynamical system*.

Lemma 3.2. *If a two dimensional subshift X has the diagonal property, then for $x \in X$ and $(i, j) \in \mathbb{Z}^2$, the configuration x is determined by the diagonal line $(x_{i+n,j-n})_{n \in \mathbb{Z}}$ through (i, j) .*

Proof. By the diagonal property, the sequence $(x_{i+n,j-n})_{n \in \mathbb{Z}}$ determines both the sequences $(x_{i+1+n,j-n})_{n \in \mathbb{Z}}$ and $(x_{i-1+n,j-n})_{n \in \mathbb{Z}}$. Repeating this way, the sequence $(x_{i+n,j-n})_{n \in \mathbb{Z}}$ determines the whole configuration x . \square

Let $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ be a C^* -textile dynamical system. It consists of two C^* -symbolic dynamical systems $(\mathcal{A}, \rho, \Sigma^\rho)$ and $(\mathcal{A}, \eta, \Sigma^\eta)$ with common unital C^* -algebra \mathcal{A} and commutation relations between their endomorphisms $\rho_\alpha, \alpha \in \Sigma^\rho, \eta_a, a \in \Sigma^\eta$ through a bijection κ between the following sets $\Sigma^{\rho\eta}$ and $\Sigma^{\eta\rho}$, where

$$\begin{aligned} \Sigma^{\rho\eta} &= \{(\alpha, b) \in \Sigma^\rho \times \Sigma^\eta \mid \eta_b \circ \rho_\alpha \neq 0\}, \\ \Sigma^{\eta\rho} &= \{(a, \beta) \in \Sigma^\eta \times \Sigma^\rho \mid \rho_\beta \circ \eta_a \neq 0\}. \end{aligned}$$

The given bijection $\kappa : \Sigma^{\rho\eta} \longrightarrow \Sigma^{\eta\rho}$ is called a specification. The required commutation relations are

$$(3.1) \quad \eta_b \circ \rho_\alpha = \rho_\beta \circ \eta_a \quad \text{if } \kappa(\alpha, b) = (a, \beta).$$

A C^* -textile dynamical system will yield a two-dimensional subshift $X_{\rho,\eta}^\kappa$. We set

$$\Sigma_\kappa = \{\omega = (\alpha, b, a, \beta) \in \Sigma^\rho \times \Sigma^\eta \times \Sigma^\eta \times \Sigma^\rho \mid \kappa(\alpha, b) = (a, \beta)\}.$$

For $\omega = (\alpha, b, a, \beta)$, since $\eta_b \circ \rho_\alpha = \rho_\beta \circ \eta_a$ as endomorphisms on \mathcal{A} , one may identify the quadruplet (α, b, a, β) with the endomorphism $\eta_b \circ \rho_\alpha (= \rho_\beta \circ \eta_a)$ on \mathcal{A} which we will denote by simply ω . Define maps $t(= \text{top}), b(= \text{bottom}) : \Sigma_\kappa \longrightarrow \Sigma^\rho$ and $l(= \text{left}), r(= \text{right}) : \Sigma_\kappa \longrightarrow \Sigma^\rho$ by setting

$$t(\omega) = \alpha, \quad b(\omega) = \beta, \quad l(\omega) = a, \quad r(\omega) = b.$$

$$\begin{array}{ccc} \cdot & \xrightarrow{\alpha=t(\omega)} & \cdot \\ a=l(\omega) \downarrow & & \downarrow b=r(\omega) \\ \cdot & \xrightarrow{\beta=b(\omega)} & \cdot \end{array}$$

A configuration $(\omega_{i,j})_{(i,j) \in \mathbb{Z}^2} \in \Sigma_{\kappa}^{\mathbb{Z}^2}$ is said to be *paved* if the conditions

$$\begin{aligned} t(\omega_{i,j}) &= b(\omega_{i,j+1}), & r(\omega_{i,j}) &= l(\omega_{i+1,j}), \\ l(\omega_{i,j}) &= r(\omega_{i-1,j}), & b(\omega_{i,j}) &= t(\omega_{i,j-1}) \end{aligned}$$

hold for all $(i, j) \in \mathbb{Z}^2$. We set

$$\begin{aligned} X_{\rho,\eta}^{\kappa} &= \{(\omega_{i,j})_{(i,j) \in \mathbb{Z}^2} \in \Sigma_{\kappa}^{\mathbb{Z}^2} \mid (\omega_{i,j})_{(i,j) \in \mathbb{Z}^2} \text{ is paved and} \\ &\quad \omega_{i+n,j-n} \circ \omega_{i+n-1,j-n+1} \circ \cdots \circ \omega_{i+1,j-1} \circ \omega_{i,j} \neq 0 \\ &\quad \text{for all } (i, j) \in \mathbb{Z}^2, n \in \mathbb{N}\}, \end{aligned}$$

where $\omega_{i+n,j-n} \circ \omega_{i+n-1,j-n+1} \circ \cdots \circ \omega_{i+1,j-1} \circ \omega_{i,j}$ is the compositions as endomorphisms on \mathcal{A} .

Lemma 3.3. *Suppose that a configuration $(\omega_{i,j})_{(i,j) \in \mathbb{Z}^2} \in \Sigma_{\kappa}^{\mathbb{Z}^2}$ is paved. Then $(\omega_{i,j})_{(i,j) \in \mathbb{Z}^2} \in X_{\rho,\eta}^{\kappa}$ if and only if*

$$\rho b(\omega_{i+n,j-m}) \circ \cdots \circ \rho b(\omega_{i+1,j-m}) \circ \rho b(\omega_{i,j-m}) \circ \eta l(\omega_{i,j-m}) \circ \cdots \circ \eta l(\omega_{i,j-1}) \circ \eta l(\omega_{i,j}) \neq 0$$

for all $(i, j) \in \mathbb{Z}^2, n, m \in \mathbb{Z}_+$.

$$\begin{array}{ccccccc} & & \cdot & & & & \\ & & \downarrow & & & & \\ & & l(\omega_{i,j}) & & & & \\ & & \cdot & & & & \\ & & \downarrow & & & & \\ & & l(\omega_{i,j-1}) & & & & \\ & & \cdot & & & & \\ & & \vdots & & & & \\ & & \cdot & & & & \\ & & \downarrow & & & & \\ & & l(\omega_{i,j-m}) & & & & \\ & \cdot & \xrightarrow{b(\omega_{i,j-m})} & \cdot & \xrightarrow{b(\omega_{i+1,j-m})} & \cdots & \xrightarrow{b(\omega_{i+n,j-m})} & \cdot \end{array}$$

Proof. Suppose that $(\omega_{i,j})_{(i,j) \in \mathbb{Z}^2} \in X_{\rho,\eta}^{\kappa}$. For $(i, j) \in \mathbb{Z}^2, n, m \in \mathbb{Z}_+$, we may assume that $m \geq n$. Since

$$\begin{aligned} &0 \neq \omega_{i+m,j-m} \circ \cdots \circ \omega_{i+n+1,j-m} \circ \omega_{i+n,j-m} \circ \cdots \circ \omega_{i,j-m} \\ &\quad \circ \cdots \circ \omega_{i+1,j-1} \circ \omega_{i,j} \\ &= \omega_{i+m,j-m} \circ \cdots \circ \omega_{i+n+1,j-m} \circ \rho b(\omega_{i+n,j-m}) \circ \cdots \circ \rho b(\omega_{i+1,j-m}) \circ \rho b(\omega_{i,j-m}) \\ &\quad \circ \eta l(\omega_{i,j-m}) \circ \cdots \circ \eta l(\omega_{i,j-1}) \circ \eta l(\omega_{i,j}), \end{aligned}$$

one has

$$\rho b(\omega_{i+n,j-m}) \circ \cdots \circ \rho b(\omega_{i+1,j-m}) \circ \rho b(\omega_{i,j-m}) \circ \eta l(\omega_{i,j-m}) \circ \cdots \circ \eta l(\omega_{i,j-1}) \circ \eta l(\omega_{i,j}) \neq 0.$$

The converse implication is clear by the equality:

$$\begin{aligned} &\omega_{i+n,j-n} \circ \cdots \circ \omega_{i,j-n} \circ \cdots \circ \omega_{i,j-1} \circ \omega_{i,j} \\ &= \rho_{b(\omega_{i+n,j-n})} \circ \cdots \circ \rho_{b(\omega_{i,j-n})} \circ \eta_{l(\omega_{i,j-n})} \cdots \circ \eta_{l(\omega_{i,j-1})} \circ \eta_{l(\omega_{i,j})}. \quad \square \end{aligned}$$

Proposition 3.4. $X_{\rho,\eta}^\kappa$ is a two-dimensional subshift having diagonal property, that is, $X_{\rho,\eta}^\kappa$ is a textile dynamical system.

Proof. It is easy to see that the set

$$E = \{(\omega_{i,j})_{(i,j) \in \mathbb{Z}^2} \in \Sigma_\kappa^{\mathbb{Z}^2} \mid (\omega_{i,j})_{(i,j) \in \mathbb{Z}^2} \text{ is paved}\}$$

is closed, because its complement is open in $\Sigma_\kappa^{\mathbb{Z}^2}$. The following set

$$\begin{aligned} U = \{(\omega_{i,j})_{(i,j) \in \mathbb{Z}^2} \in \Sigma_\kappa^{\mathbb{Z}^2} \mid &\omega_{k+n,l-n} \circ \omega_{k+n-1,l-n+1} \\ &\circ \cdots \circ \omega_{k+1,l-1} \circ \omega_{k,l} = 0 \text{ for some } (k,l) \in \mathbb{Z}^2, n \in \mathbb{N}\} \end{aligned}$$

is open in $\Sigma_\kappa^{\mathbb{Z}^2}$. As the equality $X_{\rho,\eta}^\kappa = E \cap U^c$ holds, the set $X_{\rho,\eta}^\kappa$ is closed. It is also obvious that $X_{\rho,\eta}^\kappa$ is translation invariant so that $X_{\rho,\eta}^\kappa$ is a two-dimensional subshift. It is easy to see that $X_{\rho,\eta}^\kappa$ has diagonal property. \square

We call $X_{\rho,\eta}^\kappa$ the textile dynamical system associated with

$$(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa).$$

Let us now define a (one-dimensional) subshift X_{δ^κ} over Σ_κ , which consists of diagonal sequences of $X_{\rho,\eta}^\kappa$ as follows:

$$X_{\delta^\kappa} = \{(\omega_{n,-n})_{n \in \mathbb{Z}} \in \Sigma_\kappa^{\mathbb{Z}} \mid (\omega_{i,j})_{(i,j) \in \mathbb{Z}^2} \in X_{\rho,\eta}^\kappa\}.$$

By Lemma 3.2, an element $(\omega_{n,-n})_{n \in \mathbb{Z}}$ of X_{δ^κ} may be extended to

$$(\omega_{i,j})_{(i,j) \in \mathbb{Z}^2} \in X_{\rho,\eta}^\kappa$$

in a unique way. Hence the one-dimensional subshift X_{δ^κ} determines the two-dimensional subshift $X_{\rho,\eta}^\kappa$. Therefore we have:

Lemma 3.5. *The two-dimensional subshift $X_{\rho,\eta}^\kappa$ is not empty if and only if the one-dimensional subshift X_{δ^κ} is not empty.*

For $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$, we will have a C^* -symbolic dynamical system $(\mathcal{A}, \delta^\kappa, \Sigma_\kappa)$ in Section 4. It presents the subshift X_{δ^κ} . Since a subshift presented by a C^* -symbolic dynamical system is always not empty, one sees

Proposition 3.6. *The two-dimensional subshift $X_{\rho,\eta}^\kappa$ is not empty.*

4. C^* -textile dynamical systems and their C^* -algebras

The C^* -algebra $\mathcal{O}_{\rho,\eta}^\kappa$ is defined to be the universal C^* -algebra

$$C^*(x, S_\alpha, T_a; x \in \mathcal{A}, \alpha \in \Sigma^\rho, a \in \Sigma^\eta)$$

generated by $x \in \mathcal{A}$ and partial isometries $S_\alpha, \alpha \in \Sigma^\rho, T_a, a \in \Sigma^\eta$ subject to the following relations called $(\rho, \eta; \kappa)$:

$$(4.1) \quad \sum_{\beta \in \Sigma^\rho} S_\beta S_\beta^* = 1, \quad x S_\alpha S_\alpha^* = S_\alpha S_\alpha^* x, \quad S_\alpha^* x S_\alpha = \rho_\alpha(x),$$

$$(4.2) \quad \sum_{b \in \Sigma^\eta} T_b T_b^* = 1, \quad x T_a T_a^* = T_a T_a^* x, \quad T_a^* x T_a = \eta_a(x),$$

$$(4.3) \quad S_\alpha T_b = T_a S_\beta \quad \text{if} \quad \kappa(\alpha, b) = (a, \beta)$$

for all $x \in \mathcal{A}$ and $\alpha \in \Sigma^\rho, a \in \Sigma^\eta$. We will study the algebra $\mathcal{O}_{\rho,\eta}^\kappa$. For $(\alpha, b, a, \beta) \in \Sigma^\rho \times \Sigma^\eta \times \Sigma^\eta \times \Sigma^\rho$, we set

$$\begin{aligned} RB(\alpha, a) &= \{(b, \beta) \in \Sigma^\eta \times \Sigma^\rho \mid \kappa(\alpha, b) = (a, \beta)\}, \\ R(\alpha, a, \beta) &= \{b \in \Sigma^\eta \mid \kappa(\alpha, b) = (a, \beta)\}, \\ R(\alpha, a) &= \bigcup_{\beta \in \Sigma^\rho} R(\alpha, a, \beta). \end{aligned}$$

Lemma 4.1. *For $\alpha \in \Sigma^\rho, a \in \Sigma^\eta$, one has $T_a^* S_\alpha \neq 0$ if and only if $RB(\alpha, a) \neq \emptyset$.*

Proof. Suppose that $T_a^* S_\alpha \neq 0$. As $T_a^* S_\alpha = \sum_{b' \in \Sigma^\eta} T_a^* S_\alpha T_{b'} T_{b'}^*$, there exists $b' \in \Sigma^\eta$ such that $T_a^* S_\alpha T_{b'} \neq 0$. Hence $\eta_{b'} \circ \rho_\alpha \neq 0$ so that $(\alpha, b') \in \Sigma^{\rho\eta}$. Then one may find $(a', \beta') \in \Sigma^\rho$ such that $\kappa(\alpha, b') = (a', \beta')$ and hence $S_\alpha T_{b'} = T_{a'} S_{\beta'}$. Since $0 \neq T_a^* S_\alpha T_{b'} = T_a^* T_{a'} S_{\beta'}$, one sees that $a = a'$ so that $(b', \beta') \in RB(\alpha, a)$.

Suppose next that $\kappa(\alpha, b) = (a, \beta)$ for some $(b, \beta) \in \Sigma^\eta \times \Sigma^\rho$. Since $\eta_b \circ \rho_\alpha = \rho_\beta \circ \eta_a \neq 0$, one has $0 \neq S_\alpha T_b = T_a S_\beta$. It follows that

$$S_\beta^* T_a^* S_\alpha T_b = (T_a S_\beta)^* T_a S_\beta$$

so that $T_a^* S_\alpha \neq 0$. □

Lemma 4.2. *For $\alpha \in \Sigma^\rho, a \in \Sigma^\eta$, we have*

$$(4.4) \quad T_a^* S_\alpha = \sum_{(b,\beta) \in RB(\alpha,a)} S_\beta \eta_b(\rho_\alpha(1)) T_b^*$$

and hence

$$(4.5) \quad S_\alpha^* T_a = \sum_{(b,\beta) \in RB(\alpha,a)} T_b \rho_\beta(\eta_a(1)) S_\beta^*.$$

Proof. We may assume that $T_a^* S_\alpha \neq 0$. One has

$$T_a^* S_\alpha = \sum_{b' \in \Sigma^\eta} T_a^* S_\alpha T_{b'} T_{b'}^*.$$

For $b' \in \Sigma^\eta$ with $(\alpha, b') \in \Sigma^{\rho\eta}$, take $(a', \beta') \in \Sigma^{\eta\rho}$ such that $\kappa(\alpha, b') = (a', \beta')$ so that

$$T_a^* S_\alpha T_{b'} T_{b'}^* = T_a^* T_{a'} S_{\beta'} T_{b'}^*.$$

Hence $T_a^* S_\alpha T_{b'} T_{b'}^* \neq 0$ implies $a = a'$. Since $T_a^* T_a = \eta_\alpha(1)$ which commutes with $S_{\beta'} S_{\beta'}^*$, we have

$$T_a^* T_a S_{\beta'} T_{b'}^* = S_{\beta'} S_{\beta'}^* T_a^* T_a S_{\beta'} T_{b'}^* = S_{\beta'} \rho_{\beta'}(\eta_\alpha(1)) T_{b'}^* = S_{\beta'} \eta_{b'}(\rho_\alpha(1)) T_{b'}^*.$$

It follows that

$$T_a^* S_\alpha = \sum_{(b', \beta') \in RB(\alpha, a)} T_a^* T_a S_{\beta'} T_{b'}^* = \sum_{(b', \beta') \in RB(\alpha, a)} S_{\beta'} \eta_{b'}(\rho_\alpha(1)) T_{b'}^*. \quad \square$$

Hence we have:

Lemma 4.3. For $\alpha \in \Sigma^\rho, a \in \Sigma^\eta$, we have

$$T_a T_a^* S_\alpha S_\alpha^* = \sum_{b \in R(\alpha, a)} S_\alpha T_b T_b^* S_\alpha^*.$$

Hence $T_a T_a^*$ commutes with $S_\alpha S_\alpha^*$.

Proof. By (4.4), we have

$$\begin{aligned} T_a T_a^* S_\alpha S_\alpha^* &= \sum_{(b, \beta) \in RB(\alpha, a)} T_a S_\beta \eta_b(\rho_\alpha(1)) T_b^* S_\alpha^* \\ &= \sum_{b \in R(\alpha, a)} S_\alpha T_b \eta_b(\rho_\alpha(1)) T_b^* S_\alpha^* \\ &= \sum_{b \in R(\alpha, a)} S_\alpha \rho_\alpha(1) T_b T_b^* S_\alpha^* \\ &= \sum_{b \in R(\alpha, a)} S_\alpha T_b T_b^* S_\alpha^*. \quad \square \end{aligned}$$

Recall that $Z_{\mathcal{A}}$ denotes the center of \mathcal{A} which consists of elements of \mathcal{A} commuting with all elements of \mathcal{A} .

Lemma 4.4. For $\alpha \in \Sigma^\rho, a \in \Sigma^\eta$ and $x, y \in Z_{\mathcal{A}}$, $T_a y T_a^*$ commutes with $S_\alpha x S_\alpha^*$.

Proof. By (4.4), we have

$$\begin{aligned}
 T_a y T_a^* S_\alpha x S_\alpha^* &= T_a y \sum_{(b,\beta) \in RB(\alpha,a)} S_\beta \eta_b(\rho_\alpha(1)) T_b^* x S_\alpha^* \\
 &= \sum_{(b,\beta) \in RB(\alpha,a)} T_a S_\beta S_\beta^* y S_\beta \eta_b(\rho_\alpha(1)) T_b^* x T_b T_b^* S_\alpha^* \\
 &= \sum_{(b,\beta) \in RB(\alpha,a)} S_\alpha T_b \rho_\beta(y) \eta_b(\rho_\alpha(1)) \eta_b(x) S_\beta^* T_a^* \\
 &= \sum_{(b,\beta) \in RB(\alpha,a)} S_\alpha T_b \eta_b(x) \eta_b(\rho_\alpha(1)) \rho_\beta(y) S_\beta^* T_a^* \\
 &= \sum_{(b,\beta) \in RB(\alpha,a)} S_\alpha x \rho_\alpha(1) T_b S_\beta^* y T_a^* \\
 &= \sum_{(b,\beta) \in RB(\alpha,a)} S_\alpha x S_\alpha^* S_\alpha T_b S_\beta^* T_a^* T_a y T_a^* \\
 &= \sum_{b \in R(\alpha,a)} S_\alpha x \cdot S_\alpha^* S_\alpha T_b T_b^* S_\alpha^* T_a \cdot y T_a^*.
 \end{aligned}$$

Now if $(\alpha, b') \notin \Sigma^{\rho,\eta}$, then $S_\alpha T_{b'} = 0$. Hence

$$\sum_{b \in R(\alpha,a)} S_\alpha^* S_\alpha T_b T_b^* S_\alpha^* T_a = \sum_{b \in \Sigma^\eta} S_\alpha^* S_\alpha T_b T_b^* S_\alpha^* T_a = S_\alpha^* T_a.$$

Therefore we have

$$T_a y T_a^* S_\alpha x S_\alpha^* = S_\alpha x S_\alpha^* T_a y T_a^*. \quad \square$$

For words $\mu = (\mu_1, \dots, \mu_j) \in B_j(\Lambda_\rho), \zeta = (\zeta_1, \dots, \zeta_k) \in B_k(\Lambda_\eta)$, we set

$$S_\mu = S_{\mu_1} \cdots S_{\mu_j}, \quad T_\zeta = T_{\zeta_1} \cdots T_{\zeta_k}.$$

For a subset F of $\mathcal{O}_{\rho,\eta}^\kappa$, denote by $C^*(F)$ the C^* -subalgebra of $\mathcal{O}_{\rho,\eta}^\kappa$ generated by the elements of F . We define C^* -subalgebras $\mathcal{D}_{\rho,\eta}, \mathcal{D}_{j,k}$ of $\mathcal{O}_{\rho,\eta}^\kappa$ by

$$\begin{aligned}
 \mathcal{D}_{\rho,\eta} &= C^*(S_\mu T_\zeta x T_\zeta^* S_\mu^* : \mu \in B_*(\Lambda_\rho), \zeta \in B_*(\Lambda_\eta), x \in \mathcal{A}), \\
 \mathcal{D}_{j,k} &= C^*(S_\mu T_\zeta x T_\zeta^* S_\mu^* : \mu \in B_j(\Lambda_\rho), \zeta \in B_k(\Lambda_\eta), x \in \mathcal{A}) \quad \text{for } j, k \in \mathbb{Z}_+.
 \end{aligned}$$

By the commutation relation (4.3), one sees that

$$\mathcal{D}_{j,k} = C^*(T_\xi S_\nu x S_\nu^* T_\xi^* : \nu \in B_j(\Lambda_\rho), \xi \in B_k(\Lambda_\eta), x \in \mathcal{A}).$$

The identities

$$\begin{aligned}
 S_\mu T_\zeta x T_\zeta^* S_\mu^* &= \sum_{a \in \Sigma^\eta} S_\mu T_{\zeta a} \eta_a(x) T_{\zeta a}^* S_\mu^*, \\
 T_\xi S_\nu x S_\nu^* T_\xi^* &= \sum_{\alpha \in \Sigma^\rho} T_\xi S_{\nu \alpha} \rho_\alpha(x) S_{\nu \alpha}^* T_\xi^*
 \end{aligned}$$

for $x \in \mathcal{A}$ and $\mu, \nu \in B_j(\Lambda_\rho), \zeta, \xi \in B_k(\Lambda_\eta)$ yield the embeddings

$$\mathcal{D}_{j,k} \hookrightarrow \mathcal{D}_{j,k+1}, \quad \mathcal{D}_{j,k} \hookrightarrow \mathcal{D}_{j+1,k}$$

respectively such that $\cup_{j,k \in \mathbb{Z}_+} \mathcal{D}_{j,k}$ is dense in $\mathcal{D}_{\rho,\eta}$.

Proposition 4.5. *If \mathcal{A} is commutative, so is $\mathcal{D}_{\rho,\eta}$.*

Proof. The preceding lemma tells us that $\mathcal{D}_{1,1}$ is commutative. Suppose that the algebra $\mathcal{D}_{j,k}$ is commutative for fixed $j, k \in \mathbb{N}$. We will show that the both algebras $\mathcal{D}_{j+1,k}$ and $\mathcal{D}_{j,k+1}$ are commutative. The algebra $\mathcal{D}_{j+1,k}$ consists of the linear span of elements of the form:

$$S_\alpha x S_\alpha^* \quad \text{for } x \in \mathcal{D}_{j,k}, \alpha \in \Sigma^\rho.$$

For $x, y \in \mathcal{D}_{j,k}, \alpha, \beta \in \Sigma^\rho$, we will show that $S_\alpha x S_\alpha^*$ commutes with both $S_\beta y S_\beta^*$ and y . If $\alpha = \beta$, it is easy to see that $S_\alpha x S_\alpha^*$ commutes with $S_\alpha y S_\alpha^*$, because $\rho_\alpha(1) \in \mathcal{A} \subset \mathcal{D}_{j,k}$. If $\alpha \neq \beta$, both $S_\alpha x S_\alpha^* S_\beta y S_\beta^*$ and $S_\beta y S_\beta^* S_\alpha x S_\alpha^*$ are zeros. Since $S_\alpha^* y S_\alpha \in \mathcal{D}_{j-1,k} \subset \mathcal{D}_{j,k}$, one sees $S_\alpha^* y S_\alpha$ commutes with x . One also sees that $S_\alpha S_\alpha^* \in \mathcal{D}_{j,k}$ commutes with y . It follows that

$$S_\alpha x S_\alpha^* y = S_\alpha x S_\alpha^* y S_\alpha S_\alpha^* = S_\alpha S_\alpha^* y S_\alpha x S_\alpha^* = y S_\alpha x S_\alpha^*.$$

Hence the algebra $\mathcal{D}_{j+1,k}$ is commutative, and similarly so is $\mathcal{D}_{j,k+1}$. By induction, the algebras $\mathcal{D}_{j,k}$ are all commutative for all $j, k \in \mathbb{N}$. Since $\cup_{j,k \in \mathbb{N}} \mathcal{D}_{j,k}$ is dense in $\mathcal{D}_{\rho,\eta}$, $\mathcal{D}_{\rho,\eta}$ is commutative. \square

Proposition 4.6. *Let $\mathcal{O}_{\rho,\eta}^{alg}$ be the dense *-subalgebra of $\mathcal{O}_{\rho,\eta}^\kappa$ algebraically generated by elements $x \in \mathcal{A}, S_\alpha, \alpha \in \Sigma^\rho$ and $T_a, a \in \Sigma^\eta$. Then each element of $\mathcal{O}_{\rho,\eta}^{alg}$ is a finite linear combination of elements of the form:*

$$(4.6) \quad S_\mu T_\zeta x T_\xi^* S_\nu^* \quad \text{for } x \in \mathcal{A}, \mu, \nu \in B_*(\Lambda_\rho), \zeta, \xi \in B_*(\Lambda_\eta).$$

Proof. For $\alpha, \beta \in \Sigma^\rho, a, b \in \Sigma^\eta$ and $x \in \mathcal{A}$, we have

$$\begin{aligned} S_\alpha^* S_\beta &= \begin{cases} \rho_\alpha(1) \in \mathcal{A} & \text{if } \alpha = \beta, \\ 0 & \text{otherwise,} \end{cases} & T_a^* T_b &= \begin{cases} \eta_a(1) \in \mathcal{A} & \text{if } a = b, \\ 0 & \text{otherwise,} \end{cases} \\ S_\alpha^* T_a &= \sum_{(b,\beta) \in RB(\alpha,a)} T_b \rho_\beta(\eta_a(1)) S_\beta^*, & T_a^* S_\alpha &= \sum_{(b,\beta) \in RB(\alpha,a)} S_\beta \eta_b(\rho_\alpha(1)) T_b^*, \\ S_\alpha^* x &= \rho_\alpha(x) S_\alpha, & T_a^* x &= \eta_a(x) T_a^*. \end{aligned}$$

And also

$$S_\beta^* T_a^* = \begin{cases} T_b^* S_\alpha^* & \text{if } (a, \beta) \in \Sigma^{\eta\rho} \text{ and } (a, \beta) = \kappa(\alpha, b), \\ 0 & \text{if } (a, \beta) \notin \Sigma^{\eta\rho}. \end{cases}$$

Therefore we conclude that any element of $\mathcal{O}_{\rho,\eta}^{alg}$ is a finite linear combination of elements of the form of (4.6). \square

Similarly we have:

Proposition 4.7. *Each element of $\mathcal{O}_{\rho,\eta}^{alg}$ is a finite linear combination of elements of the form:*

$$(4.7) \quad T_\zeta S_\mu x S_\nu^* T_\xi^* \quad \text{for } x \in \mathcal{A}, \mu, \nu \in B_*(\Lambda_\rho), \zeta, \xi \in B_*(\Lambda_\eta).$$

In the rest of this section, we will have a C^* -symbolic dynamical system $(\mathcal{A}, \delta^\kappa, \Sigma_\kappa)$ from $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$, which presents the one-dimensional subshift X_{δ^κ} described in the previous section. For $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$, define an endomorphism δ_ω^κ on \mathcal{A} for $\omega \in \Sigma_\kappa$ by setting

$$\delta_\omega^\kappa(x) = \eta_b(\rho_\alpha(x)) (= \rho_\beta(\eta_a(x))), \quad x \in \mathcal{A}, \quad \omega = (\alpha, b, a, \beta) \in \Sigma_\kappa.$$

Lemma 4.8. $(\mathcal{A}, \delta^\kappa, \Sigma_\kappa)$ is a C^* -symbolic dynamical system that presents X_{δ^κ} .

Proof. We will show that δ^κ is essential and faithful. Now both C^* -symbolic dynamical systems $(\mathcal{A}, \eta, \Sigma^\eta)$ and $(\mathcal{A}, \rho, \Sigma^\rho)$ are essential. Since $\rho_\alpha(Z_{\mathcal{A}}) \subset Z_{\mathcal{A}}$ and $\eta_a(Z_{\mathcal{A}}) \subset Z_{\mathcal{A}}$, it is clear that $\delta_\omega^\kappa(Z_{\mathcal{A}}) \subset Z_{\mathcal{A}}$. By the inequalities

$$\sum_{\omega \in \Sigma_\kappa} \delta_\omega^\kappa(1) = \sum_{b \in \Sigma^\eta} \sum_{\alpha \in \Sigma^\rho} \eta_b(\rho_\alpha(1)) \geq \sum_{b \in \Sigma^\eta} \eta_b(1) \geq 1$$

$\{\delta^\kappa\}_{\omega \in \Sigma_\kappa}$ is essential. For any nonzero $x \in \mathcal{A}$, there exists $\alpha \in \Sigma^\rho$ such that $\rho_\alpha(x) \neq 0$ and there exists $b \in \Sigma^\eta$ such that $\eta_b(\rho_\alpha(x)) \neq 0$. Hence δ^κ is faithful so that $(\mathcal{A}, \delta^\kappa, \Sigma_\kappa)$ is a C^* -symbolic dynamical system. It is obvious that the subshift presented by $(\mathcal{A}, \delta^\kappa, \Sigma_\kappa)$ is X_{δ^κ} . \square

Put

$$\widehat{X}_{\rho, \eta}^\kappa = \{(\omega_{i, -j})_{(i, j) \in \mathbb{N}^2} \in \Sigma_\kappa^{\mathbb{N}^2} \mid (\omega_{i, j})_{(i, j) \in \mathbb{Z}^2} \in X_{\rho, \eta}^\kappa\}$$

and

$$\widehat{X}_{\delta^\kappa} = \{(\omega_{n, -n})_{n \in \mathbb{N}} \in \Sigma_\kappa^{\mathbb{N}} \mid (\omega_{i, j})_{(i, j) \in \mathbb{N}^2} \in \widehat{X}_{\rho, \eta}^\kappa\}.$$

The latter set $\widehat{X}_{\delta^\kappa}$ is the right one-sided subshift for X_{δ^κ} .

Lemma 4.9. A configuration $(\omega_{i, -j})_{(i, j) \in \mathbb{N}^2} \in \widehat{X}_{\rho, \eta}^\kappa$ extends to a whole configuration $(\omega_{i, j})_{(i, j) \in \mathbb{Z}^2} \in X_{\rho, \eta}^\kappa$.

Proof. For $(\omega_{i, -j})_{(i, j) \in \mathbb{N}^2} \in \widehat{X}_{\rho, \eta}^\kappa$, put $x_i = \omega_{i, -i}, i \in \mathbb{N}$ so that $x = (x_i)_{i \in \mathbb{N}} \in \widehat{X}_{\delta^\kappa}$. Since $\widehat{X}_{\delta^\kappa}$ is a one-sided subshift, there exists an extension $\tilde{x} \in X_{\delta^\kappa}$ to two-sided sequence such that $\tilde{x}_i = x_i$ for $i \in \mathbb{N}$. By the diagonal property, \tilde{x} determines a whole configuration $\tilde{\omega}$ to \mathbb{Z}^2 such that $\tilde{\omega} \in X_{\delta, \eta}^\kappa$ and $(\tilde{\omega}_{i, -i})_{i \in \mathbb{N}} = \tilde{x}$. Hence $\tilde{\omega}_{i, -j} = \omega_{i, -j}$ for all $i, j \in \mathbb{N}$. \square

Let $\mathfrak{D}_{\rho, \eta}$ be the C^* -subalgebra of $\mathcal{D}_{\rho, \eta}$ defined by

$$\begin{aligned} \mathfrak{D}_{\rho, \eta} &= C^*(S_\mu T_\zeta T_\zeta^* S_\mu^* : \mu \in B_*(\Lambda_\rho), \zeta \in B_*(\Lambda_\eta)) \\ &= C^*(T_\xi S_\nu S_\nu^* T_\xi^* : \nu \in B_*(\Lambda_\rho), \xi \in B_*(\Lambda_\eta)) \end{aligned}$$

which is a commutative C^* -subalgebra of $\mathcal{D}_{\rho, \eta}$. Put for $\mu = (\mu_1, \dots, \mu_n) \in B_*(\Lambda_\rho), \zeta = (\zeta_1, \dots, \zeta_m) \in B_*(\Lambda_\eta)$ the cylinder set

$$\begin{aligned} U_{\mu, \zeta} &= \{(\omega_{i, -j})_{(i, j) \in \mathbb{N}^2} \in \widehat{X}_{\rho, \eta}^\kappa \mid \\ &\quad t(\omega_{i, -1}) = \mu_i, i = 1, \dots, n, r(\omega_{n, -j}) = \zeta_j, j = 1, \dots, m\}. \end{aligned}$$

The following lemma is direct.

Lemma 4.10. $\mathfrak{D}_{\rho,\eta}$ is isomorphic to $C(\widehat{X}_{\rho,\eta}^\kappa)$ through the correspondence such that $S_\mu T_\zeta T_\zeta^* S_\mu^*$ goes to $\chi_{U_{\mu,\zeta}}$, where $\chi_{U_{\mu,\zeta}}$ is the characteristic function for the cylinder set $U_{\mu,\zeta}$ on $\widehat{X}_{\rho,\eta}^\kappa$.

5. Condition (I) for C*-textile dynamical systems

The notion of condition (I) for finite square matrices with entries in $\{0, 1\}$ has been introduced in [8]. The condition has been generalized by many authors to corresponding conditions for generalizations of the Cuntz–Krieger algebras (cf. [12], [15], [20], [41], etc.). The condition (I) for C*-symbolic dynamical systems (including λ -graph systems) has been also defined in [29] (cf. [25], [26]). All of these conditions give rise to the uniqueness of the associated C*-algebras subject to some operator relations among certain generating elements.

In this section, we will introduce the notion of condition (I) for C*-textile dynamical systems to prove the uniqueness of the C*-algebras $\mathcal{O}_{\rho,\eta}^\kappa$ under the relation $(\rho, \eta; \kappa)$.

Let $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ be a C*-symbolic dynamical system over Σ and $X_{\rho,\eta}^\kappa$ the associated two-dimensional subshift. Denote by $\Lambda_\rho, \Lambda_\eta$ the associated subshifts to the C*-symbolic dynamical systems $(\mathcal{A}, \rho, \Sigma^\rho), (\mathcal{A}, \eta, \Sigma^\eta)$ respectively. For $\mu = (\mu_1, \dots, \mu_j) \in B_j(\Lambda_\rho), \zeta = (\zeta_1, \dots, \zeta_k) \in B_k(\Lambda_\eta)$, we put $\rho_\mu = \rho_{\mu_j} \circ \dots \circ \rho_{\mu_1}, \eta_\zeta = \eta_{\zeta_k} \circ \dots \circ \eta_{\zeta_1}$ respectively. Recall that $|\mu|, |\zeta|$ denotes the lengths j, k respectively. In the algebra $\mathcal{O}_{\rho,\eta}^\kappa$, we set the subalgebras

$$\begin{aligned} \mathcal{F}_{\rho,\eta} &= C^*(S_\mu T_\zeta x T_\xi^* S_\nu^* : \mu, \nu \in B_*(\Lambda_\rho), \zeta, \xi \in B_*(\Lambda_\eta), |\mu| = |\nu|, |\zeta| = |\xi|, x \in \mathcal{A}) \end{aligned}$$

and for $j, k \in \mathbb{Z}_+$,

$$\mathcal{F}_{j,k} = C^*(S_\mu T_\zeta x T_\xi^* S_\nu^* : \mu, \nu \in B_j(\Lambda_\rho), \zeta, \xi \in B_k(\Lambda_\eta), x \in \mathcal{A}).$$

We notice that

$$\mathcal{F}_{j,k} = C^*(T_\zeta S_\mu x S_\nu^* T_\xi^* : \mu, \nu \in B_j(\Lambda_\rho), \zeta, \xi \in B_k(\Lambda_\eta), x \in \mathcal{A}).$$

The identities

$$(5.1) \quad S_\mu T_\zeta x T_\xi^* S_\nu^* = \sum_{a \in \Sigma^\eta} S_\mu T_{\zeta a} \eta_a(x) T_{\xi a}^* S_\nu^*,$$

$$(5.2) \quad T_\zeta S_\mu x S_\nu^* T_\xi^* = \sum_{\alpha \in \Sigma^\rho} T_\zeta S_{\mu \alpha} \rho_\alpha(x) S_{\nu \alpha}^* T_\xi^*$$

for $x \in \mathcal{A}$ and $\mu, \nu \in B_j(\Lambda_\rho), \zeta, \xi \in B_k(\Lambda_\eta)$ yield the embeddings

$$(5.3) \quad \iota_{*,+1} : \mathcal{F}_{j,k} \hookrightarrow \mathcal{F}_{j,k+1}, \quad \iota_{+1,*} : \mathcal{F}_{j,k} \hookrightarrow \mathcal{F}_{j+1,k}$$

respectively, such that $\cup_{j,k \in \mathbb{Z}_+} \mathcal{F}_{j,k}$ is dense in $\mathcal{F}_{\rho,\eta}$.

By the universality of $\mathcal{O}_{\rho,\eta}^\kappa$ subject to the relations $(\rho, \eta; \kappa)$, we may define an action $\theta : \mathbb{T}^2 \rightarrow \text{Aut}(\mathcal{O}_{\rho,\eta}^\kappa)$ of the two-dimensional torus group

$$\mathbb{T}^2 = \{(z, w) \in \mathbb{C}^2 \mid |z| = |w| = 1\}$$

to $\mathcal{O}_{\rho,\eta}^\kappa$ by setting

$$\theta_{z,w}(S_\alpha) = zS_\alpha, \quad \theta_{z,w}(T_a) = wT_a, \quad \theta_{z,w}(x) = x$$

for $\alpha \in \Sigma^\rho, a \in \Sigma^\eta, x \in \mathcal{A}$ and $z, w \in \mathbb{T}$. We call the action $\theta : \mathbb{T}^2 \rightarrow \text{Aut}(\mathcal{O}_{\rho,\eta}^\kappa)$ the gauge action of \mathbb{T}^2 on $\mathcal{O}_{\rho,\eta}^\kappa$. The fixed point algebra of $\mathcal{O}_{\rho,\eta}^\kappa$ under θ is denoted by $(\mathcal{O}_{\rho,\eta}^\kappa)^\theta$. Let $\mathcal{E}_{\rho,\eta} : \mathcal{O}_{\rho,\eta}^\kappa \rightarrow (\mathcal{O}_{\rho,\eta}^\kappa)^\theta$ be the conditional expectation defined by

$$\mathcal{E}_{\rho,\eta}(X) = \int_{(z,w) \in \mathbb{T}^2} \theta_{z,w}(X) dzdw, \quad X \in \mathcal{O}_{\rho,\eta}^\kappa$$

where $dzdw$ means the normalized Haar measure on \mathbb{T}^2 . The following lemma is routine.

Lemma 5.1. $(\mathcal{O}_{\rho,\eta}^\kappa)^\theta = \mathcal{F}_{\rho,\eta}$.

Define homomorphisms $\phi_\rho, \phi_\eta : \mathcal{D}_{\rho,\eta} \rightarrow \mathcal{D}_{\rho,\eta}$ by setting

$$\phi_\rho(X) = \sum_{\alpha \in \Sigma^\rho} S_\alpha X S_\alpha^*, \quad \phi_\eta(X) = \sum_{a \in \Sigma^\eta} T_a X T_a^*, \quad X \in \mathcal{D}_{\rho,\eta}.$$

It is easy to see that by (4.3)

$$\phi_\rho \circ \phi_\eta = \phi_\eta \circ \phi_\rho \quad \text{on } \mathcal{D}_{\rho,\eta}.$$

Definition 5.2. A C^* -textile dynamical system $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ is said to satisfy *condition (I)* if there exists a unital increasing sequence

$$\mathcal{A}_0 \subset \mathcal{A}_1 \subset \dots \subset \mathcal{A}$$

of C^* -subalgebras of \mathcal{A} such that:

- (1) $\rho_\alpha(\mathcal{A}_l) \subset \mathcal{A}_{l+1}, \eta_a(\mathcal{A}_l) \subset \mathcal{A}_{l+1}$ for all $l \in \mathbb{Z}_+, \alpha \in \Sigma^\rho, a \in \Sigma^\eta$.
- (2) $\cup_{l \in \mathbb{Z}_+} \mathcal{A}_l$ is dense in \mathcal{A} .
- (3) For $\epsilon > 0, j, k, l \in \mathbb{N}$ with $j + k \leq l$ and

$$X_0 \in \mathcal{F}_{j,k}^l = C^*(S_\mu T_\zeta x T_\xi^* S_\nu^* : \mu, \nu \in B_j(\Lambda_\rho), \zeta, \xi \in B_k(\Lambda_\eta), x \in \mathcal{A}_l),$$

there exists an element

$$g \in \mathcal{D}_{\rho,\eta} \cap \mathcal{A}_l' (= \{y \in \mathcal{D}_{\rho,\eta} \mid ya = ay \text{ for } a \in \mathcal{A}_l\})$$

with $0 \leq g \leq 1$ such that:

- (i) $\|X_0 \phi_\rho^j \circ \phi_\eta^k(g)\| \geq \|X_0\| - \epsilon,$
- (ii) $\phi_\rho^n(g) \phi_\eta^m(g) = \phi_\rho^n(\phi_\eta^m(g))g = \phi_\rho^n(g)g = \phi_\eta^m(g)g = 0$ for all $n = 1, 2, \dots, j, m = 1, 2, \dots, k.$

If in particular, one may take the above subalgebras $\mathcal{A}_l \subset \mathcal{A}, l = 0, 1, 2, \dots$ to be of finite dimensional, then $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ is said to satisfy *AF-condition (I)*. In this case, $\mathcal{A} = \overline{\cup_{l=0}^\infty \mathcal{A}_l}$ is an AF-algebra.

As the element g above belongs to the diagonal subalgebra $\mathcal{D}_{\rho,\eta}$ of $\mathcal{F}_{\rho,\eta}$, the condition (I) of $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ is intrinsically determined by itself by virtue of Lemma 5.5 below.

We will also introduce the following condition called *free*, which will be stronger than condition (I) but easier to confirm than condition (I).

Definition 5.3. A C^* -textile dynamical system $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ is said to be *free* if there exists a unital increasing sequence $\mathcal{A}_0 \subset \mathcal{A}_1 \subset \dots \subset \mathcal{A}$ of C^* -subalgebras of \mathcal{A} such that:

- (1) $\rho_\alpha(\mathcal{A}_l) \subset \mathcal{A}_{l+1}, \eta_a(\mathcal{A}_l) \subset \mathcal{A}_{l+1}$ for all $l \in \mathbb{Z}_+, \alpha \in \Sigma^\rho, a \in \Sigma^\eta$.
- (2) $\cup_{l \in \mathbb{Z}_+} \mathcal{A}_l$ is dense in \mathcal{A} .
- (3) For $j, k, l \in \mathbb{N}$ with $j + k \leq l$ there exists a projection $q \in \mathcal{D}_{\rho,\eta} \cap \mathcal{A}_l'$ such that:
 - (i) $qa \neq 0$ for $0 \neq a \in \mathcal{A}_l$.
 - (ii) $\phi_\rho^n(q)\phi_\eta^m(q) = \phi_\rho^n(\phi_\eta^m(q))q = \phi_\rho^n(q)q = \phi_\eta^m(q)q = 0$ for all $n = 1, 2, \dots, j, m = 1, 2, \dots, k$.

If in particular, one may take the above subalgebras $\mathcal{A}_l \subset \mathcal{A}, l = 0, 1, 2, \dots$ to be of finite dimensional, then $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ is said to be *AF-free*.

Proposition 5.4. *If a C^* -textile dynamical system $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ is free (resp. AF-free), then it satisfies condition (I) (resp. AF-condition (I)).*

Proof. Assume that $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ is free. Take an increasing sequence $\mathcal{A}_l, l \in \mathbb{N}$ of C^* -subalgebras of \mathcal{A} satisfying the above conditions (1), (2), (3) of freeness. For $j, k, l \in \mathbb{N}$ with $j + k \leq l$ there exists a projection $q \in \mathcal{D}_{\rho,\eta} \cap \mathcal{A}_l'$ satisfying the above two conditions (3i) and (3ii). Put

$$Q_{j,k}^l = \phi_\rho^j(\phi_\eta^k(q)).$$

For $x \in \mathcal{A}_l, \mu, \nu \in B_j(\Lambda_\rho), \xi, \zeta \in B_k(\Lambda_\eta)$, one has the equality

$$Q_{j,k}^l S_\mu T_\zeta x T_\xi^* S_\nu^* = S_\mu T_\zeta x T_\xi^* S_\nu^*$$

so that $Q_{j,k}^l$ commutes with all of elements of $\mathcal{F}_{j,k}^l$. By using the condition (3i) for q one directly sees that $S_\mu T_\zeta x T_\xi^* S_\nu^* \neq 0$ if and only if

$$Q_{j,k}^l S_\mu T_\zeta x T_\xi^* S_\nu^* \neq 0.$$

Hence the map

$$X \in \mathcal{F}_{j,k}^l \longrightarrow XQ_{j,k}^l \in \mathcal{F}_{j,k}^l Q_{j,k}^l$$

defines a homomorphism, that is proved to be injective by a similar proof to the proof of [30, Proposition 3.7]. Hence we have $\|XQ_{j,k}^l\| = \|X\| \geq \|X\| - \epsilon$ for all $X \in \mathcal{F}_{j,k}^l$. □

Let \mathcal{B} be a unital C^* -algebra. Suppose that there exist an injective $*$ -homomorphism $\pi : \mathcal{A} \longrightarrow \mathcal{B}$ preserving their units and two families

$$s_\alpha \in \mathcal{B}, \alpha \in \Sigma^\rho \quad \text{and} \quad t_a \in \mathcal{B}, a \in \Sigma^\eta$$

of partial isometries satisfying

$$\begin{aligned} \sum_{\beta \in \Sigma^\rho} s_\beta s_\beta^* &= 1, & \pi(x) s_\alpha s_\alpha^* &= s_\alpha s_\alpha^* \pi(x), & s_\alpha^* \pi(x) s_\alpha &= \pi(\rho_\alpha(x)), \\ \sum_{b \in \Sigma^\eta} t_b t_b^* &= 1, & \pi(x) t_a t_a^* &= t_a t_a^* \pi(x), & t_a^* \pi(x) t_a &= \pi(\eta_a(x)), \\ s_\alpha t_b &= t_a s_\beta & \text{if } \kappa(\alpha, b) &= (a, \beta) \end{aligned}$$

for all $x \in \mathcal{A}$ and $\alpha \in \Sigma^\rho$, $a \in \Sigma^\eta$. Put $\tilde{\mathcal{A}} = \pi(\mathcal{A})$ and

$$\tilde{\rho}_\alpha(\pi(x)) = \pi(\rho_\alpha(x)), \quad \tilde{\eta}_a(\pi(x)) = \pi(\eta_a(x)), \quad x \in \mathcal{A}.$$

It is easy to see that $(\tilde{\mathcal{A}}, \tilde{\rho}, \tilde{\eta}, \Sigma^\rho, \Sigma^\eta, \kappa)$ is a C^* -textile dynamical system such that the presented textile dynamical system $X_{\tilde{\rho}, \tilde{\eta}}^\kappa$ is the same as the one $X_{\rho, \eta}^\kappa$ presented by $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$. Let $\mathcal{O}_{\pi, s, t}$ be the C^* -subalgebra of \mathcal{B} generated by $\pi(x)$ and s_α, t_a for $x \in \mathcal{A}, \alpha \in \Sigma^\rho, a \in \Sigma^\eta$. Let $\mathcal{F}_{\pi, s, t}$ be the C^* -subalgebra of $\mathcal{O}_{\pi, s, t}$ generated by $s_\mu t_\zeta \pi(x) t_\xi^* s_\nu^*$ for $x \in \mathcal{A}$ and $\mu, \nu \in B_*(\Lambda_\rho), \zeta, \xi \in B_*(\Lambda_\eta)$ with $|\mu| = |\nu|, |\zeta| = |\xi|$. By the universality of the algebra $\mathcal{O}_{\rho, \eta}^\kappa$, the correspondence

$$x \in \mathcal{A} \longrightarrow \pi(x) \in \tilde{\mathcal{A}}, \quad S_\alpha \longrightarrow s_\alpha, \quad \alpha \in \Sigma^\rho, \quad T_a \longrightarrow t_a, \quad a \in \Sigma^\eta$$

extends to a surjective $*$ -homomorphism $\tilde{\pi} : \mathcal{O}_{\rho, \eta}^\kappa \longrightarrow \mathcal{O}_{\pi, s, t}$.

Lemma 5.5. *The restriction of $\tilde{\pi}$ to the subalgebra $\mathcal{F}_{\rho, \eta}$ is a $*$ -isomorphism from $\mathcal{F}_{\rho, \eta}$ to $\mathcal{F}_{\pi, s, t}$. Hence if $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ satisfies condition (I) (resp. is free), $(\tilde{\mathcal{A}}, \tilde{\rho}, \tilde{\eta}, \Sigma^\rho, \Sigma^\eta, \kappa)$ satisfies condition (I) (resp. is free).*

Proof. It suffices to show that $\tilde{\pi}$ is injective on $\mathcal{F}_{j, k}$ for all $j, k \in \mathbb{Z}$. Suppose

$$\sum_{\mu, \nu \in B_j(\Lambda_\rho), \zeta, \xi \in B_k(\Lambda_\eta)} s_\mu t_\zeta \pi(x_{\mu, \zeta, \xi, \nu}) t_\xi^* s_\nu^* = 0$$

with $x_{\mu, \zeta, \xi, \nu} \in \mathcal{A}$. For $\mu', \nu' \in B_j(\Lambda_\rho), \zeta', \xi' \in B_k(\Lambda_\eta)$, one has

$$\begin{aligned} &\pi(\eta_{\zeta'}(\rho_{\mu'}(1)) x_{\mu', \zeta', \xi', \nu'} \eta_{\xi'}(\rho_{\nu'}(1))) \\ &= t_{\zeta'}^* s_{\mu'}^* \left(\sum_{\mu, \nu \in B_j(\Lambda_\rho), \zeta, \xi \in B_k(\Lambda_\eta)} s_\mu t_\zeta \pi(x_{\mu, \zeta, \xi, \nu}) t_\xi^* s_\nu^* \right) s_{\nu'} t_{\xi'} = 0. \end{aligned}$$

As $\pi : \mathcal{A} \longrightarrow \mathcal{B}$ is injective, one sees

$$\eta_{\zeta'}(\rho_{\mu'}(1)) x_{\mu', \zeta', \xi', \nu'} \eta_{\xi'}(\rho_{\nu'}(1)) = 0$$

so that

$$S_{\mu'} T_{\zeta'} x_{\mu', \zeta', \xi', \nu'} T_{\xi'}^* S_{\nu'}^* = 0.$$

Hence we have

$$\sum_{\mu, \nu \in B_j(\Lambda_\rho), \zeta, \xi \in B_k(\Lambda_\eta)} S_\mu T_\zeta x_{\mu, \zeta, \xi, \nu} T_\xi^* S_\nu^* = 0.$$

Therefore $\tilde{\pi}$ is injective on $\mathcal{F}_{j, k}$. □

We henceforth assume that $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ satisfies condition (I) defined above. Take a unital increasing sequence $\{\mathcal{A}_l\}_{l \in \mathbb{Z}_+}$ of C^* -subalgebras of \mathcal{A} as in the definition of condition (I). Recall that the algebra $\mathcal{F}_{j,k}^l$ for $j, k \leq l$ is defined by

$$\mathcal{F}_{j,k}^l = C^*(S_\mu T_\zeta x T_\xi^* S_\nu^* : \mu, \nu \in B_j(\Lambda_\rho), \zeta, \xi \in B_k(\Lambda_\eta), x \in \mathcal{A}_l).$$

There exists an inclusion relation $\mathcal{F}_{j,k}^l \subset \mathcal{F}_{j',k'}^{l'}$ for $j \leq j', k \leq k'$ and $l \leq l'$ through the identities (5.1), (5.2). Let $\mathcal{P}_{\pi,s,t}$ be the $*$ -subalgebra of $\mathcal{O}_{\pi,s,t}$ algebraically generated by $\pi(x), s_\alpha, t_a$ for $x \in \mathcal{A}_l, l \in \mathbb{Z}_+, \alpha \in \Sigma^\rho, a \in \Sigma^\eta$.

Lemma 5.6. *Any element $x \in \mathcal{P}_{\pi,s,t}$ can be expressed in a unique way as*

$$\begin{aligned} x = & \sum_{|\nu|, |\xi| \geq 1} x_{-\xi, -\nu} t_\xi^* s_\nu^* + \sum_{|\zeta|, |\nu| \geq 1} t_\zeta x_{\zeta, -\nu} s_\nu^* + \sum_{|\mu|, |\xi| \geq 1} s_\mu x_{\mu, -\xi} t_\xi^* \\ & + \sum_{|\mu|, |\zeta| \geq 1} s_\mu t_\zeta x_{\mu, \zeta} + \sum_{|\xi| \geq 1} x_{-\xi} t_\xi^* + \sum_{|\nu| \geq 1} x_{-\nu} s_\nu^* \\ & + \sum_{|\mu| \geq 1} s_\mu x_\mu + \sum_{|\zeta| \geq 1} t_\zeta x_\zeta + x_0 \end{aligned}$$

where the above summations Σ are all finite sums and the elements

$$x_{-\xi, -\nu}, x_{\zeta, -\nu}, x_{\mu, -\xi}, x_{\mu, \zeta}, x_{-\xi}, x_{-\nu}, x_\mu, x_\zeta, x_0$$

for $\mu, \nu \in B_*(\Lambda_\rho), \zeta, \xi \in B_*(\Lambda_\eta)$ all belong to the dense subalgebra

$$\mathcal{P}_{\pi,s,t} \cap \mathcal{F}_{\pi,s,t}$$

which satisfy

$$\begin{aligned} x_{-\xi, -\nu} &= x_{-\xi, -\nu} \eta_\xi(\rho_\nu(1)), & x_{\zeta, -\nu} &= \eta_\zeta(1) x_{\zeta, -\nu} \rho_\nu(1), \\ x_{\mu, -\xi} &= \rho_\mu(1) x_{\mu, -\xi} \eta_\xi(1), & x_{\mu, \zeta} &= \eta_\zeta(\rho_\mu(1)) x_{\mu, \zeta}, \\ x_{-\xi} &= x_{-\xi} \eta_\xi(1), & x_{-\nu} &= x_{-\nu} \rho_\nu(1), \\ x_\mu &= \rho_\mu(1) x_\mu, & x_\zeta &= \eta_\zeta(1) x_\zeta. \end{aligned}$$

Proof. Put

$$\begin{aligned} x_{-\xi, -\nu} &= \mathcal{E}_{\rho, \eta}(x s_\nu t_\xi), & x_{\zeta, -\nu} &= \mathcal{E}_{\rho, \eta}(t_\zeta^* x s_\nu), \\ x_{\mu, -\xi} &= \mathcal{E}_{\rho, \eta}(s_\mu^* x t_\xi), & x_{\mu, \zeta} &= \mathcal{E}_{\rho, \eta}(t_\zeta^* s_\mu^* x), \\ x_{-\xi} &= \mathcal{E}_{\rho, \eta}(x t_\xi), & x_{-\nu} &= \mathcal{E}_{\rho, \eta}(x s_\nu), \\ x_\mu &= \mathcal{E}_{\rho, \eta}(s_\mu^* x), & x_\zeta &= \mathcal{E}_{\rho, \eta}(t_\zeta^* x), \\ x_0 &= \mathcal{E}_{\rho, \eta}(x). \end{aligned}$$

Then we have the desired expression of x . The elements

$$x_{-\xi, -\nu}, x_{\zeta, -\nu}, x_{\mu, -\xi}, x_{\mu, \zeta}, x_{-\xi}, x_{-\nu}, x_\mu, x_\zeta, x_0$$

for $\mu, \nu \in B_*(\Lambda_\rho), \zeta, \xi \in B_*(\Lambda_\eta)$ are automatically determined by the above formulae so that the expression is unique. \square

Lemma 5.7. For $h \in \mathcal{D}_{\rho,\eta} \cap \mathcal{A}'_l$ and $j, k \in \mathbb{Z}$ with $j + k \leq l$, put

$$h^{j,k} = \phi_\rho^j \circ \phi_\eta^k(h).$$

Then we have

- (i) $h^{j,k} s_\mu = s_\mu h^{j-|\mu|,k}$ for $\mu \in B_*(\Lambda_\rho)$ with $|\mu| \leq j$.
- (ii) $h^{j,k} t_\zeta = t_\zeta h^{j,k-|\zeta|}$ for $\zeta \in B_*(\Lambda_\eta)$ with $|\zeta| \leq k$.
- (iii) $h^{j,k}$ commutes with any element of $\mathcal{F}_{j,k}^l$.

Proof. (i) It follows that for $\mu \in B_*(\Lambda_\rho)$ with $|\mu| \leq j$

$$h^{j,k} s_\mu = \sum_{|\mu'|=|\mu|} s_{\mu'} \phi_\rho^{j-|\mu|}(\phi_\eta^k(h)) s_{\mu'}^* s_\mu = s_\mu \phi_\rho^{j-|\mu|}(\phi_\eta^k(h)) s_\mu^* s_\mu.$$

Since $h \in \mathcal{A}'_l$ and $\mathcal{A}_{j+k} \subset \mathcal{A}_l$, one has

$$\begin{aligned} \phi_\rho^{j-|\mu|}(\phi_\eta^k(h)) s_\mu^* s_\mu &= \sum_{\nu \in B_{j-|\mu|}(\Lambda_\rho)} \sum_{\xi \in B_k(\Lambda_\eta)} s_\nu t_\xi h t_\xi^* s_\nu^* s_\mu^* s_\mu \\ &= \sum_{\nu \in B_{j-|\mu|}(\Lambda_\rho)} \sum_{\xi \in B_k(\Lambda_\eta)} s_\nu t_\xi h t_\xi^* s_\nu^* s_\mu^* s_\nu s_\nu t_\xi t_\xi^* s_\nu^* \\ &= \sum_{\nu \in B_{j-|\mu|}(\Lambda_\rho)} \sum_{\xi \in B_k(\Lambda_\eta)} s_\nu t_\xi \eta_\xi(\rho_{\mu\nu}(1)) h t_\xi^* s_\nu^* \\ &= \sum_{\nu \in B_{j-|\mu|}(\Lambda_\rho)} \sum_{\xi \in B_k(\Lambda_\eta)} s_\nu \rho_{\mu\nu}(1) t_\xi h t_\xi^* s_\nu^* \\ &= s_\mu^* s_\mu \phi_\rho^{j-|\mu|}(\phi_\eta^k(h)) = s_\mu^* s_\mu h^{j-|\mu|,k} \end{aligned}$$

so that $h^{j,k} s_\mu = s_\mu h^{j-|\mu|,k}$.

- (ii) Similarly we have $h^{j,k} t_\zeta = t_\zeta h^{j,k-|\zeta|}$ for $\zeta \in B_*(\Lambda_\eta)$ with $|\zeta| \leq k$.
- (iii) For $x \in \mathcal{A}_l$, $\mu, \nu \in B_j(\Lambda_\rho)$, $\zeta, \xi \in B_k(\Lambda_\eta)$, we have

$$h^{j,k} s_\mu t_\zeta = s_\mu h^{0,k} t_\zeta = s_\mu t_\zeta h^{0,0} = s_\mu t_\zeta h.$$

It follows that

$$h^{j,k} s_\mu t_\zeta x t_\xi^* s_\nu^* = s_\mu t_\zeta h x t_\xi^* s_\nu^* = s_\mu t_\zeta x h t_\xi^* s_\nu^* = s_\mu t_\zeta x t_\xi^* s_\nu^* h^{j,k}.$$

Hence $h^{j,k}$ commutes with any element of $\mathcal{F}_{j,k}^l$. \square

Lemma 5.8. Assume that $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ satisfies condition (I). For $x \in \mathcal{P}_{\pi,s,t}$, let $x_0 = \mathcal{E}_{\rho,\eta}(x)$ as in Lemma 5.6. Then we have

$$\|x_0\| \leq \|x\|.$$

Proof. We may assume that the elements for $x \in \mathcal{P}_{\pi,s,t}$

$$x_{-\xi, -\nu}, x_{\zeta, -\nu}, x_{\mu, -\xi}, x_{\mu, \zeta}, x_{-\xi}, x_{-\nu}, x_\mu, x_\zeta, x_0$$

in Lemma 5.6 belong to $\tilde{\pi}(\mathcal{F}_{j_1, k_1}^{l_1})$ for some j_1, k_1, l_1 and $\mu, \nu \in \cup_{n=0}^{j_0} B_n(\Lambda_\rho)$, $\zeta, \xi \in \cup_{n=0}^{k_0} B_n(\Lambda_\eta)$ for some j_0, k_0 . Take $j, k, l \in \mathbb{Z}_+$ such as

$$j \geq j_0 + j_1, \quad k \geq k_0 + k_1, \quad l \geq \max\{j + k, l_1\}.$$

By Lemma 5.5, $(\tilde{\mathcal{A}}, \tilde{\rho}, \tilde{\eta}, \Sigma^\rho, \Sigma^\eta, \kappa)$ satisfies condition (I). For any $\epsilon > 0$, the numbers j, k, l , and the element $x_0 \in \tilde{\pi}(\mathcal{F}_{j_1, k_1}^{l_1})$, one may find

$$g \in \tilde{\pi}(\mathcal{D}_{\rho, \eta}) \cap \pi(\mathcal{A}_l)'$$

with $0 \leq g \leq 1$ such that:

- (i) $\|x_0 \phi_\rho^j \circ \phi_\eta^k(g)\| \geq \|x_0\| - \epsilon$.
- (ii) $\phi_\rho^n(g) \phi_\eta^m(g) = \phi_\rho^n(\phi_\eta^m(g))g = \phi_\rho^n(g)g = \phi_\eta^m(g)g = 0$ for all $n = 1, 2, \dots, j, m = 1, 2, \dots, k$.

Put $h = g^{\frac{1}{2}}$ and $h^{j,k} = \phi_\rho^j \circ \phi_\eta^k(h)$. It follows that $\|x\| \geq \|h^{j,k} x h^{j,k}\|$ and

$$\|h^{j,k} x h^{j,k}\| = \|(1) + (2) + (3) + (4) + (5) + (6)\|$$

where the summands are given by

- (1) $\sum_{|\nu|, |\xi| \geq 1} h^{j,k} x_{-\xi, -\nu} t_\xi^* s_\nu^* h^{j,k}$
- (2) $\sum_{|\zeta|, |\nu| \geq 1} h^{j,k} t_\zeta x_{\zeta, -\nu} s_\nu^* h^{j,k}$
- (3) $\sum_{|\mu|, |\xi| \geq 1} h^{j,k} s_\mu x_{\mu, -\xi} t_\xi^* h^{j,k}$
- (4) $\sum_{|\mu|, |\zeta| \geq 1} h^{j,k} s_\mu t_\zeta x_{\mu, \zeta} h^{j,k}$
- (5) $\sum_{|\xi| \geq 1} h^{j,k} x_{-\xi} t_\xi^* h^{j,k} + \sum_{|\nu| \geq 1} h^{j,k} x_{-\nu} s_\nu^* h^{j,k} + \sum_{|\mu| \geq 1} h^{j,k} s_\mu x_\mu h^{j,k} + \sum_{|\zeta| \geq 1} h^{j,k} t_\zeta x_\zeta h^{j,k}$
- (6) $h^{j,k} x_0 h^{j,k}$.

For (1), as $x_{-\xi, -\nu} \in \tilde{\pi}(\mathcal{F}_{j_1, k_1}^{l_1}) \subset \tilde{\pi}(\mathcal{F}_{j, k}^l)$, one sees that $x_{-\xi, -\nu}$ commutes with $h^{j,k}$. Hence we have

$$h^{j,k} x_{-\xi, -\nu} t_\xi^* s_\nu^* h^{j,k} = x_{-\xi, -\nu} h^{j,k} t_\xi^* s_\nu^* h^{j,k} = x_{-\xi, -\nu} h^{j,k} h^{j-|\nu|, k-|\xi|} t_\xi^* s_\nu^*$$

and

$$\begin{aligned} h^{j,k} h^{j-|\nu|, k-|\xi|} (h^{j,k} h^{j-|\nu|, k-|\xi|})^* &= \phi_\rho^j(\phi_\eta^k(g)) \cdot \phi_\rho^{j-|\nu|}(\phi_\eta^{k-|\xi|}(g)) \\ &= \phi_\rho^{j-|\nu|} \circ \phi_\eta^{k-|\xi|}(\phi_\eta^{|\xi|}(\phi_\rho^{|\nu|}(g))) = 0 \end{aligned}$$

so that

$$h^{j,k} x_{-\xi, -\nu} t_\xi^* s_\nu^* h^{j,k} = 0.$$

For (2), as $x_{\xi, -\nu} \in \tilde{\pi}(\mathcal{F}_{j_1, k_1}^{l_1}) \subset \tilde{\pi}(\mathcal{F}_{j, k-|\xi|}^l)$, one sees that $x_{\xi, -\nu}$ commutes with $h^{j, k-|\xi|}$. Hence we have

$$h^{j,k} t_\xi x_{\xi, -\nu} s_\nu^* h^{j,k} = t_\xi h^{j, k-|\xi|} x_{\xi, -\nu} h^{j-|\nu|, k} s_\nu^* = t_\xi x_{\xi, -\nu} h^{j, k-|\xi|} h^{j-|\nu|, k} s_\nu^*$$

and

$$\begin{aligned} h^{j,k-|\xi|} h^{j-|\nu|,k} (h^{j,k-|\xi|} h^{j-|\nu|,k})^* &= \phi_\rho^j(\phi_\eta^{k-|\zeta|}(g)) \cdot \phi_\rho^{j-|\nu|}(\phi_\eta^k(g)) \\ &= \phi_\rho^{j-|\nu|} \circ \phi_\eta^{k-|\zeta|}(\phi_\rho^{|\nu|}(g) \phi_\eta^{|\zeta|}(g)) = 0 \end{aligned}$$

so that

$$h^{j,k} t_\xi x_{\xi, -\nu} s_\nu^* h^{j,k} = 0.$$

For (3), as $x_{\mu, -\xi} \in \tilde{\pi}(\mathcal{F}_{j_1, k_1}^{l_1}) \subset \tilde{\pi}(\mathcal{F}_{j-|\mu|, k}^l)$, one sees that $x_{\mu, -\xi}$ commutes with $h^{j-|\mu|, k}$. Hence we have

$$h^{j,k} s_\mu x_{\mu, -\xi} t_\xi^* h^{j,k} = s_\mu h^{j-|\mu|, k} x_{\mu, -\xi} h^{j, k-|\xi|} t_\xi^* = s_\mu x_{\mu, -\xi} h^{j-|\mu|, k} h^{j, k-|\xi|} t_\xi^*$$

and

$$\begin{aligned} h^{j-|\mu|, k} h^{j, k-|\xi|} (h^{j-|\mu|, k} h^{j, k-|\xi|})^* &= \phi_\rho^{j-|\mu|}(\phi_\eta^k(g)) \cdot \phi_\rho^j(\phi_\eta^{k-|\xi|}(g)) \\ &= \phi_\rho^{j-|\mu|} \circ \phi_\eta^{k-|\xi|}(\phi_\eta^{|\xi|}(g) \phi_\rho^{|\mu|}(g)) = 0 \end{aligned}$$

so that

$$h^{j,k} s_\mu x_{\mu, -\xi} t_\xi^* h^{j,k} = 0.$$

For (4), as $x_{\mu, \zeta} \in \tilde{\pi}(\mathcal{F}_{j_1, k_1}^{l_1}) \subset \tilde{\pi}(\mathcal{F}_{j-|\mu|, k-|\zeta|}^l)$, one sees that $x_{\mu, \zeta}$ commutes with $h^{j-|\mu|, k-|\zeta|}$. Hence we have

$$h^{j,k} s_\mu t_\zeta x_{\mu, \zeta} h^{j,k} = s_\mu t_\zeta h^{j-|\mu|, k-|\zeta|} x_{\mu, \zeta} h^{j,k} = s_\mu t_\zeta x_{\mu, \zeta} h^{j-|\mu|, k-|\zeta|} h^{j,k}$$

and

$$\begin{aligned} h^{j-|\mu|, k-|\zeta|} h^{j,k} (h^{j-|\mu|, k-|\zeta|} h^{j,k})^* &= \phi_\rho^{j-|\mu|}(\phi_\eta^{k-|\zeta|}(g)) \cdot \phi_\rho^j(\phi_\eta^k(g)) \\ &= \phi_\rho^{j-|\mu|} \circ \phi_\eta^{k-|\zeta|}(g \phi_\rho^{|\mu|}(\phi_\eta^{|\zeta|}(g))) = 0 \end{aligned}$$

so that

$$h^{j,k} s_\mu t_\zeta x_{\mu, \zeta} h^{j,k} = 0.$$

For (5), as $x_{-\xi}$ commutes with $h^{j,k}$, we have

$$h^{j,k} x_{-\xi} t_\xi^* h^{j,k} = x_{-\xi} h^{j,k} h^{j, k-|\xi|} t_\xi^*$$

and

$$\begin{aligned} h^{j,k} h^{j, k-|\xi|} (h^{j,k} h^{j, k-|\xi|})^* &= \phi_\rho^j(\phi_\eta^k(g)) \cdot \phi_\rho^j(\phi_\eta^{k-|\xi|}(g)) \\ &= \phi_\rho^j \circ \phi_\eta^{k-|\xi|}(\phi_\eta^{|\xi|}(g) g) = 0 \end{aligned}$$

so that

$$h^{j,k} x_{-\xi} t_\xi^* h^{j,k} = 0.$$

We similarly see that

$$h^{j,k} x_{-\nu} s_\nu^* h^{j,k} = h^{j,k} s_\mu x_\mu h^{j,k} = h^{j,k} t_\zeta x_\zeta h^{j,k} = 0.$$

Therefore we have

$$\|x\| \geq \|h^{j,k} x_0 h^{j,k}\| = \|x_0 (h^{j,k})^2\| = \|x_0 \phi_\rho^j \circ \phi_\eta^k(g)\| \geq \|x_0\| - \epsilon. \quad \square$$

By a similar argument to [8, 2.8 Proposition], one sees:

Corollary 5.9. *Assume $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ satisfies condition (I). There exists a conditional expectation $\mathcal{E}_{\pi,s,t} : \mathcal{O}_{\pi,s,t} \rightarrow \mathcal{F}_{\pi,s,t}$ such that*

$$\mathcal{E}_{\pi,s,t} \circ \tilde{\pi} = \tilde{\pi} \circ \mathcal{E}_{\rho,\eta}.$$

Therefore we have

Proposition 5.10. *Assume that $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ satisfies condition (I). The *-homomorphism $\tilde{\pi} : \mathcal{O}_{\rho,\eta}^\kappa \rightarrow \mathcal{O}_{\pi,s,t}$ defined by*

$$\tilde{\pi}(x) = \pi(x), \quad x \in \mathcal{A}, \quad \tilde{\pi}(S_\alpha) = s_\alpha, \quad \alpha \in \Sigma^\rho, \quad \tilde{\pi}(T_a) = t_a, \quad a \in \Sigma^\eta$$

*becomes a surjective *-isomorphism, and hence the C*-algebras $\mathcal{O}_{\rho,\eta}^\kappa$ and $\mathcal{O}_{\pi,s,t}$ are canonically *-isomorphic through $\tilde{\pi}$.*

Proof. The map $\tilde{\pi} : \mathcal{F}_{\rho,\eta} \rightarrow \mathcal{F}_{\pi,s,t}$ is *-isomorphic and satisfies $\mathcal{E}_{\pi,s,t} \circ \tilde{\pi} = \tilde{\pi} \circ \mathcal{E}_{\rho,\eta}$. Since $\mathcal{E}_{\rho,\eta} : \mathcal{O}_{\rho,\eta}^\kappa \rightarrow \mathcal{F}_{\rho,\eta}$ is faithful, a routine argument shows that the *-homomorphism $\tilde{\pi} : \mathcal{O}_{\rho,\eta}^\kappa \rightarrow \mathcal{O}_{\pi,s,t}$ is actually a *-isomorphism. \square

Hence the following uniqueness of the C*-algebra $\mathcal{O}_{\rho,\eta}^\kappa$ holds.

Theorem 5.11. *Assume that $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ satisfies condition (I). The C*-algebra $\mathcal{O}_{\rho,\eta}^\kappa$ is the unique C*-algebra subject to the relation $(\rho, \eta; \kappa)$. This means that if there exist a unital C*-algebra \mathcal{B} , an injective *-homomorphism $\pi : \mathcal{A} \rightarrow \mathcal{B}$ and two families of partial isometries $s_\alpha, \alpha \in \Sigma^\rho, t_a, a \in \Sigma^\eta$ satisfying the following relations :*

$$\begin{aligned} \sum_{\beta \in \Sigma^\rho} s_\beta s_\beta^* &= 1, & \pi(x) s_\alpha s_\alpha^* &= s_\alpha s_\alpha^* \pi(x), & s_\alpha^* \pi(x) s_\alpha &= \pi(\rho_\alpha(x)), \\ \sum_{b \in \Sigma^\eta} t_b t_b^* &= 1, & \pi(x) t_a t_a^* &= t_a t_a^* \pi(x), & t_a^* \pi(x) t_a &= \pi(\eta_a(x)) \\ s_\alpha t_b &= t_a s_\beta & \text{if } \kappa(\alpha, b) &= (a, \beta) \end{aligned}$$

for $(\alpha, b) \in \Sigma^{\rho\eta}, (a, \beta) \in \Sigma^{\eta\rho}$ and $x \in \mathcal{A}, \alpha \in \Sigma^\rho, a \in \Sigma^\eta$, then the correspondence

$$x \in \mathcal{A} \rightarrow \pi(x) \in \mathcal{B}, \quad S_\alpha \rightarrow s_\alpha \in \mathcal{B}, \quad T_a \rightarrow t_a \in \mathcal{B}$$

*extends to a *-isomorphism $\tilde{\pi}$ from $\mathcal{O}_{\rho,\eta}^\kappa$ onto the C*-subalgebra $\mathcal{O}_{\pi,s,t}$ of \mathcal{B} generated by $\pi(x), x \in \mathcal{A}$ and $s_\alpha, \alpha \in \Sigma, t_a, a \in \Sigma^\eta$.*

For a C*-textile dynamical system $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$, let $\lambda_{\rho,\eta} : \mathcal{A} \rightarrow \mathcal{A}$ be the positive map on \mathcal{A} defined by

$$\lambda_{\rho,\eta}(x) = \sum_{\alpha \in \Sigma^\rho, a \in \Sigma^\eta} \eta_a \circ \rho_\alpha(x), \quad x \in \mathcal{A}.$$

Then $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ is said to be *irreducible* if there exists no nontrivial ideal of \mathcal{A} invariant under $\lambda_{\rho,\eta}$.

Corollary 5.12. *If $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ satisfies condition (I) and is irreducible, the C*-algebra $\mathcal{O}_{\rho,\eta}^\kappa$ is simple.*

Proof. Assume that there exists a nontrivial ideal \mathcal{I} of $\mathcal{O}_{\rho,\eta}^\kappa$. Now suppose that $\mathcal{I} \cap \mathcal{A} = \{0\}$. As $S_\alpha^* S_\alpha = \rho_\alpha(1), T_a^* T_a = \eta_a(1) \in \mathcal{A}$, one knows that $S_\alpha, T_a \notin \mathcal{I}$ for all $\alpha \in \Sigma^\rho, a \in \Sigma^\eta$. By the above theorem, the quotient map $q : \mathcal{O}_{\rho,\eta}^\kappa \rightarrow \mathcal{O}_{\rho,\eta}^\kappa/\mathcal{I}$ must be injective so that \mathcal{I} is trivial. Hence one sees that $\mathcal{I} \cap \mathcal{A} \neq \{0\}$ and it is invariant under $\lambda_{\rho,\eta}$. \square

6. Concrete realization

In this section we will realize the C^* -algebra $\mathcal{O}_{\rho,\eta}^\kappa$ for $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ in a concrete way as a C^* -algebra constructed from a Hilbert C^* -bimodule. For $\gamma_i \in \Sigma^\rho \cup \Sigma^\eta$, put

$$\xi_{\gamma_i} = \begin{cases} \rho_{\gamma_i} & \text{if } \gamma_i \in \Sigma^\rho, \\ \eta_{\gamma_i} & \text{if } \gamma_i \in \Sigma^\eta. \end{cases}$$

A finite sequence of labels $(\gamma_1, \gamma_2, \dots, \gamma_k) \in (\Sigma^\rho \cup \Sigma^\eta)^k$ is said to be *concatenated labeled path* if $\xi_{\gamma_k} \circ \dots \circ \xi_{\gamma_2} \circ \xi_{\gamma_1}(1) \neq 0$. For $m, n \in \mathbb{Z}_+$, let $L_{(n,m)}$ be the set of concatenated labeled paths $(\gamma_1, \gamma_2, \dots, \gamma_{m+n})$ such that symbols in Σ^ρ appear in $(\gamma_1, \gamma_2, \dots, \gamma_{m+n})$ n -times and symbols in Σ^η appear in $(\gamma_1, \gamma_2, \dots, \gamma_{m+n})$ m -times. We define a relation in $L_{(n,m)}$ for $i = 1, 2, \dots, n + m - 1$. We write

$$\begin{aligned} &(\gamma_1, \dots, \gamma_{i-1}, \gamma_i, \gamma_{i+1}, \gamma_{i+2}, \dots, \gamma_{m+n}) \\ &\approx_i (\gamma_1, \dots, \gamma_{i-1}, \gamma'_i, \gamma'_{i+1}, \gamma_{i+2}, \dots, \gamma_{m+n}) \end{aligned}$$

if one of the following two conditions holds:

- (1) $(\gamma_i, \gamma_{i+1}) \in \Sigma^{\rho\eta}, (\gamma'_i, \gamma'_{i+1}) \in \Sigma^{\eta\rho}$ and $\kappa(\gamma_i, \gamma_{i+1}) = (\gamma'_i, \gamma'_{i+1})$,
- (2) $(\gamma_i, \gamma_{i+1}) \in \Sigma^{\eta\rho}, (\gamma'_i, \gamma'_{i+1}) \in \Sigma^{\rho\eta}$ and $\kappa(\gamma'_i, \gamma'_{i+1}) = (\gamma_i, \gamma_{i+1})$.

Denote by \approx the equivalence relation in $L_{(n,m)}$ generated by the relations $\approx_i, i = 1, 2, \dots, n + m - 1$. Let $\mathfrak{T}_{(n,m)} = L_{(n,m)}/\approx$ be the set of equivalence classes of $L_{(n,m)}$ under \approx . Denote by $[\gamma] \in \mathfrak{T}_{(n,m)}$ the equivalence class of $\gamma \in L_{(n,m)}$. Put the vectors $e = (1, 0), f = (0, -1)$ in \mathbb{R}^2 . Consider the set of all paths consisting of sequences of vectors e, f starting at the point $(-n, m) \in \mathbb{R}^2$ for $n, m \in \mathbb{Z}_+$ and ending at the origin. Such a path consists of n e -vectors and m f -vectors. Let $\mathfrak{P}_{(n,m)}$ be the set of all such paths from $(-n, m)$ to the origin. We consider the correspondence

$$\rho_\alpha \longrightarrow e \quad (\alpha \in \Sigma^\rho), \quad \eta_a \longrightarrow f \quad (a \in \Sigma^\eta),$$

denoted by π . It extends a surjective map from $L_{(n,m)}$ to $\mathfrak{P}_{(n,m)}$ in a natural way. For a concatenated labeled path $\gamma = (\gamma_1, \gamma_2, \dots, \gamma_{n+m}) \in L_{(n,m)}$, put the projection in \mathcal{A}

$$P_\gamma = (\xi_{\gamma_{n+m}} \circ \dots \circ \xi_{\gamma_2} \circ \xi_{\gamma_1})(1).$$

We note that $P_\gamma \neq 0$ for all $\gamma \in L_{(n,m)}$.

Lemma 6.1. For $\gamma, \gamma' \in L_{(n,m)}$, if $\gamma \approx \gamma'$, we have $P_\gamma = P_{\gamma'}$. Hence the projection $P_{[\gamma]}$ for $[\gamma] \in \mathfrak{T}_{(n,m)}$ is well-defined.

Proof. If $\kappa(\alpha, b) = (a, \beta)$, one has $\eta_b \circ \rho_\alpha(1) = \rho_\beta \circ \eta_a(1) \neq 0$. Hence the assertion is obvious. \square

Denote by $|\mathfrak{T}_{(n,m)}|$ the cardinal number of the finite set $\mathfrak{T}_{(n,m)}$. Let $e_t, t \in \mathfrak{T}_{(n,m)}$ be the standard complete orthonormal basis of $\mathbb{C}^{|\mathfrak{T}_{(n,m)}|}$. Define

$$H_{(n,m)} = \sum_{t \in \mathfrak{T}_{(n,m)}} \oplus \mathbb{C}e_t \otimes P_t \mathcal{A} \left(= \sum_{t \in \mathfrak{T}_{(n,m)}} \oplus \text{Span}\{ce_t \otimes P_t x \mid c \in \mathbb{C}, x \in \mathcal{A}\} \right)$$

the direct sum of $\mathbb{C}e_t \otimes P_t \mathcal{A}$ over $t \in \mathfrak{T}_{(n,m)}$. $H_{(n,m)}$ has a structure of C^* -bimodule over \mathcal{A} by setting

$$(e_t \otimes P_t x)y := e_t \otimes P_t xy, \\ \phi(y)(e_t \otimes P_t x) := e_t \otimes \xi_\gamma(y)x (= e_t \otimes P_t \xi_\gamma(y)x) \quad \text{for } x, y \in \mathcal{A}$$

where $t = [\gamma]$ for $\gamma = (\gamma_1, \dots, \gamma_{n+m})$ and $\xi_\gamma(y) = (\xi_{\gamma_{n+m}} \circ \dots \circ \xi_{\gamma_2} \circ \xi_{\gamma_1})(y)$. Define an \mathcal{A} -valued inner product on $H_{(n,m)}$ by setting

$$\langle e_t \otimes P_t x \mid e_s \otimes P_s y \rangle := \begin{cases} x^* P_t y & \text{if } t = s, \\ 0 & \text{otherwise} \end{cases}$$

for $t, s \in \mathfrak{T}_{(n,m)}$ and $x, y \in \mathcal{A}$. Then $H_{(n,m)}$ becomes a Hilbert C^* -bimodule over \mathcal{A} . Put $H_{(0,0)} = \mathcal{A}$. Denote by F_κ the Hilbert C^* -bimodule over \mathcal{A} defined by the direct sum:

$$F_\kappa = \sum_{(n,m) \in \mathbb{Z}_+^2} \oplus H_{(n,m)}.$$

For $\alpha \in \Sigma^\rho, a \in \Sigma^\eta$, the creation operators s_α, t_a on F_κ :

$$s_\alpha : H_{(n,m)} \longrightarrow H_{(n+1,m)}, \quad t_a : H_{(n,m)} \longrightarrow H_{(n,m+1)}$$

are defined by

$$s_\alpha x = e_{[\alpha]} \otimes P_{[\alpha]} x, \quad \text{for } x \in H_{(0,0)} (= \mathcal{A}), \\ s_\alpha(e_{[\gamma]} \otimes P_{[\gamma]} x) = \begin{cases} e_{[\alpha\gamma]} \otimes P_{[\alpha\gamma]} x & \text{if } \alpha\gamma \in L_{(n+1,m)}, \\ 0 & \text{otherwise,} \end{cases} \\ t_a x = e_{[a]} \otimes P_{[a]} x, \quad \text{for } x \in H_{(0,0)} (= \mathcal{A}), \\ t_a(e_{[\gamma]} \otimes P_{[\gamma]} x) = \begin{cases} e_{[a\gamma]} \otimes P_{[a\gamma]} x & \text{if } a\gamma \in L_{(n,m+1)}, \\ 0 & \text{otherwise.} \end{cases}$$

For $y \in \mathcal{A}$ an operator $i_{F_\kappa}(y)$ on F_κ :

$$i_{F_\kappa}(y) : H_{(n,m)} \longrightarrow H_{(n,m)}$$

is defined by

$$i_{F_\kappa}(y)x = yx \quad \text{for } x \in H_{(0,0)} (= \mathcal{A}),$$

$$i_{F_\kappa}(y)(e_{[\gamma]} \otimes P_{[\gamma]}x) = \phi(y)(e_{[\gamma]} \otimes P_{[\gamma]}x) (= e_{[\gamma]} \otimes \xi_\gamma(y)x).$$

Define the Cuntz–Toeplitz C^* -algebra for $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ by

$$\mathcal{T}_{\rho,\eta}^\kappa = C^*(s_\alpha, t_a, i_{F_\kappa}(y) \mid \alpha \in \Sigma^\rho, a \in \Sigma^\eta, y \in \mathcal{A})$$

as the C^* -algebra on F_κ generated by $s_\alpha, t_a, i_{F_\kappa}(y)$ for $\alpha \in \Sigma^\rho, a \in \Sigma^\eta, y \in \mathcal{A}$.

Lemma 6.2. *For $\alpha \in \Sigma^\rho, a \in \Sigma^\eta$, we have*

$$(i) \quad s_\alpha^*(e_{[\gamma]} \otimes P_{[\gamma]}x) = \begin{cases} \phi(\rho_\alpha(1))(e_{[\gamma']} \otimes P_{[\gamma']}x) & \text{if } \gamma \approx \alpha\gamma', \\ 0 & \text{otherwise.} \end{cases}$$

$$(ii) \quad t_a^*(e_{[\gamma]} \otimes P_{[\gamma]}x) = \begin{cases} \phi(\eta_a(1))(e_{[\gamma']} \otimes P_{[\gamma']}x) & \text{if } \gamma \approx a\gamma', \\ 0 & \text{otherwise.} \end{cases}$$

Proof. (i) For $\gamma \in L_{(n,m)}, \gamma' \in L_{(n-1,m)}$ and $\alpha \in \Sigma^\rho$, we have

$$\begin{aligned} \langle s_\alpha^*(e_{[\gamma]} \otimes P_{[\gamma]}x) \mid e_{[\gamma']} \otimes P_{[\gamma']}x' \rangle &= \langle e_{[\gamma]} \otimes P_{[\gamma]}x \mid e_{[\alpha\gamma']} \otimes P_{[\alpha\gamma']}x' \rangle \\ &= \begin{cases} x^*P_{[\alpha\gamma']}x & \text{if } \gamma \approx \alpha\gamma', \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

On the other hand,

$$\phi(\rho_\alpha(1))(e_{[\gamma']} \otimes P_{[\gamma']}x) = e_{[\gamma']} \otimes P_{[\alpha\gamma']}P_{\gamma'}x = e_{[\gamma']} \otimes P_{[\alpha\gamma']}x$$

so that

$$\langle \phi(\rho_\alpha(1))(e_{[\gamma']} \otimes P_{[\gamma']}x) \mid e_{[\gamma']} \otimes P_{[\gamma']}x' \rangle = x^*P_{[\alpha\gamma']}x'.$$

Hence we obtain the desired equality. Similarly we see (ii). □

The following lemma is straightforward.

Lemma 6.3. *For $\alpha \in \Sigma^\rho, a \in \Sigma^\eta$ and $\gamma \in L_{(n,m)}, x \in \mathcal{A}$, we have:*

$$(i) \quad s_\alpha s_\alpha^*(e_{[\gamma]} \otimes P_{[\gamma]}x) = \begin{cases} e_{[\gamma]} \otimes P_{[\gamma]}x & \text{if } \gamma \approx \alpha\gamma' \text{ for some } \gamma' \in L_{(n-1,m)}, \\ 0 & \text{otherwise.} \end{cases}$$

$$(ii) \quad t_a t_a^*(e_{[\gamma]} \otimes P_{[\gamma]}x) = \begin{cases} e_{[\gamma]} \otimes P_{[\gamma]}x & \text{if } \gamma \approx a\gamma' \text{ for some } \gamma' \in L_{(n,m-1)}, \\ 0 & \text{otherwise.} \end{cases}$$

Hence we see:

Lemma 6.4.

$$(i) \quad 1 - \sum_{\alpha \in \Sigma^\rho} s_\alpha s_\alpha^* = \text{the projection onto the subspace spanned by the vectors } e_{[\gamma]} \otimes P_{[\gamma]}x \text{ for } \gamma \in \cup_{m=0}^\infty L_{(0,m)}, x \in \mathcal{A}.$$

- (ii) $1 - \sum_{a \in \Sigma^\eta} t_a t_a^* =$ the projection onto the subspace spanned by the vectors $e_{[\gamma]} \otimes P_{[\gamma]}x$ for $\gamma \in \cup_{n=0}^\infty L(n,0), x \in \mathcal{A}$.

Lemma 6.5. For $\alpha \in \Sigma^\rho, a \in \Sigma^\eta$ and $x \in \mathcal{A}$, we have:

- (i) $s_\alpha^* x s_\alpha = \phi(\rho_\alpha(x))$ and in particular $s_\alpha^* s_\alpha = \phi(\rho_\alpha(1))$.
- (ii) $t_a^* x t_a = \phi(\eta_a(x))$ and in particular $t_a^* t_a = \phi(\eta_a(1))$.

Proof. (i) It follows that for $\gamma \in L(n, m)$ with $\alpha\gamma \in L(n+1, m)$ and $y \in \mathcal{A}$,

$$\begin{aligned} s_\alpha^* x s_\alpha (e_{[\gamma]} \otimes P_{[\gamma]}y) &= s_\alpha^* (e_{[\alpha\gamma]} \otimes P_{[\alpha\gamma]}y \xi_{\alpha\gamma}(x)) \\ &= e_{[\gamma]} \otimes P_{[\gamma]}y \xi_\gamma(\rho_\alpha(x)) \\ &= \phi(\rho_\alpha(x))(e_{[\gamma]} \otimes P_{[\gamma]}y). \end{aligned}$$

If $\alpha\gamma \notin L(n+1, m)$, we have

$$s_\alpha (e_{[\gamma]} \otimes P_{[\gamma]}y) = 0, \quad \phi(\rho_\alpha(x))(e_{[\gamma]} \otimes P_{[\gamma]}y) = 0.$$

Hence we see that $s_\alpha^* x s_\alpha = \phi(\rho_\alpha(x))$. Similarly we see (ii). □

Lemma 6.6. For $\alpha, \beta \in \Sigma^\rho, a, b \in \Sigma^\eta$ we have:

$$(6.1) \quad s_\alpha t_b = t_a s_\beta \quad \text{if } \kappa(\alpha, b) = (a, \beta).$$

Proof. For $\gamma \in L(n, m)$ with $\alpha b\gamma, a\beta\gamma \in L(n+1, m+1)$ and $x \in \mathcal{A}$, we have

$$\begin{aligned} s_\alpha t_b (e_{[\gamma]} \otimes P_{[\gamma]}x) &= e_{[\alpha b\gamma]} \otimes P_{[\alpha b\gamma]}y, \\ t_a s_\beta (e_{[\gamma]} \otimes P_{[\gamma]}x) &= (e_{[a\beta\gamma]} \otimes P_{[a\beta\gamma]}x). \end{aligned}$$

Since $\kappa(\alpha, b) = (a, \beta)$, the condition $\alpha b\gamma \in L(n+1, m+1)$ is equivalent to the condition $a\beta\gamma \in L(n+1, m+1)$. We then have $[\alpha b\gamma] = [a\beta\gamma]$ and $P_{[\alpha b\gamma]} = P_{[a\beta\gamma]}$. □

Let $\mathcal{I}_{\rho, \eta}^\kappa$ be the ideal of $\mathcal{T}_{\rho, \eta}^\kappa$ generated by the two projections:

$$1 - \sum_{\alpha \in \Sigma^\rho} s_\alpha s_\alpha^* \quad \text{and} \quad 1 - \sum_{a \in \Sigma^\eta} t_a t_a^*.$$

Let $\widehat{\mathcal{O}}_{\rho, \eta}^\kappa$ be the quotient C^* -algebra

$$\widehat{\mathcal{O}}_{\rho, \eta}^\kappa = \mathcal{T}_{\rho, \eta}^\kappa / \mathcal{I}_{\rho, \eta}^\kappa.$$

Let $\pi_{\rho, \eta} : \mathcal{T}_{\rho, \eta}^\kappa \rightarrow \widehat{\mathcal{O}}_{\rho, \eta}^\kappa$ be the quotient map. Put

$$\widehat{S}_\alpha = \pi_{\rho, \eta}(s_\alpha), \quad \widehat{T}_a = \pi_{\rho, \eta}(t_a), \quad \widehat{i}(x) = \pi_{\rho, \eta}(i_{(F_\kappa)}(x))$$

for $\alpha \in \Sigma^\rho, a \in \Sigma^\eta$ and $x \in \mathcal{A}$. By the above discussions, the following relations hold:

$$\begin{aligned} \sum_{\beta \in \Sigma^\rho} \widehat{S}_\beta \widehat{S}_\beta^* &= 1, & \widehat{i}(x) \widehat{S}_\alpha \widehat{S}_\alpha^* &= \widehat{S}_\alpha \widehat{S}_\alpha^* \widehat{i}(x), & \widehat{S}_\alpha^* \widehat{i}(x) \widehat{S}_\alpha &= \widehat{i}(\rho_\alpha(x)), \\ \sum_{b \in \Sigma^\eta} \widehat{T}_b \widehat{T}_b^* &= 1, & \widehat{i}(x) \widehat{T}_a \widehat{T}_a^* &= \widehat{T}_a \widehat{T}_a^* \widehat{i}(x), & \widehat{T}_a^* \widehat{i}(x) \widehat{T}_a &= \widehat{i}(\eta_a(x)), \\ \widehat{S}_\alpha \widehat{T}_b &= \widehat{T}_a \widehat{S}_\beta & \text{if } \kappa(\alpha, b) &= (a, \beta) \end{aligned}$$

for all $x \in \mathcal{A}$ and $\alpha \in \Sigma^\rho, a \in \Sigma^\eta$.

For $(z, w) \in \mathbb{T}^2$, the correspondence

$$e_{[\gamma]} \otimes P_{[\gamma]}x \in H_{(n,m)} \longrightarrow z^n w^m e_{[\gamma]} \otimes P_{[\gamma]}x \in H_{(n,m)}$$

yields a unitary representation of \mathbb{T}^2 on $H_{(n,m)}$, which extends to F_κ , denoted by $u_{(z,w)}$. Since

$$u_{(z,w)} \mathcal{T}_{\rho,\eta}^\kappa u_{(z,w)}^* = \mathcal{T}_{\rho,\eta}^\kappa, \quad u_{(z,w)} \mathcal{I}_{\rho,\eta}^\kappa u_{(z,w)}^* = \mathcal{I}_{\rho,\eta}^\kappa,$$

The map

$$X \in \mathcal{T}_{\rho,\eta}^\kappa \longrightarrow u_{(z,w)} X u_{(z,w)}^* \in \mathcal{T}_{\rho,\eta}^\kappa$$

yields an action of \mathbb{T}^2 on the C^* -algebra $\widehat{\mathcal{O}}_{\rho,\eta}^\kappa$, which we denote by $\widehat{\theta}$. Similarly to the action θ on $\mathcal{O}_{\rho,\eta}^\kappa$, we may define the conditional expectation $\widehat{\mathcal{E}}_{\rho,\eta}$ from $\widehat{\mathcal{O}}_{\rho,\eta}^\kappa$ to the fixed point algebra $(\widehat{\mathcal{O}}_{\rho,\eta}^\kappa)^{\widehat{\theta}}$ by taking the integration of the function $\widehat{\theta}_{(z,w)}(X)$ over $(z, w) \in \mathbb{T}^2$ for $X \in \widehat{\mathcal{O}}_{\rho,\eta}^\kappa$. Then as in the proof of Proposition 5.10, one may prove the following theorem.

Theorem 6.7. *The algebra $\widehat{\mathcal{O}}_{\rho,\eta}^\kappa$ is canonically $*$ -isomorphic to the C^* -algebra $\mathcal{O}_{\rho,\eta}^\kappa$ through the correspondences:*

$$S_\alpha \longrightarrow \widehat{S}_\alpha, \quad T_a \longrightarrow \widehat{T}_a, \quad x \longrightarrow \widehat{i}(x)$$

for $\alpha \in \Sigma^\rho, a \in \Sigma^\eta$ and $x \in \mathcal{A}$.

7. K-Theory machinery

Let us denote by \mathcal{K} the C^* -algebra of compact operators on a separable infinite dimensional Hilbert space. For a C^* -algebra \mathcal{B} , we denote by $M(\mathcal{B})$ its multiplier algebra. In this section, we will study K-theory groups $K_*(\mathcal{O}_{\rho,\eta}^\kappa)$ for the C^* -algebra $\mathcal{O}_{\rho,\eta}^\kappa$. We fix a C^* -textile dynamical system $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$. We define two actions

$$\widehat{\rho} : \mathbb{T} \longrightarrow \text{Aut}(\mathcal{O}_{\rho,\eta}^\kappa), \quad \widehat{\eta} : \mathbb{T} \longrightarrow \text{Aut}(\mathcal{O}_{\rho,\eta}^\kappa)$$

of the circle group $\mathbb{T} = \{z \in \mathbb{C} \mid |z| = 1\}$ to $\mathcal{O}_{\rho,\eta}^\kappa$ by setting

$$\widehat{\rho}_z = \theta_{(z,1)}, \quad \widehat{\eta}_w = \theta_{(1,w)}, \quad z, w \in \mathbb{T}.$$

They satisfy

$$\widehat{\rho}_z \circ \widehat{\eta}_w = \widehat{\eta}_w \circ \widehat{\rho}_z = \theta_{(z,w)}, \quad z, w \in \mathbb{T}.$$

Set the fixed point algebras

$$\begin{aligned} (\mathcal{O}_{\rho,\eta}^\kappa)^{\widehat{\rho}} &= \{x \in \mathcal{O}_{\rho,\eta}^\kappa \mid \widehat{\rho}_z(x) = x \text{ for all } z \in \mathbb{T}\}, \\ (\mathcal{O}_{\rho,\eta}^\kappa)^{\widehat{\eta}} &= \{x \in \mathcal{O}_{\rho,\eta}^\kappa \mid \widehat{\eta}_w(x) = x \text{ for all } w \in \mathbb{T}\}. \end{aligned}$$

For $x \in (\mathcal{O}_{\rho,\eta}^\kappa)^{\widehat{\rho}}$, define the $\mathcal{O}_{\rho,\eta}^\kappa$ -valued constant function

$$\widehat{x} \in L^1(\mathbb{T}, \mathcal{O}_{\rho,\eta}^\kappa) \subset \mathcal{O}_{\rho,\eta}^\kappa \times_{\widehat{\rho}} \mathbb{T}$$

from \mathbb{T} by setting $\hat{x}(z) = x, z \in \mathbb{T}$. Put $p_0 = \hat{1}$. By [45], the algebra $(\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}}$ is canonically isomorphic to $p_0(\mathcal{O}_{\rho,\eta}^\kappa \times_{\hat{\rho}} \mathbb{T})p_0$ through the map

$$j_\rho : x \in (\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}} \longrightarrow \hat{x} \in p_0(\mathcal{O}_{\rho,\eta}^\kappa \times_{\hat{\rho}} \mathbb{T})p_0$$

which induces an isomorphism

$$(7.1) \quad j_{\rho^*} : K_i((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}}) \longrightarrow K_i(p_0(\mathcal{O}_{\rho,\eta}^\kappa \times_{\hat{\rho}} \mathbb{T})p_0), \quad i = 0, 1$$

on their K-groups. By a similar manner to the proofs given in [23, Section 4], one may prove the following lemma.

Lemma 7.1.

(i) *There exists an isometry*

$$v \in M((\mathcal{O}_{\rho,\eta}^\kappa \times_{\hat{\rho}} \mathbb{T}) \otimes \mathcal{K})$$

such that $vv^* = p_0 \otimes 1, v^*v = 1$.

(ii) $\mathcal{O}_{\rho,\eta}^\kappa \times_{\hat{\rho}} \mathbb{T}$ is stably isomorphic to $(\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}}$, and similarly $\mathcal{O}_{\rho,\eta}^\kappa \times_{\hat{\eta}} \mathbb{T}$ is stably isomorphic to $(\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\eta}}$.

(iii) The inclusion $\iota_{\hat{\rho}} : p_0(\mathcal{O}_{\rho,\eta}^\kappa \times_{\hat{\rho}} \mathbb{T})p_0 \hookrightarrow \mathcal{O}_{\rho,\eta}^\kappa \times_{\hat{\rho}} \mathbb{T}$ induces an isomorphism

$$\iota_{\hat{\rho}^*} : K_i(p_0(\mathcal{O}_{\rho,\eta}^\kappa \times_{\hat{\rho}} \mathbb{T})p_0) \cong K_i(\mathcal{O}_{\rho,\eta}^\kappa \times_{\hat{\rho}} \mathbb{T}), \quad i = 0, 1$$

on their K-groups.

Thanks to the lemma above, the isomorphism

$$\text{Ad}(v^*) : x \in p_0(\mathcal{O}_{\rho,\eta}^\kappa \times_{\hat{\rho}} \mathbb{T})p_0 \otimes \mathcal{K} \longrightarrow v^*xv \in (\mathcal{O}_{\rho,\eta}^\kappa \times_{\hat{\rho}} \mathbb{T}) \otimes \mathcal{K}$$

induces isomorphisms

$$(7.2) \quad \text{Ad}(v^*)_* : K_i(p_0(\mathcal{O}_{\rho,\eta}^\kappa \times_{\hat{\rho}} \mathbb{T})p_0) \longrightarrow K_i(\mathcal{O}_{\rho,\eta}^\kappa \times_{\hat{\rho}} \mathbb{T}), \quad i = 0, 1.$$

Let $\hat{\rho}$ be the automorphism on $\mathcal{O}_{\rho,\eta}^\kappa \times_{\hat{\rho}} \mathbb{T}$ for the positive generator of \mathbb{Z} for the dual action of $\hat{\rho}$. By (7.1) and (7.2), we may define an isomorphism

$$\beta_{\rho,i} = j_{\rho^*}^{-1} \circ \text{Ad}(v^*)_*^{-1} \circ \hat{\rho}_* \circ \text{Ad}(v^*)_* \circ j_{\rho^*} : K_i((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}}) \longrightarrow K_i((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}})$$

for $i = 0, 1$, so that the diagram is commutative:

$$\begin{array}{ccc} K_i(\mathcal{O}_{\rho,\eta}^\kappa \times_{\hat{\rho}} \mathbb{T}) & \xrightarrow{\hat{\rho}_*} & K_i(\mathcal{O}_{\rho,\eta}^\kappa \times_{\hat{\rho}} \mathbb{T}) \\ \uparrow \text{Ad}(v^*)_* & & \uparrow \text{Ad}(v^*)_* \\ K_i(p_0(\mathcal{O}_{\rho,\eta}^\kappa \times_{\hat{\rho}} \mathbb{T})p_0) & & K_i(p_0(\mathcal{O}_{\rho,\eta}^\kappa \times_{\hat{\rho}} \mathbb{T})p_0) \\ \uparrow j_{\rho^*} & & \uparrow j_{\rho^*} \\ K_i((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}}) & \xrightarrow{\beta_{\rho,i}} & K_i((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}}). \end{array}$$

By [39] (cf. [15]), one has the six term exact sequence of K-theory:

$$\begin{array}{ccccc} K_0(\mathcal{O}_{\rho,\eta}^\kappa \times_{\hat{\rho}} \mathbb{T}) & \xrightarrow{\text{id}-\hat{\rho}_*} & K_0(\mathcal{O}_{\rho,\eta}^\kappa \times_{\hat{\rho}} \mathbb{T}) & \xrightarrow{\iota_*} & K_0((\mathcal{O}_{\rho,\eta}^\kappa \times_{\hat{\rho}} \mathbb{T}) \times_{\hat{\rho}} \mathbb{Z}) \\ \delta \uparrow & & & & \exp \downarrow \\ K_1((\mathcal{O}_{\rho,\eta}^\kappa \times_{\hat{\rho}} \mathbb{T}) \times_{\hat{\rho}} \mathbb{Z}) & \xleftarrow{\iota_*} & K_1(\mathcal{O}_{\rho,\eta}^\kappa \times_{\hat{\rho}} \mathbb{T}) & \xleftarrow{\text{id}-\hat{\rho}_*} & K_1(\mathcal{O}_{\rho,\eta}^\kappa \times_{\hat{\rho}} \mathbb{T}). \end{array}$$

Since $(\mathcal{O}_{\rho,\eta}^\kappa \times_{\hat{\rho}} \mathbb{T}) \times_{\hat{\rho}} \mathbb{Z} \cong \mathcal{O}_{\rho,\eta}^\kappa \otimes \mathcal{K}$ and $K_i(\mathcal{O}_{\rho,\eta}^\kappa \times_{\hat{\rho}} \mathbb{T}) \cong K_i((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}})$, one has:

Lemma 7.2. *The following six term exact sequence of K-theory holds:*

$$\begin{array}{ccccc} K_0((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}}) & \xrightarrow{\text{id}-\beta_{\rho,0}} & K_0((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}}) & \xrightarrow{\iota_*} & K_0(\mathcal{O}_{\rho,\eta}^\kappa) \\ \delta \uparrow & & & & \exp \downarrow \\ K_1(\mathcal{O}_{\rho,\eta}^\kappa) & \xleftarrow{\iota_*} & K_1((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}}) & \xleftarrow{\text{id}-\beta_{\rho,1}} & K_1((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}}). \end{array}$$

Hence there exist short exact sequences for $i = 0, 1$:

$$\begin{aligned} 0 &\longrightarrow \text{Coker}(\text{id} - \beta_{\rho,i}) \text{ in } K_i((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}}) \\ &\longrightarrow K_i(\mathcal{O}_{\rho,\eta}^\kappa) \\ &\longrightarrow \text{Ker}(\text{id} - \beta_{\rho,i+1}) \text{ in } K_{i+1}((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}}) \\ &\longrightarrow 0. \end{aligned}$$

In the rest of this section, we will study the groups

$$\text{Coker}(\text{id} - \beta_{\rho,i}) \text{ in } K_i((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}}), \quad \text{Ker}(\text{id} - \beta_{\rho,i+1}) \text{ in } K_{i+1}((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}}).$$

The action $\hat{\eta}$ acts on the subalgebra $(\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}}$, which we still denote by $\hat{\eta}$. Then the fixed point algebra $((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}})^{\hat{\eta}}$ of $(\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}}$ under $\hat{\eta}$ coincides with $\mathcal{F}_{\rho,\eta}$. The above discussions for the action $\hat{\rho} : \mathbb{T} \rightarrow \mathcal{O}_{\rho,\eta}^\kappa$ works for the action $\hat{\eta} : \mathbb{T} \rightarrow (\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}}$ as in the following way. For $y \in ((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}})^{\hat{\eta}}$, define the constant function $\hat{y} \in L^1(\mathbb{T}, (\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}}) \subset (\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}} \times_{\hat{\eta}} \mathbb{T}$ by setting $\hat{y}(w) = y, w \in \mathbb{T}$. Putting $q_0 = \hat{1}$, the algebra $((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}})^{\hat{\eta}}$ is canonically isomorphic to $q_0((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}} \times_{\hat{\eta}} \mathbb{T})q_0$ through the map

$$j_\eta^\rho : y \in ((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}})^{\hat{\eta}} \longrightarrow \hat{y} \in q_0((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}} \times_{\hat{\eta}} \mathbb{T})q_0$$

which induces an isomorphism

$$j_{\eta_*}^\rho : K_i(((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}})^{\hat{\eta}}) \longrightarrow K_i(q_0((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}} \times_{\hat{\eta}} \mathbb{T})q_0), \quad i = 0, 1$$

on their K-groups. Similarly to Lemma 7.1, we have:

Lemma 7.3.

(i) *There exists an isometry*

$$u \in M(((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}} \times_{\hat{\eta}} \mathbb{T}) \otimes \mathcal{K})$$

such that $uu^* = q_0 \otimes 1, u^*u = 1$.

- (ii) $(\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}} \times_{\hat{\eta}} \mathbb{T}$ is stably isomorphic to $((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}})^{\hat{\eta}}$.
- (iii) The inclusion

$$\iota_{\hat{\eta}}^{\hat{\rho}} : q_0((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}} \times_{\hat{\eta}} \mathbb{T})q_0 = ((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}})^{\hat{\eta}} = \mathcal{F}_{\rho,\eta} \hookrightarrow (\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}} \times_{\hat{\eta}} \mathbb{T}$$

induces an isomorphism

$$\iota_{\hat{\eta}*}^{\hat{\rho}} : K_i(q_0((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}} \times_{\hat{\eta}} \mathbb{T})q_0) \cong K_i((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}} \times_{\hat{\eta}} \mathbb{T}), \quad i = 0, 1$$

on their K -groups.

The isomorphism

$$\text{Ad}(u^*) : y \in q_0((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}} \times_{\hat{\eta}} \mathbb{T})q_0 \longrightarrow u^* y u \in (\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}} \times_{\hat{\eta}} \mathbb{T}$$

induces isomorphisms

$$\text{Ad}(u^*)_* : K_i(q_0((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}} \times_{\hat{\eta}} \mathbb{T})q_0) \cong K_i((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}} \times_{\hat{\eta}} \mathbb{T}), \quad i = 0, 1.$$

Let $\hat{\eta}_\rho$ be the automorphism on $(\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}} \times_{\hat{\eta}} \mathbb{T}$ for the positive generator of \mathbb{Z} for the dual action of $\hat{\eta}$. Define an isomorphism

$$\gamma_{\eta,i} = j_{\hat{\eta}*}^{\rho-1} \circ \text{Ad}(u^*)_*^{-1} \circ \hat{\eta}_{\rho*} \circ \text{Ad}(u^*)_* \circ j_{\hat{\eta}*}^\rho : K_i(\mathcal{F}_{\rho,\eta}) \longrightarrow K_i(\mathcal{F}_{\rho,\eta}), \quad i = 0, 1$$

such that the diagram is commutative for $i = 0, 1$:

$$\begin{array}{ccc} K_i((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}} \times_{\hat{\eta}} \mathbb{T}) & \xrightarrow{\hat{\eta}_{\rho*}} & K_i((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}} \times_{\hat{\eta}} \mathbb{T}) \\ \uparrow \text{Ad}(u^*)_* & & \uparrow \text{Ad}(u^*)_* \\ K_i(q_0((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}} \times_{\hat{\eta}} \mathbb{T})q_0) & & K_i(q_0((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}} \times_{\hat{\eta}} \mathbb{T})q_0) \\ \uparrow j_{\hat{\eta}*}^\rho & & \uparrow j_{\hat{\eta}*}^\rho \\ K_i(((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}})^{\hat{\eta}}) & & K_i(((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}})^{\hat{\eta}}) \\ \parallel & & \parallel \\ K_i(\mathcal{F}_{\rho,\eta}) & \xrightarrow{\gamma_{\eta,i}} & K_i(\mathcal{F}_{\rho,\eta}). \end{array}$$

We similarly define an endomorphism $\gamma_{\rho,i} : K_i(\mathcal{F}_{\rho,\eta}) \longrightarrow K_i(\mathcal{F}_{\rho,\eta})$ by exchanging the rôles of ρ and η .

Under the equality $((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}})^{\hat{\eta}} = \mathcal{F}_{\rho,\eta}$, we have the following lemma which is similar to Lemma 7.2

Lemma 7.4. *The following six term exact sequence of K -theory holds:*

$$\begin{array}{ccccc} K_0(\mathcal{F}_{\rho,\eta}) & \xrightarrow{\text{id}-\gamma_{\eta,0}} & K_0(\mathcal{F}_{\rho,\eta}) & \xrightarrow{\iota_*} & K_0((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}}) \\ \delta \uparrow & & & & \exp \downarrow \\ K_1((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}}) & \xleftarrow{\iota_*} & K_1(\mathcal{F}_{\rho,\eta}) & \xleftarrow{\text{id}-\gamma_{\eta,1}} & K_1(\mathcal{F}_{\rho,\eta}). \end{array}$$

In particular, if $K_1(\mathcal{F}_{\rho,\eta}) = 0$, we have

$$\begin{aligned} K_0((\mathcal{O}_{\rho,\eta}^\kappa)^\hat{\rho}) &= \text{Coker}(\text{id} - \gamma_{\eta,0}) \quad \text{in } K_0(\mathcal{F}_{\rho,\eta}), \\ K_1((\mathcal{O}_{\rho,\eta}^\kappa)^\hat{\rho}) &= \text{Ker}(\text{id} - \gamma_{\eta,0}) \quad \text{in } K_0(\mathcal{F}_{\rho,\eta}). \end{aligned}$$

Denote by $M_n(\mathcal{B})$ the $n \times n$ matrix algebra over a C^* -algebra \mathcal{B} , which is identified with the tensor product $\mathcal{B} \otimes M_n(\mathbb{C})$. The following lemmas hold.

Lemma 7.5. *For a projection $q \in M_n((\mathcal{O}_{\rho,\eta}^\kappa)^\rho)$ and a partial isometry $S \in \mathcal{O}_{\rho,\eta}^\kappa$ such that*

$$\hat{\rho}_z(S) = zS \quad \text{for } z \in \mathbb{T}, \quad q(SS^* \otimes 1_n) = (SS^* \otimes 1_n)q,$$

we have

$$\beta_{\rho,0}^{-1}([(SS^* \otimes 1_n)q]) = [(S^* \otimes 1_n)q(S \otimes 1_n)] \quad \text{in } K_0((\mathcal{O}_{\rho,\eta}^\kappa)^\hat{\rho}).$$

Proof. As q commutes with $SS^* \otimes 1_n$, $p = (S^* \otimes 1_n)q(S \otimes 1_n)$ is a projection in $(\mathcal{O}_{\rho,\eta}^\kappa)^\hat{\rho}$. Since $p \leq S^*S \otimes 1_n$, By a similar argument to the proof of [23, Lemma 4.5], one sees that $\beta_{\rho,0}([p]) = [(S \otimes 1_n)p(S^* \otimes 1_n)]$ in $K_0((\mathcal{O}_{\rho,\eta}^\kappa)^\hat{\rho})$. \square

Lemma 7.6.

(i) *For a projection $q \in M_n(\mathcal{F}_{\rho,\eta})$ and a partial isometry $T \in (\mathcal{O}_{\rho,\eta}^\kappa)^\hat{\rho}$ such that*

$$\hat{\eta}_w(T) = wT \quad \text{for } w \in \mathbb{T}, \quad q(TT^* \otimes 1_n) = (TT^* \otimes 1_n)q,$$

we have

$$\gamma_{\eta,0}^{-1}([(TT^* \otimes 1_n)q]) = [(T^* \otimes 1_n)q(T \otimes 1_n)] \quad \text{in } K_0(\mathcal{F}_{\rho,\eta}).$$

(ii) *For a projection $q \in M_n(\mathcal{F}_{\rho,\eta})$ and a partial isometry $S \in (\mathcal{O}_{\rho,\eta}^\kappa)^\hat{\eta}$ such that*

$$\hat{\rho}_z(S) = zS \quad \text{for } z \in \mathbb{T}, \quad q(SS^* \otimes 1_n) = (SS^* \otimes 1_n)q,$$

we have

$$\gamma_{\rho,0}^{-1}([(SS^* \otimes 1_n)q]) = [(S^* \otimes 1_n)q(S \otimes 1_n)] \quad \text{in } K_0(\mathcal{F}_{\rho,\eta}).$$

Hence we have

Lemma 7.7. *The diagram*

$$\begin{array}{ccc} K_0(\mathcal{F}_{\rho,\eta}) & \xrightarrow{\text{id} - \gamma_{\rho,0}} & K_0(\mathcal{F}_{\rho,\eta}) \\ \downarrow \iota_* & & \downarrow \iota_* \\ K_0((\mathcal{O}_{\rho,\eta}^\kappa)^\hat{\rho}) & \xrightarrow{\text{id} - \beta_{\rho,0}} & K_0((\mathcal{O}_{\rho,\eta}^\kappa)^\hat{\rho}) \end{array}$$

is commutative.

Proof. By [35, Proposition 3.3], the map $\iota_* : K_0(\mathcal{F}_{\rho,\eta}) \rightarrow K_0((\mathcal{O}_{\rho,\eta}^\kappa)^\hat{\rho})$ is induced by the natural inclusion $\mathcal{F}_{\rho,\eta} (= ((\mathcal{O}_{\rho,\eta}^\kappa)^\hat{\rho})^\eta) \hookrightarrow (\mathcal{O}_{\rho,\eta}^\kappa)^\hat{\rho}$. For an element $[q] \in K_0(\mathcal{F}_{\rho,\eta})$ one may assume that $q \in M_n(\mathcal{F}_{\rho,\eta})$ for some $n \in \mathbb{N}$ so that one has

$$\begin{aligned} \gamma_{\rho,0}^{-1}([q]) &= \sum_{\alpha \in \Sigma^\rho} [(S_\alpha S_\alpha^* \otimes 1_n)q] \\ &= \sum_{\alpha \in \Sigma^\rho} [(S_\alpha^* \otimes 1_n)q(S_\alpha \otimes 1_n)] \\ &= \sum_{\alpha \in \Sigma^\rho} \beta_{\rho,0}^{-1}([q(S_\alpha S_\alpha^* \otimes 1_n)]) = \beta_{\rho,0}^{-1}([q]) \end{aligned}$$

so that $\beta_{\rho,0}|_{K_0(\mathcal{F}_{\rho,\eta})} = \gamma_{\rho,0}$. □

In the rest of this section, we assume that $K_1(\mathcal{F}_{\rho,\eta}) = 0$. The following lemma is crucial in our further discussions.

Lemma 7.8. *In the six term exact sequence in Lemma 7.4 with $K_1(\mathcal{F}_{\rho,\eta}) = 0$, we have the following commutative diagrams:*

$$(7.3) \quad \begin{array}{ccc} 0 & & 0 \\ \downarrow & & \downarrow \\ K_1((\mathcal{O}_{\rho,\eta}^\kappa)^\hat{\rho}) & \xrightarrow{\text{id}-\beta_{\rho,1}} & K_1((\mathcal{O}_{\rho,\eta}^\kappa)^\hat{\rho}) \\ \delta \downarrow & & \delta \downarrow \\ K_0(\mathcal{F}_{\rho,\eta}) & \xrightarrow{\text{id}-\gamma_{\rho,0}} & K_0(\mathcal{F}_{\rho,\eta}) \\ \text{id}-\gamma_{\eta,0} \downarrow & & \text{id}-\gamma_{\eta,0} \downarrow \\ K_0(\mathcal{F}_{\rho,\eta}) & \xrightarrow{\text{id}-\gamma_{\rho,0}} & K_0(\mathcal{F}_{\rho,\eta}) \\ \iota_* \downarrow & & \iota_* \downarrow \\ K_0((\mathcal{O}_{\rho,\eta}^\kappa)^\hat{\rho}) & \xrightarrow{\text{id}-\beta_{\rho,0}} & K_0((\mathcal{O}_{\rho,\eta}^\kappa)^\hat{\rho}) \\ \downarrow & & \downarrow \\ 0 & & 0 \end{array}$$

Proof. It is well-known that δ -map is functorial (see [48, Theorem 7.2.5], [4, p.266 (LX)]). Hence the diagram of the upper square

$$\begin{array}{ccc} K_1((\mathcal{O}_{\rho,\eta}^\kappa)^\hat{\rho}) & \xrightarrow{\text{id}-\beta_{\rho,1}} & K_1((\mathcal{O}_{\rho,\eta}^\kappa)^\hat{\rho}) \\ \delta \downarrow & & \delta \downarrow \\ K_0(\mathcal{F}_{\rho,\eta}) & \xrightarrow{\text{id}-\gamma_{\rho,0}} & K_0(\mathcal{F}_{\rho,\eta}) \end{array}$$

is commutative. Since $\gamma_{\rho,0} \circ \gamma_{\eta,0} = \gamma_{\eta,0} \circ \gamma_{\rho,0}$, the diagram of the middle square

$$(7.4) \quad \begin{array}{ccc} K_0(\mathcal{F}_{\rho,\eta}) & \xrightarrow{\text{id}-\gamma_{\rho,0}} & K_0(\mathcal{F}_{\rho,\eta}) \\ \downarrow \text{id}-\gamma_{\eta,0} & & \downarrow \text{id}-\gamma_{\eta,0} \\ K_0(\mathcal{F}_{\rho,\eta}) & \xrightarrow{\text{id}-\gamma_{\rho,0}} & K_0(\mathcal{F}_{\rho,\eta}) \end{array}$$

is commutative. The commutativity of the lower square comes from the preceding lemma. \square

We will describe the K-groups $K_*(\mathcal{O}_{\rho,\eta}^\kappa)$ in terms of the kernels and cokernels of the homomorphisms $\text{id} - \gamma_{\rho,0}$ and $\text{id} - \gamma_{\eta,0}$ on $K_0(\mathcal{F}_{\rho,\eta})$. Recall that there exist two short exact sequences by Lemma 7.2:

$$\begin{aligned} 0 &\longrightarrow \text{Coker}(\text{id} - \beta_{\rho,0}) \text{ in } K_0((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}}) \\ &\longrightarrow K_0(\mathcal{O}_{\rho,\eta}^\kappa) \\ &\longrightarrow \text{Ker}(\text{id} - \beta_{\rho,1}) \text{ in } K_1((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}}) \\ &\longrightarrow 0 \end{aligned}$$

and

$$\begin{aligned} 0 &\longrightarrow \text{Coker}(\text{id} - \beta_{\rho,1}) \text{ in } K_1((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}}) \\ &\longrightarrow K_1(\mathcal{O}_{\rho,\eta}^\kappa) \\ &\longrightarrow \text{Ker}(\text{id} - \beta_{\rho,0}) \text{ in } K_0((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}}) \\ &\longrightarrow 0. \end{aligned}$$

As $\gamma_{\eta,0} \circ \gamma_{\rho,0} = \gamma_{\rho,0} \circ \gamma_{\eta,0}$ on $K_0(\mathcal{F}_{\rho,\eta})$, the homomorphisms $\gamma_{\rho,0}$ and $\gamma_{\eta,0}$ naturally act on $\text{Coker}(\text{id} - \gamma_{\eta,0}) = K_0(\mathcal{F}_{\rho,\eta})/(\text{id} - \gamma_{\eta,0})K_0(\mathcal{F}_{\rho,\eta})$ and $\text{Coker}(\text{id} - \gamma_{\rho,0}) = K_0(\mathcal{F}_{\rho,\eta})/(\text{id} - \gamma_{\rho,0})K_0(\mathcal{F}_{\rho,\eta})$ as endomorphisms respectively, which we denote by $\bar{\gamma}_{\rho,0}$ and $\bar{\gamma}_{\eta,0}$ respectively.

Lemma 7.9.

(i) For $K_0(\mathcal{O}_{\rho,\eta}^\kappa)$, we have

$$\begin{aligned} &\text{Coker}(\text{id} - \beta_{\rho,0}) \text{ in } K_0((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}}) \\ &\cong \text{Coker}(\text{id} - \bar{\gamma}_{\rho,0}) \text{ in } K_0(\mathcal{F}_{\rho,\eta})/(\text{id} - \gamma_{\eta,0})K_0(\mathcal{F}_{\rho,\eta}) \\ &\cong K_0(\mathcal{F}_{\rho,\eta})/((\text{id} - \gamma_{\rho,0})K_0(\mathcal{F}_{\rho,\eta}) + (\text{id} - \gamma_{\eta,0})K_0(\mathcal{F}_{\rho,\eta})) \end{aligned}$$

and

$$\begin{aligned} &\text{Ker}(\text{id} - \beta_{\rho,1}) \text{ in } K_1((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}}) \\ &\cong \text{Ker}(\text{id} - \gamma_{\rho,0}) \text{ in } (\text{Ker}(\text{id} - \gamma_{\eta,0}) \text{ in } K_0(\mathcal{F}_{\rho,\eta})) \\ &\cong \text{Ker}(\text{id} - \gamma_{\rho,0}) \cap \text{Ker}(\text{id} - \gamma_{\eta,0}) \text{ in } K_0(\mathcal{F}_{\rho,\eta}). \end{aligned}$$

(ii) For $K_1(\mathcal{O}_{\rho,\eta}^\kappa)$, we have

$$\begin{aligned} & \text{Coker}(\text{id} - \beta_{\rho,1}) \text{ in } K_1((\mathcal{O}_{\rho,\eta}^\kappa)^\hat{\rho}) \\ & \cong (\text{Ker}(\text{id} - \gamma_{\eta,0}) \text{ in } K_0(\mathcal{F}_{\rho,\eta})) / (\text{id} - \gamma_{\rho,0})(\text{Ker}(\text{id} - \gamma_{\eta,0}) \text{ in } K_0(\mathcal{F}_{\rho,\eta})) \end{aligned}$$

and

$$\begin{aligned} & \text{Ker}(\text{id} - \beta_{\rho,0}) \text{ in } K_0((\mathcal{O}_{\rho,\eta}^\kappa)^\hat{\rho}) \\ & \cong \text{Ker}(\text{id} - \bar{\gamma}_{\rho,0}) \text{ in } (K_0(\mathcal{F}_{\rho,\eta}) / (\text{id} - \gamma_{\eta,0})K_0(\mathcal{F}_{\rho,\eta})). \end{aligned}$$

Proof. (i) We will first prove the assertions for the group

$$\text{Coker}(\text{id} - \beta_{\rho,0}) \text{ in } K_0((\mathcal{O}_{\rho,\eta}^\kappa)^\hat{\rho}).$$

In the diagram (7.3), the exactness of the vertical arrows implies that ι_* is surjective so that

$$K_0((\mathcal{O}_{\rho,\eta}^\kappa)^\hat{\rho}) \cong \iota_*(K_0(\mathcal{F}_{\rho,\eta})) \cong K_0(\mathcal{F}_{\rho,\eta}) / \text{Ker}(\text{id} - \gamma_{\eta,0}) \text{ in } K_0(\mathcal{F}_{\rho,\eta}).$$

By the commutativity in the lower square in the diagram (7.3), one has

$$\begin{aligned} & \text{Coker}(\text{id} - \beta_{\rho,0}) \text{ in } K_0((\mathcal{O}_{\rho,\eta}^\kappa)^\hat{\rho}) \\ & \cong \text{Coker}(\text{id} - \bar{\gamma}_{\rho,0}) \text{ in } (\text{Coker}(\text{id} - \gamma_{\eta,0}) \text{ in } K_0(\mathcal{F}_{\rho,\eta})). \end{aligned}$$

The latter group will be proved to be isomorphic to the group

$$K_0(\mathcal{F}_{\rho,\eta}) / ((\text{id} - \gamma_{\eta,0})K_0(\mathcal{F}_{\rho,\eta}) + (\text{id} - \gamma_{\rho,0})K_0(\mathcal{F}_{\rho,\eta})).$$

Put $H_{\rho,\eta} = (\text{id} - \gamma_{\eta,0})K_0(\mathcal{F}_{\rho,\eta}) + (\text{id} - \gamma_{\rho,0})K_0(\mathcal{F}_{\rho,\eta})$ the subgroup of $K_0(\mathcal{F}_{\rho,\eta})$ generated by $(\text{id} - \gamma_{\eta,0})K_0(\mathcal{F}_{\rho,\eta})$ and $(\text{id} - \gamma_{\rho,0})K_0(\mathcal{F}_{\rho,\eta})$. Set the quotient maps

$$\begin{aligned} K_0(\mathcal{F}_{\rho,\eta}) & \xrightarrow{q_\eta} K_0(\mathcal{F}_{\rho,\eta}) / (\text{id} - \gamma_{\eta,0})K_0(\mathcal{F}_{\rho,\eta}) \\ & \xrightarrow{q(\text{id} - \gamma_{\rho,0})} \text{Coker}(\text{id} - \bar{\gamma}_{\rho,0}) \text{ in } K_0(\mathcal{F}_{\rho,\eta}) / (\text{id} - \gamma_{\eta,0})K_0(\mathcal{F}_{\rho,\eta}) \end{aligned}$$

and

$$\begin{aligned} \Phi & = q(\text{id} - \gamma_{\rho,0}) \circ q_\eta : K_0(\mathcal{F}_{\rho,\eta}) \\ & \longrightarrow \text{Coker}(\text{id} - \bar{\gamma}_{\rho,0}) \text{ in } K_0(\mathcal{F}_{\rho,\eta}) / (\text{id} - \gamma_{\eta,0})K_0(\mathcal{F}_{\rho,\eta}). \end{aligned}$$

It suffices to show the equality $\text{Ker}(\Phi) = H_{\rho,\eta}$. As $(\text{id} - \gamma_{\rho,0})$ commutes with $(\text{id} - \gamma_{\eta,0})$, one has

$$(\text{id} - \gamma_{\eta,0})K_0(\mathcal{F}_{\rho,\eta}) \subset \text{Ker}(\Phi), \quad (\text{id} - \gamma_{\rho,0})K_0(\mathcal{F}_{\rho,\eta}) \subset \text{Ker}(\Phi).$$

Hence we have $H_{\rho,\eta} \subset \text{Ker}(\Phi)$. On the other hand, for $g \in \text{Ker}(\Phi)$, we have $g \in (\text{id} - \bar{\gamma}_{\rho,0})(K_0(\mathcal{F}_{\rho,\eta}) / (\text{id} - \gamma_{\eta,0})K_0(\mathcal{F}_{\rho,\eta}))$ so that $g = (\text{id} - \gamma_{\rho,0})[h]$ for some $[h] \in K_0(\mathcal{F}_{\rho,\eta}) / (\text{id} - \gamma_{\eta,0})K_0(\mathcal{F}_{\rho,\eta})$. Hence

$$g = (\text{id} - \gamma_{\rho,0})h + (\text{id} - \gamma_{\rho,0})(\text{id} - \gamma_{\eta,0})K_0(\mathcal{F}_{\rho,\eta})$$

so that $g \in H_{\rho,\eta}$. Hence we have $\text{Ker}(\Phi) \subset H_{\rho,\eta}$ and $\text{Ker}(\Phi) = H_{\rho,\eta}$.

We will second prove the assertions for the group

$$\text{Ker}(\text{id} - \beta_{\rho,1}) \text{ in } K_1((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}}).$$

In the diagram (7.3), the exactness of the vertical arrows implies that δ is injective and $\text{Im}(\delta) = \text{Ker}(\text{id} - \gamma_{\eta,0})$ so that we have

$$(7.5) \quad K_1((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}}) \cong \text{Ker}(\text{id} - \gamma_{\eta,0}) \text{ in } K_0(\mathcal{F}_{\rho,\eta}).$$

By the commutativity in the upper square in the diagram (7.3), one has

$$\text{Ker}(\text{id} - \beta_{\rho,1}) \text{ in } K_1((\mathcal{O}_{\rho,\eta}^\kappa)^{\hat{\rho}}) \cong \text{Ker}(\text{id} - \gamma_{\rho,0}) \text{ in } (\text{Ker}(\text{id} - \gamma_{\eta,0}) \text{ in } K_0(\mathcal{F}_{\rho,\eta})).$$

Since $\gamma_{\eta,0}$ commutes with $\gamma_{\rho,0}$ in $K_0(\mathcal{F}_{\rho,\eta})$, we have

$$\begin{aligned} & \text{Ker}(\text{id} - \gamma_{\rho,0}) \text{ in } (\text{Ker}(\text{id} - \gamma_{\eta,0}) \text{ in } K_0(\mathcal{F}_{\rho,\eta})) \\ & \cong \text{Ker}(\text{id} - \gamma_{\rho,0}) \cap \text{Ker}(\text{id} - \gamma_{\eta,0}) \text{ in } K_0(\mathcal{F}_{\rho,\eta}). \end{aligned}$$

(ii) The assertions are similarly shown as in (i). □

Therefore we have:

Theorem 7.10. *Assume that $K_1(\mathcal{F}_{\rho,\eta}) = 0$. There exist short exact sequences:*

$$\begin{aligned} 0 & \longrightarrow K_0(\mathcal{F}_{\rho,\eta}) / ((\text{id} - \gamma_{\rho,0})K_0(\mathcal{F}_{\rho,\eta}) + (\text{id} - \gamma_{\eta,0})K_0(\mathcal{F}_{\rho,\eta})) \\ & \longrightarrow K_0(\mathcal{O}_{\rho,\eta}^\kappa) \\ & \longrightarrow \text{Ker}(\text{id} - \gamma_{\rho,0}) \cap \text{Ker}(\text{id} - \gamma_{\eta,0}) \text{ in } K_0(\mathcal{F}_{\rho,\eta}) \\ & \longrightarrow 0 \end{aligned}$$

and

$$\begin{aligned} 0 & \longrightarrow (\text{Ker}(\text{id} - \gamma_{\eta,0}) \text{ in } K_0(\mathcal{F}_{\rho,\eta})) / (\text{id} - \gamma_{\rho,0})(\text{Ker}(\text{id} - \gamma_{\eta,0}) \text{ in } K_0(\mathcal{F}_{\rho,\eta})) \\ & \longrightarrow K_1(\mathcal{O}_{\rho,\eta}^\kappa) \\ & \longrightarrow \text{Ker}(\text{id} - \bar{\gamma}_{\rho,0}) \text{ in } (K_0(\mathcal{F}_{\rho,\eta}) / (\text{id} - \gamma_{\eta,0})K_0(\mathcal{F}_{\rho,\eta})) \\ & \longrightarrow 0. \end{aligned}$$

We may describe the above formulae as follows.

Corollary 7.11. *Suppose $K_1(\mathcal{F}_{\rho,\eta}) = 0$. There exist short exact sequences:*

$$\begin{aligned} 0 & \longrightarrow \text{Coker}(\text{id} - \bar{\gamma}_{\rho,0}) \text{ in } (\text{Coker}(\text{id} - \gamma_{\eta,0}) \text{ in } K_0(\mathcal{F}_{\rho,\eta})) \\ & \longrightarrow K_0(\mathcal{O}_{\rho,\eta}^\kappa) \\ & \longrightarrow \text{Ker}(\text{id} - \gamma_{\rho,0}) \text{ in } (\text{Ker}(\text{id} - \gamma_{\eta,0}) \text{ in } K_0(\mathcal{F}_{\rho,\eta})) \\ & \longrightarrow 0 \end{aligned}$$

and

$$\begin{aligned} 0 &\longrightarrow \text{Coker}(\text{id} - \gamma_{\rho,0}) \text{ in } ((\text{Ker}(\text{id} - \gamma_{\eta,0}) \text{ in } K_0(\mathcal{F}_{\rho,\eta})) \\ &\longrightarrow K_1(\mathcal{O}_{\rho,\eta}^\kappa) \\ &\longrightarrow \text{Ker}(\text{id} - \bar{\gamma}_{\rho,0}) \text{ in } (\text{Coker}(\text{id} - \gamma_{\eta,0}) \text{ in } K_0(\mathcal{F}_{\rho,\eta})) \\ &\longrightarrow 0. \end{aligned}$$

8. K-Theory formulae

In this section, we will present more useful formulae to compute the K-groups $K_i(\mathcal{O}_{\rho,\eta}^\kappa)$ under a certain additional assumption on $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$. The additional condition on $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ is the following:

Definition 8.1. A C^* -textile dynamical system $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ is said to *form square* if the C^* -subalgebra $C^*(\rho_\alpha(1) : \alpha \in \Sigma^\rho)$ of \mathcal{A} generated by the projections $\rho_\alpha(1), \alpha \in \Sigma^\rho$ coincides with the C^* -subalgebra $C^*(\eta_a(1) : a \in \Sigma^\eta)$ of \mathcal{A} generated by the projections $\eta_a(1), a \in \Sigma^\eta$.

Lemma 8.2. Assume that $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ forms square. Put for $l \in \mathbb{Z}_+$

$$\mathcal{A}_l^\rho = C^*(\rho_\mu(1) : \mu \in B_l(\Lambda_\rho)), \quad \mathcal{A}_l^\eta = C^*(\eta_\xi(1) : \xi \in B_l(\Lambda_\eta)).$$

Then $\mathcal{A}_l^\rho = \mathcal{A}_l^\eta$.

Proof. By the assumption, we have $\mathcal{A}_1^\rho = \mathcal{A}_1^\eta$. Hence the desired equality for $l = 1$ holds. Suppose that the equalities hold for all $l \leq k$ for some $k \in \mathbb{N}$. For $\mu = (\mu_1, \mu_2, \dots, \mu_k, \mu_{k+1}) \in B_{k+1}(\Lambda_\rho)$ we have $\rho_\mu(1) = \rho_{\mu_{k+1}}(\rho_{\mu_1\mu_2\cdots\mu_k}(1))$ so that $\rho_\mu(1) \in \rho_{\mu_{k+1}}(\mathcal{A}_k^\rho)$. By the commutation relation (3.1), one sees that

$$\rho_{\mu_{k+1}}(\mathcal{A}_k^\rho) \subset C^*(\eta_\xi(\rho_\alpha(1)) : \xi \in B_k(\Lambda_\eta), \alpha \in \Sigma^\rho).$$

Since $C^*(\rho_\alpha(1) : \alpha \in \Sigma^\rho) = C^*(\eta_a(1) : a \in \Sigma^\eta)$, the algebra $C^*(\eta_\xi(\rho_\alpha(1)) : \xi \in B_k(\Lambda_\eta), \alpha \in \Sigma^\rho)$ is contained in \mathcal{A}_{k+1}^η so that $\rho_{\mu_{k+1}}(\mathcal{A}_k^\rho) \subset \mathcal{A}_{k+1}^\eta$. This implies $\rho_\mu(1) \in \mathcal{A}_{k+1}^\eta$ so that $\mathcal{A}_{k+1}^\rho \subset \mathcal{A}_{k+1}^\eta$ and hence $\mathcal{A}_{k+1}^\rho = \mathcal{A}_{k+1}^\eta$. \square

Therefore we have

Lemma 8.3. Assume that $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ forms square. Put for $j, k \in \mathbb{Z}_+$

$$\begin{aligned} \mathcal{A}_{j,k} &= C^*(\rho_\mu(\eta_\zeta(1)) : \mu \in B_j(\Lambda_\rho), \zeta \in B_k(\Lambda_\eta)) \\ &= C^*(\eta_\xi(\rho_\nu(1)) : \xi \in B_k(\Lambda_\eta), \nu \in B_j(\Lambda_\rho)). \end{aligned}$$

Then $\mathcal{A}_{j,k}$ is commutative and of finite dimensional such that

$$\mathcal{A}_{j,k} = \mathcal{A}_{j+k}^\rho (= \mathcal{A}_{j+k}^\eta).$$

Hence $\mathcal{A}_{j,k} = \mathcal{A}_{j',k'}$ if $j + k = j' + k'$.

Proof. Since $\eta_\zeta(1) \in Z_{\mathcal{A}}$ and $\rho_\mu(Z_{\mathcal{A}}) \subset Z_{\mathcal{A}}$, the algebra $\mathcal{A}_{j,k}$ belongs to the center $Z_{\mathcal{A}}$ of \mathcal{A} . By the preceding lemma, we have

$$\mathcal{A}_{j,k} = C^*(\rho_\mu(\rho_\nu(1)) : \mu \in B_j(\Lambda_\rho), \nu \in B_k(\Lambda_\rho)) = \mathcal{A}_{j+k}^\rho. \quad \square$$

For $j, k \in \mathbb{Z}_+$, put $l = j + k$. We denote by \mathcal{A}_l the commutative finite dimensional algebra $\mathcal{A}_{j,k}$. Put $m(l) = \dim \mathcal{A}_l$. Take the finite sequence of minimal projections $E_i^l, i = 1, 2, \dots, m(l)$ in \mathcal{A}_l such that $\sum_{i=1}^{m(l)} E_i^l = 1$ and hence $\mathcal{A}_l = \bigoplus_{i=1}^{m(l)} \mathbb{C}E_i^l$. Since $\rho_\alpha(\mathcal{A}_l) \subset \mathcal{A}_{l+1}$, there exists $A_{l,l+1}^\rho(i, \alpha, n)$, which takes 0 or 1, such that

$$\rho_\alpha(E_i^l) = \sum_{n=1}^{m(l+1)} A_{l,l+1}^\rho(i, \alpha, n) E_n^{l+1}, \quad \alpha \in \Sigma^\rho, i = 1, \dots, m(l).$$

Similarly, there exists $A_{l,l+1}^\eta(i, a, n)$, which takes 0 or 1, such that

$$\eta_a(E_i^l) = \sum_{n=1}^{m(l+1)} A_{l,l+1}^\eta(i, a, n) E_n^{l+1}, \quad a \in \Sigma^\eta, i = 1, \dots, m(l).$$

Set for $i = 1, \dots, m(l)$

$$\begin{aligned} \mathcal{F}_{j,k}(i) &= C^*(S_\mu T_\zeta E_i^l x E_i^l T_\xi^* S_\nu^* \mid \mu, \nu \in B_j(\Lambda_\rho), \zeta, \xi \in B_k(\Lambda_\eta), x \in \mathcal{A}), \\ &= C^*(T_\zeta S_\mu E_i^l x E_i^l S_\nu^* T_\xi^* \mid \mu, \nu \in B_j(\Lambda_\rho), \zeta, \xi \in B_k(\Lambda_\eta), x \in \mathcal{A}). \end{aligned}$$

Let $N_{j,k}(i)$ be the cardinal number of the finite set

$$\{(\mu, \zeta) \in B_j(\Lambda_\rho) \times B_k(\Lambda_\eta) \mid \rho_\mu(\eta_\zeta(1)) \geq E_i^l\}.$$

Since E_i^l is a central projection in \mathcal{A} , we have

Lemma 8.4. *For $j, k \in \mathbb{Z}_+$, put $l = j + k$. Then we have:*

(i) $\mathcal{F}_{j,k}(i)$ is isomorphic to the matrix algebra

$$M_{N_{j,k}(i)}(E_i^l \mathcal{A} E_i^l) (= M_{N_{j,k}(i)}(\mathbb{C}) \otimes E_i^l \mathcal{A} E_i^l)$$

over $E_i^l \mathcal{A} E_i^l$ for $i = 1, \dots, m(l)$.

(ii) $\mathcal{F}_{j,k} = \mathcal{F}_{j,k}(1) \oplus \dots \oplus \mathcal{F}_{j,k}(m(l))$.

Proof. (i) For $(\mu, \zeta) \in B_j(\Lambda_\rho) \times B_k(\Lambda_\eta)$ with $S_\mu T_\zeta E_i^l \neq 0$, one has

$$\eta_\zeta(\rho_\mu(1)) E_i^l \neq 0$$

so that $\eta_\zeta(\rho_\mu(1)) \geq E_i^l$. Hence $(S_\mu T_\zeta E_i^l)^* S_\mu T_\zeta E_i^l = E_i^l$. One sees that the set

$$\{S_\mu T_\zeta E_i^l \mid (\mu, \zeta) \in B_j(\Lambda_\rho) \times B_k(\Lambda_\eta); S_\mu T_\zeta E_i^l \neq 0\}$$

consist of partial isometries which give rise to matrix units of $\mathcal{F}_{j,k}(i)$ such that $\mathcal{F}_{j,k}(i)$ is isomorphic to $M_{N_{j,k}(i)}(E_i^l \mathcal{A} E_i^l)$.

(ii) Since $\mathcal{A} = E_1^l \mathcal{A} E_1^l \oplus \dots \oplus E_{m(l)}^l \mathcal{A} E_{m(l)}^l$, the assertion is easy. □

Define homomorphisms $\lambda_\rho, \lambda_\eta : K_0(\mathcal{A}) \rightarrow K_0(\mathcal{A})$ by setting

$$\lambda_\rho([p]) = \sum_{\alpha \in \Sigma^\rho} [(\rho_\alpha \otimes 1_n)(p)], \quad \lambda_\eta([p]) = \sum_{a \in \Sigma^\eta} [(\eta_a \otimes 1_n)(p)]$$

for a projection $p \in M_n(\mathcal{A})$ for some $n \in \mathbb{N}$. Recall that the identities (5.1), (5.2) give rise to the embeddings (5.3), which induce homomorphisms

$$K_0(\mathcal{F}_{j,k}) \longrightarrow K_0(\mathcal{F}_{j,k+1}), \quad K_0(\mathcal{F}_{j,k}) \longrightarrow K_0(\mathcal{F}_{j+1,k}).$$

We still denote them by $\iota_{*,+1}, \iota_{+1,*}$ respectively.

Lemma 8.5. *Assume that $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ forms square. There exists an isomorphism*

$$\Phi_{j,k} : K_0(\mathcal{F}_{j,k}) \longrightarrow K_0(\mathcal{A})$$

such that the following diagrams are commutative:

(i)

$$\begin{CD} K_0(\mathcal{F}_{j,k}) @>\iota_{+1,*}>> K_0(\mathcal{F}_{j+1,k}) \\ @V\Phi_{j,k}VV @VV\Phi_{j+1,k}V \\ K_0(\mathcal{A}) @>\lambda_\rho>> K_0(\mathcal{A}) \end{CD}$$

(ii)

$$\begin{CD} K_0(\mathcal{F}_{j,k}) @>\iota_{*,+1}>> K_0(\mathcal{F}_{j,k+1}) \\ @V\Phi_{j,k}VV @VV\Phi_{j,k+1}V \\ K_0(\mathcal{A}) @>\lambda_\eta>> K_0(\mathcal{A}). \end{CD}$$

Proof. Put for $i = 1, 2, \dots, m(l)$

$$P_i = \sum_{\mu \in B_j(\Lambda_\rho), \zeta \in B_k(\Lambda_\eta)} S_\mu T_\zeta E_i^l T_\zeta^* S_\mu^*.$$

Then P_i is a central projection in $\mathcal{F}_{j,k}$ such that $\sum_{i=1}^{m(l)} P_i = 1$. For $X \in \mathcal{F}_{j,k}$, one has $P_i X P_i \in \mathcal{F}_{j,k}(i)$ such that

$$X = \sum_{i=1}^{m(l)} P_i X P_i \in \bigoplus_{i=1}^{m(l)} \mathcal{F}_{j,k}(i).$$

Define an isomorphism

$$\varphi_{j,k} : X \in \mathcal{F}_{j,k} \longrightarrow \sum_{i=1}^{m(l)} P_i X P_i \in \bigoplus_{i=1}^{m(l)} \mathcal{F}_{j,k}(i)$$

which induces an isomorphism on their K-groups

$$\varphi_{j,k*} : K_0(\mathcal{F}_{j,k}) \longrightarrow \bigoplus_{i=1}^{m(l)} K_0(\mathcal{F}_{j,k}(i)).$$

Take and fix $\nu(i), \mu(i) \in B_j(\Lambda_\rho)$ and $\zeta(i), \xi(i) \in B_k(\Lambda_\eta)$ such that

$$(8.1) \quad T_{\xi(i)} S_{\nu(i)} = S_{\mu(i)} T_{\zeta(i)} \quad \text{and} \quad T_{\xi(i)} S_{\nu(i)} E_i^l \neq 0.$$

Hence $S_{\nu(i)}^* T_{\xi(i)}^* T_{\xi(i)} S_{\nu(i)} \geq E_i^l$. Since $\mathcal{F}_{j,k}(i)$ is isomorphic to

$$M_{N_{j,k(i)}}(\mathbb{C}) \otimes E_i^l \mathcal{A} E_i^l,$$

the embedding

$$\iota_{j,k}(i) : x \in E_i^l \mathcal{A} E_i^l \longrightarrow T_{\xi(i)} S_{\nu(i)} x S_{\nu(i)}^* T_{\xi(i)}^* \in \mathcal{F}_{j,k}(i)$$

induces an isomorphism on their K-groups

$$\iota_{j,k}(i)_* : K_0(E_i^l \mathcal{A} E_i^l) \longrightarrow K_0(\mathcal{F}_{j,k}(i)).$$

Put

$$\psi_{j,k} = \bigoplus_{i=1}^{m(l)} \iota_{j,k}(i) : \bigoplus_{i=1}^{m(l)} E_i^l \mathcal{A} E_i^l \longrightarrow \bigoplus_{i=1}^{m(l)} \mathcal{F}_{j,k}(i)$$

and hence we have an isomorphism

$$\psi_{j,k*} = \bigoplus_{i=1}^{m(l)} \iota_{j,k}(i)_* : \bigoplus_{i=1}^{m(l)} K_0(E_i^l \mathcal{A} E_i^l) \longrightarrow \bigoplus_{i=1}^{m(l)} K_0(\mathcal{F}_{j,k}(i)).$$

Since $K_0(\mathcal{A}) = \bigoplus_{i=1}^{m(l)} K_0(E_i^l \mathcal{A} E_i^l)$, we have an isomorphism

$$\Phi_{j,k} = \psi_{j,k*}^{-1} \circ \varphi_{j,k*} : K_0(\mathcal{F}_{j,k}) \xrightarrow{\varphi_{j,k*}} \bigoplus_{i=1}^{m(l)} K_0(\mathcal{F}_{j,k}(i)) \xrightarrow{\psi_{j,k*}^{-1}} K_0(\mathcal{A}).$$

(i) It suffices to show the following diagram

$$\begin{array}{ccc} K_0(\mathcal{F}_{j,k}) & \xrightarrow{\iota_{+1,*}} & K_0(\mathcal{F}_{j+1,k}) \\ \varphi_{j,k*} \downarrow & & \varphi_{j+1,k*} \downarrow \\ \bigoplus_{i=1}^{m(l)} K_0(\mathcal{F}_{j,k}(i)) & & \bigoplus_{i=1}^{m(l)} K_0(\mathcal{F}_{j+1,k}(i)) \\ \psi_{j,k*} \uparrow & & \psi_{j+1,k*} \uparrow \\ K_0(\mathcal{A}) & \xrightarrow{\lambda_\rho} & K_0(\mathcal{A}) \end{array}$$

is commutative. For $x = \sum_{i=1}^{m(l)} E_i^l x E_i^l \in \mathcal{A}$, we have

$$\psi_{j,k}(x) = \sum_{i=1}^{m(l)} T_{\xi(i)} S_{\nu(i)} E_i^l x E_i^l S_{\nu(i)}^* T_{\xi(i)}^* = \sum_{i=1}^{m(l)} S_{\mu(i)} T_{\zeta(i)} E_i^l x E_i^l T_{\zeta(i)}^* S_{\mu(i)}^*.$$

Since $P_i T_{\xi(i)} S_{\nu(i)} E_i^l x E_i^l S_{\nu(i)}^* T_{\xi(i)}^* P_i = T_{\xi(i)} S_{\nu(i)} E_i^l x E_i^l S_{\nu(i)}^* T_{\xi(i)}^*$, we have

$$\varphi_{j,k}^{-1} \circ \psi_{j,k}(x) = \sum_{i=1}^{m(l)} T_{\xi(i)} S_{\nu(i)} E_i^l x E_i^l S_{\nu(i)}^* T_{\xi(i)}^*$$

so that

$$\iota_{+1,*} \circ \varphi_{j,k}^{-1} \circ \psi_{j,k}(x) = \sum_{\alpha \in \Sigma^\rho} \sum_{i=1}^{m(l)} T_{\xi(i)} S_{\nu(i)\alpha} \rho_\alpha (E_i^l x E_i^l) S_{\nu(i)\alpha}^* T_{\xi(i)}^*.$$

Since

$$S_{\nu(i)\alpha}\rho_\alpha(E_i^l x E_i^l)S_{\nu(i)\alpha}^* = \sum_{n=1}^{m(l+1)} A_{l,l+1}^\rho(i, \alpha, n)S_{\nu(i)\alpha}E_n^{l+1}\rho_\alpha(x)E_n^{l+1}S_{\nu(i)\alpha}^*$$

and $A_{l,l+1}^\rho(i, \alpha, n)S_{\nu(i)\alpha}E_n^{l+1} = S_{\nu(i)\alpha}E_n^{l+1}$, we have

$$\sum_{\alpha \in \Sigma^\rho} S_{\nu(i)\alpha}\rho_\alpha(E_i^l x E_i^l)S_{\nu(i)\alpha}^* = \sum_{n=1}^{m(l+1)} \sum_{\alpha \in \Sigma^\rho} S_{\nu(i)\alpha}E_n^{l+1}\rho_\alpha(x)E_n^{l+1}S_{\nu(i)\alpha}^*$$

so that

$$\iota_{+1,*} \circ \varphi_{j,k}^{-1} \circ \psi_{j,k}(x) = \sum_{\alpha \in \Sigma^\rho} \sum_{i=1}^{m(l)} \sum_{n=1}^{m(l+1)} T_{\xi(i)}S_{\nu(i)\alpha}E_n^{l+1}\rho_\alpha(x)E_n^{l+1}S_{\nu(i)\alpha}^*T_{\xi(i)}^*.$$

On the other hand,

$$\begin{aligned} \psi_{j,k}(\lambda_\rho(x)) &= \psi_{j,k} \left(\sum_{n=1}^{m(l+1)} \sum_{\alpha \in \Sigma^\rho} E_n^{l+1}\rho_\alpha(x)E_n^{l+1} \right) \\ &= \sum_{\alpha \in \Sigma^\rho} \sum_{i=1}^{m(l)} \sum_{n=1}^{m(l+1)} T_{\xi(i)}S_{\nu(i)\alpha}E_n^{l+1}\rho_\alpha(x)E_n^{l+1}S_{\nu(i)\alpha}^*T_{\xi(i)}^*. \end{aligned}$$

Therefore we have

$$\iota_{+1,*} \circ \varphi_{j,k}^{-1} \circ \psi_{j,k}(x) = \psi_{j,k}(\lambda_\rho(x)).$$

(ii) is symmetric to (i). □

Define the abelian groups of the inductive limits:

$$G_\rho = \lim\{\lambda_\rho : K_0(\mathcal{A}) \longrightarrow K_0(\mathcal{A})\}, \quad G_\eta = \lim\{\lambda_\eta : K_0(\mathcal{A}) \longrightarrow K_0(\mathcal{A})\}.$$

Put the subalgebras of $\mathcal{F}_{\rho,\eta}$ for $j, k \in \mathbb{Z}_+$

$$\begin{aligned} \mathcal{F}_{\rho,k} &= C^*(T_\zeta S_\mu x S_\nu^* T_\xi^* \mid \mu, \nu \in B_*(\Lambda_\rho), |\mu| = |\nu|, \zeta, \xi \in B_k(\Lambda_\eta), x \in \mathcal{A}) \\ &= C^*(T_\zeta y T_\xi^* \mid \zeta, \xi \in B_k(\Lambda_\eta), y \in \mathcal{F}_\rho), \\ \mathcal{F}_{j,\eta} &= C^*(S_\mu T_\zeta x T_\xi^* S_\nu^* \mid \mu, \nu \in B_j(\Lambda_\rho), \zeta, \xi \in B_*(\Lambda_\eta), |\zeta| = |\xi|, x \in \mathcal{A}) \\ &= C^*(S_\mu y S_\nu^* \mid \mu, \nu \in B_j(\Lambda_\rho), y \in \mathcal{F}_\eta). \end{aligned}$$

By the preceding lemma, we have:

Lemma 8.6. *For $j, k \in \mathbb{Z}_+$, there exist isomorphisms*

$$\Phi_{\rho,k} : K_0(\mathcal{F}_{\rho,k}) \longrightarrow G_\rho, \quad \Phi_{j,\eta} : K_0(\mathcal{F}_{j,\eta}) \longrightarrow G_\eta$$

such that the following diagrams are commutative:

(i)

$$\begin{array}{ccccccc}
 K_0(\mathcal{F}_{j,k}) & \xrightarrow{\iota_{+1,*}} & K_0(\mathcal{F}_{j+1,k}) & \xrightarrow{\iota_{+1,*}} & \cdots & \xrightarrow{\iota_{+1,*}} & K_0(\mathcal{F}_{\rho,k}) \\
 \Phi_{j,k} \downarrow & & \Phi_{j+1,k} \downarrow & & & & \Phi_{\rho,k} \downarrow \\
 K_0(\mathcal{A}) & \xrightarrow{\lambda_\rho} & K_0(\mathcal{A}) & \xrightarrow{\lambda_\rho} & \cdots & \xrightarrow{\lambda_\rho} & G_\rho
 \end{array}$$

(ii)

$$\begin{array}{ccccccc}
 K_0(\mathcal{F}_{j,k}) & \xrightarrow{\iota_{*,+1}} & K_0(\mathcal{F}_{j,k+1}) & \xrightarrow{\iota_{*,+1}} & \cdots & \xrightarrow{\iota_{*,+1}} & K_0(\mathcal{F}_{j,\eta}) \\
 \Phi_{j,k} \downarrow & & \Phi_{j,k+1} \downarrow & & & & \Phi_{j,\eta} \downarrow \\
 K_0(\mathcal{A}) & \xrightarrow{\lambda_\eta} & K_0(\mathcal{A}) & \xrightarrow{\lambda_\eta} & \cdots & \xrightarrow{\lambda_\eta} & G_\eta.
 \end{array}$$

Lemma 8.7. *If $\xi = (\xi_1, \dots, \xi_k) \in B_k(\Lambda_\eta), \nu = (\nu_1, \dots, \nu_j) \in B_j(\Lambda_\rho)$ satisfy the condition $\rho_\nu(\eta_\xi(1)) \geq E_i^l$ for some $i = 1, \dots, m(l)$ with $l = j + k$, then $T_{\xi_1}^* T_\xi S_\nu E_i^l = T_{\bar{\xi}} S_\nu E_i^l$ where $\bar{\xi} = (\xi_2, \dots, \xi_k)$.*

Proof. Since $T_{\xi_1}^* T_\xi = T_{\xi_1}^* T_{\xi_1} T_\xi T_{\bar{\xi}}^* T_{\bar{\xi}} = T_{\bar{\xi}} T_{\bar{\xi}}^* T_{\xi_1}^* T_{\xi_1} T_\xi = T_{\bar{\xi}} T_\xi^* T_\xi$, we have

$$T_{\xi_1}^* T_\xi S_\nu E_i^l = T_{\bar{\xi}} S_\nu S_\nu^* T_\xi^* T_\xi S_\nu E_i^l = T_{\bar{\xi}} S_\nu \rho_\nu(\eta_\xi(1)) E_i^l = T_{\bar{\xi}} S_\nu E_i^l. \quad \square$$

Let us denote by γ_ρ, γ_η the endomorphisms $\gamma_{\rho,0}, \gamma_{\eta,0}$ on $K_0(\mathcal{F}_{\rho,\eta})$ appeared in Lemma 7.6, respectively.

Lemma 8.8. *For $k, j \in \mathbb{Z}_+$, we have:*

(i) *The restriction of γ_η^{-1} to $K_0(\mathcal{F}_{j,k})$ makes the following diagram commutative:*

$$\begin{array}{ccc}
 K_0(\mathcal{F}_{j,k}) & \xrightarrow{\gamma_\eta^{-1}} & K_0(\mathcal{F}_{j,k-1}) & \xrightarrow{\iota_{*,+1}} & K_0(\mathcal{F}_{j,k}) \\
 \Phi_{j,k} \downarrow & & & & \Phi_{j,k} \downarrow \\
 K_0(\mathcal{A}) & & \xrightarrow{\lambda_\eta} & & K_0(\mathcal{A}).
 \end{array}$$

(ii) *The restriction of γ_ρ^{-1} to $K_0(\mathcal{F}_{j,k})$ makes the following diagram commutative:*

$$\begin{array}{ccc}
 K_0(\mathcal{F}_{j,k}) & \xrightarrow{\gamma_\rho^{-1}} & K_0(\mathcal{F}_{j-1,k}) & \xrightarrow{\iota_{+1,*}} & K_0(\mathcal{F}_{j,k}) \\
 \Phi_{j,k} \downarrow & & & & \Phi_{j,k} \downarrow \\
 K_0(\mathcal{A}) & & \xrightarrow{\lambda_\rho} & & K_0(\mathcal{A}).
 \end{array}$$

Proof. (i) Put $l = j + k$. Take a projection $p \in M_n(\mathcal{A})$ for some $n \in \mathbb{N}$. Since $\mathcal{A} \otimes M_n(\mathbb{C}) = \sum_{i=1}^{m(l)} (E_i^l \otimes 1)(\mathcal{A} \otimes M_n)(E_i^l \otimes 1)$, by putting

$$p_i^l = (E_i^l \otimes 1)p(E_i^l \otimes 1) \in M_n(E_i^l \mathcal{A} E_i^l),$$

we have $p = \sum_{i=1}^{m(l)} p_i^l$. Take

$$\xi(i) = (\xi_1(i), \dots, \xi_k(i)) \in B_k(\Lambda_\eta), \quad \nu(i) = (\nu_1(i), \dots, \nu_j(i)) \in B_j(\Lambda_\rho)$$

as in (8.1) so that $\rho_{\nu(i)}(\eta_{\xi(i)}(1)) \geq E_i^l$ and put $\bar{\xi}(i) = (\xi_2(i), \dots, \xi_k(i))$ so that $\xi(i) = \xi_1(i)\bar{\xi}(i)$. We have

$$\psi_{j,k^*}([p]) = \sum_{i=1}^{m(l)} \oplus [(T_{\xi(i)}S_{\nu(i)} \otimes 1_n)p_i^l(S_{\nu(i)}^*T_{\xi(i)}^* \otimes 1_n)] \in \bigoplus_{i=1}^{m(l)} K_0(\mathcal{F}_{j,k}(i)).$$

As

$$(T_{\xi(i)}S_{\nu(i)} \otimes 1_n)p_i^l(S_{\nu(i)}^*T_{\xi(i)}^* \otimes 1_n) \leq T_{\xi_1(i)}T_{\xi_1(i)}^* \otimes 1_n,$$

by the preceding lemma we have

$$T_{\xi_1(i)}^*T_{\xi(i)}S_{\nu(i)}E_i^l = T_{\bar{\xi}(i)}S_{\nu(i)}E_i^l$$

so that by Lemma 7.6

$$\gamma_\eta^{-1}([(T_{\xi(i)}S_{\nu(i)} \otimes 1_n)p_i^l(S_{\nu(i)}^*T_{\xi(i)}^* \otimes 1_n)]) = [(T_{\bar{\xi}(i)}S_{\nu(i)} \otimes 1_n)p_i^l(S_{\nu(i)}^*T_{\bar{\xi}(i)}^* \otimes 1_n)].$$

Hence $K_0(\mathcal{F}_{j,k})$ goes to $K_0(\mathcal{F}_{j,k-1})$ by the homomorphism γ_η^{-1} . Take $\mu(i) \in B_j(\Lambda_\rho)$, $\bar{\zeta}(i) \in B_{k-1}(\Lambda_\eta)$ such that $T_{\bar{\xi}(i)}S_{\nu(i)} = S_{\mu(i)}T_{\bar{\zeta}(i)}$ for $i = 1, \dots, m(l)$. The element

$$\begin{aligned} & \sum_{i=1}^{m(l)} [(T_{\bar{\xi}(i)}S_{\nu(i)} \otimes 1_n)p_i^l(S_{\nu(i)}^*T_{\bar{\xi}(i)}^* \otimes 1_n)] \\ &= \sum_{i=1}^{m(l)} [(S_{\mu(i)}T_{\bar{\zeta}(i)} \otimes 1_n)p_i^l(T_{\bar{\zeta}(i)}^*S_{\mu(i)}^* \otimes 1_n)] \in K_0(\mathcal{F}_{j,k-1}) \end{aligned}$$

goes to

$$\sum_{i=1}^{m(l)} \sum_{a \in \Sigma^\eta} [(S_{\mu(i)}T_{\bar{\zeta}(i)a} \otimes 1_n)(T_a^* \otimes 1_n)p_i^l(T_a \otimes 1_n)(T_{\bar{\zeta}(i)a}^*S_{\mu(i)}^* \otimes 1_n)] \in K_0(\mathcal{F}_{j,k})$$

by $\iota_{*,+1}$. The latter one is expressed as

$$(8.2) \quad \sum_{h=1}^{m(l)} \oplus \sum_{i=1}^{m(l)} \sum_{a \in \Sigma^\eta} [(S_{\mu(i)}T_{\bar{\zeta}(i)a} \otimes 1_n)E_h^l(T_a^* \otimes 1_n)p_i^l(T_a \otimes 1_n)E_h^l(T_{\bar{\zeta}(i)a}^*S_{\mu(i)}^* \otimes 1_n)]$$

in $\bigoplus_{h=1}^{m(l)} K_0(\mathcal{F}_{j,k}(h))$. On the other hand, we have

$$\begin{aligned} \lambda_\eta([p]) &= \sum_{a \in \Sigma^\eta} [(T_a^* \otimes 1_n)p(T_a \otimes 1_n)] \\ &= \sum_{h=1}^{m(l)} \oplus \sum_{a \in \Sigma^\eta} [E_h^l(T_a^* \otimes 1_n)p(T_a \otimes 1_n)E_h^l] \in \bigoplus_{h=1}^{m(l)} K_0(E_h^l \mathcal{A} E_h^l), \end{aligned}$$

which is expressed as

$$\begin{aligned} & \sum_{h=1}^{m(l)} \oplus \sum_{a \in \Sigma^\eta} [(T_{\xi(h)} S_{\nu(h)} E_h^l \otimes 1_n)(T_a^* \otimes 1_n) p(T_a \otimes 1_n)(E_h^l S_{\nu(h)}^* T_{\xi(h)}^* \otimes 1_n)] \\ &= \sum_{h=1}^{m(l)} \oplus \sum_{a \in \Sigma^\eta} \sum_{i=1}^{m(l)} \left[(T_{\xi(h)} S_{\nu(h)} E_h^l \otimes 1_n)(T_a^* \otimes 1_n) \right. \\ & \quad \left. \cdot p_i^l(T_a \otimes 1_n)(E_h^l S_{\nu(h)}^* T_{\xi(h)}^* \otimes 1_n) \right] \end{aligned}$$

in $\bigoplus_{h=1}^{m(l)} K_0(\mathcal{F}_{j,k}(h))$. Take $\mu'(h) \in B_j(\Lambda_\rho), \zeta'(h) \in B_k(\Lambda_\eta)$ such that $T_{\xi(h)} S_{\nu(h)} = S_{\mu'(h)} T_{\zeta'(h)}$ so that the above element is

(8.3)

$$\sum_{h=1}^{m(l)} \oplus \sum_{i=1}^{m(l)} \sum_{a \in \Sigma^\eta} [(S_{\mu'(h)} T_{\zeta'(h)} E_h^l \otimes 1_n)(T_a^* \otimes 1_n) p_i^l(T_a \otimes 1_n)(E_h^l T_{\zeta'(h)}^* S_{\nu'(h)}^* \otimes 1_n)]$$

in $\bigoplus_{h=1}^{m(l)} K_0(\mathcal{F}_{j,k}(h))$. Since for $h, i = 1, \dots, m(l), a \in \Sigma^\eta$ their classes of the K-groups coincide such as

$$\begin{aligned} & [(S_{\mu(i)} T_{\zeta(i)a} \otimes 1_n) E_h^l (T_a^* \otimes 1_n) p_i^l(T_a \otimes 1_n) E_h^l (T_{\zeta(i)a}^* S_{\mu(i)}^* \otimes 1_n)] \\ &= [(S_{\mu'(h)} T_{\zeta'(h)} E_h^l \otimes 1_n)(T_a^* \otimes 1_n) p_i^l(T_a \otimes 1_n)(E_h^l T_{\zeta'(h)}^* S_{\nu'(h)}^* \otimes 1_n)] \\ & \in K_0(\mathcal{F}_{j,k}(h)), \end{aligned}$$

the element of (8.2) is equal to the element of (8.3) in $K_0(\mathcal{F}_{j,k})$. Thus (i) holds.

(ii) is similar to (i). □

We note that for $j, k \in \mathbb{Z}_+$,

$$\begin{aligned} K_0(\mathcal{F}_{\rho,k}) &= \lim_j \{ \iota_{+1,*} : K_0(\mathcal{F}_{j,k}) \longrightarrow K_0(\mathcal{F}_{j+1,k}) \}, \\ K_0(\mathcal{F}_{j,\eta}) &= \lim_k \{ \iota_{*,+1} : K_0(\mathcal{F}_{j,k}) \longrightarrow K_0(\mathcal{F}_{j,k+1}) \}. \end{aligned}$$

The following lemma is direct.

Lemma 8.9. *For $k, j \in \mathbb{Z}_+$, the following diagrams are commutative:*

(i)

$$\begin{array}{ccc} K_0(\mathcal{F}_{j,k}) & \xrightarrow{\gamma_\eta^{-1}} & K_0(\mathcal{F}_{j,k-1}) \\ \iota_{+1,*} \downarrow & & \iota_{+1,*} \downarrow \\ K_0(\mathcal{F}_{j+1,k}) & \xrightarrow{\gamma_\eta^{-1}} & K_0(\mathcal{F}_{j+1,k-1}). \end{array}$$

Hence γ_η^{-1} yields a homomorphism from $K_0(\mathcal{F}_{\rho,k})$ to $K_0(\mathcal{F}_{\rho,k-1})$.

(ii)

$$\begin{CD} K_0(\mathcal{F}_{j,k}) @>\gamma_\rho^{-1}>> K_0(\mathcal{F}_{j-1,k}) \\ @V\iota_{*,+1}VV @VV\iota_{*,+1}V \\ K_0(\mathcal{F}_{j,k+1}) @>\gamma_\rho^{-1}>> K_0(\mathcal{F}_{j-1,k+1}). \end{CD}$$

Hence γ_ρ^{-1} yields a homomorphism from $K_0(\mathcal{F}_{j,\eta})$ to $K_0(\mathcal{F}_{j-1,\eta})$.

The homomorphisms

$$\iota_{+1,*} : K_0(\mathcal{F}_{j,k}) \longrightarrow K_0(\mathcal{F}_{j+1,k}), \quad \iota_{*,+1} : K_0(\mathcal{F}_{j,k}) \longrightarrow K_0(\mathcal{F}_{j,k+1})$$

are naturally induce homomorphisms

$$K_0(\mathcal{F}_{j,\eta}) \longrightarrow K_0(\mathcal{F}_{j+1,\eta}), \quad \iota_{*,+1} : K_0(\mathcal{F}_{\rho,k}) \longrightarrow K_0(\mathcal{F}_{\rho,k+1})$$

which we denote by $\iota_{+1,\eta}$, $\iota_{\rho,+1}$ respectively. They are also induced by the identities (5.1), (5.2) respectively.

Lemma 8.10. *For $k, j \in \mathbb{Z}_+$, the following diagrams are commutative:*

(i)

$$\begin{CD} K_0(\mathcal{F}_{\rho,k}) @>\gamma_\eta^{-1}>> K_0(\mathcal{F}_{\rho,k-1}) \\ @V\iota_{\rho,+1}VV @VV\iota_{\rho,+1}V \\ K_0(\mathcal{F}_{\rho,k+1}) @>\gamma_\eta^{-1}>> K_0(\mathcal{F}_{\rho,k}). \end{CD}$$

(ii)

$$\begin{CD} K_0(\mathcal{F}_{j,\eta}) @>\gamma_\rho^{-1}>> K_0(\mathcal{F}_{j-1,\eta}) \\ @V\iota_{+1,\eta}VV @VV\iota_{+1,\eta}V \\ K_0(\mathcal{F}_{j+1,\eta}) @>\gamma_\rho^{-1}>> K_0(\mathcal{F}_{j,\eta}). \end{CD}$$

Proof. (i) As in the proof of Lemma 8.9, one may take an element of $K_0(\mathcal{F}_{\rho,k})$ as in the following form:

$$\sum_{i=1}^{m(l)} \oplus [(T_{\xi(i)} S_{\nu(i)} \otimes 1_n) p_i^l (S_{\nu(i)}^* T_{\xi(i)}^* \otimes 1_n)] \in \bigoplus_{i=1}^{m(l)} K_0(\mathcal{F}_{j,k}(i))$$

for some projection $p \in M_n(\mathcal{A})$ and j, l with $l = j + k$, where

$$p_i^l = (E_i^l \otimes 1) p (E_i^l \otimes 1) \in M_n(E_i^l \mathcal{A} E_i^l).$$

Let $\xi(i) = \xi_1(i) \bar{\xi}(i)$ with $\xi_1(i) \in \Sigma^\eta$, $\bar{\xi}(i) \in B_{k-1}(\Lambda_\eta)$. One may assume that $T_{\xi(i)} S_{\nu(i)} \neq 0$ so that $T_{\xi(i)} S_{\nu(i)} = S_{\nu(i)'} T_{\bar{\xi}(i)'}$ for some $\nu(i)' \in B_j(\Lambda_\rho)$, $\bar{\xi}(i)' \in$

$B_{k-1}(\Lambda_\eta)$. As in the proof of Lemma 8.9, one has

$$\begin{aligned} & \gamma_\eta^{-1}([(T_{\xi(i)}S_{\nu(i)} \otimes 1_n)p_i^l(S_{\nu(i)}^*T_{\xi(i)}^* \otimes 1_n)]) \\ &= [(T_{\bar{\xi}(i)}S_{\nu(i)} \otimes 1_n)p_i^l(S_{\nu(i)}^*T_{\bar{\xi}(i)}^* \otimes 1_n)] \\ &= [(S_{\nu(i)'}T_{\bar{\xi}(i)'} \otimes 1_n)p_i^l(S_{\nu(i)'}^*T_{\bar{\xi}(i)'}^* \otimes 1_n)]. \end{aligned}$$

Hence we have

$$\begin{aligned} & \iota_{*,+1} \circ \gamma_\eta^{-1}([(T_{\xi(i)}S_{\nu(i)} \otimes 1_n)p_i^l(S_{\nu(i)}^*T_{\xi(i)}^* \otimes 1_n)]) \\ &= \iota_{*,+1}([(S_{\nu(i)'}T_{\bar{\xi}(i)'} \otimes 1_n)p_i^l(T_{\bar{\xi}(i)'}^*S_{\nu(i)'}^* \otimes 1_n)]) \\ &= \sum_{b \in \Sigma^n} [(S_{\nu(i)'}T_{\bar{\xi}(i)'}b \otimes 1_n)(T_b^* \otimes 1_n)p_i^l(T_b \otimes 1_n)(T_{\bar{\xi}(i)'}^*bS_{\nu(i)'}^* \otimes 1_n)]. \end{aligned}$$

On the other hand, the equality $T_{\xi(i)}S_{\nu(i)} = T_{\xi(i)_1}S_{\nu(i)'}T_{\bar{\xi}(i)'}$ implies

$$\begin{aligned} & \iota_{*,+1}([(T_{\xi(i)}S_{\nu(i)} \otimes 1_n)p_i^l(S_{\nu(i)}^*T_{\xi(i)}^* \otimes 1_n)]) \\ &= \sum_{b \in \Sigma^n} [(T_{\xi(i)_1}S_{\nu(i)'}T_{\bar{\xi}(i)'}b \otimes 1_n)(T_b^* \otimes 1_n)p_i^l(T_b \otimes 1_n)(T_{\bar{\xi}(i)'}^*bS_{\nu(i)'}^*T_{\xi(i)_1}^* \otimes 1_n)] \end{aligned}$$

and hence

$$\begin{aligned} & \gamma_\eta^{-1} \circ \iota_{*,+1}([(T_{\xi(i)}S_{\nu(i)} \otimes 1_n)p_i^l(S_{\nu(i)}^*T_{\xi(i)}^* \otimes 1_n)]) \\ &= \sum_{b \in \Sigma^n} \gamma_\eta^{-1} \left([(T_{\xi(i)_1}S_{\nu(i)'}T_{\bar{\xi}(i)'}b \otimes 1_n)(T_b^* \otimes 1_n) \right. \\ & \quad \left. \cdot p_i^l(T_b \otimes 1_n)(T_{\bar{\xi}(i)'}^*bS_{\nu(i)'}^*T_{\xi(i)_1}^* \otimes 1_n)] \right) \\ &= \sum_{b \in \Sigma^n} [(S_{\nu(i)'}T_{\bar{\xi}(i)'}b \otimes 1_n)(T_b^* \otimes 1_n)p_i^l(T_b \otimes 1_n)(T_{\bar{\xi}(i)'}^*bS_{\nu(i)'}^* \otimes 1_n)]. \end{aligned}$$

(ii) The proof is completely symmetric to the above proof. □

Since the homomorphisms $\lambda_\rho, \lambda_\eta : K_0(\mathcal{A}) \rightarrow K_0(\mathcal{A})$ are mutually commutative, the map λ_η induces a homomorphism on the inductive limit $G_\rho = \lim\{\lambda_\rho : K_0(\mathcal{A}) \rightarrow K_0(\mathcal{A})\}$ and similarly λ_ρ does on the inductive limit G_η . They are still denoted by $\lambda_\rho, \lambda_\eta$ respectively.

Lemma 8.11. *For $k, j \in \mathbb{Z}_+$, the following diagrams are commutative:*

(i)

$$\begin{array}{ccccc} K_0(\mathcal{F}_{\rho,k}) & \xrightarrow{\gamma_\eta^{-1}} & K_0(\mathcal{F}_{\rho,k-1}) & \xrightarrow{\iota_{\rho,+1}} & K_0(\mathcal{F}_{\rho,k}) \\ \Phi_{\rho,k} \downarrow & & & & \Phi_{\rho,k} \downarrow \\ G_\rho & & \xrightarrow{\lambda_\eta} & & G_\rho. \end{array}$$

(ii)

$$\begin{array}{ccccc}
 K_0(\mathcal{F}_{j,\eta}) & \xrightarrow{\gamma_\rho^{-1}} & K_0(\mathcal{F}_{j-1,\eta}) & \xrightarrow{\iota_{+1,\eta}} & K_0(\mathcal{F}_{j,\eta}) \\
 \Phi_{j,\eta} \downarrow & & & & \Phi_{j,\eta} \downarrow \\
 G_\eta & & \xrightarrow{\lambda_\rho} & & G_\eta.
 \end{array}$$

Proof. (i) As in the proof of Lemma 8.8 and Lemma 8.10 one may take an element of $K_0(\mathcal{F}_{\rho,k})$ as in the following form:

$$\sum_{i=1}^{m(l)} \oplus [(T_{\xi(i)} S_{\nu(i)} \otimes 1_n) p_i^l (S_{\nu(i)}^* T_{\xi(i)}^* \otimes 1_n)] \in \bigoplus_{i=1}^{m(l)} K_0(\mathcal{F}_{j,k}(i))$$

for some projection $p \in M_n(\mathcal{A})$ and j, l with $l = j + k$, where

$$p_i^l = (E_i^l \otimes 1) p (E_i^l \otimes 1).$$

Keep the notations as in the proof of Lemma 8.8, we have

$$\begin{aligned}
 & \iota_{*,+1} \circ \gamma_\eta^{-1}([(T_{\xi(i)} S_{\nu(i)} \otimes 1_n) p_i^l (S_{\nu(i)}^* T_{\xi(i)}^* \otimes 1_n)]) \\
 &= \sum_{b \in \Sigma^n} [(S_{\nu(i)'} T_{\bar{\xi}(i)'}^* \otimes 1_n) (T_b^* \otimes 1_n) p_i^l (T_b \otimes 1_n) (T_{\bar{\xi}(i)'}^* S_{\nu(i)'}^* \otimes 1_n)]
 \end{aligned}$$

so that

$$\begin{aligned}
 & \Phi_{\rho,k} \circ \iota_{*,+1} \circ \gamma_\eta^{-1}([(T_{\xi(i)} S_{\nu(i)} \otimes 1_n) p_i^l (S_{\nu(i)}^* T_{\xi(i)}^* \otimes 1_n)]) \\
 &= \sum_{b \in \Sigma^n} \Phi_{\rho,k}([(S_{\nu(i)'} T_{\bar{\xi}(i)'}^* \otimes 1_n) (T_b^* \otimes 1_n) p_i^l (T_b \otimes 1_n) (T_{\bar{\xi}(i)'}^* S_{\nu(i)'}^* \otimes 1_n)]) \\
 &= \sum_{b \in \Sigma^n} [(T_b^* \otimes 1_n) p_i^l (T_b \otimes 1_n)] \\
 &= \lambda_\eta([p_i^l]) = (\lambda_\eta \circ \Phi_{\rho,k})([(T_{\xi(i)} S_{\nu(i)} \otimes 1_n) p_i^l (S_{\nu(i)}^* T_{\xi(i)}^* \otimes 1_n)]).
 \end{aligned}$$

Therefore we have $\Phi_{\rho,k} \circ \iota_{\rho,+1} \circ \gamma_\eta^{-1} = \lambda_\eta \circ \Phi_{\rho,k}$.

(ii) The proof is completely symmetric to the above proof. □

Put for $j, k \in \mathbb{Z}_+$

$$\begin{aligned}
 G_{\rho,k} &= K_0(\mathcal{F}_{\rho,k}) (\cong G_\rho = \lim\{\lambda_\rho : K_0(\mathcal{A}) \longrightarrow K_0(\mathcal{A})\}), \\
 G_{j,\eta} &= K_0(\mathcal{F}_{j,\eta}) (\cong G_\eta = \lim\{\lambda_\eta : K_0(\mathcal{A}) \longrightarrow K_0(\mathcal{A})\}).
 \end{aligned}$$

The map $\lambda_\eta : K_0(\mathcal{A}) \longrightarrow K_0(\mathcal{A})$ naturally gives rise to a homomorphism from $G_{\rho,k}$ to $G_{\rho,k+1}$ which we will still denote by λ_η . Similarly we have a homomorphism λ_ρ from $G_{j,\eta}$ to $G_{j+1,\eta}$.

Lemma 8.12. For $k, j \in \mathbb{Z}_+$, the following diagrams are commutative:

$$\begin{array}{ccc}
 \text{(i)} & & \\
 & & K_0(\mathcal{F}_{\rho,k}) \xrightarrow{\iota_{\rho,+1}} K_0(\mathcal{F}_{\rho,k+1}) \\
 & & \parallel \qquad \qquad \qquad \parallel \\
 & & G_{\rho,k} \xrightarrow{\lambda_\eta} G_{\rho,k+1} \\
 \text{(ii)} & & \\
 & & K_0(\mathcal{F}_{j,\eta}) \xrightarrow{\iota_{+1,\eta}} K_0(\mathcal{F}_{j+1,\eta}) \\
 & & \parallel \qquad \qquad \qquad \parallel \\
 & & G_{j,\eta} \xrightarrow{\lambda_\rho} G_{j+1,\eta}
 \end{array}$$

We denote the abelian group $K_0(\mathcal{F}_{\rho,\eta})$ by $G_{\rho,\eta}$. Since

$$\begin{aligned}
 K_0(\mathcal{F}_{\rho,\eta}) &= \lim_k \{ \iota_{\rho,+1} : K_0(\mathcal{F}_{\rho,k}) \longrightarrow K_0(\mathcal{F}_{\rho,k+1}) \} \\
 &= \lim_j \{ \iota_{+1,\eta} : K_0(\mathcal{F}_{j,\eta}) \longrightarrow K_0(\mathcal{F}_{j+1,\eta}) \},
 \end{aligned}$$

one has

$$G_{\rho,\eta} = \lim_k \{ \lambda_\eta : G_{\rho,k} \longrightarrow G_{\rho,k+1} \} = \lim_j \{ \lambda_\rho : G_{j,\eta} \longrightarrow G_{j+1,\eta} \}.$$

Define two endomorphisms

$$\begin{aligned}
 \sigma_\eta \text{ on } G_{\rho,\eta} &= \lim_k \{ \lambda_\eta : G_{\rho,k} \longrightarrow G_{\rho,k+1} \} \quad \text{and} \\
 \sigma_\rho \text{ on } G_{\rho,\eta} &= \lim_j \{ \lambda_\rho : G_{j,\eta} \longrightarrow G_{j+1,\eta} \}
 \end{aligned}$$

by setting

$$\begin{aligned}
 \sigma_\rho : [g, k] \in G_{\rho,k} &\longrightarrow [g, k - 1] \in G_{\rho,k-1} \text{ for } g \in G_\rho \text{ and} \\
 \sigma_\eta : [h, j] \in G_{j,\eta} &\longrightarrow [h, j - 1] \in G_{j-1,\eta} \text{ for } h \in G_\eta.
 \end{aligned}$$

Therefore we have:

Lemma 8.13.

(i) *There exists an isomorphism $\Phi_{\rho,\infty} : K_0(\mathcal{F}_{\rho,\eta}) \longrightarrow G_{\rho,\eta}$ such that the following diagrams are commutative:*

$$\begin{array}{ccc}
 K_0(\mathcal{F}_{\rho,\eta}) & \xrightarrow{\gamma_\eta^{-1}} & K_0(\mathcal{F}_{\rho,\eta}) \\
 \Phi_{\rho,\infty} \downarrow & & \Phi_{\rho,\infty} \downarrow \\
 G_{\rho,\eta} & \xrightarrow{\sigma_\eta} & G_{\rho,\eta}
 \end{array}$$

and hence

$$\begin{array}{ccc}
 K_0(\mathcal{F}_{\rho,\eta}) & \xrightarrow{id - \gamma_\eta^{-1}} & K_0(\mathcal{F}_{\rho,\eta}) \\
 \Phi_{\rho,\infty} \downarrow & & \Phi_{\rho,\infty} \downarrow \\
 G_{\rho,\eta} & \xrightarrow{id - \sigma_\eta} & G_{\rho,\eta}
 \end{array}$$

(ii) *There exists an isomorphism $\Phi_{\infty,\eta} : K_0(\mathcal{F}_{\rho,\eta}) \longrightarrow G_{\rho,\eta}$ such that the following diagrams are commutative:*

$$\begin{CD} K_0(\mathcal{F}_{\rho,\eta}) @>\gamma_\rho^{-1}>> K_0(\mathcal{F}_{\rho,\eta}) \\ @V\Phi_{\infty,\eta}VV @VV\Phi_{\infty,\eta}V \\ G_{\rho,\eta} @>\sigma_\rho>> G_{\rho,\eta} \end{CD}$$

and hence

$$\begin{CD} K_0(\mathcal{F}_{\rho,\eta}) @>id-\gamma_\rho^{-1}>> K_0(\mathcal{F}_{\rho,\eta}) \\ @V\Phi_{\infty,\eta}VV @VV\Phi_{\infty,\eta}V \\ G_{\rho,\eta} @>id-\sigma_\rho>> G_{\rho,\eta}. \end{CD}$$

Let us denote by $J_{\mathcal{A}}$ the natural embedding $\mathcal{A} = \mathcal{F}_{0,0} \hookrightarrow \mathcal{F}_{\rho,\eta}$, which induces a homomorphism $J_{\mathcal{A}*} : K_0(\mathcal{A}) \longrightarrow K_0(\mathcal{F}_{\rho,\eta})$.

Lemma 8.14. *The homomorphism $J_{\mathcal{A}*} : K_0(\mathcal{A}) \longrightarrow K_0(\mathcal{F}_{\rho,\eta})$ is injective such that*

$$J_{\mathcal{A}*} \circ \lambda_\rho = \gamma_\rho^{-1} \circ J_{\mathcal{A}*} \quad \text{and} \quad J_{\mathcal{A}*} \circ \lambda_\eta = \gamma_\eta^{-1} \circ J_{\mathcal{A}*}.$$

Proof. We will first show that the endomorphisms $\lambda_\rho, \lambda_\eta$ on $K_0(\mathcal{A})$ are both injective. Put a projection $Q_\alpha = S_\alpha S_\alpha^*$ and a subalgebra $\mathcal{A}_\alpha = \rho_\alpha(\mathcal{A})$ of \mathcal{A} for $\alpha \in \Sigma^\rho$. Then the endomorphism ρ_α on \mathcal{A} extends to an isomorphism from $\mathcal{A}Q_\alpha$ onto \mathcal{A}_α by setting $\rho_\alpha(x) = S_\alpha^* x S_\alpha, x \in \mathcal{A}Q_\alpha$ whose inverse is $\phi_\alpha : \mathcal{A}_\alpha \longrightarrow \mathcal{A}Q_\alpha$ defined by $\phi_\alpha(y) = S_\alpha y S_\alpha^*, y \in \mathcal{A}_\alpha$. Hence the induced homomorphism $\rho_{\alpha*} : K_0(\mathcal{A}Q_\alpha) \longrightarrow K_0(\mathcal{A}_\alpha)$ is an isomorphism. Since $\mathcal{A} = \bigoplus_{\alpha \in \Sigma^\rho} Q_\alpha \mathcal{A}$, the homomorphism

$$\sum_{\alpha \in \Sigma^\rho} \phi_{\alpha*} \circ \rho_{\alpha*} : K_0(\mathcal{A}) \longrightarrow \bigoplus_{\alpha \in \Sigma^\rho} K_0(Q_\alpha \mathcal{A})$$

is an isomorphism, one may identify $K_0(\mathcal{A}) = \bigoplus_{\alpha \in \Sigma^\rho} K_0(Q_\alpha \mathcal{A})$. Let $g \in K_0(\mathcal{A})$ satisfy $\lambda_\rho(g) = 0$. Put $g_\alpha = \phi_{\alpha*} \circ \rho_{\alpha*}(g) \in K_0(Q_\alpha \mathcal{A})$ for $\alpha \in \Sigma^\rho$ so that $g = \sum_{\alpha \in \Sigma^\rho} g_\alpha$. As $\rho_{\beta*} \circ \phi_{\alpha*} = 0$ for $\beta \neq \alpha$, one sees $\rho_{\beta*}(g_\alpha) = 0$ for $\beta \neq \alpha$. Hence

$$0 = \lambda_\rho(g) = \sum_{\beta \in \Sigma^\rho} \sum_{\alpha \in \Sigma^\rho} \rho_{\beta*}(g_\alpha) = \sum_{\alpha \in \Sigma^\rho} \rho_{\alpha*}(g_\alpha) \in \bigoplus_{\alpha \in \Sigma^\rho} K_0(\mathcal{A}_\alpha).$$

It follows that $\rho_{\alpha*}(g_\alpha) = 0$ in $K_0(\mathcal{A}_\alpha)$. Since $\rho_{\alpha*} : K_0(Q_\alpha \mathcal{A}) \longrightarrow K_0(\mathcal{A}_\alpha)$ is isomorphic, one sees that $g_\alpha = 0$ in $K_0(Q_\alpha \mathcal{A})$ for all $\alpha \in \Sigma^\rho$. This implies that $g = \sum_{\alpha \in \Sigma^\rho} g_\alpha = 0$ in $K_0(\mathcal{A})$. Therefore the endomorphism λ_ρ on $K_0(\mathcal{A})$ is injective, and similarly so is λ_η .

By the previous lemma, there exists an isomorphism $\Phi_{j,k} : K_0(\mathcal{F}_{j,k}) \rightarrow K_0(\mathcal{A})$ such that the diagram

$$\begin{array}{ccc} K_0(\mathcal{F}_{j,k}) & \xrightarrow{\iota_{+1,*}} & K_0(\mathcal{F}_{j+1,k}) \\ \Phi_{j,k} \downarrow & & \Phi_{j+1,k} \downarrow \\ K_0(\mathcal{A}) & \xrightarrow{\lambda_\rho} & K_0(\mathcal{A}) \end{array}$$

is commutative so that the embedding $\iota_{+1,*} : K_0(\mathcal{F}_{j,k}) \rightarrow K_0(\mathcal{F}_{j+1,k})$ is injective, and similarly $\iota_{*,+1} : K_0(\mathcal{F}_{j,k}) \rightarrow K_0(\mathcal{F}_{j,k+1})$ is injective. Hence for $n, m \in \mathbb{N}$, the homomorphism

$$\iota_{n,m} : K_0(\mathcal{A}) = K_0(\mathcal{F}_{0,0}) \rightarrow K_0(\mathcal{F}_{n,m})$$

defined by the compositions of $\iota_{+1,*}$ and $\iota_{*,+1}$ is injective. By [44, Theorem 6.3.2 (iii)], one knows $\text{Ker}(J_{\mathcal{A}*}) = \cup_{n,m \in \mathbb{N}} \text{Ker}(\iota_{n,m})$, so that $\text{Ker}(J_{\mathcal{A}*}) = 0$. □

We henceforth identify the group $K_0(\mathcal{A})$ with its image $J_{\mathcal{A}*}(K_0(\mathcal{A}))$ in $K_0(\mathcal{F}_{\rho,\eta})$. As in the above proof, not only $K_0(\mathcal{A})(= K_0(\mathcal{F}_{0,0}))$ but also the groups $K_0(\mathcal{F}_{j,k})$ for j, k are identified with subgroups of $K_0(\mathcal{F}_{\rho,\eta})$ via injective homomorphisms from $K_0(\mathcal{F}_{j,k})$ to $K_0(\mathcal{F}_{\rho,\eta})$ induced by the embeddings of $\mathcal{F}_{j,k}$ into $\mathcal{F}_{\rho,\eta}$. We note that

$$\begin{aligned} (\text{id} - \gamma_\eta)K_0(\mathcal{F}_{\rho,\eta}) &= (\text{id} - \gamma_\eta^{-1})K_0(\mathcal{F}_{\rho,\eta}), \\ (\text{id} - \gamma_\rho)K_0(\mathcal{F}_{\rho,\eta}) &= (\text{id} - \gamma_\rho^{-1})K_0(\mathcal{F}_{\rho,\eta}) \end{aligned}$$

and

$$\begin{aligned} &\text{Ker}(\text{id} - \gamma_\rho) \cap \text{Ker}(\text{id} - \gamma_\eta) \text{ in } K_0(\mathcal{F}_{\rho,\eta}) \\ &= \text{Ker}(\text{id} - \gamma_\rho^{-1}) \cap \text{Ker}(\text{id} - \gamma_\eta^{-1}) \text{ in } K_0(\mathcal{F}_{\rho,\eta}). \end{aligned}$$

Denote by $(\text{id} - \gamma_\rho)K_0(\mathcal{F}_{\rho,\eta}) + (\text{id} - \gamma_\eta)K_0(\mathcal{F}_{\rho,\eta})$ the subgroup of $K_0(\mathcal{F}_{\rho,\eta})$ generated by $(\text{id} - \gamma_\rho)K_0(\mathcal{F}_{\rho,\eta})$ and $(\text{id} - \gamma_\eta)K_0(\mathcal{F}_{\rho,\eta})$.

Lemma 8.15. *Any element in $K_0(\mathcal{F}_{\rho,\eta})$ is equivalent to some element of $K_0(\mathcal{A})$ modulo the subgroup $(\text{id} - \gamma_\rho)K_0(\mathcal{F}_{\rho,\eta}) + (\text{id} - \gamma_\eta)K_0(\mathcal{F}_{\rho,\eta})$.*

Proof. For $g \in K_0(\mathcal{F}_{\rho,\eta})$, we may assume that $g \in K_0(\mathcal{F}_{j,k})$ for some $j, k \in \mathbb{Z}_+$. As γ_ρ^{-1} commutes with γ_η^{-1} , one sees that $(\gamma_\rho^{-1})^j \circ (\gamma_\eta^{-1})^k(g) \in K_0(\mathcal{A})$. Put $g_1 = \gamma_\rho^{-1}(g)$ so that

$$g - (\gamma_\rho^{-1})^j \circ (\gamma_\eta^{-1})^k(g) = g - \gamma_\rho^{-1}(g) + g_1 - (\gamma_\rho^{-1})^{j-1} \circ (\gamma_\eta^{-1})^k(g_1).$$

We inductively see that $g - (\gamma_\rho^{-1})^j \circ (\gamma_\eta^{-1})^k(g)$ belongs to the subgroup

$$(\text{id} - \gamma_\rho)K_0(\mathcal{F}_{\rho,\eta}) + (\text{id} - \gamma_\eta)K_0(\mathcal{F}_{\rho,\eta}). \quad \square$$

Denote by $(\text{id} - \lambda_\rho)K_0(\mathcal{A}) + (\text{id} - \lambda_\eta)K_0(\mathcal{A})$ the subgroup of $K_0(\mathcal{A})$ generated by $(\text{id} - \lambda_\rho)K_0(\mathcal{A})$ and $(\text{id} - \lambda_\eta)K_0(\mathcal{A})$.

Lemma 8.16. *If $g \in K_0(\mathcal{A})$ belongs to*

$$(\text{id} - \gamma_\rho^{-1})K_0(\mathcal{F}_{\rho,\eta}) + (\text{id} - \gamma_\eta^{-1})K_0(\mathcal{F}_{\rho,\eta}),$$

then g belongs to $(\text{id} - \lambda_\rho)K_0(\mathcal{A}) + (\text{id} - \lambda_\eta)K_0(\mathcal{A})$.

Proof. By the assumption that $g \in (\text{id} - \gamma_\rho^{-1})K_0(\mathcal{F}_{\rho,\eta}) + (\text{id} - \gamma_\eta^{-1})K_0(\mathcal{F}_{\rho,\eta})$, there exist $h_1, h_2 \in K_0(\mathcal{F}_{\rho,\eta})$ such that $g = (\text{id} - \gamma_\rho^{-1})(h_1) + (\text{id} - \gamma_\eta^{-1})(h_2)$. We may assume that $h_1, h_2 \in K_0(\mathcal{F}_{j,k})$ for large enough $j, k \in \mathbb{Z}_+$. Put $e_i = (\gamma_\rho^{-1})^j \circ (\gamma_\eta^{-1})^k(h_i)$ which belongs to $K_0(\mathcal{F}_{0,0}) (= K_0(\mathcal{A}))$ for $i = 0, 1$. It follows that

$$\lambda_\rho^j \circ \lambda_\eta^k(g) = (\text{id} - \lambda_\eta)(e_1) + (\text{id} - \lambda_\rho)(e_2).$$

Since $g \in K_0(\mathcal{A})$ and $\lambda_\rho^j \circ \lambda_\eta^k(g) \in (\text{id} - \lambda_\eta)K_0(\mathcal{A}) + (\text{id} - \lambda_\rho)K_0(\mathcal{A})$, as in the proof of Lemma 8.15, by putting $g^{(n)} = \lambda_\rho^n(g), g^{(n,m)} = \lambda_\eta^m(g^{(n)}) \in K_0(\mathcal{A})$ we have

$$\begin{aligned} &g - \lambda_\rho^j \circ \lambda_\eta^k(g) \\ &= g - \lambda_\rho(g) + g^{(1)} - \lambda_\rho(g^{(1)}) + g^{(2)} - \lambda_\rho(g^{(2)}) + \dots + g^{(j-1)} - \lambda_\rho(g^{(j-1)}) \\ &\quad + g^{(j)} - \lambda_\eta(g^{(j)}) + g^{(j,1)} - \lambda_\eta(g^{(j,1)}) + g^{(j,2)} - \lambda_\eta(g^{(j,2)}) + \dots \\ &\quad + g^{(j,k-1)} - \lambda_\eta(g^{(j,k-1)}) \\ &= (\text{id} - \lambda_\rho)(g + g^{(1)} + \dots + g^{(j-1)}) + (\text{id} - \lambda_\eta)(g^{(j)} + g^{(j,1)} + \dots + g^{(j,k-1)}) \end{aligned}$$

so that g belongs to the subgroup $(\text{id} - \lambda_\eta)K_0(\mathcal{A}) + (\text{id} - \lambda_\rho)K_0(\mathcal{A})$. □

Hence we obtain the following lemma for the cokernel.

Lemma 8.17. *The quotient group*

$$K_0(\mathcal{F}_{\rho,\eta}) / ((\text{id} - \gamma_\eta^{-1})K_0(\mathcal{F}_{\rho,\eta}) + (\text{id} - \gamma_\rho^{-1})K_0(\mathcal{F}_{\rho,\eta}))$$

is isomorphic to the quotient group

$$K_0(\mathcal{A}) / ((\text{id} - \lambda_\eta)K_0(\mathcal{A}) + (\text{id} - \lambda_\rho)K_0(\mathcal{A})).$$

Proof. Surjectivity of the quotient map

$$K_0(\mathcal{A}) \longrightarrow K_0(\mathcal{F}_{\rho,\eta}) / ((\text{id} - \gamma_\eta^{-1})K_0(\mathcal{F}_{\rho,\eta}) + (\text{id} - \gamma_\rho^{-1})K_0(\mathcal{F}_{\rho,\eta}))$$

comes from Lemma 8.15. Its kernel coincides with

$$(\text{id} - \lambda_\eta)K_0(\mathcal{A}) + (\text{id} - \lambda_\rho)K_0(\mathcal{A})$$

by the preceding lemma. □

For the kernel, we have:

Lemma 8.18. *The subgroup*

$$\text{Ker}(\text{id} - \gamma_\eta^{-1}) \cap \text{Ker}(\text{id} - \gamma_\rho^{-1}) \text{ in } K_0(\mathcal{F}_{\rho,\eta})$$

is isomorphic to the subgroup

$$\text{Ker}(\text{id} - \lambda_\eta) \cap \text{Ker}(\text{id} - \lambda_\rho) \text{ in } K_0(\mathcal{A})$$

through $J_{\mathcal{A}^*}$.

Proof. For $g \in \text{Ker}(\text{id} - \gamma_\eta^{-1}) \cap \text{Ker}(\text{id} - \gamma_\rho^{-1})$ in $K_0(\mathcal{F}_{\rho,\eta})$, one may assume that $g \in K_0(\mathcal{F}_{j,k})$ for some $j, k \in \mathbb{Z}_+$ so that $g = (\gamma_\rho^{-1})^j \circ (\gamma_\eta^{-1})^k(g) \in K_0(\mathcal{A})$. Since $\lambda_\eta = \gamma_\eta^{-1}$ and $\lambda_\rho = \gamma_\rho^{-1}$ on $K_0(\mathcal{A})$ under the identification between $J_{\mathcal{A}^*}(K_0(\mathcal{A}))$ and $K_0(\mathcal{A})$ via $J_{\mathcal{A}^*}$, one has that $g \in \text{Ker}(\text{id} - \lambda_\eta) \cap \text{Ker}(\text{id} - \lambda_\rho)$ in $K_0(\mathcal{A})$. The converse inclusion relation

$$\text{Ker}(\text{id} - \lambda_\eta) \cap \text{Ker}(\text{id} - \lambda_\rho) \subset \text{Ker}(\text{id} - \gamma_\eta^{-1}) \cap \text{Ker}(\text{id} - \gamma_\rho^{-1})$$

is clear through the above identification. □

Therefore the short exact sequence for $K_0(\mathcal{O}_{\rho,\eta}^\kappa)$ in Theorem 7.10 is restated as the following proposition.

Proposition 8.19. *Assume that $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ forms square and*

$$K_1(\mathcal{F}_{\rho,\eta}) = \{0\}.$$

Then there exists a short exact sequence:

$$\begin{aligned} 0 &\longrightarrow K_0(\mathcal{A})/((\text{id} - \lambda_\eta)K_0(\mathcal{A}) + (\text{id} - \lambda_\rho)K_0(\mathcal{A})) \\ &\longrightarrow K_0(\mathcal{O}_{\rho,\eta}^\kappa) \\ &\longrightarrow \text{Ker}(\text{id} - \lambda_\eta) \cap \text{Ker}(\text{id} - \lambda_\rho) \text{ in } K_0(\mathcal{A}) \\ &\longrightarrow 0. \end{aligned}$$

Let \mathcal{F}_ρ be the fixed point algebra $(\mathcal{O}_\rho)^{\hat{\rho}}$ of the C^* -algebra \mathcal{O}_ρ by the gauge action $\hat{\rho}$ for the C^* -symbolic dynamical system $(\mathcal{A}, \rho, \Sigma^\rho)$. The algebra \mathcal{F}_ρ is isomorphic to the subalgebra $\mathcal{F}_{\rho,0}$ of $\mathcal{F}_{\rho,\eta}$ in a natural way. As in the proof of Lemma 8.15, the group $K_0(\mathcal{F}_{\rho,0})$ is regarded as a subgroup of $K_0(\mathcal{F}_{\rho,\eta})$ and the restriction of γ_η^{-1} to $K_0(\mathcal{F}_{\rho,0})$ satisfies $\gamma_\eta^{-1}(K_0(\mathcal{F}_{\rho,0})) \subset K_0(\mathcal{F}_{\rho,0})$ so that γ_η^{-1} yields an endomorphism on $K_0(\mathcal{F}_\rho)$, which we still denote by γ_η^{-1} .

For the group $K_1(\mathcal{O}_{\rho,\eta}^\kappa)$, we provide several lemmas.

Lemma 8.20.

- (i) *Any element in $K_0(\mathcal{F}_{\rho,\eta})$ is equivalent to some element of $K_0(\mathcal{F}_{\rho,0})$ (= $K_0(\mathcal{F}_\rho)$) modulo the subgroup $(\text{id} - \gamma_\eta)K_0(\mathcal{F}_{\rho,\eta})$.*
- (ii) *If $g \in K_0(\mathcal{F}_{\rho,0})$ (= $K_0(\mathcal{F}_\rho)$) belongs to $(\text{id} - \gamma_\eta)K_0(\mathcal{F}_{\rho,\eta})$, then g belongs to $(\text{id} - \gamma_\eta)K_0(\mathcal{F}_\rho)$.*

As γ_ρ commutes with γ_η on $K_0(\mathcal{F}_{\rho,\eta})$, it naturally acts on the quotient group $K_0(\mathcal{F}_{\rho,\eta})/(\text{id} - \gamma_\eta^{-1})K_0(\mathcal{F}_{\rho,\eta})$. We denote it by $\bar{\gamma}_\rho$. Similarly λ_ρ naturally induces an endomorphism on $K_0(\mathcal{A})/(\text{id} - \lambda_\eta)K_0(\mathcal{A})$. We denote it by $\bar{\lambda}_\rho$.

Lemma 8.21.

- (i) *The quotient group $K_0(\mathcal{F}_{\rho,\eta})/(\text{id} - \gamma_\eta^{-1})K_0(\mathcal{F}_{\rho,\eta})$ is isomorphic to the quotient group $K_0(\mathcal{F}_\rho)/(\text{id} - \gamma_\eta^{-1})K_0(\mathcal{F}_\rho)$, that is also isomorphic to the quotient group $K_0(\mathcal{A})/(\text{id} - \lambda_\eta)K_0(\mathcal{A})$.*

(ii) The kernel of $\text{id} - \bar{\gamma}_\rho$ in $K_0(\mathcal{F}_{\rho,\eta})/(\text{id} - \gamma_\eta^{-1})K_0(\mathcal{F}_{\rho,\eta})$ is isomorphic to the kernel of $\text{id} - \bar{\lambda}_\rho$ in $K_0(\mathcal{A})/(\text{id} - \lambda_\eta)K_0(\mathcal{A})$.

Proof. (i) The fact that the three quotient groups

$$\begin{aligned} &K_0(\mathcal{F}_{\rho,\eta})/(\text{id} - \gamma_\eta^{-1})K_0(\mathcal{F}_{\rho,\eta}), \\ &K_0(\mathcal{F}_\rho)/(\text{id} - \gamma_\eta^{-1})K_0(\mathcal{F}_\rho), \\ &K_0(\mathcal{A})/(\text{id} - \lambda_\eta)K_0(\mathcal{A}), \end{aligned}$$

are naturally isomorphic is similarly proved to the previous discussions.

(ii) The kernel $\text{Ker}(\text{id} - \bar{\gamma}_\rho)$ in $K_0(\mathcal{F}_{\rho,\eta})/(\text{id} - \gamma_\eta^{-1})K_0(\mathcal{F}_{\rho,\eta})$ is isomorphic to the kernel $\text{Ker}(\text{id} - \bar{\gamma}_\rho)$ in $K_0(\mathcal{F}_\rho)/(\text{id} - \gamma_\eta^{-1})K_0(\mathcal{F}_\rho)$ which is isomorphic to the kernel $\text{Ker}(\text{id} - \bar{\lambda}_\rho)$ in $K_0(\mathcal{A})/(\text{id} - \lambda_\eta)K_0(\mathcal{A})$. \square

Lemma 8.22. *The kernel of $\text{id} - \gamma_\rho$ in $K_0(\mathcal{F}_{\rho,\eta})$ is isomorphic to the kernel of $\text{id} - \gamma_\rho$ in $K_0(\mathcal{F}_\rho)$ that is also isomorphic to the kernel of $\text{id} - \lambda_\eta$ in $K_0(\mathcal{A})$ such that the quotient group*

$$(\text{Ker}(\text{id} - \gamma_\eta) \text{ in } K_0(\mathcal{F}_{\rho,\eta}))/(\text{id} - \gamma_\rho)(\text{Ker}(\text{id} - \gamma_\eta) \text{ in } K_0(\mathcal{F}_{\rho,\eta}))$$

is isomorphic to the quotient group

$$(\text{Ker}(\text{id} - \lambda_\eta) \text{ in } K_0(\mathcal{A}))/(\text{id} - \lambda_\rho)(\text{Ker}(\text{id} - \lambda_\eta) \text{ in } K_0(\mathcal{A})).$$

Proof. The proofs are similar to the previous discussions. \square

Therefore the short exact sequence for $K_1(\mathcal{O}_{\rho,\eta}^\kappa)$ in Theorem 7.10 is restated as the following proposition.

Proposition 8.23. *Assume that $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ forms square and*

$$K_1(\mathcal{F}_{\rho,\eta}) = \{0\}.$$

Then there exists a short exact sequence:

$$\begin{aligned} 0 &\longrightarrow (\text{Ker}(\text{id} - \lambda_\eta) \text{ in } K_0(\mathcal{A}))/(\text{id} - \lambda_\rho)(\text{Ker}(\text{id} - \lambda_\eta) \text{ in } K_0(\mathcal{A})) \\ &\longrightarrow K_1(\mathcal{O}_{\rho,\eta}^\kappa) \\ &\longrightarrow \text{Ker}(\text{id} - \bar{\lambda}_\rho) \text{ in } (K_0(\mathcal{A})/(\text{id} - \lambda_\eta)K_0(\mathcal{A})) \\ &\longrightarrow 0. \end{aligned}$$

We give a condition on $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ which makes $K_1(\mathcal{F}_{\rho,\eta}) = \{0\}$.

Lemma 8.24. *Suppose that a C*-textile dynamical system*

$$(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$$

forms square and satisfies $K_1(\mathcal{A}) = \{0\}$. Then $K_1(\mathcal{F}_{\rho,\eta}) = \{0\}$.

Proof. The algebra $\mathcal{F}_{\rho,\eta}$ is an inductive limit C*-algebra of subalgebras $\mathcal{F}_{j,k}$ with inclusion maps (5.3). Let $E_i^l, i = 1, \dots, m(l)$ be the minimal projections

in \mathcal{A}_l as in Lemma 8.4, which are central in \mathcal{A} such that $\sum_{i=1}^{m(l)} E_i^l = 1$. By Lemma 8.4, we have

$$K_1(\mathcal{F}_{j,k}) = \bigoplus_{i=1}^{m(l)} K_1(\mathcal{F}_{j,k}(i)) = \bigoplus_{i=1}^{m(l)} K_1(E_i^l \mathcal{A} E_i^l) = K_1(\mathcal{A})$$

so that the condition $K_1(\mathcal{A}) = \{0\}$ implies $K_1(\mathcal{F}_{\rho,\eta}) = \{0\}$. □

A C^* -textile dynamical system $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ is said to *have trivial K_1* if $K_1(\mathcal{A}) = \{0\}$.

Consequently we reach the following K-theory formulae for the C^* -algebra $\mathcal{O}_{\rho,\eta}^\kappa$ by Proposition 8.19 and Proposition 8.23.

Theorem 8.25. *Suppose that a C^* -textile dynamical system*

$$(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$$

forms square having trivial K_1 . Then there exist short exact sequences for their K -groups as in the following way:

$$\begin{aligned} 0 &\longrightarrow K_0(\mathcal{A}) / ((\text{id} - \lambda_\eta)K_0(\mathcal{A}) + (\text{id} - \lambda_\rho)K_0(\mathcal{A})) \\ &\longrightarrow K_0(\mathcal{O}_{\rho,\eta}^\kappa) \\ &\longrightarrow \text{Ker}(\text{id} - \lambda_\eta) \cap \text{Ker}(\text{id} - \lambda_\rho) \text{ in } K_0(\mathcal{A}) \\ &\longrightarrow 0 \end{aligned}$$

and

$$\begin{aligned} 0 &\longrightarrow (\text{Ker}(\text{id} - \lambda_\eta) \text{ in } K_0(\mathcal{A})) / (\text{id} - \lambda_\rho)(\text{Ker}(\text{id} - \lambda_\eta) \text{ in } K_0(\mathcal{A})) \\ &\longrightarrow K_1(\mathcal{O}_{\rho,\eta}^\kappa) \\ &\longrightarrow \text{Ker}(\text{id} - \bar{\lambda}_\rho) \text{ in } (K_0(\mathcal{A}) / (\text{id} - \lambda_\eta)K_0(\mathcal{A})) \\ &\longrightarrow 0 \end{aligned}$$

where the endomorphisms $\lambda_\rho, \lambda_\eta : K_0(\mathcal{A}) \longrightarrow K_0(\mathcal{A})$ are defined by

$$\begin{aligned} \lambda_\rho([p]) &= \sum_{\alpha \in \Sigma^\rho} [\rho_\alpha(p)] \in K_0(\mathcal{A}) \text{ for } [p] \in K_0(\mathcal{A}), \\ \lambda_\eta([p]) &= \sum_{a \in \Sigma^\eta} [\eta_a(p)] \in K_0(\mathcal{A}) \text{ for } [p] \in K_0(\mathcal{A}). \end{aligned}$$

9. Examples

9.1. LR-textile λ -graph systems. A symbolic matrix

$$\mathcal{M} = [\mathcal{M}(i, j)]_{i,j=1}^N$$

is a matrix whose components consist of formal sums of elements of an alphabet Σ , such as

$$\mathcal{M} = \begin{bmatrix} a & a + c \\ c & 0 \end{bmatrix} \quad \text{where } \Sigma = \{a, b, c\}.$$

\mathcal{M} is said to be essential if there is no zero column or zero row. \mathcal{M} is said to be left-resolving if for each column a symbol does not appear in two different rows. For example, $\begin{bmatrix} a & a+b \\ c & 0 \end{bmatrix}$ is left-resolving, but $\begin{bmatrix} a & a+b \\ c & b \end{bmatrix}$ is not left-resolving because of b at the second column. We assume that symbolic matrices are always essential and left-resolving. We denote by $\Sigma^{\mathcal{M}}$ the alphabet Σ of the symbolic matrix \mathcal{M} .

Let $\mathcal{M} = [\mathcal{M}(i, j)]_{i, j=1}^N$ and $\mathcal{M}' = [\mathcal{M}'(i, j)]_{i, j=1}^N$ be $N \times N$ symbolic matrices over $\Sigma^{\mathcal{M}}$ and $\Sigma^{\mathcal{M}'}$ respectively. Suppose that there is a bijection $\kappa : \Sigma^{\mathcal{M}} \rightarrow \Sigma^{\mathcal{M}'}$. Following Nasu's terminology [34] we say that \mathcal{M} and \mathcal{M}' are equivalent under specification κ , or simply, specified equivalent if \mathcal{M}' can be obtained from \mathcal{M} by replacing every symbol $\alpha \in \Sigma^{\mathcal{M}}$ by $\kappa(\alpha) \in \Sigma^{\mathcal{M}'}$. That is if $\mathcal{M}(i, j) = \alpha_1 + \dots + \alpha_n$, then $\mathcal{M}'(i, j) = \kappa(\alpha_1) + \dots + \kappa(\alpha_n)$. We write this situation as $\mathcal{M} \stackrel{\kappa}{\cong} \mathcal{M}'$ (see [34]).

For a symbolic matrix $\mathcal{M} = [\mathcal{M}(i, j)]_{i, j=1}^N$ over $\Sigma^{\mathcal{M}}$, we set for $\alpha \in \Sigma^{\mathcal{M}}, i, j = 1, \dots, N$

$$A^{\mathcal{M}}(i, \alpha, j) = \begin{cases} 1 & \text{if } \alpha \text{ appears in } \mathcal{M}(i, j), \\ 0 & \text{otherwise.} \end{cases}$$

Put an $N \times N$ nonnegative matrix $A^{\mathcal{M}} = [A^{\mathcal{M}}(i, j)]_{i, j=1}^N$ by setting

$$A^{\mathcal{M}}(i, j) = \sum_{\alpha \in \Sigma^{\mathcal{M}}} A^{\mathcal{M}}(i, \alpha, j).$$

Let \mathcal{A} be an N -dimensional commutative C^* -algebra \mathbb{C}^N with minimal projections E_1, \dots, E_N such that

$$\mathcal{A} = \mathbb{C}E_1 \oplus \dots \oplus \mathbb{C}E_N.$$

We set for $\alpha \in \Sigma^{\mathcal{M}}$:

$$\rho_{\alpha}^{\mathcal{M}}(E_i) = \sum_{j=1}^N A^{\mathcal{M}}(i, \alpha, j)E_j, \quad i = 1, \dots, N.$$

Then we have a C^* -symbolic dynamical system $(\mathcal{A}, \rho^{\mathcal{M}}, \Sigma^{\mathcal{M}})$.

Let $\mathcal{M} = [\mathcal{M}(i, j)]_{i, j=1}^N$ and $\mathcal{N} = [\mathcal{N}(i, j)]_{i, j=1}^N$ be $N \times N$ symbolic matrices over $\Sigma^{\mathcal{M}}$ and $\Sigma^{\mathcal{N}}$ respectively. We have two C^* -symbolic dynamical systems $(\mathcal{A}, \rho^{\mathcal{M}}, \Sigma^{\mathcal{M}})$ and $(\mathcal{A}, \rho^{\mathcal{N}}, \Sigma^{\mathcal{N}})$. Put

$$\begin{aligned} \Sigma^{\mathcal{M}\mathcal{N}} &= \{(\alpha, b) \in \Sigma^{\mathcal{M}} \times \Sigma^{\mathcal{N}} \mid \rho_b^{\mathcal{N}} \circ \rho_{\alpha}^{\mathcal{M}} \neq 0\}, \\ \Sigma^{\mathcal{N}\mathcal{M}} &= \{(a, \beta) \in \Sigma^{\mathcal{N}} \times \Sigma^{\mathcal{M}} \mid \rho_{\beta}^{\mathcal{M}} \circ \rho_a^{\mathcal{N}} \neq 0\}. \end{aligned}$$

Suppose that there is a bijection κ from $\Sigma^{\mathcal{M}\mathcal{N}}$ to $\Sigma^{\mathcal{N}\mathcal{M}}$ such that κ yields a specified equivalence

$$(9.1) \quad \mathcal{M}\mathcal{N} \stackrel{\kappa}{\cong} \mathcal{N}\mathcal{M}$$

and fix it.

Proposition 9.1. *Keep the above situations. The specified equivalence (9.1) induces a specification $\kappa : \Sigma^{\mathcal{MN}} \rightarrow \Sigma^{\mathcal{NM}}$ such that*

$$(9.2) \quad \rho_b^{\mathcal{N}} \circ \rho_\alpha^{\mathcal{M}} = \rho_\beta^{\mathcal{M}} \circ \rho_a^{\mathcal{N}} \quad \text{if} \quad \kappa(\alpha, b) = (a, \beta).$$

Hence $(\mathcal{A}, \rho^{\mathcal{M}}, \rho^{\mathcal{N}}, \Sigma^{\mathcal{M}}, \Sigma^{\mathcal{N}}, \kappa)$ gives rise to a C^* -textile dynamical system which forms square having trivial K_1 .

Proof. Since $\mathcal{MN} \cong^{\kappa} \mathcal{NM}$, one sees that for $i, j = 1, 2, \dots, N$,

$$\kappa(\mathcal{MN}(i, j)) = \mathcal{NM}(i, j).$$

For $(\alpha, b) \in \Sigma^{\mathcal{MN}}$, there exists $i, k = 1, 2, \dots, N$ such that

$$\rho_b^{\mathcal{N}} \circ \rho_\alpha^{\mathcal{M}}(E_i) \geq E_k.$$

As $\kappa(\alpha, b)$ appears in $\mathcal{NM}(i, k)$, by putting $(a, \beta) = \kappa(\alpha, b)$, we have

$$\rho_\beta^{\mathcal{M}} \circ \rho_a^{\mathcal{N}}(E_i) \geq E_k.$$

Hence $\kappa(\alpha, b) \in \Sigma^{\mathcal{NM}}$. One indeed sees that $\rho_b^{\mathcal{N}} \circ \rho_\alpha^{\mathcal{M}} = \rho_\beta^{\mathcal{M}} \circ \rho_a^{\mathcal{N}}$ by the relation $\mathcal{MN} \cong^{\kappa} \mathcal{NM}$. □

Two symbolic matrices satisfying (9.1) give rise to an LR textile system that has been introduced by Nasu (see [34]). Textile systems introduced by Nasu give a strong tool to analyze automorphisms and endomorphisms of topological Markov shifts. The author has generalized LR-textile systems to LR-textile λ -graph systems which consist of two pairs of sequences $(\mathcal{M}, I) = (\mathcal{M}_{l,l+1}, I_{l,l+1})_{l \in \mathbb{Z}_+}$ and $(\mathcal{N}, I) = (\mathcal{N}_{l,l+1}, I_{l,l+1})_{l \in \mathbb{Z}_+}$ such that

$$(9.3) \quad \mathcal{M}_{l,l+1} \mathcal{N}_{l+1,l+2} \cong^{\kappa} \mathcal{N}_{l,l+1} \mathcal{M}_{l+1,l+2}, \quad l \in \mathbb{Z}_+$$

through a specification κ ([28]). We denote the LR-textile λ -graph system by $\mathcal{T}_{\mathcal{K}_{\mathcal{N}}^{\mathcal{M}}}$. Denote by $\mathfrak{L}^{\mathcal{M}}$ and $\mathfrak{L}^{\mathcal{N}}$ the associated λ -graph systems respectively. Since $\mathfrak{L}^{\mathcal{M}}$ and $\mathfrak{L}^{\mathcal{N}}$ have common sequences $V_l^{\mathcal{M}} = V_l^{\mathcal{N}}, l \in \mathbb{Z}_+$ of vertices which denoted by $V_l, l \in \mathbb{Z}_+$, and its common inclusion matrices $I_{l,l+1}, l \in \mathbb{Z}_+$. Hence $\mathfrak{L}^{\mathcal{M}}$ and $\mathfrak{L}^{\mathcal{N}}$ form square in the sense of [28, p.170]. Let $(\mathcal{A}_{\mathcal{M}}, \rho^{\mathcal{M}}, \Sigma^{\mathcal{M}})$ and $(\mathcal{A}_{\mathcal{N}}, \rho^{\mathcal{N}}, \Sigma^{\mathcal{N}})$ be the associated C^* -symbolic dynamical systems with the λ -graph systems $\mathfrak{L}^{\mathcal{M}}$ and $\mathfrak{L}^{\mathcal{N}}$ respectively. Since both the algebras $\mathcal{A}_{\mathcal{M}}$ and $\mathcal{A}_{\mathcal{N}}$ are the C^* -algebras of inductive limit of the system $I_{l,l+1}^* : C(V_l) \rightarrow C(V_{l+1}), l \in \mathbb{Z}_+$, they are identical, which is denoted by \mathcal{A} . It is easy to see that the relation (9.3) implies

$$(9.4) \quad \rho_\alpha^{\mathcal{M}} \circ \rho_b^{\mathcal{N}} = \rho_a^{\mathcal{N}} \circ \rho_\beta^{\mathcal{M}} \quad \text{if} \quad \kappa(\alpha, b) = (a, \beta).$$

Proposition 9.2. *An LR-textile λ -graph system $\mathcal{T}_{\mathcal{K}_{\mathcal{N}}^{\mathcal{M}}}$ yields a C^* -textile dynamical system $(\mathcal{A}, \rho^{\mathcal{M}}, \rho^{\mathcal{N}}, \Sigma^{\mathcal{M}}, \Sigma^{\mathcal{N}}, \kappa)$ which forms square. Conversely, a C^* -textile dynamical system $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ which forms square yields*

an LR-textile λ -graph system $\mathcal{T}_{\mathcal{K}^{\mathcal{M}^\rho}}^{\mathcal{M}^\eta}$ such that the associated C*-textile dynamical system written $(\mathcal{A}_{\rho,\eta}, \rho^{\mathcal{M}^\rho}, \rho^{\mathcal{M}^\eta}, \Sigma^{\mathcal{M}^\rho}, \Sigma^{\mathcal{M}^\eta}, \kappa)$ is a subsystem of $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ in the sense that the relations:

$$\mathcal{A}_{\rho,\eta} \subset \mathcal{A}, \quad \rho|_{\mathcal{A}_{\rho,\eta}} = \rho^{\mathcal{M}^\rho}, \quad \eta|_{\mathcal{A}_{\rho,\eta}} = \rho^{\mathcal{M}^\eta}$$

hold.

Proof. Let $\mathcal{T}_{\mathcal{K}^{\mathcal{M}}}$ be an LR-textile λ -graph system. As in the above discussions, we have a C*-textile dynamical system $(\mathcal{A}, \rho^{\mathcal{M}}, \rho^{\mathcal{N}}, \Sigma^{\mathcal{M}}, \Sigma^{\mathcal{N}}, \kappa)$. Conversely, let $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ be a C*-textile dynamical system which forms square. Put for $l \in \mathbb{N}$

$$\mathcal{A}_l^\rho = C^*(\rho_\mu(1) : \mu \in B_l(\Lambda_\rho)), \quad \mathcal{A}_l^\eta = C^*(\eta_\xi(1) : \xi \in B_l(\Lambda_\eta)).$$

Since $\mathcal{A}_l^\rho = \mathcal{A}_l^\eta$ and they are commutative and of finite dimensional, the algebra

$$\mathcal{A}_{\rho,\eta} = \overline{\cup_{l \in \mathbb{Z}_+} \mathcal{A}_l^\rho} = \overline{\cup_{l \in \mathbb{Z}_+} \mathcal{A}_l^\eta}$$

is a commutative AF-subalgebra of \mathcal{A} . It is easy to see that both $(\mathcal{A}_{\rho,\eta}, \rho, \Sigma^\rho)$ and $(\mathcal{A}_{\rho,\eta}, \eta, \Sigma^\eta)$ are C*-symbolic dynamical systems such that

$$(9.5) \quad \eta_b \circ \rho_\alpha = \rho_\beta \circ \eta_a \quad \text{if} \quad \kappa(\alpha, b) = (a, \beta)$$

By [27], there exist λ -graph systems \mathcal{L}^ρ and \mathcal{L}^η whose C*-symbolic dynamical systems are $(\mathcal{A}_{\rho,\eta}, \rho, \Sigma^\rho)$ and $(\mathcal{A}_{\rho,\eta}, \eta, \Sigma^\eta)$ respectively. Let $(\mathcal{M}^\rho, I^\rho)$ and $(\mathcal{M}^\eta, I^\eta)$ be the associated symbolic matrix systems. It is easy to see that the relation (9.5) implies

$$\mathcal{M}_{l,l+1}^\rho \mathcal{M}_{l+1,l+2}^\eta \stackrel{\kappa}{\cong} \mathcal{M}_{l,l+1}^\eta \mathcal{M}_{l+1,l+2}^\rho, \quad l \in \mathbb{Z}_+.$$

Hence we have an LR-textile λ -graph system $\mathcal{T}_{\mathcal{K}^{\mathcal{M}^\rho}}^{\mathcal{M}^\eta}$. It is direct to see that the associated C*-textile dynamical system is $(\mathcal{A}_{\rho,\eta}, \rho|_{\mathcal{A}_{\rho,\eta}}, \eta|_{\mathcal{A}_{\rho,\eta}}, \Sigma^\rho, \Sigma^\eta, \kappa)$. \square

Let A be an $N \times N$ matrix with entries in nonnegative integers. We may consider a directed graph $G_A = (V_A, E_A)$ with vertex set V_A and edge set E_A . The vertex set V_A consists of N vertices which we denote by $\{v_1, \dots, v_N\}$. We equip $A(i, j)$ edges from the vertex v_i to the vertex v_j . Denote by E_A the set of the edges. Let $\Sigma^A = E_A$ and the labeling map $\lambda_A : E_A \rightarrow \Sigma^A$ be defined as the identity map. Then we have a labeled directed graph denoted by G_A as well as a symbolic matrix $\mathcal{M}_A = [\mathcal{M}_A(i, j)]_{i,j=1}^N$ by setting

$$\mathcal{M}_A(i, j) = \begin{cases} e_1 + \dots + e_n & \text{if } e_1, \dots, e_n \text{ are edges from } v_i \text{ to } v_j, \\ 0 & \text{if there is no edge from } v_i \text{ to } v_j. \end{cases}$$

Let B be an $N \times N$ matrix with entries in nonnegative integers such that

$$(9.6) \quad AB = BA.$$

The equality (9.6) implies that the cardinal numbers of the sets of the pairs of directed edges

$$\begin{aligned}\Sigma^{AB}(i, j) &= \{(e, f) \in E_A \times E_B \mid s(e) = v_i, t(e) = s(f), t(f) = v_j\} \text{ and} \\ \Sigma^{BA}(i, j) &= \{(f, e) \in E_B \times E_A \mid s(f) = v_i, t(f) = s(e), t(e) = v_j\}\end{aligned}$$

coincide with each other for each v_i and v_j . We put $\Sigma^{AB} = \cup_{i,j=1}^N \Sigma^{AB}(i, j)$ and $\Sigma^{BA} = \cup_{i,j=1}^N \Sigma^{BA}(i, j)$ so that one may take a bijection $\kappa : \Sigma^{AB} \rightarrow \Sigma^{BA}$ which gives rise to a specified equivalence $\mathcal{M}_A \mathcal{M}_B \stackrel{\kappa}{\cong} \mathcal{M}_B \mathcal{M}_A$. We then have a C^* -textile dynamical system

$$(\mathcal{A}, \rho^{\mathcal{M}_A}, \rho^{\mathcal{M}_B}, \Sigma^A, \Sigma^B, \kappa)$$

which we denote by

$$(\mathcal{A}, \rho^A, \rho^B, \Sigma^A, \Sigma^B, \kappa).$$

The associated C^* -algebra is denoted by $\mathcal{O}_{A,B}^\kappa$. The algebra $\mathcal{O}_{A,B}^\kappa$ depends on the choice of a specification $\kappa : \Sigma^{AB} \rightarrow \Sigma^{BA}$. The algebras are 2-graph algebras of Kumjian and Pask [19]. They are also C^* -algebras associated to textile systems studied by V. Deaconu [9]. By Theorem 8.25, we have:

Proposition 9.3. *Keep the above situations. There exist short exact sequences:*

$$\begin{aligned}0 &\longrightarrow \mathbb{Z}^N / ((1 - A)\mathbb{Z}^N + (1 - B)\mathbb{Z}^N) \\ &\longrightarrow K_0(\mathcal{O}_{A,B}^\kappa) \\ &\longrightarrow \text{Ker}(1 - A) \cap \text{Ker}(1 - B) \text{ in } \mathbb{Z}^N \longrightarrow 0\end{aligned}$$

and

$$\begin{aligned}0 &\longrightarrow (\text{Ker}(1 - B) \text{ in } \mathbb{Z}^N) / (1 - A)(\text{Ker}(1 - B) \text{ in } \mathbb{Z}^N) \\ &\longrightarrow K_1(\mathcal{O}_{A,B}^\kappa) \\ &\longrightarrow \text{Ker}(1 - A) \text{ in } \mathbb{Z}^N / (1 - B)\mathbb{Z}^N \longrightarrow 0.\end{aligned}$$

We consider 1×1 matrices $[N]$ and $[M]$ with its entries N and M respectively for $1 < N, M \in \mathbb{N}$. Let G_N be a directed graph with one vertex and N directed self-loops. Similarly we consider a directed graph G_M with M directed self-loops at the vertex. The self-loops are denoted by $\Sigma^N = \{e_1, \dots, e_N\}$ and $\Sigma^M = \{f_1, \dots, f_M\}$ respectively. As a specification κ , we take the exchanging map $(e, f) \in \Sigma^N \times \Sigma^M \rightarrow (f, e) \in \Sigma^M \times \Sigma^N$ which we will fix. Put

$$\rho_{e_i}^N(1) = 1, \quad \rho_{f_j}^M(1) = 1 \quad \text{for } i = 1, \dots, N, \quad j = 1, \dots, M.$$

Then we have a C^* -textile dynamical system

$$(\mathbb{C}, \rho^N, \rho^M, \Sigma^N, \Sigma^M, \kappa).$$

The associated C^* -algebra is denoted by $\mathcal{O}_{N,M}^\kappa$.

Lemma 9.4. $\mathcal{O}_{N,M}^\kappa = \mathcal{O}_N \otimes \mathcal{O}_M$.

Proof. Let $s_i, i = 1, \dots, N$ and $t_j, i = 1, \dots, M$ be the generating isometries of the Cuntz algebra \mathcal{O}_N and those of \mathcal{O}_M respectively which satisfy

$$\sum_{i=1}^N s_i s_i^* = 1, \quad \sum_{j=1}^M t_j t_j^* = 1.$$

Let $S_i, i = 1, \dots, N$ and $T_j, i = 1, \dots, M$ be the generating isometries of $\mathcal{O}_{N,M}^\kappa$ satisfying

$$\sum_{i=1}^N S_i S_i^* = 1, \quad \sum_{j=1}^M T_j T_j^* = 1$$

and

$$S_i T_j = T_j S_i, \quad i = 1, \dots, N, \quad j = 1, \dots, M.$$

The universality of $\mathcal{O}_{N,M}^\kappa$ subject to the relations and that of the tensor product $\mathcal{O}_N \otimes \mathcal{O}_M$ ensure us that the correspondence $\Phi : \mathcal{O}_{N,M} \rightarrow \mathcal{O}_N \otimes \mathcal{O}_M$ given by $\Phi(S_i) = s_i \otimes 1, \Phi(T_j) = 1 \otimes t_j$ yields an isomorphism. \square

Although we may easily compute the K-groups $K_*(\mathcal{O}_{M,N}^\kappa)$ by using the Künneth formula for $K_i(\mathcal{O}_N \otimes \mathcal{O}_M)$ ([46]), we will compute them by Proposition 9.3 as in the following way.

Proposition 9.5 (cf. [19]). *For $1 < N, M \in \mathbb{N}$, the C*-algebra $\mathcal{O}_{N,M}^\kappa$ is simple, purely infinite, such that*

$$K_0(\mathcal{O}_{N,M}^\kappa) \cong K_1(\mathcal{O}_{N,M}^\kappa) \cong \mathbb{Z}/d\mathbb{Z}$$

where $d = \text{gcd}(N - 1, M - 1)$ the greatest common divisor of $N - 1, M - 1$.

Proof. It is easy to see that the group $\mathbb{Z}/((N - 1)\mathbb{Z} + (M - 1)\mathbb{Z})$ is isomorphic to $\mathbb{Z}/d\mathbb{Z}$. As $\text{Ker}(N - 1) = \text{Ker}(M - 1) = 0$ in \mathbb{Z} , we see that

$$K_0(\mathcal{O}_{N,M}^\kappa) \cong \mathbb{Z}/d\mathbb{Z}.$$

It is elementary to see that the subgroup

$$\{[k] \in \mathbb{Z}/(M - 1)\mathbb{Z} \mid (N - 1)k \in (M - 1)\mathbb{Z}\}$$

of $\mathbb{Z}/(M - 1)\mathbb{Z}$ is isomorphic to $\mathbb{Z}/d\mathbb{Z}$. Hence we have

$$K_1(\mathcal{O}_{N,M}^\kappa) \cong \mathbb{Z}/d\mathbb{Z}. \quad \square$$

We will generalize the above examples from the view point of tensor products.

9.2. Tensor products. Let $(\mathcal{A}^\rho, \rho, \Sigma^\rho)$ and $(\mathcal{A}^\eta, \eta, \Sigma^\eta)$ be C^* -symbolic dynamical systems. We will construct a C^* -textile dynamical system by taking tensor product. Put

$$\bar{\mathcal{A}} = \mathcal{A}^\rho \otimes \mathcal{A}^\eta, \quad \bar{\rho}_\alpha = \rho_\alpha \otimes \text{id}, \quad \bar{\eta}_a = \text{id} \otimes \eta_a, \quad \Sigma^{\bar{\rho}} = \Sigma^\rho, \quad \Sigma^{\bar{\eta}} = \Sigma^\eta$$

for $\alpha \in \Sigma^\rho, a \in \Sigma^\eta$, where \otimes means the minimal C^* -tensor product \otimes_{\min} . For $(\alpha, a) \in \Sigma^\rho \times \Sigma^\eta$, we see $\eta_b \circ \rho_\alpha(1) \neq 0$ if and only if $\eta_b(1) \neq 0, \rho_\alpha(1) \neq 0$, so that

$$\Sigma^{\bar{\rho}\bar{\eta}} = \Sigma^\rho \times \Sigma^\eta \quad \text{and similarly} \quad \Sigma^{\bar{\eta}\bar{\rho}} = \Sigma^\eta \times \Sigma^\rho.$$

Define $\bar{\kappa} : \Sigma^{\bar{\rho}\bar{\eta}} \rightarrow \Sigma^{\bar{\eta}\bar{\rho}}$ by setting $\bar{\kappa}(\alpha, b) = (b, \alpha)$.

Lemma 9.6. $(\bar{\mathcal{A}}, \bar{\rho}, \bar{\eta}, \Sigma^{\bar{\rho}}, \Sigma^{\bar{\eta}}, \bar{\kappa})$ is a C^* -textile dynamical system.

Proof. By [2], we have $Z_{\bar{\mathcal{A}}} = Z_{\mathcal{A}^\rho} \otimes Z_{\mathcal{A}^\eta}$ so that

$$\bar{\rho}_\alpha(Z_{\bar{\mathcal{A}}}) \subset Z_{\bar{\mathcal{A}}}, \quad \alpha \in \Sigma^{\bar{\rho}} \quad \text{and} \quad \bar{\eta}_a(Z_{\bar{\mathcal{A}}}) \subset Z_{\bar{\mathcal{A}}}, \quad a \in \Sigma^{\bar{\eta}}.$$

We also have $\sum_{\alpha \in \Sigma^{\bar{\rho}}} \bar{\rho}_\alpha(1) = \sum_{\alpha \in \Sigma^\rho} \rho_\alpha(1) \otimes 1 \geq 1$, and similarly

$$\sum_{a \in \Sigma^{\bar{\eta}}} \bar{\eta}(1) \geq 1$$

so that both families $\{\bar{\rho}_\alpha\}_{\alpha \in \Sigma^{\bar{\rho}}}$ and $\{\bar{\eta}_a\}_{a \in \Sigma^{\bar{\eta}}}$ of endomorphisms are essential. Since $\{\rho_\alpha\}_{\alpha \in \Sigma^\rho}$ is faithful on \mathcal{A}^ρ , the homomorphism

$$x \in \mathcal{A}^\rho \longrightarrow \sum_{\alpha \in \Sigma^\rho} \oplus \rho_\alpha(x) \in \sum_{\alpha \in \Sigma^\rho} \oplus \mathcal{A}^\rho$$

is injective so that the homomorphism

$$x \otimes y \in \mathcal{A}^\rho \otimes \mathcal{A}^\eta \longrightarrow \sum_{\alpha \in \Sigma^\rho} \oplus \rho_\alpha(x) \otimes y \in \sum_{\alpha \in \Sigma^\rho} \oplus \mathcal{A}^\rho \otimes \mathcal{A}^\eta$$

is injective. This implies that $\{\bar{\rho}_\alpha\}_{\alpha \in \Sigma^{\bar{\rho}}}$ is faithful. Similarly, so is $\{\bar{\eta}_a\}_{a \in \Sigma^{\bar{\eta}}}$. Hence $(\bar{\mathcal{A}}, \bar{\rho}, \Sigma^{\bar{\rho}})$ and $(\bar{\mathcal{A}}, \bar{\eta}, \Sigma^{\bar{\eta}})$ are both C^* -symbolic dynamical systems. It is direct to see that $\bar{\eta}_b \circ \bar{\rho}_\alpha = \bar{\rho}_\alpha \circ \bar{\eta}_b$ for $(\alpha, b) \in \Sigma^{\bar{\rho}\bar{\eta}}$. Therefore $(\bar{\mathcal{A}}, \bar{\rho}, \bar{\eta}, \Sigma^{\bar{\rho}}, \Sigma^{\bar{\eta}}, \bar{\kappa})$ is a C^* -textile dynamical system. \square

We call $(\bar{\mathcal{A}}, \bar{\rho}, \bar{\eta}, \Sigma^{\bar{\rho}}, \Sigma^{\bar{\eta}}, \bar{\kappa})$ the tensor product between $(\mathcal{A}^\rho, \rho, \Sigma^\rho)$ and $(\mathcal{A}^\eta, \eta, \Sigma^\eta)$. Denote by $S_\alpha, \alpha \in \Sigma^{\bar{\rho}}, T_a, a \in \Sigma^{\bar{\eta}}$ the generating partial isometries of the C^* -algebra $\mathcal{O}_{\bar{\rho}, \bar{\eta}}^{\bar{\kappa}}$ for the C^* -textile dynamical system

$$(\bar{\mathcal{A}}, \bar{\rho}, \bar{\eta}, \Sigma^{\bar{\rho}}, \Sigma^{\bar{\eta}}, \bar{\kappa}).$$

By the universality for the algebra $\mathcal{O}_{\bar{\rho}, \bar{\eta}}^{\bar{\kappa}}$ subject to the relations $(\bar{\rho}, \bar{\eta}; \bar{\kappa})$, the algebra $\mathcal{D}_{\bar{\rho}, \bar{\eta}}$ is isomorphic to the tensor product $\mathcal{D}_\rho \otimes \mathcal{D}_\eta$ through the correspondence

$$S_\mu T_\xi (x \otimes y) T_\xi^* S_\mu^* \longleftrightarrow S_\mu x S_\mu^* \otimes T_\xi y T_\xi^*$$

for $\mu \in B_*(\Lambda_\rho), \xi \in B_*(\Lambda_\eta), x \in \mathcal{A}^\rho, y \in \mathcal{A}^\eta$.

Lemma 9.7. Suppose that $(\mathcal{A}^\rho, \rho, \Sigma^\rho)$ and $(\mathcal{A}^\eta, \eta, \Sigma^\eta)$ are both free (resp. AF-free). Then the tensor product $(\bar{\mathcal{A}}, \bar{\rho}, \bar{\eta}, \Sigma^{\bar{\rho}}, \Sigma^{\bar{\eta}}, \bar{\kappa})$ is free (resp. AF-free).

Proof. Suppose that $(\mathcal{A}^\rho, \rho, \Sigma^\rho)$ and $(\mathcal{A}^\eta, \eta, \Sigma^\eta)$ are both free. There exist increasing sequences $\mathcal{A}_l^\rho, l \in \mathbb{Z}_+$ and $\mathcal{A}_l^\eta, l \in \mathbb{Z}_+$ of C^* -subalgebras of \mathcal{A}^ρ and \mathcal{A}^η satisfying the conditions of their freeness respectively. Put

$$\bar{\mathcal{A}}_l = \mathcal{A}_l^\rho \otimes \mathcal{A}_l^\eta, \quad l \in \mathbb{Z}_+.$$

It is clear that:

- (1) $\bar{\rho}_\alpha(\bar{\mathcal{A}}_l) \subset \bar{\mathcal{A}}_{l+1}, \alpha \in \Sigma^{\bar{\rho}}$ and $\bar{\eta}_a(\bar{\mathcal{A}}_l) \subset \bar{\mathcal{A}}_{l+1}, a \in \Sigma^{\bar{\eta}}$ for $l \in \mathbb{Z}_+$.
- (2) $\cup_{l \in \mathbb{Z}_+} \bar{\mathcal{A}}_l$ is dense in $\bar{\mathcal{A}}$.

We will show that the condition (3) for $\bar{\mathcal{A}}$ in Definition 5.3 holds. Take and fix arbitrary $j, k, l \in \mathbb{N}$ with $j + k \leq l$. For $j \leq l$, one may take a projection $q_\rho \in \mathcal{D}_\rho \cap \mathcal{A}_j^{\rho'}$ satisfying the condition (3) of the freeness of $(\mathcal{A}^\rho, \rho, \Sigma^\rho)$, and similarly for $k \leq l$, one may take a projection $q_\eta \in \mathcal{D}_\eta \cap \mathcal{A}_k^{\eta'}$. Put $q = q_\rho \otimes q_\eta \in \mathcal{D}_\rho \otimes \mathcal{D}_\eta (= \mathcal{D}_{\bar{\rho}, \bar{\eta}})$ so that $q \in \mathcal{D}_{\bar{\rho}, \bar{\eta}} \cap \bar{\mathcal{A}}_l^i$. As the maps $\Phi_l^\rho : x \in \mathcal{A}_l^\rho \rightarrow q_\rho x \in q_\rho \mathcal{A}_l^\rho$ and $\Phi_l^\eta : y \in \mathcal{A}_l^\eta \rightarrow q_\eta y \in q_\eta \mathcal{A}_l^\eta$ are both isomorphisms, the tensor product

$$\Phi_l^\rho \otimes \Phi_l^\eta : x \otimes y \in \mathcal{A}_l^\rho \otimes \mathcal{A}_l^\eta \rightarrow (q_\rho \otimes q_\eta)(x \otimes y) \in (q_\rho \otimes q_\eta)(\mathcal{A}_l^\rho \otimes \mathcal{A}_l^\eta)$$

is isomorphic. Hence $qa \neq 0$ for $0 \neq a \in \bar{\mathcal{A}}_l$. It is straightforward to see that q satisfies the condition (3) (ii) of Definition 5.3. Therefore the tensor product $(\bar{\mathcal{A}}, \bar{\rho}, \bar{\eta}, \Sigma^{\bar{\rho}}, \Sigma^{\bar{\eta}}, \bar{\kappa})$ is free. It is obvious to see that if both $(\mathcal{A}^\rho, \rho, \Sigma^\rho)$ and $(\mathcal{A}^\eta, \eta, \Sigma^\eta)$ are AF-free, then $(\bar{\mathcal{A}}, \bar{\rho}, \bar{\eta}, \Sigma^{\bar{\rho}}, \Sigma^{\bar{\eta}}, \bar{\kappa})$ is AF-free. \square

Proposition 9.8. *Suppose that $(\mathcal{A}^\rho, \rho, \Sigma^\rho)$ and $(\mathcal{A}^\eta, \eta, \Sigma^\eta)$ are both free. Then the C^* -algebra $\mathcal{O}_{\bar{\rho}, \bar{\eta}}^{\bar{\kappa}}$ for the tensor product C^* -textile dynamical system $(\bar{\mathcal{A}}, \bar{\rho}, \bar{\eta}, \Sigma^{\bar{\rho}}, \Sigma^{\bar{\eta}}, \bar{\kappa})$ is isomorphic to the minimal tensor product $\mathcal{O}_\rho \otimes \mathcal{O}_\eta$ of the C^* -algebras between \mathcal{O}_ρ and \mathcal{O}_η . If in particular, $(\mathcal{A}^\rho, \rho, \Sigma^\rho)$ and $(\mathcal{A}^\eta, \eta, \Sigma^\eta)$ are both irreducible, the C^* -algebra $\mathcal{O}_{\bar{\rho}, \bar{\eta}}^{\bar{\kappa}}$ is simple.*

Proof. Suppose that $(\mathcal{A}^\rho, \rho, \Sigma^\rho)$ and $(\mathcal{A}^\eta, \eta, \Sigma^\eta)$ are both free. By the preceding lemma, the tensor product $(\bar{\mathcal{A}}, \bar{\rho}, \bar{\eta}, \Sigma^{\bar{\rho}}, \Sigma^{\bar{\eta}}, \bar{\kappa})$ is free and hence satisfies condition (I). Let $s_\alpha, \alpha \in \Sigma^\rho$ and $t_a, a \in \Sigma^\eta$ be the generating partial isometries of the C^* -algebras \mathcal{O}_ρ and \mathcal{O}_η respectively. Let $S_\alpha, \alpha \in \Sigma^{\bar{\rho}}$ and $T_a, a \in \Sigma^{\bar{\eta}}$ be the generating partial isometries of the C^* -algebra $\mathcal{O}_{\bar{\rho}, \bar{\eta}}^{\bar{\kappa}}$. By the uniqueness of the algebra $\mathcal{O}_{\bar{\rho}, \bar{\eta}}^{\bar{\kappa}}$ with respect to the relations $(\bar{\rho}, \bar{\eta}; \bar{\kappa})$, the correspondence

$$S_\alpha \rightarrow s_\alpha \otimes 1 \in \mathcal{O}_\rho \otimes \mathcal{O}_\eta, \quad T_a \rightarrow 1 \otimes t_a \in \mathcal{O}_\rho \otimes \mathcal{O}_\eta$$

naturally gives rise to an isomorphism from $\mathcal{O}_{\bar{\rho}, \bar{\eta}}^{\bar{\kappa}}$ onto the tensor product $\mathcal{O}_\rho \otimes \mathcal{O}_\eta$.

If in particular, $(\mathcal{A}^\rho, \rho, \Sigma^\rho)$ and $(\mathcal{A}^\eta, \eta, \Sigma^\eta)$ are both irreducible, the C^* -algebras \mathcal{O}_ρ and \mathcal{O}_η are both simple so that $\mathcal{O}_{\bar{\rho}, \bar{\eta}}^{\bar{\kappa}}$ is simple. \square

We remark that the tensor product $(\bar{\mathcal{A}}, \bar{\rho}, \bar{\eta}, \Sigma^{\bar{\rho}}, \Sigma^{\bar{\eta}}, \bar{\kappa})$ does not necessarily form square. The K-theory groups $K_*(\mathcal{O}_{\bar{\rho}, \bar{\eta}}^{\bar{\kappa}})$ are computed from the Künneth formulae for $K_*(\mathcal{O}_\rho \otimes \mathcal{O}_\eta)$ [46].

10. Concluding remark

In [31], a different construction of C^* -algebra written $\mathcal{O}_{\mathcal{H}_\kappa}$ from C^* -textile dynamical system $(\mathcal{A}, \rho, \eta, \Sigma^\rho, \Sigma^\eta, \kappa)$ is studied by using a 2-dimensional analogue of Hilbert C^* -bimodule. The C^* -algebra $\mathcal{O}_{\mathcal{H}_\kappa}$ is different from the C^* -algebra $\mathcal{O}_{\rho, \eta}^\kappa$ in the present paper (see also [33], [32]).

References

- [1] ARANDA PINO, GONZALO; CLARK, JOHN; AN HUEF, ASTRID; RAEBURN, IAIN. Kumjian–Pask algebras of higher-rank graphs. *Trans. Amer. Math. Soc.* **365** (2013), no. 7, 3613–3641. MR3042597, Zbl 1282.16006, doi:10.1090/S0002-9947-2013-05717-0.
- [2] ARCHBOLD, R. J. On the centre of a tensor product of C^* -algebras. *J. London Math. Soc.* (2) **10** (1975), 257–262. MR0402512 (53 # 6331), Zbl 0303.46057, doi:10.1112/jlms/s2-10.3.257.
- [3] BATES, TERESA; PASK, DAVID. C^* -algebras of labelled graphs. *J. Operator Theory* **57** (2007), no. 1, 207–226. MR2304922 (2008c:46079), Zbl 1113.46049, arXiv:math/0503374.
- [4] BLACKADAR, BRUCE. K -theory for operator algebras. Second edition. Mathematical Sciences Research Institute Publications, 5. *Cambridge University Press, Cambridge*, 1998. xx+300 pp. ISBN: 0-521-63532-2. MR1656031 (99g:46104), Zbl 0913.46054.
- [5] BRATTELI, OLA. Inductive limits of finite-dimensional C^* -algebras. *Trans. Amer. Math. Soc.* **171** (1972), 195–234. MR0312282 (47 #844), Zbl 0264.46057, doi:10.2307/1996380.
- [6] BROWN, LAWRENCE G. Stable isomorphism of hereditary subalgebras of C^* -algebras. *Pacific J. Math.* **71** (1977), no. 2, 335–348. MR0454645 (56 #12894), Zbl 0362.46042, doi:10.2140/pjm.1977.71.335.
- [7] CUNTZ, JOACHIM. Simple C^* -algebras generated by isometries. *Comm. Math. Phys.* **57** (1977), no. 2, 173–185. MR0467330 (57 7189), Zbl 0399.46045, doi:10.1007/BF01625776.
- [8] CUNTZ, JOACHIM; KRIEGER, WOLFGANG. A class of C^* -algebras and topological Markov chains. *Invent. Math.* **56** (1980), no. 3, 251–268. MR561974 (82f:46073a), Zbl 0434.46045, doi:10.1007/BF01390048.
- [9] DEACONU, VALENTIN. C^* -algebras and Fell bundles associated to a textile system. *J. Math. Anal. Appl.* **372** (2010), no. 2, 515–524. MR2678879 (2012a:46092), Zbl 1205.37020, arXiv:1001.0037, doi:10.1016/j.jmaa.2010.07.034.
- [10] EVANS, D. GWION. On the K -theory of higher rank graph C^* -algebras. *New York J. Math.* **14** (2008), 1–31. MR2383584 (2009e:46065), Zbl 1146.46048, arXiv:math/0406458.
- [11] EXEL, RUY; GONÇALVES, DANIEL; STARLING, CHARLES. The tiling C^* -algebra viewed as a tight inverse semigroup algebra. *Semigroup Forum* **84** (2012), no. 2, 229–240. MR2898758 (2000i:46064), Zbl 1256.46034, arXiv:1106.4535, doi:10.1007/s00233-011-9359-x.
- [12] EXEL, RUY; LACA, MARCELO. Cuntz–Krieger algebras for infinite matrices. *J. Reine Angew. Math.* **512** (1999), 119–172. MR1703078 (2000i:46064), Zbl 0932.47053, arXiv:funct-an/9712008, doi:10.1515/crll.1999.051.
- [13] FARTHING, CYNTHIA; MUHLY, PAUL S.; YEEND, TRENT. Higher-rank graph C^* -algebras: an inverse semigroup and groupoid approach. *Semigroup Forum* **71** (2005), no. 2, 159–187. MR2184052 (2006h:46052), Zbl 1099.46036, arXiv:math/0409505, doi:10.1007/s00233-005-0512-2.

- [14] FISCHER, ROLAND. Sofic systems and graphs. *Monatsh. Math.* **80** (1975), no. 3, 179–186. MR0407235 (53 #11018), Zbl 0314.54043, doi:10.1007/BF01319913.
- [15] KAJIWARA, TSUYOSHI; PINZARI, CLAUDIA; WATATANI, YASUO. Ideal structure and simplicity of the C^* -algebras generated by Hilbert bimodules. *J. Funct. Anal.* **159** (1998), no. 2, 295–322. MR1658088 (2000a:46094), Zbl 0942.46035, doi:10.1006/jfan.1998.3306.
- [16] KANG, SOORAN; PASK, DAVID; WEBSTER, SAMUEL B. G. Textile systems, coloured graphs and their applications to higher-rank graphs. Preprint, 2013. arXiv:1310.7651.
- [17] KRIEGER, WOLFGANG. On sofic systems. I. *Israel J. Math.* **48** (1984), no. 4, 305–330. MR0776312 (86j:54074), Zbl 0573.54032, doi:10.1007/BF02760631.
- [18] KRIEGER, W. On sofic systems II. *Israel J. Math.* **60** (1987), 167–176. MR0931874 (89k:54098), Zbl 0638.54034, doi:10.1007/BF02790789.
- [19] KUMJIAN, ALEX; PASK, DAVID. Higher rank graph C^* -algebras. *New York J. Math.* **6** (2000), 1–20. MR1745529 (2001b:46102), Zbl 0946.46044, arXiv:math/0007029.
- [20] KUMJIAN, ALEX; PASK, DAVID; RAEBURN, IAIN; RENAULT, JEAN. Graphs, groupoids and Cuntz–Krieger algebras. *J. Funct. Anal.* **144** (1997), no. 2, 505–541. MR1432596 (98g:46083), Zbl 0929.46055, doi:10.1006/jfan.1996.3001.
- [21] LIND, DOUGLAS; MARCUS, BRIAN. An introduction to symbolic dynamics and coding. *Cambridge University Press, Cambridge*, 1995. xvi+495 pp. ISBN: 0-521-55124-2; 0-521-55900-6. MR1369092 (97a:58050), Zbl 1106.37301, doi:10.1017/CBO9780511626302.
- [22] MARKLEY, NELSON G.; PAUL, MICHAEL E. Matrix subshifts for \mathbb{Z}^{ν} symbolic dynamics. *Proc. London Math. Soc.* (3) **43** (1981), no. 2, 251–272. MR0628277 (82i:54073), Zbl 0479.54022, doi:10.1112/plms/s3-43.2.251.
- [23] MATSUMOTO, KENGO. K -theory for C^* -algebras associated with subshifts. *Math. Scand.* **82** (1998), no. 2, 237–255. MR1646513 (2000e:46087), Zbl 0921.46078.
- [24] MATSUMOTO, KENGO. Presentations of subshifts and their topological conjugacy invariants. *Doc. Math.* **4** (1999), 285–340. MR1710375 (2000h:37013), Zbl 0926.37002.
- [25] MATSUMOTO, KENGO. C^* -algebras associated with presentations of subshifts. *Doc. Math.* **7** (2002), 1–30. MR1911208 (2004j:46083), Zbl 1007.46051.
- [26] MATSUMOTO, KENGO. Construction and pure infiniteness of C^* -algebras associated with lambda-graph systems. *Math. Scand.* **97** (2005), no. 1, 73–88. MR2179590 (2006e:46063), Zbl 1094.46041.
- [27] MATSUMOTO, KENGO. Actions of symbolic dynamical systems on C^* -algebras. *J. Reine Angew. Math.* **605** (2007), 23–49. MR2338126 (2009g:46131), Zbl 1123.46050, doi:10.1515/CRELLE.2007.026.
- [28] MATSUMOTO, KENGO. Textile systems on lambda-graph systems. *Yokohama Math. J.* **54** (2008), no. 2, 121–206. MR2488282 (2010e:37011), Zbl 1175.37017, arXiv:math/0607520.
- [29] MATSUMOTO, KENGO. Orbit equivalence in C^* -algebras defined by actions of symbolic dynamical systems. *Operator structures and dynamical systems*, 121–140, Contemporary Math., 503. *Amer. Math. Soc., Providence, RI*, 2009. MR2590619 (2012d:46166), Zbl 1194.46093,
- [30] MATSUMOTO, KENGO. Actions of symbolic dynamical systems on C^* -algebras. II. Simplicity of C^* -symbolic crossed products and some examples. *Math. Z.* **265** (2010), no. 4, 735–760. MR2652533 (2011g:46123), Zbl 1208.46057, arXiv:0705.3283, doi:10.1007/s00209-009-0538-3.
- [31] MATSUMOTO, KENGO. C^* -algebras associated with Hilbert C^* -quad modules of C^* -textile dynamical systems. Preprint, 2011. arXiv:1111.3091v1.
- [32] MATSUMOTO, KENGO. Cuntz–Krieger algebras associated with Hilbert C^* -quad modules of commuting matrices. *Math. Scand.* **117** (2015), no. 1, 126–149. arXiv:1201.1056v1.

- [33] MATSUMOTO, KENGO. C^* -algebras associated with Hilbert C^* -quad modules of finite type. *Int. J. Math. Math. Sci.* **2014** Art. ID 952068, 21 pp. MR3166746, Zbl 1286.46060, arXiv:1309.7704, doi:10.1155/2014/952068.
- [34] NASU, MASAKAZU. Textile systems for endomorphisms and automorphisms of the shift. *Mem. Amer. Math. Soc.* **114** (1995), no. 546, viii+215 pp. MR1234883 (95i:54051), Zbl 0845.54031, doi:10.1090/memo/0546.
- [35] PASCHKE, W. L. K -theory for actions of the circle group on C^* -algebras. *J. Operator Theory* **6** (1981), no. 1, 125–133. MR0637006 (82m:46074), Zbl 0478.46057.
- [36] PASK, DAVID; RAEBURN, IAIN; WEAVER, NATASHA A. A family of 2-graphs arising from two-dimensional subshifts. *Ergodic Theory Dynam. Systems* **29** (2009), no. 5, 1613–1639. MR2545020 (2010g:46115), Zbl 1187.46040. arXiv:0804.3447, doi:10.1017/S0143385708000795.
- [37] PASK, DAVID; RAEBURN, IAIN; WEAVER, NATASHA A. Periodic 2-graphs arising from subshifts. *Bull. Aust. Math. Soc.* **82** (2010), no. 1, 120–138. MR2669150 (2011f:46083), Zbl 1200.46060, arXiv:0911.0730, doi:10.1017/S0004972709001269.
- [38] PATERSON, ALAN L. T. Groupoids, inverse semigroups, and their operator algebras. Progress in Mathematics 170. *Birkhäuser Boston Inc., Boston, MA*, 1999. xvi+274 pp. ISBN: 0-8176-4051-7. MR1724106 (2001a:22003), Zbl 0913.22001, doi:10.1007/978-1-4612-1774-9.
- [39] PIMSNER, MICHAEL V. A class of C^* -algebras generalizing both Cuntz–Krieger algebras and crossed product by \mathbb{Z} . *Free probability theory* (Waterloo, ON, 1995), 189–212, Fields Inst. Commun., 12. *Amer. Math. Soc., Providence, RI*, 1997. MR1426840 (97k:46069), Zbl 0871.46028.
- [40] PIMSNER, M.; VOICULESCU, D. Exact sequences for K -groups and Ext-groups of certain cross-product C^* -algebras. *J. Operator Theory* **4** (1980), no. 1, 93–118. MR0587369 (82c:46074), Zbl 0474.46059.
- [41] RENAULT, JEAN. A groupoid approach to C^* -algebras. Lecture Notes in Mathematics, 793. *Springer, Berlin*, 1980. ii+160 pp. ISBN: 3-540-09977-8. MR0584266 (82h:46075), Zbl 0433.46049, doi:10.1007/BFb0091072.
- [42] ROBERTSON, DAVID I.; SIMS, AIDAN. Simplicity of C^* -algebras associated to higher-rank graphs. *Bull. Lond. Math. Soc.* **39** (2007), no. 2, 337–344. MR2323468 (2008g:46099), Zbl 1125.46045, arXiv:math/0602120, doi:10.1112/blms/bdm006.
- [43] ROBERTSON, GUYAN; STEGER, TIM. Affine buildings, tiling systems and higher rank Cuntz–Krieger algebras. *J. Reine Angew. Math.* **513** (1999), 115–144. MR1713322 (2000j:46109), Zbl 1064.46504, arXiv:1302.5593, doi:10.1515/crll.1999.057.
- [44] RØRDAM, M.; LARSEN, F.; LAUSTSEN, N. An introduction to K -theory for C^* -algebras. London Mathematical Society Student Texts, 49. *Cambridge University Press, Cambridge*, 2000. xii+242 pp. ISBN: 0-521-78334-8; 0-521-78944-3. MR1783408 (2001g:46001), Zbl 0967.19001.
- [45] ROSENBERG, JONATHAN. Appendix to: “Crossed products of UHF algebras by product type actions” by O. Bratteli. *Duke Math. J.* **46** (1979), no. 1, 25–26. MR0523599 (82a:46064), Zbl 0395.46049, doi:10.1215/S0012-7094-79-04602-7.
- [46] ROSENBERG, JONATHAN; SCHOCHET, CLAUDE. The Künneth theorem and the universal coefficient theorem for Kasparov’s generalized K -functor. *Duke Math. J.* **55** (1987), no. 2, 431–474. MR0894590 (88i:46091), Zbl 0644.46051, doi:10.1215/S0012-7094-87-05524-4.
- [47] SCHMIDT, KLAUS. Dynamical systems of algebraic origin. Progress in Mathematics, 128. *Birkhäuser, Verlag, Basel*, 1995. xviii+310 pp. ISBN: 3-7643-5174-8. MR1345152 (97c:28041), Zbl 0833.28001, doi:10.1007/978-3-0348-0277-2.
- [48] WEGGE-OLSEN, N. E. K -theory and C^* -algebras. A friendly approach. Oxford Science Publications. *The Clarendon Press, Oxford University Press, New York*, 1993. xii+370 pp. ISBN: 0-19-859694-4. MR1222415 (95c:46116), Zbl 0780.46038.

(Kengo Matsumoto) DEPARTMENT OF MATHEMATICS, JOETSU UNIVERSITY OF EDUCATION, JOETSU 943-8512, JAPAN
kengo@juen.ac.jp

This paper is available via <http://nyjm.albany.edu/j/2015/21-54.html>.