

Homogeneous holomorphic hermitian principal bundles over hermitian symmetric spaces

Indranil Biswas and Harald Upmeyer

ABSTRACT. We give a complete characterization of invariant integrable complex structures on principal bundles defined over hermitian symmetric spaces, using the Jordan algebraic approach for the curvature computations. In view of possible generalizations, the general setup of invariant holomorphic principal fibre bundles is described in a systematic way.

CONTENTS

1. Introduction	21
2. Homogeneous H -bundles	22
3. Connexions and complexions	26
4. Homogeneous connexions and complexions	32
5. Curvature and Integrability	36
6. The symmetric case	40
7. Concluding remarks	45
References	46

1. Introduction

The classification of hermitian holomorphic vector bundles, or more general holomorphic principal fibre bundles, over a complex manifold M is a central problem in algebraic geometry and quantization theory, e.g., for a compact Riemann surface M . In geometric quantization, where $M = G/K$ is a co-adjoint orbit, G -invariant principal fibre bundles have been investigated from various points of view [Bo, Ra, OR]. In case M is a hermitian symmetric space of noncompact type, a complete characterization of invariant integrable complex structures on principal bundles over M was obtained in [BiM] (for the unit disk) and [Bi] (for all bounded symmetric domains).

Received August 5, 2015.

2010 *Mathematics Subject Classification.* 32M10, 14M17, 32L05.

Key words and phrases. Irreducible hermitian symmetric space; principal bundle; homogeneous complex structure; hermitian structure.

The main objective of this paper is to treat the dual case of compact hermitian symmetric spaces, and to show that the compact case (as well as the flat case) leads to exactly the same characterization, resulting in an explicit duality correspondence for invariant integrable complex principal bundles (with hermitian structure) between the noncompact type, the compact type and the flat type as well. The proof is carried out using the Jordan theoretic approach towards hermitian symmetric spaces [Lo, FK]. Of course, the traditional Lie triple system approach could be used instead, but Jordan triple systems (essentially the hermitian polarization of the underlying Lie triple system) make things more transparent and somewhat more elementary.

More importantly, the Jordan triple approach leads in a natural way to more general complex homogeneous (nonsymmetric) manifolds G/C where C is a proper subgroup of K . These manifolds are fibre bundles over G/K with compact fibres given by Jordan theoretic flag manifolds. Again, there exists a duality between such spaces of compact, noncompact and flat type, and in a subsequent paper [BiU] the duality correspondence for invariant fibre bundles, proved here in the hermitian symmetric case $C = K$, will be studied in the more general setting. In view of these more general situations, the current paper describes the general setup for homogeneous holomorphic principal fibre bundles in a careful way, specializing to the symmetric case only in the last section.

As a next step beyond the classification, its dependence on the underlying complex structure on M is of fundamental importance. While the hermitian symmetric case G/K has a unique G -invariant complex structure, the more general flag manifold bundles G/C have an interesting moduli space of invariant complex structures. It is a challenging problem whether this moduli space carries a projectively flat connexion (with values in the vector bundle of holomorphic sections) similar to the case of abelian varieties, [Mu], [We], or Chern–Simons theory [ADW], [Wi].

From the geometric quantization point of view, it is also of interest to describe the spaces of holomorphic sections, given by suitable Dolbeault operators, in an explicit way.

2. Homogeneous H -bundles

Let M be a manifold and G a connected real Lie group, with Lie algebra \mathfrak{g} , acting smoothly on M . Denoting the action $G \times M \rightarrow M$ by $(a, x) \mapsto a(x)$, we define $R_x^M : G \rightarrow M$, $x \in M$, by

$$R_x^M(a) = a(x) \quad \forall (a, x) \in G \times M.$$

Let H be a connected complex Lie group; its Lie algebra will be denoted by \mathfrak{h} . Fix a maximal compact subgroup $L \subset H$; all such subgroups are conjugate in H . The Lie algebra of L is denoted by \mathfrak{l} . Let Q be a C^∞ principal H -bundle over M , with projection $\pi : Q \rightarrow M = Q/H$. The free action $Q \times H \rightarrow Q$ is

written as $(q, b) \mapsto qb$. Define $R_b^Q : Q \rightarrow Q$ and $L_q^Q : H \rightarrow Q$ by

$$R_b^Q q = L_q^Q b = qb \quad \forall (q, b) \in Q \times H.$$

For notational convenience, we have omitted the parentheses.

Then $\pi \circ R_b^Q = \pi$ and $\ker(d\pi)_q \subset T_q Q$ is the “vertical” subspace of the tangent space $T_q Q$ at $q \in Q$. We call Q an *equivariant H -bundle* if there is a C^∞ action $G \times Q \rightarrow Q$, denoted by $(a, q) \mapsto aq$, such that $\pi(aq) = a\pi(q)$ and

$$a(qb) = (aq)b.$$

Define $L_a^Q : Q \rightarrow Q$ and $R_q^Q : G \rightarrow Q$ by

$$L_a^Q q = R_q^Q a = aq \quad \forall (a, q) \in G \times Q.$$

Then $L_a^Q \circ R_b^Q = R_b^Q \circ L_a^Q$ and $\pi \circ L_a^Q = L_a^M \circ \pi$ for all $a \in G, b \in H$. A *hermitian structure* on a principal H -bundle Q is a principal subbundle $P \subset Q$ with structure group L , i.e., we have

$$R_b^Q : P \rightarrow P \quad \forall b \in L$$

and the action of L on the fibre P_x is transitive for all $x \in M$. We also say that (Q, P) is a *hermitian H -bundle*. An *equivariant hermitian H -bundle* is an equivariant H -bundle Q endowed with a hermitian structure $P \subset Q$ such that $G \cdot P = P$, i.e., we have

$$L_a^Q P = P \quad \forall a \in G.$$

In this case P becomes an equivariant L -bundle. When G acts transitively on M , we call Q a *homogeneous H -bundle*. In the homogeneous case, fix a base point $o \in M$ and put

$$K = \{k \in G \mid k(o) = o\}.$$

Then $M = G/K$. The Lie algebra of K will be denoted by \mathfrak{k} .

The following proposition is straightforward to prove.

Proposition 2.1. *Let $f : K \rightarrow H$ be a (real-analytic) homomorphism. Consider the quotient manifold*

$$Q := G \times_{K, f} H$$

consisting of all equivalence classes

$$(2.1) \quad \langle g|h \rangle = \langle gk^{-1}|f(k)h \rangle,$$

with $g \in G, h \in H$ and $k \in K$. (This bracket notation is used to avoid confusion with the commutator bracket.) Then Q becomes a homogeneous H -bundle, with projection $\pi(g|h) = g(o) = R_o^M(g)$. The action of $(a, b) \in G \times H$ is given by

$$a\langle g|h \rangle b = \langle ag|hb \rangle.$$

If in addition, $f(K) \subset L$, we obtain a hermitian homogeneous principal H -bundle

$$P := G \times_{K, f} L \subset Q := G \times_{K, f} H.$$

In the set-up of Proposition 2.1, the maps $L_a^Q : Q \rightarrow Q$ and $R_{g|h}^Q : G \rightarrow Q$ have the form

$$L_a^Q \langle g|h \rangle = R_{g|h}^Q(a) = \langle ag|h \rangle.$$

For any $\ell \in H$, let

$$I_\ell^H h = \ell h \ell^{-1}$$

be the inner automorphism induced by ℓ . Then $d_e I_\ell^H = \text{Ad}_\ell^{\mathfrak{h}}$. The conjugate homomorphism

$$I_\ell^H \circ f : K \rightarrow H$$

induces the H -bundle isomorphism

$$G \times_{K,f} H \rightarrow G \times_{K, I_\ell^H \circ f} H$$

mapping the equivalence class $\langle g|h \rangle_f$ to $\langle g|I_\ell^H h \rangle_{I_\ell^H \circ f}$.

Theorem 2.2. *Every homogeneous principal H -bundle Q on $M = G/K$ is isomorphic to $G \times_{K,f} H$ for a homomorphism $f : K \rightarrow H$, which is unique up to conjugation by elements $\ell \in H$. Similarly, every hermitian homogeneous H -bundle (Q, P) is isomorphic to the pair*

$$G \times_{K,f} L \subset G \times_{K,f} H$$

for a homomorphism $f : K \rightarrow L \subset H$, which is unique up to conjugation by an element in L . More precisely, for any base point $\mathfrak{o} \in Q_{\mathfrak{o}}$ (respectively, $\mathfrak{o} \in P_{\mathfrak{o}}$) there exists a unique homomorphism $f_{\mathfrak{o}} : K \rightarrow H$ (respectively, $f_{\mathfrak{o}} : K \rightarrow L$) such that

$$(2.2) \quad k\mathfrak{o} = \mathfrak{o}f_{\mathfrak{o}}(k) \quad \forall k \in K,$$

and

$$(2.3) \quad \langle g|h \rangle \mapsto g\mathfrak{o}h$$

defines an isomorphism $G \times_{K, f_{\mathfrak{o}}} H \rightarrow Q$ of equivariant H -bundles (respectively, an isomorphism $G \times_{K, f_{\mathfrak{o}}} L \rightarrow P$ of hermitian equivariant H -bundles). Another base point $\mathfrak{o}' = \mathfrak{o}\ell^{-1}$, with $\ell \in H$ (respectively, $\ell \in L$), corresponds to the homomorphism $f_{\mathfrak{o}'} = I_\ell \circ f_{\mathfrak{o}}$.

Proof. Let Q be a homogeneous H -bundle. Choose $\mathfrak{o} \in Q$ with $\pi(\mathfrak{o}) = o$. Since the fibre $Q_{\mathfrak{o}}$ is preserved by K , and H acts freely on $Q_{\mathfrak{o}}$, there exists a unique map $f_{\mathfrak{o}} : K \rightarrow H$ such that (2.2) holds for all $k \in K$. Then

$$\begin{aligned} \mathfrak{o}f_{\mathfrak{o}}(k_1 k_2) &= (k_1 k_2)\mathfrak{o} = k_1(k_2\mathfrak{o}) = k_1(\mathfrak{o}f_{\mathfrak{o}}(k_2)) = (k_1\mathfrak{o})f_{\mathfrak{o}}(k_2) \\ &= (\mathfrak{o}f_{\mathfrak{o}}(k_1))f_{\mathfrak{o}}(k_2) = \mathfrak{o}(f_{\mathfrak{o}}(k_1)f_{\mathfrak{o}}(k_2)) \end{aligned}$$

for all $k_1, k_2 \in K$. Since H operates freely on Q , this implies that $f_{\mathfrak{o}}(k_1 k_2) = f_{\mathfrak{o}}(k_1)f_{\mathfrak{o}}(k_2)$, and hence $f_{\mathfrak{o}} : K \rightarrow H$ is a (real-analytic) homomorphism. For all $b \in H$ we have

$$k\mathfrak{o}b = (\mathfrak{o}f_{\mathfrak{o}}(k))b = \mathfrak{o}(f_{\mathfrak{o}}(k)b).$$

The resulting identity

$$(ak)\mathfrak{o}b = a(k\mathfrak{o}b) = a(\mathfrak{o}(f_{\mathfrak{o}}(k)b)) = a\mathfrak{o}(f_{\mathfrak{o}}(k)b)$$

shows that (2.3) defines an isomorphism $G \times_{K, f_o} H \rightarrow Q$ of equivariant H -bundles, mapping the “base point” $\langle e|e \rangle$ to \mathfrak{o} . If $\mathfrak{o}' \in Q_o$ is another base point, there exists a unique $\ell \in H$ such that $\mathfrak{o}' = \mathfrak{o}\ell^{-1}$. It follows that

$$k\mathfrak{o}' = k\mathfrak{o}\ell^{-1} = \mathfrak{o}f_o(k)\ell^{-1} = \mathfrak{o}'\ell f_o(k)\ell^{-1}.$$

Thus the new base point \mathfrak{o}' corresponds to the conjugate homomorphism

$$f_{\mathfrak{o}'\ell^{-1}}(k) = \ell f_o(k)\ell^{-1} = I_\ell^H f_o(k).$$

In the hermitian case, for any base point $\mathfrak{o} \in P_o \subset Q_o$ the defining identity (2.2) implies $f_o(k) \in L$ for all $k \in K$. Another base point $\mathfrak{o}' = \mathfrak{o}\ell^{-1} \in P_o$ differs by a unique element $\ell \in L$. In both cases, since G acts transitively on M , the G -invariance condition implies that the entire construction is independent of the choice of base point $o \in M$. \square

In view of (2.2), the homomorphism f_o could be denoted by $f_o = I_o^{-1}$, i.e., $k\mathfrak{o} = \mathfrak{o}I_o^{-1}(k)$. In this notation, the identity $I_{\mathfrak{o}'\ell^{-1}}^{-1} = I_\ell \circ I_o^{-1}$ is obvious. It will be convenient to express the tangent spaces in an explicit manner using equivalence classes. Note that the differential $d_e f : \mathfrak{k} \rightarrow \mathfrak{h}$ of f at the unit element $e \in K$ is a Lie algebra homomorphism.

Lemma 2.3. *For a given class $\langle g|h \rangle \in Q := G \times_{K, f} H$ the tangent space $T_{g|h}Q$ consists of all equivalence classes*

$$\langle \dot{g}|\dot{h} \rangle = \left\langle (d_g R_{k^{-1}}^G)\dot{g} - (d_e L_{gk^{-1}}^G)\kappa | (d_h L_{f(k)}^H)\dot{h} + (d_e R_{f(k)h}^H)(d_e f)\kappa \right\rangle,$$

where $\dot{g} \in T_g G$, $\dot{h} \in T_h H$, $k \in K$ and $\kappa \in \mathfrak{k}$.

Here we regard TG and TH as the disjoint union of the respective tangent spaces, so that the first expression is evaluated at $\langle g|h \rangle$ whereas the second expression is evaluated at the same class written as $\langle gk^{-1}|f(k)h \rangle$. The identity follows from differentiating the relation $\langle g_t|h_t \rangle = \langle g_t k_t^{-1}|f(k_t)h_t \rangle$ at $t = 0$, where $k_t \in K$ satisfies $k_0 = k$ and $\partial_t^0 k_t = \kappa$. As special cases we obtain

$$(2.4) \quad \begin{aligned} \langle \dot{g}|\dot{h} \rangle &= \left\langle (d_g R_{k^{-1}}^G)\dot{g} | (d_h L_{f(k)}^H)\dot{h} \right\rangle \\ &= \left\langle \dot{g} - (d_e L_g^G)\kappa | \dot{h} + (d_e R_h^H)(d_e f)\kappa \right\rangle \end{aligned}$$

putting $\kappa = 0$ or $k = e$, respectively. The projection π has the differential

$$(d_{g|h}\pi)\langle \dot{g}|\dot{h} \rangle = \left(\frac{d}{dt}(g_t)(0) \right) (o) = (d_g R_o^M)\dot{g}.$$

Thus the vertical subspace is given by

$$\ker(d_{g|h}\pi) = \{ \langle \dot{g}|\dot{h} \rangle \mid \dot{g} \in \ker(d_g R_o^M) \subset T_g G, \dot{h} \in T_h H \}.$$

For $\beta \in \mathfrak{h}$, the fundamental vector field ρ_β^Q has the form

$$(\rho_\beta^Q)_{g|h} = \partial_t^0 \langle g|hb_t \rangle = \langle 0_g | (d_e L_h^H)\beta \rangle = \langle 0_g | (d_e R_h^H) \text{Ad}_h^H \beta \rangle.$$

3. Connexions and complexions

A *connexion* on a principal H -bundle Q is a smooth distribution $q \mapsto T_q^\Theta Q$ of “horizontal” subspaces of $T_q Q$ such that $T_q Q = T_q^\Theta Q \oplus \ker(d_q \pi)$ and

$$T_{qb}^\Theta Q = (d_q R_b^Q)(T_q^\Theta Q) \quad \forall (q, b) \in Q \times H.$$

We use the same symbol for the associated connexion 1-form $\Theta_q : T_q Q \rightarrow \mathfrak{h}$ on Q , uniquely determined by the condition that $X \in T_q Q$ has the horizontal projection

$$X^\Theta = X - (d_e L_q^Q)(\Theta_q(X)).$$

A connexion Θ on an equivariant H -bundle Q is called *invariant* if

$$(d_q L_a^Q)(T_q^\Theta Q) = T_{aq}^\Theta Q \quad \forall a \in G.$$

In this case the associated connexion 1-form satisfies

$$\Theta_q = \Theta_{aq}(d_q L_a^Q).$$

Let $Q \times_H \mathfrak{h}$ denote the associated bundle of type Ad_H , with fibres

$$(Q \times_H \mathfrak{h})_x = \{[q : \beta] = [qh : \text{Ad}_h^{-1} \beta] \mid q \in Q_x, \beta \in \mathfrak{h}\}$$

for $x \in M$, with $h \in H$ being arbitrary. By [KN, Section II.5] every tensorial i -form on Q is given by

$$(\mathbf{C}_q^Q \circ d_q \pi)(X_q^1, \dots, X_q^i) := \mathbf{C}_q^Q((d_q \pi)X_q^1, \dots, (d_q \pi)X_q^i)$$

for $X_q^1, \dots, X_q^i \in T_q Q$, where

$$\mathbf{C}_x(v_1 \wedge \dots \wedge v_i) = [q : \mathbf{C}_q^Q(v_1 \wedge \dots \wedge v_i)] \quad \forall q \in Q_x$$

is an i -form of type Ad_H (on M), with homogeneous lift $\mathbf{C}_q^Q : \bigwedge^i T_x M \rightarrow \mathfrak{h}$ having the right invariance property

$$\mathbf{C}_{qb}^Q = \text{Ad}_{b^{-1}}^H \mathbf{C}_q^Q \quad \forall b \in H.$$

An i -form \mathbf{C} of type Ad_H is called *invariant* if

$$(3.1) \quad \mathbf{C}_{aq}^Q(d_x L_a^M) = \mathbf{C}_q^Q \quad \forall (a, q) \in G \times Q.$$

Proposition 3.1.

(i) Let Θ^0 be a connexion on Q . If \mathbf{C} is a 1-form of type Ad_H , then

$$(3.2) \quad \Theta_q = \Theta_q^0 + \mathbf{C}_q^Q \circ d_q \pi \quad \forall q \in Q$$

is a connexion 1-form on Q . Every connexion 1-form Θ on Q arises this way.

(ii) In the equivariant case, let Θ^0 be an invariant connexion on Q . Then Θ is invariant if and only if \mathbf{C} is invariant, i.e.,

$$(3.3) \quad \mathbf{C}_{aq}^Q(d_x L_a^M) = \mathbf{C}_q^Q \quad \forall (a, q) \in G \times Q.$$

Proof. Part (i) is well-known. For part (ii) let the connexion Θ be invariant. The condition $\pi \circ L_a^Q = L_a^M \circ \pi$ implies that $(d_{aq}\pi)(d_q L_a^Q) = (d_x L_a^M)(d_q \pi)$, and hence we have

$$\begin{aligned} \mathbf{C}_q^Q(d_q \pi) &= \Theta_q - \Theta_q^0 = (\Theta_{aq} - \Theta_{aq}^0)(d_q L_a^Q) \\ &= \mathbf{C}_{aq}^Q(d_{aq}\pi)(d_q L_a^Q) = \mathbf{C}_{aq}^Q(d_x L_a^M)(d_q \pi). \end{aligned}$$

Since $d_q \pi$ is surjective, (3.3) follows. The converse is proved in a similar way. \square

If (3.2) holds, we say that Θ is related to Θ^0 via \mathbf{C} . For a hermitian H -bundle (Q, P) a connexion Ξ on P is called *invariant* if

$$(d_p L_a^P)T_p^\Xi P = T_{ap}^\Xi P \quad \forall (a, p) \in G \times P.$$

In this case the associated connexion 1-form satisfies

$$\Xi_p = \Xi_{ap}(d_p L_a^P).$$

By [KN, Proposition II.6.2] every connexion Ξ on P has a unique extension to a connexion $\iota \Xi$ on Q such that

$$T_p^\Xi P = T_p^{\iota \Xi} Q \quad \forall p \in P \subset Q.$$

Equivalently, the connexion forms satisfy

$$(\iota \Xi)_p|_{T_p P} = \Xi_p \quad \forall p \in P.$$

A connexion Θ on Q is called *hermitian* if $\Theta = \iota \Xi$ for a (unique) connexion Ξ on P . Thus hermitian connexions on Q are in 1-1 correspondence with connexions on P . A connexion Ξ on P is invariant if and only if its extension $\iota \Xi$ is invariant.

For hermitian H -bundles (P, Q) , we have i -forms of type Ad_L written as

$$\mathbf{A}_x(v_1 \wedge \cdots \wedge v_i) = [p : \mathbf{A}_p^P(v_1 \wedge \cdots \wedge v_i)] \quad \forall p \in P_x,$$

with homogeneous lift $\mathbf{A}_p^P : \bigwedge^i T_x M \rightarrow \mathfrak{l}$ having the right invariance property

$$(3.4) \quad \mathbf{A}_{p\ell}^P = \text{Ad}_{\ell^{-1}}^L \mathbf{A}_p^P \quad \forall \ell \in L.$$

Let \mathfrak{l} be the Lie algebra of L . For each $x \in M$, there is a linear injection of fibres

$$\iota_x : (P \times_L \mathfrak{l})_x \rightarrow (Q \times_H \mathfrak{h})_x, \quad [p, \beta] \mapsto [p, \iota \beta],$$

where $p \in P_x, \beta \in \mathfrak{l}$ and $\iota : \mathfrak{l} \rightarrow \mathfrak{h}$ is the inclusion map. The map ι_x is well-defined because

$$[p_1, \beta_1] = [p_2, \beta_2] \in (P \times_L \mathfrak{l})_x$$

implying that $p_2 = p_1 \ell$ and $\beta_2 = \text{Ad}_{\ell^{-1}}^L \beta_1$ for some $\ell \in L$. Therefore, we also have $\iota \beta_2 = \text{Ad}_{\ell^{-1}}^H \iota \beta_1$. To show that ι_x is injective, suppose that

$$[p_1, \iota \beta_1] = [p_2, \iota \beta_2] \in (Q \times_H \mathfrak{h})_x.$$

Then we have $p_2 = p_1 h$ and $\iota\beta_2 = \text{Ad}_{h^{-1}}^H \iota\beta_1$ for some $h \in H$. Since $p_1, p_2 \in P_x$ it follows that $h \in L$ and hence $\beta_2 = \text{Ad}_{h^{-1}}^L \beta_1$.

As a consequence, an i -form \mathbf{A} of type Ad_L induces an i -form $\iota\mathbf{A}$ of type Ad_H , with homogeneous lift

$$(\iota\mathbf{A})_{ph}^Q := \text{Ad}_{h^{-1}}^H \mathbf{A}_p^P : \bigwedge^i T_o M \rightarrow \mathfrak{h} \quad \forall (p, h) \in P \times H.$$

Then \mathbf{A} is invariant in the sense that

$$\mathbf{A}_{ap}^P(d_x L_a^M) = \mathbf{A}_p^P \quad \forall (a, p) \in G \times P$$

if and only if $\iota\mathbf{A}$ is invariant as in (3.1).

Proposition 3.2.

- (i) For a hermitian bundle (P, Q) , let Ξ^0 be a connexion on P . If \mathbf{A} is a 1-form of type Ad_L , then

$$(3.5) \quad \Xi_p = \Xi_p^0 + \mathbf{A}_p^P \circ d_p \pi \quad \forall p \in P$$

is a connexion 1-form on P . Every connexion 1-form Ξ on P arises this way. We also have

$$(\iota\Xi)_q = (\iota\Xi)_q^0 + (\iota\mathbf{A})_q^Q \circ d_q \pi \quad \forall q \in Q.$$

- (ii) In the equivariant case, let Ξ^0 be an invariant connexion on P . Then Ξ is invariant if and only if \mathbf{A} is invariant, i.e.,

$$(3.6) \quad \mathbf{A}_{ap}^P(d_x L_a^M) = \mathbf{A}_p^P \quad \forall (a, p) \in G \times P.$$

If (3.5) holds, we say that Ξ is related to Ξ^0 via \mathbf{A} . In this case, $\iota\Xi$ is related to $\iota\Xi^0$ via $\iota\mathbf{A}$. Now suppose that (M, j) is an almost complex manifold, with almost complex structure $j_x \in \text{End}(T_x M)$, $x \in M$, having the left invariance property

$$(d_x L_g^M)j_x = j_{gx}(d_x L_g^M) \quad \forall (g, x) \in G \times M.$$

Let H be a complex Lie group. Consider the bi-invariant complex structure $i_h \in \text{End}(T_h H)$ on H such that $i_e \in \text{End}(\mathfrak{h})$ is multiplication by $\sqrt{-1}$. See [At], [Kos] for complex structures on principal bundles.

Definition 3.3. An almost complex structure $J_q \in \text{End}(T_q Q)$ on an H -bundle Q is called a *complexion* if

$$(3.7) \quad (d_q \pi)J_q = j_x(d_q \pi)$$

and the map $Q \times H \rightarrow Q$ is almost-holomorphic. Writing

$$qb = L_q^Q(b) = R_b^Q(q)$$

for $q \in Q$, $b \in H$, this means that

$$(3.8) \quad (d_q R_b^Q)J_q = J_{qb}(d_q R_b^Q)$$

$$(3.9) \quad J_{qb}(d_b L_q^Q) = (d_b L_q^Q)i_b.$$

In the equivariant case, a complexation J is called *invariant* if in addition

$$(d_q L_a^Q) J_q = J_{aq} (d_q L_a^Q) \quad \forall a \in G.$$

By right invariance the condition (3.9) is equivalent to

$$J_q (d_e L_q^Q) \beta = (d_e L_q^Q) (\sqrt{-1} \beta) \quad \forall \beta \in \mathfrak{h}.$$

Since the fibre $(Q \times_H \mathfrak{h})_x$ is a complex vector space, the notion of (p, q) -forms of type Ad_H makes sense.

Proposition 3.4.

(i) *Let J^0 be a complexation on Q . If \mathbf{B} is a $(0, 1)$ -form of type Ad_H , then*

$$(3.10) \quad J_q = J_q^0 + (d_e L_q^Q) (\mathbf{B}_q^Q \circ d_q \pi)$$

defines a complexation J on Q . Every complexation J on Q arises this way.

(ii) *Let J^0 be an invariant complexation on an equivariant bundle Q . Then J is invariant if and only if \mathbf{B} is invariant, i.e.,*

$$(3.11) \quad \mathbf{B}_{aq}^Q (d_x L_a^M) = \mathbf{B}_q^Q \quad \forall (a, q) \in G \times Q.$$

Proof. By (3.7) we have

$$(d_q \pi) (J_q - J_q^0) = (j_x - j_x) (d_q \pi) = 0.$$

Thus $\text{Image}(J_q - J_q^0) \subset \text{Ker}(d_q \pi)$. Since $d_e L_q^Q : \mathfrak{h} \rightarrow \text{Ker}(d_q \pi)$ is an isomorphism, there exists a 1-form $\Psi_q : T_q Q \rightarrow \mathfrak{h}$ such that

$$J_q - J_q^0 = (d_e L_q^Q) \Psi_q.$$

For $b \in H$ we have $R_b^Q \circ L_q^Q = L_{qb}^Q \circ I_{b^{-1}}^H$, and hence

$$(d_q R_b^Q) (d_e L_q^Q) = (d_e L_{qb}^Q) \text{Ad}_{b^{-1}}^H.$$

It follows that

$$\begin{aligned} (d_e L_{qb}^Q) \Psi_{qb} (d_q R_b^Q) &= (J_{qb} - J_{qb}^0) (d_q R_b^Q) = (d_q R_b^Q) (J_q - J_q^0) \\ &= (d_q R_b^Q) (d_e L_q^Q) \Psi_q = (d_e L_{qb}^Q) \text{Ad}_{b^{-1}}^H \Psi_q. \end{aligned}$$

Therefore, we have $\Psi_{qb} (d_q R_b^Q) = \text{Ad}_{b^{-1}}^H \Psi_q$, so Ψ is pseudo-tensorial. For $\beta \in \mathfrak{h}$ we have

$$J_q (d_e L_q^Q) \beta = (d_e L_q^Q) (\sqrt{-1} \beta) = J_q^0 (d_e L_q^Q) \beta.$$

Therefore, we have $(J_q - J_q^0)|_{\text{Ker}(d_q \pi)} = 0$, which implies that Ψ is tensorial. Hence there exists a unique 1-form \mathbf{B} of type Ad_H such that $\Psi_q = \mathbf{B}_q^Q (d_q \pi)$.

We also have $(J_q - J_q^0)^2 = 0$ and hence

$$\begin{aligned}
(d_e L_q^Q)(\mathbf{B}_q^Q j_x + \sqrt{-1}\mathbf{B}_q^Q)(d_q \pi) &= (d_e L_q^Q)\mathbf{B}_q^Q j_x(d_q \pi) + (d_e L_q^Q)\sqrt{-1}\mathbf{B}_q^Q(d_q \pi) \\
&= (d_e L_q^Q)\mathbf{B}_q^Q(d_q \pi)J_q^0 + J_q^0(d_e L_q^Q)\mathbf{B}_q^Q(d_q \pi) \\
&= (J_q - J_q^0)J_q^0 + J_q^0(J_q - J_q^0) \\
&= J_q^2 - (J_q - J_q^0)^2 - (J_q^0)^2 = J_q^2 - (J_q^0)^2 \\
&= -id + id = 0.
\end{aligned}$$

Since $d_e L_q^Q$ is invertible, we obtain

$$\mathbf{B}_q^Q j_x + \sqrt{-1}\mathbf{B}_q^Q = 0.$$

Thus \mathbf{B} is of type $(0, 1)$.

For part (ii), let J be invariant. Using $L_a^Q \circ L_q^Q = L_{aq}^Q$ and $L_a^M \circ \pi = \pi \circ L_a^Q$ we obtain

$$\begin{aligned}
(d_e L_{aq}^Q)\mathbf{B}_{aq}^Q(d_x L_a^M)(d_q \pi) &= (d_e L_{aq}^Q)\mathbf{B}_{aq}^Q(d_{aq}\pi)(d_q L_a^Q) = (J_{aq} - J_{aq}^0)(d_q L_a^Q) \\
&= (d_q L_a^Q)(J_q - J_q^0) = (d_q L_a^Q)(d_e L_q^Q)\mathbf{B}_q^Q(d_q \pi) \\
&= (d_e L_{aq}^Q)\mathbf{B}_q^Q(d_q \pi).
\end{aligned}$$

Since $d_e L_{aq}^Q$ is invertible, (3.11) follows. The converse is proved in a similar way. \square

If (3.10) holds, we say that J is related to J^0 via \mathbf{B} . There is a close relationship between (invariant) connexions and complexions: Every connexion Θ on Q induces a unique complexion J^Θ which is ‘‘horizontal’’ in the sense that

$$J_q^\Theta : T_q^\Theta Q \rightarrow T_q^\Theta Q \quad \forall q \in Q.$$

In fact, for vertical tangent vectors we have

$$(3.12) \quad J_q^\Theta(d_e L_q^Q)\beta = (d_e L_q^Q)(\sqrt{-1}\beta) \quad \forall \beta \in \mathfrak{h},$$

and if $Y \in T_q^\Theta Q$ is horizontal, then $J_q^\Theta(Y) \in T_q^\Theta Q$ is uniquely determined by the condition $(d_q \pi)J_q^\Theta(Y) = j_x(d_q \pi)(Y)$. If Θ is invariant, then the associated complexion J^Θ is also invariant.

Proposition 3.5.

- (i) *If a connexion Θ is related to Θ^0 via \mathbf{C} , then the induced complexion J^Θ is related to J^0 via \mathbf{B} , where*

$$(3.13) \quad \mathbf{B}_x = \sqrt{-1}\mathbf{C}_x - \mathbf{C}_x j_x \quad \forall x \in M.$$

- (ii) *Θ and Θ^0 induce the same complexion if and only if \mathbf{C} is a $(1, 0)$ -form, i.e., $\mathbf{C}_q^Q : T_x M \rightarrow \mathfrak{h}$ is \mathbb{C} -linear.*

- (iii) If J^0 is induced by a connexion Θ^0 and J is related to J^0 via \mathbf{B} , then J is induced by the connexion

$$\Theta_q = \Theta_q^0 - \frac{\sqrt{-1}}{2} \mathbf{B}_q^Q(d_q\pi)$$

which is related to Θ^0 via $-\frac{\sqrt{-1}}{2}\mathbf{B}$.

Proof. In order to check (3.10) on a tangent vector $X \in T_qQ$, we may assume that X is Θ^0 -horizontal, since for vertical vectors of the form $X = (d_eL_q^Q)\beta$, $\beta \in \mathfrak{h}$, both sides of (3.10) agree. Thus assume that $\Theta_q^0 X = 0$. Then $\Theta_q X = \mathbf{C}_q^Q(d_q\pi)X$. Since J^0 is induced by Θ^0 , it follows that $J_q^0 X$ is also Θ^0 -horizontal, and hence $\Theta_q J_q^0 X = \mathbf{C}_q^Q(d_q\pi)J_q^0 X$. Therefore,

$$X - (d_eL_q^Q)\mathbf{C}_q^Q(d_q\pi)X = X - (d_eL_q^Q)\Theta_q X$$

is Θ -horizontal. It follows that

$$U := J_q^\Theta X - (d_eL_q^Q)\mathbf{C}_q^Q(d_q\pi)X = J_q^\Theta \left(X - (d_eL_q^Q)\mathbf{C}_q^Q(d_q\pi)X \right),$$

$$V := J_q^0 X - (d_eL_q^Q)\mathbf{C}_q^Q(d_q\pi)J_q^0 X = J_q^0 X - (d_eL_q^Q)\Theta_q J_q^0 X$$

are both Θ -horizontal. Since $(d_q\pi)U = j_x(d_q\pi)X = (d_q\pi)V$, it follows that $U = V$, i.e.,

$$J_q^\Theta X - (d_eL_q^Q)\sqrt{-1}\mathbf{C}_q^Q(d_q\pi)X = J_q^0 X - (d_eL_q^Q)\mathbf{C}_q^Q(d_q\pi)J_q^0 X.$$

Therefore, we have

$$\begin{aligned} (d_eL_q^Q)\mathbf{B}_q^Q(d_q\pi)X &= J_q^\Theta X - J_q^0 X \\ &= (d_eL_q^Q) \left(\sqrt{-1}\mathbf{C}_q^Q(d_q\pi)X - \mathbf{C}_q^Q(d_q\pi)J_q^0 X \right). \end{aligned}$$

Since $d_eL_q^Q$ is injective, using (3.7) it follows that

$$\mathbf{B}_q^Q(d_q\pi)X = \sqrt{-1}\mathbf{C}_q^Q(d_q\pi)X - \mathbf{C}_q^Q(d_q\pi)J_q^0 X = (\sqrt{-1}\mathbf{C}_q^Q - \mathbf{C}_q^Q j_x)(d_q\pi)X.$$

We now have $\mathbf{B}_q^Q = \sqrt{-1}\mathbf{C}_q^Q - \mathbf{C}_q^Q j_x$ on T_xM because $d_q\pi$ is surjective. This proves part (i).

Part (ii) is a direct consequence, since $\mathbf{B} = 0$ if and only if \mathbf{C} is \mathbb{C} -linear.

For part (iii), since $\mathbf{B}_q^Q : T_xM \rightarrow \mathfrak{h}$ is \mathbb{C} -antilinear, the 1-form

$$\mathbf{C}_x := -\frac{\sqrt{-1}}{2} \mathbf{B}_x$$

yields \mathbf{B}_x via (3.13). \square

In the hermitian case, we can sharpen the correspondence as follows:

Proposition 3.6.

- (i) For a hermitian H -bundle (Q, P) , a complexion J on Q is induced by a unique hermitian connexion $\iota\Xi$, with Ξ being a connexion on P . Thus there is a 1-1 correspondence between complexions on Q , connexions on P , and hermitian connexions on Q .

- (ii) *In the equivariant case, J is invariant if and only if Ξ (equivalently, $\iota\Xi$) is invariant.*

Proof. Let Ξ^0 be a connexion on P , and let J^0 be the complexation on Q induced by $\iota\Xi^0$. The complexation J is related to J^0 via a unique $(0, 1)$ -form \mathbf{B} of type Ad_H . Since the Lie algebra splitting

$$(3.14) \quad \mathfrak{h} = \mathfrak{l} \oplus \sqrt{-1}\mathfrak{l}$$

is Ad_L -invariant, the anti-linear map $\mathbf{B}_x : T_x M \rightarrow (Q \times_H \mathfrak{h})_x$ has a unique decomposition

$$(3.15) \quad \mathbf{B}_x v = \sqrt{-1}(\iota_x \mathbf{A}_x)v - (\iota_x \mathbf{A}_x)j_x v = \sqrt{-1}(\iota \mathbf{A})_x v - (\iota \mathbf{A})_x j_x v$$

$\forall x \in M$, where $\mathbf{A}_x : T_x M \rightarrow (P \times_L \mathfrak{l})_x$ is a 1-form of type Ad_L . Let Ξ be the connexion on P related to Ξ^0 via \mathbf{A} . By Proposition 3.5, the complexation J is induced by the hermitian connexion

$$(\iota\Xi^0)_q + (\iota \mathbf{A})_q^Q \circ d_q \pi = (\iota\Xi)_q.$$

For uniqueness, suppose that for two connexions Ξ^0 and Ξ on P , the extensions induce the same complexation. The connexion Ξ is related to Ξ^0 via a unique 1-form \mathbf{A} of type Ad_L . Then $\iota\Xi$ is related to $\iota\Xi^0$ via $\iota \mathbf{A}$, and both induce the same complexation. Proposition 3.5 implies that $\iota \mathbf{A}$ is a $(1, 0)$ -form of type Ad_H . Since \mathbf{A}_p^P is \mathfrak{l} -valued, it follows that $\mathbf{A}_p^P = 0$. Hence $\mathbf{A} = 0$ and $\Xi = \Xi^0$. As Ξ is invariant if and only if the complexation induced by $\iota\Xi$ is invariant, the assertion (ii) follows. \square

4. Homogeneous connexions and complexions

We now turn to the homogeneous case. By Theorem 2.2 we may assume that the homogeneous H -bundle Q is given by $Q = G \times_{K,f} H$ for a homomorphism $f : K \rightarrow H$, with $f(K) \subset L$ in the hermitian case. We fix an Ad_K -invariant splitting

$$(4.1) \quad \mathfrak{g} = \mathfrak{k} \oplus \mathfrak{m}.$$

Thus $[\mathfrak{k}, \mathfrak{m}] \subset \mathfrak{m}$ but not necessarily $[\mathfrak{m}, \mathfrak{m}] \subset \mathfrak{k}$. Let $P_{\mathfrak{k}} : \mathfrak{g} \rightarrow \mathfrak{k}$ and $P_{\mathfrak{m}} : \mathfrak{g} \rightarrow \mathfrak{m}$ denote the projection maps given by (4.1).

Definition 4.1.

- (i) The *tautological* connexion Θ^0 on $Q = G \times_{K,f} H$ is defined by the horizontal subspaces

$$\begin{aligned} T_{g|h}^0 Q &:= \{ \langle (d_e L_g^G) \tau | 0_h \rangle : \tau \in \mathfrak{m} \} \\ &= \{ \langle (d_e L_g^G) \gamma | (d_e R_h^H) \eta \rangle : \gamma \in \mathfrak{g}, \eta \in \mathfrak{h}, (d_e f) P_{\mathfrak{k}} \gamma + \eta = 0 \} \\ &= \{ \langle \dot{g} | \dot{h} \rangle : \dot{g} \in T_g G, \dot{h} \in T_h H, (d_e f) P_{\mathfrak{k}} (d_e L_g^G)^{-1} \dot{g} + (d_e R_h^H)^{-1} \dot{h} = 0 \}. \end{aligned}$$

- (ii) For a hermitian homogeneous H -bundle (Q, P) , the tautological connexion is the extension of the “tautological” connexion Ξ^0 on P with horizontal subspaces

$$T_{g|\ell}^0 P := \{ \langle (d_e L_g^G) \tau | 0_\ell \rangle \mid \tau \in \mathfrak{m} \} \quad \forall \langle g|\ell \rangle \in P = G \times_{K,f} L.$$

The equivalence of the various realizations of the horizontal subspaces follows from the definition. Under right translations, we have

$$(d_{g|h} R_b^Q) \langle \dot{g} | \dot{h} \rangle = \partial_t^0 \langle g_t | h_t b \rangle = \langle \dot{g} | (d_h R_b^H) \dot{h} \rangle.$$

It follows that $(T_{g|h} R_b^Q) T_{g|h}^0 Q = T_{\langle g|h \rangle}^0 Q$, showing that Θ^0 is indeed a connexion on Q . Since

$$(d_{g|h} L_a^Q) \langle \dot{g} | \dot{h} \rangle = \langle (d_g L_a^G) \dot{g} | \dot{h} \rangle,$$

it follows that

$$(d_{g|h} L_a^Q) \langle (d_e L_g^G) \tau | 0_h \rangle = \langle (d_g L_a^G) (d_e L_g^G) \tau | 0_h \rangle = (d_e L_{ag}^G \tau | 0_h).$$

Thus Θ^0 is an invariant connexion. In the hermitian case, the connexion Ξ^0 on P is also invariant.

Proposition 4.2. *The connexion 1-form of the tautological connection is given by*

$$\Theta_{g|h}^0 \langle (d_e L_g^G) \gamma | (d_e L_h^H) \eta \rangle = \text{Ad}_{h^{-1}}^H (d_e f) P_{\mathfrak{k}} \gamma + \eta \quad \forall (\gamma, \eta) \in \mathfrak{g} \times \mathfrak{h}.$$

In particular, $\Theta_{e|e}^0 \langle \gamma | \eta \rangle = (d_e f) P_{\mathfrak{k}} \gamma + \eta$.

Proof. Applying (2.4) to $\kappa = P_{\mathfrak{k}} \gamma$ we conclude that

$$\begin{aligned} \langle (d_e L_g^G) \gamma | (d_e L_h^H) \eta \rangle &= \langle (d_e L_g^G) P_{\mathfrak{m}} \gamma | (d_e R_h^H) (d_e f) P_{\mathfrak{k}} \gamma + (d_e L_h^H) \eta \rangle. \\ &= \langle (d_e L_g^G) P_{\mathfrak{m}} \gamma | 0_h \rangle + \langle 0_g | (d_e R_h^H) (d_e f) P_{\mathfrak{k}} \gamma + (d_e L_h^H) \eta \rangle. \end{aligned}$$

This gives the direct sum decomposition into horizontal and vertical components, and

$$\beta := \Theta_{g|h}^0 \langle (d_e L_g^G) \gamma | (d_e L_h^H) \eta \rangle \in \mathfrak{h}$$

is determined by the condition

$$\begin{aligned} \langle 0_g | (d_e R_h^H) (d_e f) P_{\mathfrak{k}} \gamma + (d_e L_h^H) \eta \rangle &= (d_e L_{g|h}) \beta = \partial_t^0 \langle g|h \rangle b_t \\ &= \partial_t^0 \langle g|h b_t \rangle = \langle 0 | (d_e L_h^H) \beta \rangle. \end{aligned}$$

Therefore, we have

$$\beta = (d_e L_h^H)^{-1} (d_e R_h^H) (d_e f) P_{\mathfrak{k}} \gamma + \eta = \text{Ad}_{h^{-1}}^H (d_e f) P_{\mathfrak{k}} \gamma + \eta. \quad \square$$

In the homogeneous case, invariance can be checked at the base point $o \in M$. An i -linear map

$$\mathbf{c} : \bigwedge^i T_o M \rightarrow \mathfrak{h}$$

will be called f -covariant if

$$\text{Ad}_{f(k)}^H \circ \mathbf{c} = \mathbf{c} \circ (d_o L_k^M) \quad \forall k \in K.$$

Every such map induces an invariant i -form \mathbf{C} of type Ad_H via

$$\mathbf{C}_{go}(d_oL_g^M)v = [\langle g|h \rangle, \text{Ad}_h^{-1} \mathbf{c}v]$$

for all $(g, h) \in Q = G \times_{K,f} H$ and $v \in \bigwedge^i T_oM$. Equivalently, the homogeneous lift satisfies

$$(4.2) \quad \mathbf{C}_{g|h}^Q(d_oL_g^M) = \text{Ad}_{h^{-1}}^H \circ \mathbf{c}.$$

Conversely, for any invariant i -form \mathbf{C} of type Ad_H , the map

$$\mathbf{c} := \mathbf{C}_{e|e}^Q : \bigwedge^i T_oM \rightarrow \mathfrak{h}$$

is f -covariant. If the homomorphism f is replaced by the conjugate $f' = I_\ell \circ f$, then $\mathbf{c}' : \bigwedge^i T_oM \rightarrow \mathfrak{h}$ is f' -covariant if and only if $\mathbf{c} := \text{Ad}_{\ell^{-1}}^H \circ \mathbf{c}'$ is f -covariant. In the homogeneous case Proposition 3.1 yields the following:

Proposition 4.3. *Let $\mathbf{c} : T_oM \rightarrow \mathfrak{h}$ be a linear map which is f -covariant, i.e.,*

$$\text{Ad}_{f(k)}^H \circ \mathbf{c} = \mathbf{c} \circ (d_oL_k^M)$$

for all $k \in K$. Then the 1-form \mathbf{C} of type Ad_H defined by

$$\mathbf{C}_{g|h}^Q(d_oL_g^M) = \text{Ad}_{h^{-1}}^H \circ \mathbf{c}, \quad \forall (g, h) \in G \times H$$

is invariant. Hence the connexion Θ on $Q = G \times_{K,f} H$, related to the tautological connexion Θ^0 via \mathbf{C} , is invariant, and also every invariant connexion is of this form.

In this case we say that Θ is generated by \mathbf{c} , the choice of the (invariant) tautological connexion Θ^0 being understood. In the hermitian homogeneous case Proposition 3.2 yields the following:

Proposition 4.4. *Let $\mathbf{a} : T_oM \rightarrow \mathfrak{l}$ be a linear map which is f -covariant, i.e., satisfies $\text{Ad}_{f(k)}^L \circ \mathbf{a} = \mathbf{a} \circ (d_oL_k^M)$ for all $k \in K$. Then the 1-form \mathbf{A} of type Ad_L defined by*

$$\mathbf{A}_{g|\ell}^Q(d_oL_g^M) = \text{Ad}_{\ell^{-1}}^L \circ \mathbf{a} \quad \forall (g, \ell) \in G \times L,$$

is invariant. Hence the connexion Ξ on $P = G \times_{K,f} \mathfrak{l}$, related to the tautological connexion Ξ^0 via \mathbf{A} , is invariant, and every invariant connexion is of this form.

In this case we say that Ξ is generated by \mathbf{a} , the choice of the (invariant) tautological connexion Ξ^0 being understood. Since $\iota \Xi^0 = \Theta^0$, the invariant connexion $\iota \Xi$ is generated by the f -covariant map $\iota \circ \mathbf{a} : T_oM \rightarrow \mathfrak{h}$. Combining Propositions 4.3 and 4.4 with Theorem 2.2 we obtain the following:

Theorem 4.5.

- (i) *The homogeneous H -bundles Q endowed with an invariant connexion are in 1 – 1 correspondence with pairs (f, \mathbf{c}) consisting of a homomorphism $f : K \rightarrow H$ and an f -covariant linear map $\mathbf{c} : T_oM \rightarrow \mathfrak{h}$, modulo the equivalence*

$$(f, \mathbf{c}) \sim (I_\ell^H \circ f, \text{Ad}_\ell^H \mathbf{c}),$$

where $\ell \in H$ is arbitrary.

- (ii) *The hermitian homogeneous H -bundles (P, Q) endowed with an invariant hermitian connexion are in 1 – 1 correspondence with pairs (f, \mathbf{a}) consisting of a homomorphism $f : K \rightarrow L$ and an f -covariant linear map $\mathbf{a} : T_oM \rightarrow \mathfrak{l}$, modulo the equivalence*

$$(f, \mathbf{a}) \sim (I_\ell^L \circ f, \text{Ad}_\ell^H \mathbf{a}),$$

where $\ell \in L$ is arbitrary.

In the complex homogeneous case, Proposition 3.4 yields the following:

Proposition 4.6. *Let $\mathbf{b} : T_oM \rightarrow \mathfrak{h}$ be an anti-linear map which is f -covariant, i.e.,*

$$\text{Ad}_{f(k)}^H \circ \mathbf{b} = \mathbf{b} \circ (d_o L_k^M), \quad \forall k \in K.$$

Then the $(0, 1)$ -form \mathbf{B} of type Ad_H defined by

$$\mathbf{B}_{g|h}^Q(d_o L_g^M) = \text{Ad}_{h^{-1}}^H \circ \mathbf{b} \quad \forall (g, h) \in G \times H$$

is invariant. Hence the complexation J on $Q = G \times_{K, f} H$, related to the tautological complexation J^0 via \mathbf{B} , is invariant, and every invariant complexation J is of this form.

In this case we say that J is generated by \mathbf{b} , the choice of the (invariant) tautological complexation J^0 being understood. Using the K -invariant inverse map $T_oM \ni v \mapsto \tilde{v} \in \mathfrak{m}$ of $d_e R_o^M$, the tautological complexation J^0 has the value

$$J_{e|e}^0(\langle \tilde{v} | 0_e \rangle) = \langle (j_o v)^\sim | 0_e \rangle$$

at the base point. For a general invariant complexation J generated by $\mathbf{b} : T_oM \rightarrow \mathfrak{h}$ we have

$$L_{e|e}^Q h = \langle e|e \rangle h = \langle e|h \rangle$$

and hence $d_{e|e} L_{e|e}^Q \eta = \langle 0_e | \eta \rangle$. Therefore

$$\begin{aligned} J_{e|e}(\langle \tilde{v} | 0_e \rangle) - J_{e|e}^0(\langle \tilde{v} | 0_e \rangle) &= (d_{e|e} L_{e|e}^Q) \mathbf{b}(d_{e|e} \pi)(\langle \tilde{v} | 0_e \rangle) \\ &= (d_{e|e} L_{e|e}^Q) \mathbf{b} v = \langle 0_e | \mathbf{b} v \rangle \end{aligned}$$

and hence

$$J_{e|e}(\langle \tilde{v} | 0_e \rangle) = \langle (j_o v)^\sim | 0_e \rangle + \langle 0_e | \mathbf{b} v \rangle = \langle (j_o v)^\sim | \mathbf{b} v \rangle.$$

All other values can be computed via $G \times H$ -invariance. In the complex homogeneous case Proposition 3.5 yields the following:

Proposition 4.7.

(i) *If an invariant connexion Θ is generated by \mathbf{c} , then the induced invariant complexation J^Θ is generated by*

$$(4.3) \quad \mathbf{b} = \sqrt{-1}\mathbf{c} - \mathbf{c}j_o.$$

(ii) *Θ induces the tautological complexation if and only if $\mathbf{c} : T_oM \rightarrow \mathfrak{h}$ is \mathbb{C} -linear.*

(iii) *If the invariant complexation J is generated by \mathbf{b} , then J is induced by the invariant connexion Θ generated by $-\frac{\sqrt{-1}}{2}\mathbf{b}$.*

In the hermitian homogeneous case. there is a 1-1 correspondence

$$(4.4) \quad \mathbf{b}v = \sqrt{-1}\mathbf{a}(v) - \mathbf{a}j_ov \quad \forall v \in T_oM$$

between f -covariant anti-linear maps $\mathbf{b} : T_oM \rightarrow \mathfrak{h}$ and f -covariant maps $\mathbf{A} : T_oM \rightarrow \mathfrak{l}$, since j_o commutes with $d_oL_k^M$ for all $k \in K$. This correspondence realizes the correspondence between invariant complexations and invariant hermitian connexions addressed in Proposition 3.6. Combining Propositions 4.6 and 4.7 with Theorem 2.2, we obtain the following:

Theorem 4.8. *The homogeneous H -bundles Q (respectively, the hermitian homogeneous H -bundles (P, Q)) endowed with an invariant complexation J are in 1-1 correspondence with pairs (f, \mathbf{b}) consisting of a homomorphism $f : K \rightarrow H$ (respectively, $f : K \rightarrow L$) and an f -covariant anti-linear map*

$$\mathbf{b} : T_oM \rightarrow \mathfrak{h},$$

modulo the equivalence $(f, \mathbf{b}) \sim (I_\ell^H \circ f, \text{Ad}_\ell^H \mathbf{b})$, where $\ell \in H$ (respectively, $\ell \in L$) is arbitrary. In the hermitian case, we may equivalently consider pairs (f, \mathbf{a}) where $\mathbf{a} : T_oM \rightarrow \mathfrak{l}$ is any f -covariant \mathbb{R} -linear map.

5. Curvature and Integrability

For any connexion Θ on a principal H -bundle Q there exists a 2-form \mathbf{K} of type Ad_H giving the *curvature*

$$d\Theta(X, Y) + \frac{1}{2}[\Theta X, \Theta Y] = \mathbf{K}_q^Q((d_q\pi)X, (d_q\pi)Y) \quad \forall X, Y \in T_qQ.$$

If Θ is invariant, the associated curvature form \mathbf{K} is also invariant. For $\beta \in \mathfrak{h}$ define the right action vector field ρ_β^Q on Q by

$$(\rho_\beta^Q)_q = \partial_t^0 q \exp(t\beta) = (d_e L_q^Q)\beta \quad \forall q \in Q.$$

Let J be a complexation on Q . For vector fields X, Y the $(0, 2)$ -part of the bilinear bracket $[X, Y]$ defined by

$$N(X, Y) = [X, Y] + J[JX, Y] + J[X, JY] - [JX, JY]$$

is called the *Nijenhuis tensor*. It is well-known that J is integrable if and only if N vanishes.

Lemma 5.1. *Let J be a complexion on Q . Then $N(\rho_\beta^Q, Y) = 0$ for $\beta \in \mathfrak{h}$ and any vector field Y on Q .*

Proof. Let

$$b_t = \exp(t\beta).$$

Applying [KN, Proposition I.1.9] to $\rho_\beta^Q = \partial_t^0 \exp(R_{b_t}^Q)$ we have

$$[\rho_\beta^Q, Y]_q = \lim_{t \rightarrow 0} (Y_q - (d_{qb_t} R_{b_t}^Q) Y_{qb_t}) = \lim_{t \rightarrow 0} (Y_q - (d_{qb_t} R_{b_{-t}}^Q) Y_{qb_t})$$

for every vector field X on Q . Since J commutes with right translations R_b^Q on Q it follows that

$$\begin{aligned} J_q[\rho_\beta^Q, Y]_q &= \lim_{t \rightarrow 0} (J_q Y_q - J_q (d_{qb_t} R_{b_{-t}}^Q) Y_{qb_t}) \\ &= \lim_{t \rightarrow 0} (J_q Y_q - (d_{qb_t} R_{b_{-t}}^Q) J_{qb_t} Y_{qb_t}) = [\rho_\beta^Q, JY]_q. \end{aligned}$$

Thus $J[\rho_\beta^Q, Y] = [\rho_\beta^Q, JY]$ as vector fields. Since

$$JJY = -Y \quad \text{and} \quad J\rho_\beta^Q = \rho_{\sqrt{-1}\beta}^Q,$$

we obtain

$$\begin{aligned} N(\rho_\beta^Q, Y) &= [\rho_\beta^Q, Y] + J[\rho_\beta^Q, JY] + J[\rho_{\sqrt{-1}\beta}^Q, Y] - [\rho_{\sqrt{-1}\beta}^Q, JY] \\ &= [\rho_\beta^Q, Y] + [\rho_\beta^Q, JJY] + [\rho_{\sqrt{-1}\beta}^Q, JY] - [\rho_{\sqrt{-1}\beta}^Q, JY] = 0. \quad \square \end{aligned}$$

Theorem 5.2. *Let (M, j) be integrable. Then the complexion J^Θ has the Nijenhuis tensor*

$$N_q(X, Y) = -2(d_e L_q^Q) \bar{\mathbf{K}}_q^Q((d_q \pi)X, (d_q \pi)Y) \quad \forall X, Y \in T_q Q,$$

where the $(0, 2)$ -part $\bar{\mathbf{K}}$ of \mathbf{K} is defined by

$$\bar{\mathbf{K}}_x(u, v) = \mathbf{K}_x(u, v) + \sqrt{-1}\mathbf{K}_x(j_x u, v) + \sqrt{-1}\mathbf{K}_x(u, j_x v) - \mathbf{K}_x(j_x u, j_x v)$$

$\forall u, v \in T_x M$.

Proof. Every $X \in T_q Q$ is given by $X = (\xi^\Theta + \rho_\beta^Q)_q$ for some vector field ξ on M and $\beta \in \mathfrak{h}$. Thus it suffices to consider vector fields of the form $\xi^\Theta + \rho_\beta^Q$. By Lemma 5.1 it is enough to consider horizontal lifts of vector fields ξ, η on M . Denoting the Nijenhuis tensor of (M, j) by $n(\xi, \eta)$, the integrability assumption on (M, j) implies that

$$(5.1) \quad n(\xi, \eta) = [\xi, \eta] + j[j\xi, \eta] + j[\xi, j\eta] - [j\xi, j\eta] = 0.$$

From [KN, Corollary I.5.3] we have

$$\Theta_q[\xi^\Theta, \eta^\Theta]_q = -2\Omega_q(\xi_q^\Theta, \eta_q^\Theta) = -2\mathbf{K}_q^Q(\xi_x, \eta_x).$$

By [KN, Proposition I.1.3] the horizontal part of $[\xi^\ominus, \eta^\ominus]_q$ coincides with $[\xi, \eta]_q^\ominus$. It follows that

$$[\xi^\ominus, \eta^\ominus]_q = [\xi, \eta]_q^\ominus + (d_e L_q^Q) \Theta_q [\xi^\ominus, \eta^\ominus]_q = [\xi, \eta]_q^\ominus - 2(d_e L_q^Q) \mathbf{K}_q^Q(\xi_x, \eta_x).$$

Using (3.12) and $J^\ominus \xi^\ominus = (j\xi)^\ominus$ this implies that

$$\begin{aligned} J_q^\ominus [\xi^\ominus, \eta^\ominus]_q &= J_q^\ominus [\xi, \eta]_q^\ominus - 2J_q^\ominus (d_e L_q^Q) \mathbf{K}_q^Q(\xi_x, \eta_x) \\ &= (j[\xi, \eta])_q^\ominus - 2(d_e L_q^Q) \sqrt{-1} \mathbf{K}_q^Q(\xi_x, \eta_x) \end{aligned}$$

and $[J^\ominus \xi^\ominus, \eta^\ominus]_q = [j\xi, \eta]_q^\ominus - 2(d_e L_q^Q) \mathbf{K}_q^Q(j\xi_x, \eta_x)$. Therefore, we have

$$\begin{aligned} N_q(\xi_q^\ominus, \eta_q^\ominus) &= [\xi^\ominus, \eta^\ominus]_q + J_q^\ominus [J^\ominus \xi^\ominus, \eta^\ominus]_q + J_q^\ominus [\xi^\ominus, J^\ominus \eta^\ominus]_q - [J^\ominus \xi^\ominus, J^\ominus \eta^\ominus]_q \\ &= (n(\xi, \eta))_q^\ominus - 2(d_e L_q^Q) \left(\mathbf{K}_q^Q(\xi_x, \eta_x) + \sqrt{-1} \mathbf{K}_q^Q(j\xi_x, \eta_x) \right. \\ &\quad \left. + \sqrt{-1} \mathbf{K}_q^Q(\xi_x, j\eta_x) - \mathbf{K}_q^Q(j\xi_x, j\eta_x) \right) \\ &= (n(\xi, \eta))_q^\ominus - 2(d_e L_q^Q) \overline{\mathbf{K}}_q^Q(\xi_x, \eta_x). \end{aligned}$$

In view of (5.1) the proof is now complete. \square

Corollary 5.3. *If (M, j) is integrable, then the complexion J^\ominus is integrable on Q if and only if the curvature form \mathbf{K} of Θ has vanishing $(0, 2)$ -part.*

In the homogeneous case the curvature form \mathbf{K} of an invariant connexion Θ is induced by a unique f -covariant map $\mathbf{k} : \bigwedge^2 T_o M \rightarrow \mathfrak{h}$ as in (4.2).

Theorem 5.4. *The curvature \mathbf{K}^0 of the tautological connection Θ^0 satisfies the equation*

$$2\mathbf{k}^0(u, v) = -(d_e f) P_{\mathfrak{k}}[\tilde{u}, \tilde{v}] \quad \forall u, v \in T_o M,$$

where $\tilde{u} \in \mathfrak{m}$ is uniquely determined by $(d_e R_o^M) \tilde{u} = u$.

Proof. Let $\alpha, \gamma \in \mathfrak{g}$. Consider the left action vector field

$$(\lambda_\alpha^Q)_g|_h = \partial_t^0 \langle a_t g | h \rangle = \langle (d_e R_g^G) \alpha | 0_h \rangle$$

on Q . The identity $R_g^G = L_g^G \circ I_{g^{-1}}^G$ implies $d_e R_g^G = (d_e L_g^G) \text{Ad}_{g^{-1}}^G$. Therefore Proposition 4.2 yields

$$(\Theta^0 \lambda_\alpha^Q)_{g|e} = \Theta_{g|e}^0 \langle (d_e R_g^G) \alpha | 0_h \rangle = (d_e f) P_{\mathfrak{k}} \text{Ad}_{g^{-1}}^G \alpha.$$

Putting $g_t = \exp(t\gamma) \in G$ we conclude that

$$(\Theta^0 \lambda_\alpha^Q)_{g_t|e} = (d_e f) P_{\mathfrak{k}} \text{Ad}_{g_t^{-1}}^G \alpha = (d_e f) P_{\mathfrak{k}} \exp(-t \text{ad}_\gamma) \alpha$$

and hence

$$\begin{aligned} d_{e|e}(\Theta^0 \lambda_\alpha^Q) \langle \gamma | 0_e \rangle &= \partial_t^0 (\Theta^0 \lambda_\alpha^Q)_{g_t|e} \\ &= (d_e f) P_{\mathfrak{k}} \partial_t^0 \exp(-t \text{ad}_\gamma) \alpha \\ &= (d_e f) P_{\mathfrak{k}}[\alpha, \gamma]. \end{aligned}$$

Since $[\lambda_\alpha^Q, \lambda_\gamma^Q] = \lambda_{[\gamma, \alpha]}^Q$ for left actions we have

$$\Theta_{e|e}^0[\lambda_\alpha^Q, \lambda_\gamma^Q]_{e|e} = \Theta_{e|e}^0(\lambda_{[\gamma, \alpha]}^Q)_{e|e} = (d_e f)P_{\mathfrak{k}}[\gamma, \alpha].$$

Therefore, we have

$$\begin{aligned} 2\mathbf{K}_{e|e}^0(\langle \alpha|0_e \rangle, \langle \gamma, 0_e \rangle) &= 2(d\Theta^0)_{e|e}(\langle \alpha|0_e \rangle, \langle \gamma, 0_e \rangle) + [\Theta_{e|e}^0 \langle \alpha|0_e \rangle, \Theta_{e|e}^0 \langle \gamma|0_e \rangle] \\ &= d_{e|e}(\Theta^0 \lambda_\gamma^Q) \langle \alpha|0_e \rangle - d_{e|e}(\Theta^0 \lambda_\alpha^Q) \langle \gamma|0_e \rangle \\ &\quad - \Theta_{e|e}^0[\lambda_\alpha^Q, \lambda_\gamma^Q]_{e|e} + [\Theta_{e|e}^0 \langle \alpha|0_e \rangle, \Theta_{e|e}^0 \langle \gamma|0_e \rangle] \\ &= (d_e f)(P_{\mathfrak{k}}[\gamma, \alpha]) - (d_e f)(P_{\mathfrak{k}}[\alpha, \gamma]) - (d_e f)(P_{\mathfrak{k}}[\gamma, \alpha]) \\ &\quad + [(d_e f)(P_{\mathfrak{k}}\alpha), (d_e f)(P_{\mathfrak{k}}\gamma)] \\ &= (d_e f)([P_{\mathfrak{k}}\alpha, P_{\mathfrak{k}}\gamma] - P_{\mathfrak{k}}[\alpha, \gamma]). \end{aligned}$$

Since $P_{\mathfrak{k}}\tilde{u} = 0$ we obtain

$$2\mathbf{k}^0(u, v) = 2\Omega_{e|e}^0(\langle \tilde{u}|0 \rangle, \langle \tilde{v}|0 \rangle) = -(d_e f)(P_{\mathfrak{k}}[\tilde{u}, \tilde{v}]). \quad \square$$

Every $\alpha \in \mathfrak{g}$ induces a left action vector field λ_α^M on M by putting

$$(\lambda_\alpha^M)_x = (d_e R_x^M)\alpha = \partial_t^0(a_t x),$$

where $a_t = \exp(t\alpha)$. For left actions we have the reverse commutator identity $[\lambda_\alpha^M, \lambda_\gamma^M] = \lambda_{[\gamma, \alpha]}^M$ for $\alpha, \gamma \in \mathfrak{g}$.

Lemma 5.5. *For $\tau \in \mathfrak{m}$ the vector field λ_τ^M has the horizontal lift*

$$(5.2) \quad X_{g|h}^\tau := \langle (d_e L_g^G) P_{\mathfrak{m}} \text{Ad}_{g^{-1}}^G \tau | 0_h \rangle.$$

Proof. Let $(g, h) \in G \times H$ and $k \in K$. Then the equality

$$R_{k^{-1}}^G \circ L_g^G = L_{gk^{-1}}^G \circ I_k^G$$

implies that

$$(d_g R_{k^{-1}}^G)(d_e L_g^G) = (d_e L_{gk^{-1}}^G) \text{Ad}_k^G.$$

Since $[\text{Ad}_k^G, P_{\mathfrak{m}}] = 0$, we have

$$\text{Ad}_k^G P_{\mathfrak{m}} \text{Ad}_{g^{-1}}^G = P_{\mathfrak{m}} \text{Ad}_k^G \text{Ad}_{g^{-1}}^G = P_{\mathfrak{m}} \text{Ad}_{kg^{-1}}^G$$

and therefore

$$\begin{aligned} &\langle (d_g R_{k^{-1}}^G)(d_e L_g^G) P_{\mathfrak{m}} \text{Ad}_{g^{-1}}^G \tau | (d_h L_{f(k)}^H) 0_h \rangle \\ &= \langle (d_e L_{gk^{-1}}^G) \text{Ad}_k^G P_{\mathfrak{m}} \text{Ad}_{g^{-1}}^G \tau | 0_{f(k)h} \rangle \\ &= \langle (d_e L_{gk^{-1}}^G) P_{\mathfrak{m}} \text{Ad}_{kg^{-1}}^G \tau | 0_{f(k)h} \rangle. \end{aligned}$$

This shows that (5.2) depends only on the class of (g, h) and therefore defines a vector field on Q which is Θ^0 -horizontal by construction. Moreover the equality $R_o^M \circ L_g^G = L_g^M \circ R_o^M$ implies that

$$(d_g R_o^M)(d_e L_g^G) = (d_o L_g^M)(d_e R_o^M),$$

and the equality $L_g^M \circ R_o^M \circ I_{g^{-1}}^G = R_{go}^M$ implies that

$$(d_o L_g^M)(d_e R_o^M) \text{Ad}_{g^{-1}}^G = d_e R_{go}^M.$$

Since $(d_e R_o^M)P_m = d_e R_o^M$ it follows that

$$\begin{aligned} (X^\tau \pi)\langle g|h \rangle &= (d_{g|h} \pi)X_{g|h}^\tau = (d_g R_o^M)(d_e L_g^G) P_m \text{Ad}_{g^{-1}}^G \tau \\ &= (d_o L_g^M)(d_e R_o^M) P_m \text{Ad}_{g^{-1}}^G \tau \\ &= (d_o L_g^M)(d_e R_o^M) \text{Ad}_{g^{-1}}^G \tau \\ &= (d_e R_{go}^M)\tau = (\lambda_\tau^M)_{go}. \quad \square \end{aligned}$$

6. The symmetric case

Now we consider the special case where $M = G/K$ is a *symmetric* space. These spaces have a well-known algebraic description using the so-called Lie triple systems [He]. As discovered by M. Koecher [Koe], in the hermitian symmetric case there is a more “elementary” approach using instead the so-called *hermitian Jordan triple systems* [Lo]. These are (complex) vector spaces Z which carry a Jordan triple product

$$(u, v, w) \mapsto \{uv^*w\} \quad \forall u, v, w \in Z$$

which is symmetric bilinear in (u, w) and conjugate-linear in v . The Jacobi identity is replaced by the *Jordan triple identity*

$$[u \square v^*, z \square w^*] = \{uv^*z\} \square w^* - z \square \{wu^*v\}^* \quad \forall u, v, z, w \in Z.$$

Here $u \square v^* \in \text{End}(Z)$ is defined by

$$(u \square v^*)z = \{uv^*z\} \quad \forall z \in Z.$$

The basic example is the matrix space $Z = \mathbb{C}^{r \times s}$ with Jordan triple product

$$\{uv^*w\} = \frac{1}{2}(uv^*w + wv^*u).$$

The Jordan theoretic approach applies to all complex hermitian symmetric spaces, including the two exceptional types, and also to all classical real symmetric spaces. More generally, all real forms of complex hermitian symmetric spaces, for example symmetric convex cones [FK], and therefore also some exceptional real symmetric spaces, are included. (On the other hand, there exist nonclassical real symmetric spaces which cannot be treated this way.)

We first consider both the real and complex case. Given a (real or complex) Jordan triple Z , we put

$$Q_z w := \{zw^*z\} \quad \forall z, w \in Z$$

and define the *Bergman operator*

$$B_{z,w} = id_Z - 2z \square w^* + Q_z Q_w \quad \forall z, w \in Z,$$

acting linearly on Z . In case $B_{z,w} \in \text{GL}(Z)$ is invertible, we define the *quasi-inverse*

$$z^w := B_{z,w}^{-1}(z - Q_z w).$$

Define ϵ to be $\epsilon = -1$ for the noncompact case, $\epsilon = 1$ for the compact case, and $\epsilon = 0$ for the flat case. We define symmetric spaces M^ϵ associated with the Jordan triple Z as follows: $M^0 = Z$ is the flat model, M^- is the connected component

$$M^- \subset \{z \in Z \mid \det B(z, z) \neq 0\}$$

containing the origin $o = 0 \in Z$ (a bounded symmetric domain, more precisely a norm unit ball of Z), and

$$M^+ = (Z \times Z) / \sim$$

is a compact manifold consisting of all equivalence classes $[z, a] = [z^{b-a}, b]$, whenever $B(z, a - b)$ is invertible [Lo]. (In view of the "addition formula" $(z^u)^v = z^{u+v}$ for quasi-inverses [Lo], we may informally regard M^+ as the set of all quasi-inverses z^a , even when $B(z, a)$ is not invertible.) Thus we have natural inclusions

$$M^- \subset Z = M^0 \subset M^+,$$

under the embedding $Z \subset M^+$ given by $z \mapsto z^0 = [z, 0]$. The points at infinity are precisely the classes $[z, a]$ where $\det B(z, a) = 0$. The compact dual M^+ is also called the *conformal compactification* of Z . At the origin the tangent space $T_o M = Z$ is independent of the choice of ϵ . Let $K \subset \text{GL}(Z)$ be the identity component of the Jordan triple automorphism group of Z , i.e., all linear isomorphisms of Z preserving the Jordan triple product. The group K acts by linear transformations on every type M^ϵ . For fixed $w \in Z \cap M^\epsilon$, the (nonlinear) *transvection*, defined by

$$(6.1) \quad \mathfrak{t}_w^\epsilon(z) := w + B_{w, -\epsilon w}^{1/2} z^{\epsilon w},$$

is a birational automorphism of M^ϵ . For $\epsilon = 1, 0, -1$ let G^ϵ denote the connected real Lie group generated by K together with the transvections (6.1). In the flat case $\epsilon = 0$ we obtain the so-called Cartan motion group $G^0 := K \times Z$ which is a semi-direct product of K and the translations $\mathfrak{t}_w^0(z) = z + w$ for $w \in Z$. If $\epsilon \neq 0$, then G^ϵ is a reductive Lie group of compact type ($\epsilon = 1$) or noncompact type ($\epsilon = -1$), respectively. We treat all three cases simultaneously, using the notation M^ϵ and G^ϵ to denote the curvature type. In all three cases we have

$$K = \{k \in G^\epsilon \mid ko = o\}.$$

In the Jordan theoretic setting, the Lie algebra \mathfrak{g}^ϵ of G^ϵ can be described using polynomial vector fields

$$\xi = \xi(z) \frac{\partial}{\partial z}$$

(or degree ≤ 2) on the underlying vector space Z . More precisely, there is a Cartan decomposition

$$(6.2) \quad \mathfrak{g}^\epsilon = \mathfrak{k} \oplus \mathfrak{p}^\epsilon,$$

where the Lie algebra $\mathfrak{k} \subset \mathfrak{gl}(Z)$ of K , consisting of all Jordan triple derivations of Z , is identified with a space of linear vector fields, whereas the Lie triple system \mathfrak{p}^ϵ consists of all nonlinear vector fields

$$v^\epsilon := v + \epsilon Q_z v = (v + \epsilon \{zv^*z\}) \frac{\partial}{\partial z} \quad \forall v \in Z = T_o M^\epsilon.$$

The projection $\mathfrak{g}^\epsilon \rightarrow \mathfrak{k}$ is realized as the derivative

$$P_{\mathfrak{k}} \gamma = \gamma'(o)$$

at the origin $o = 0 \in Z$. One can show that $\exp v^\epsilon = \mathfrak{k}_w^\epsilon$, where $w := \tan_\epsilon v \in M^\epsilon$ is given by a ‘‘tangent’’ power series defined via the Jordan triple calculus [Lo].

Proposition 6.1. *In the symmetric case the tautological connection Θ^0 induced by the Cartan decomposition (6.2) satisfies*

$$(6.3) \quad \mathbf{k}^0(u, v) = -\epsilon(df)(u \square v^* - v \square u^*) \quad \forall u, v \in Z.$$

Proof. Denoting by u^∂ the (holomorphic) partial derivative in direction $u \in Z$, we have the commutator identity

$$\begin{aligned} [u^\epsilon, v^\epsilon] &= [u + \epsilon Q_z u, v + \epsilon Q_z v] = \epsilon (u^\partial Q_z v - v^\partial Q_z u) \\ &= 2\epsilon (u \square v^* - v \square u^*) \in \mathfrak{k}. \end{aligned}$$

Thus we obtain a linear vector field satisfying $[u^\epsilon, v^\epsilon]'(o) = [u^\epsilon, v^\epsilon]$, and Theorem 5.4 implies

$$2\mathbf{k}^0(u, v) = -df([u^\epsilon, v^\epsilon]'(o)) = -2\epsilon df(u \square v^* - v \square u^*). \quad \square$$

Theorem 6.2. *In the symmetric case, let Θ be an invariant connexion related to Θ^0 by an f -covariant linear map $\mathbf{c} : T_o M \rightarrow \mathfrak{h}$. Then the respective curvatures satisfy*

$$\mathbf{k}(u, v) - \mathbf{k}^0(u, v) = \frac{1}{2}[\mathbf{c}u, \mathbf{c}v] \quad \forall u, v \in T_o M.$$

Proof. For $v \in Z$ let X^v be the Θ^0 -horizontal lift of λ_v^M , as in Lemma 5.5. Consider the tensorial 1-form $\Phi := \Theta - \Theta^0$ on Q . Then

$$\Omega(X^u, X^v) - \Omega^0(X^u, X^v) = d\Phi(X^u, X^v) + \frac{1}{2}[\Phi X^u, \Phi X^v] \quad \forall u, v \in T_o M.$$

Since $L_g^M \circ R_o^M \circ I_{g^{-1}}^G = R_{go}^M$ implies $(d_o L_g^M)(d_e R_o^M) \text{Ad}_{g^{-1}}^G = d_e R_{go}^M$, it follows that

$$\begin{aligned} (\Phi X^v)\langle g|e \rangle &= \mathbf{C}_{g|e}^Q(d_{g|e} \pi) X_{g|e}^v = \mathbf{C}_{g|e}^Q(d_e R_{go}^M) v^\epsilon \\ &= \mathbf{C}_{g|e}^Q(d_o L_g^M)(d_e R_o^M) \text{Ad}_{g^{-1}}^G v^\epsilon = \mathbf{c}(d_e R_o^M)(\text{Ad}_{g^{-1}}^G) v^\epsilon, \end{aligned}$$

since $\mathbf{C}_{g|e}^Q(d_o L_g^M) = \mathbf{c}$ as a special case of (4.2). With $g_t := \exp(t u^\epsilon)$ we obtain

$$(\Phi X^v)\langle g_t|e\rangle = \mathbf{c}(d_e R_o^M)(\text{Ad}_{g_{-t}}^G)v^\epsilon = \mathbf{c}(d_e R_o^M)\exp(-t \text{ad}_{u^\epsilon})v^\epsilon.$$

Since the linear map $\mathbf{c}(d_e R_o^M)$ commutes with taking the ∂_t -derivative, it follows that

$$\begin{aligned} X^u(\Phi X^v)\langle e|e\rangle &= d_{e|e}(\Phi X^v)\langle u^\epsilon|0_e\rangle = \partial_t^0(\Phi X^v)\langle g_t|e\rangle \\ &= \mathbf{c}(d_e R_o^M)\partial_t^0 \exp(-t \text{ad}_{u^\epsilon})v^\epsilon = -\mathbf{c}(d_e R_o^M) \text{ad}_{u^\epsilon}v^\epsilon \\ &= -\mathbf{c}(d_e R_o^M)[u^\epsilon, v^\epsilon] = 0, \end{aligned}$$

since $[u^\epsilon, v^\epsilon] \in [\mathfrak{p}^\epsilon, \mathfrak{p}^\epsilon] \subset \mathfrak{k}$ implies $(d_e R_o^M)[u^\epsilon, v^\epsilon] = 0$. On the other hand,

$$\begin{aligned} X^u(X^v \pi)\langle e|e\rangle &= d_{e|e}(X^v \pi)X_{e|e}^u = d_{e|e}(X^v \pi)\langle u^\epsilon|0_e\rangle \\ &= \partial_t^0(X^v \pi)\langle g_t|e\rangle = \partial_t^0(\lambda_{v^\epsilon}^M)_{g_t o} \\ &= \partial_t^0(v + \epsilon Q_{g_t(0)}v) = 0, \end{aligned}$$

since applying the product rule to the quadratic term yields a term $g_0(0) = 0$. Thus

$$(d_{e|e}\pi)[X^u, X^v]_{e|e} = ([X^u, X^v]\pi)\langle e|e\rangle = (X^u(X^v \pi) - X^v(X^u \pi))\langle e|e\rangle = 0$$

and hence

$$\begin{aligned} d\Phi(X^u, X^v)\langle e|e\rangle &= X^u(\Phi X^v)\langle e|e\rangle - X^v(\Phi X^u)\langle e|e\rangle - \Phi_{e|e}[X^u, X^v]_{e|e} \\ &= -\mathbf{c}(d_{e|e}\pi)[X^u, X^v]_{e|e} = 0. \end{aligned}$$

It follows that

$$\begin{aligned} \mathbf{k}(u, v) - \mathbf{k}^0(u, v) &= \Omega_{e|e}(X^u, X^v) - \Omega_{e|e}^0(X^u, X^v) \\ &= d\Phi(X^u, X^v)\langle e|e\rangle + \frac{1}{2}[(\Phi X^u)\langle e|e\rangle, (\Phi X^v)\langle e|e\rangle] \\ &= \frac{1}{2}[\mathbf{c}u, \mathbf{c}v]. \quad \square \end{aligned}$$

Now we consider the case where $M = G/K$ is an (irreducible) *hermitian symmetric* space. In this case Z is a complex hermitian Jordan triple, G^- is the identity component of the real Lie group of all biholomorphic automorphisms of M^- , and G^+ is the identity component of the biholomorphic isometry group of M^+ , i.e., the biholomorphic automorphisms of M^+ that preserve the Kähler metric. Both Lie groups are semi-simple. The identity (6.3) can be polarized and hence *defines* the Jordan triple product in terms of the curvature tensor at the base point o .

Proposition 6.3. *In the hermitian symmetric case, the tautological complexion J^0 is integrable.*

Proof. By Corollary 5.3 we have to show that the curvature 2-form \mathbf{K}^0 of the tautological connexion Θ^0 has vanishing $(0, 2)$ -part. By invariance under $G \times H$ it suffices to consider the base point $\langle e|e \rangle$. The \mathbb{R} -bilinear map

$$(u, v) \mapsto D(u, v) = u \square v^* - v \square u^* \in \mathfrak{k},$$

for $u, v \in Z$, satisfies

$$D(u, \sqrt{-1}v) + D(\sqrt{-1}u, v) = 0 \quad \text{and} \quad D(\sqrt{-1}u, \sqrt{-1}v) = D(u, v).$$

It follows that

$$D(u, v) + \sqrt{-1}D(\sqrt{-1}u, v) + \sqrt{-1}D(u, \sqrt{-1}v) - D(\sqrt{-1}u, \sqrt{-1}v) = 0.$$

Therefore, Theorem 6.2 shows that

$$\mathbf{k}^0(u, v) = \epsilon(d_e f)(u \square v^* - v \square u^*) = \epsilon(d_e f)D(u, v)$$

has vanishing $(0, 2)$ -part. \square

If $\mathbf{b} : Z \rightarrow \mathfrak{h}$ is an f -covariant anti-linear map, then $[\mathbf{b} \wedge \mathbf{b}](u, v) := [\mathbf{b}u, \mathbf{b}v]$ defines an f -covariant anti-bilinear map

$$[\mathbf{b} \wedge \mathbf{b}] : Z \wedge Z \rightarrow \mathfrak{h}.$$

Theorem 6.4. *In the hermitian symmetric case, the invariant complexation J generated by \mathbf{b} is integrable if and only if $[\mathbf{b} \wedge \mathbf{b}] = 0$.*

Proof. By Proposition 4.7 the complexation J is induced by the invariant connexion Θ generated by $-\frac{\sqrt{-1}}{2}\mathbf{b}$. Applying Theorem 6.2 we have

$$\mathbf{k}(u, v) - \mathbf{k}^0(u, v) = \frac{1}{2} \left[\frac{\sqrt{-1}}{2}\mathbf{b}u, \frac{\sqrt{-1}}{2}\mathbf{b}v \right] = -\frac{1}{8}[\mathbf{b}u, \mathbf{b}v]$$

for all $u, v \in Z$. By Theorem 5.2 the Nijenhuis tensor for J satisfies

$$N_{e|e}(X, Y) = -2\bar{\mathbf{k}}((d_e \pi)X, (d_e \pi)Y) \quad \forall X, Y \in T_{e|e}Q,$$

where

$$\bar{\mathbf{k}}(u, v) = \mathbf{k}(u, v) + \sqrt{-1}\mathbf{k}(\sqrt{-1}u, v) + \sqrt{-1}\mathbf{k}(u, \sqrt{-1}v) - \mathbf{k}(\sqrt{-1}u, \sqrt{-1}v)$$

is the $(0, 2)$ -part of the curvature \mathbf{k} of Θ . Since the tautological complexation J^0 on Q is integrable by Proposition 6.3, it follows that

$$\begin{aligned} \bar{\mathbf{k}}^0(u, v) &= \mathbf{k}^0(u, v) + \sqrt{-1}\mathbf{k}^0(\sqrt{-1}u, v) + \sqrt{-1}\mathbf{k}^0(u, \sqrt{-1}v) - \mathbf{k}^0(\sqrt{-1}u, \sqrt{-1}v) \\ &= 0. \end{aligned}$$

On the other hand, $[\mathbf{b} \wedge \mathbf{b}]$ is of type $(0, 2)$ since \mathbf{b} is anti-linear by Proposition 4.6. Therefore

$$\bar{\mathbf{k}}(u, v) = -\frac{1}{8}[\mathbf{b}u, \mathbf{b}v]$$

and hence

$$N_{e|e}(X, Y) = \frac{1}{4}[\mathbf{b}(d_e \pi)X, \mathbf{b}(d_e \pi)Y] = \frac{1}{4}[\mathbf{b} \wedge \mathbf{b}](X \otimes Y) \quad \forall X, Y \in T_{e|e}Q.$$

This implies that N vanishes at the base point $\langle e|e \rangle$ if and only if $[\mathbf{b} \wedge \mathbf{b}] = 0$. By $G \times H$ -invariance, this implies that N vanishes at all points $\langle g|h \rangle \in Q$. \square

Theorem 6.5. *In the hermitian symmetric case, the hermitian homogeneous H -bundles Q endowed with an integrable invariant complexion J are in 1 – 1 correspondence with pairs (f, \mathbf{b}) consisting of a homomorphism $f : K \rightarrow L \subset H$ and an f -covariant anti-linear map $\mathbf{b} : T_oM \rightarrow \mathfrak{h}$ satisfying*

$$(6.4) \quad [\mathbf{b} \wedge \mathbf{b}] = 0,$$

modulo the equivalence $(f, \mathbf{b}) \sim (I_\ell^H \circ f, \text{Ad}_\ell^H \mathbf{b})$, where $\ell \in L$ is arbitrary. Via the relation (4.4), we may equivalently consider pairs (f, \mathbf{a}) where

$$\mathbf{a} : T_oM \rightarrow \mathfrak{l}$$

is any f -covariant \mathbb{R} -linear map such that

$$(6.5) \quad [\mathbf{a}u, \mathbf{a}v] = [\mathbf{a}j_o u, \mathbf{a}j_o v] \quad \forall u, v \in T_oM.$$

Proof. If \mathbf{a} and \mathbf{b} are related by (4.4), then the conditions (6.4) and (6.5) are equivalent. \square

Since the classification of Theorem 6.5 uses only data coming from the group K , which is the same for the hermitian symmetric spaces $M^\epsilon = G^\epsilon/K$, for $\epsilon = 0, 1, -1$, we obtain:

Corollary 6.6. *There is a canonical 1 – 1 correspondence between the sets classifying the hermitian homogeneous H -bundles (P, Q) , endowed with an integrable invariant complexion J , over the hermitian symmetric spaces of compact type, noncompact type and flat type, respectively.*

7. Concluding remarks

In order to put our results in perspective, and motivate the general treatment in the first four Sections, we outline possible generalizations of our approach. In the geometric quantization program, one considers more general nonsymmetric G -homogeneous spaces $N = G/C$, endowed with an invariant complex structure j . An interesting class of examples is obtained as follows: Let $M = G/K$ be an irreducible hermitian symmetric space of compact, noncompact or flat type (depending on the choice of G). Then the tangent space $Z = T_o(M)$ at the origin $o \in M$ is a hermitian Jordan triple. The so-called principal inner ideals $U \subset Z$ are precisely the Peirce 2-spaces $U = Z_c^2$ for a given tripotent $c \in Z$. Let F_j denote the Grassmann type manifold of all such Peirce 2-spaces of fixed rank $j \leq r$. More generally, for any increasing sequence $1 \leq j_1 < j_2 < \dots < j_\ell \leq r$ there is a flag type manifold F_{j_1, \dots, j_ℓ} consisting of all flags $U_1 \subset U_2 \subset \dots \subset U_\ell$ of Peirce 2-spaces U_i of rank j_i . Using the G -action, one can define such flag manifolds for any tangent space $T_x(M)$, $x \in M$, and obtains a fibre bundle $N_{j_1, \dots, j_\ell} \rightarrow M$ with typical fibre F_{j_1, \dots, j_ℓ} . This fibre bundle is homogeneous $N_{j_1, \dots, j_\ell} = G/K_{j_1, \dots, j_\ell}$, where

K_{j_1, \dots, j_ℓ} is a closed subgroup of K which is the same for the compact, non-compact and flat case. Our principal result, concerning the classification of holomorphic principal fibre bundles and the duality between the compact and noncompact case, may be generalized to this setting [BiU].

Another important problem, of interest in quantization theory, is the explicit construction of Dolbeault cohomology operators depending on the given complex structure, in the symmetric case or the more general setting outlined above. A first step towards this goal is a more explicit realization of the classifying space of holomorphic principal fibre bundles, described in Theorem (6.5) in the symmetric case. According to (6.4), the basic case $H = GL_n(\mathbf{C})$ involves pairwise commuting $n \times n$ -matrices B_1, \dots, B_d , where $d = \dim_{\mathbf{C}} G/K$, modulo joint conjugation; already a quite complicated object in algebraic geometry. Finally, the whole construction depends on the underlying invariant complex structure j of $N = G/C$ which may not be unique if C is a proper subgroup of K . Analogous to the Narasimhan-Seshadri Theorem for Riemann surfaces, the moduli space of invariant complex structures may carry a canonical projectively flat connexion on the bundle of holomorphic sections.

References

- [At] ATIYAH, M. F. Complex analytic connections in fibre bundles. *Trans. Amer. Math. Soc.* **85** (1957), 181–207. [MR0086359](#) (19,172c), [Zbl 0078.16002](#), doi: [10.2307/1992969](#).
- [ADW] AXELROD, SCOTT; DELLA PIETRA, STEVE; WITTEN, EDWARD. Geometric quantization of Chern–Simons gauge theory. *J. Differential Geom.* **33** (1991), no. 3, 787–902. [MR1100212](#) (92i:58064), [Zbl 0697.53061](#).
- [Bi] BISWAS, INDRANIL. Classification of homogeneous holomorphic hermitian principal bundles over G/K . *Forum Math.* **27** (2015), no. 2, 937–960. [MR3334089](#), [Zbl 1317.32044](#), doi: [10.1515/forum-2012-0131](#).
- [BiM] BISWAS, INDRANIL; MISRA, GADADHAR. $\widetilde{SL}(2, \mathbb{R})$ -homogeneous vector bundles. *Internat J. Math.* **19** (2008), no. 1, 1–19. [MR2380469](#) (2009f:32041), [Zbl 1158.53043](#), doi: [10.1142/S0129167X08004534](#).
- [BiU] BISWAS, INDRANIL; UPMEIER, HARALD. Invariant principal fibre bundles over flag domains. In preparation.
- [Bo] BOTT, RAOUL. Homogeneous vector bundles. *Ann. of Math.* (2) **66** (1957), 203–248. [MR0089473](#) (19,681d), [Zbl 0094.35701](#), doi: [10.2307/1969996](#).
- [FK] FARAUT, JACQUES; KORÁNYI, ADAM. Analysis on symmetric cones. Oxford Mathematical Monographs. Oxford Science Publications. *The Clarendon Press, Oxford University Press, New York*, 1994. xii+382 pp. ISBN: 0-19-853477-9. [MR1446489](#) (98g:17031), [Zbl 0841.43002](#).
- [He] HELGASON, SIGURDUR. Differential geometry, Lie groups, and symmetric spaces. Corrected reprint of the 1978 original. Graduate Studies in Mathematics, 34. *American Mathematical Society, Providence, RI*, 2001. xxvi+641 pp. ISBN: 0-8218-2848-7. [MR1834454](#) (2002b:53081), [Zbl 0993.53002](#).
- [KN] KOBAYASHI, SHOSHICHI; NOMIZU, KATSUMI. Foundations of differential geometry. I. *Interscience Publishers, a division of John Wiley & Sons, New York-London*, 1963. xi+329 pp. [MR0152974](#), [Zbl 0119.37502](#).

- [Koe] KOECHER, MAX. An elementary approach to bounded symmetric domains. *Rice University, Houston, Tex.*, 1969. iii+143 pp. [MR0261032](#), [Zbl 0217.10901](#).
- [Kos] KOSZUL, J.-L.. Lectures on fibre bundles and differential geometry. Notes by S. Ramanan. Tata Institute of Fundamental Research Lectures on Mathematics, 20. *Tata Institute of Fundamental Research, Bombay*, 1965. ii+130+iii pp. [MR0268801](#), [Zbl 0244.53026](#), <http://www.math.tifr.res.in/~publ/ln/tifr20.pdf>.
- [Lo] LOOS, OTTMAR. Bounded symmetric domains and Jordan pairs. University of California, Irvine, 1977.
- [Mu] MUMFORD, D. On the equations defining abelian varieties. I. *Invent. Math.* **1** (1966), 287–354. [MR0204427](#), [Zbl 0219.14024](#), doi: [10.1007/BF01389737](https://doi.org/10.1007/BF01389737).
- [OR] OTTAVIANI, GIORGIO; RUBEI, ELENA. Quivers and the cohomology of homogeneous vector bundles. *Duke Math. J.* **132** (2006), no. 3, 459–508. [MR2219264](#) (2008b:14075), [Zbl 1100.14012](#), [arXiv:math/0403307](https://arxiv.org/abs/math/0403307), doi: [10.1215/S0012-7094-06-13233-7](https://doi.org/10.1215/S0012-7094-06-13233-7).
- [Ra] RAMANAN, S. Holomorphic vector bundles on homogeneous spaces. *Topology* **5** (1966), 159–177. [MR0190947](#), [Zbl 0138.18602](#), doi: [10.1016/0040-9383\(66\)90017-6](https://doi.org/10.1016/0040-9383(66)90017-6).
- [We] WELTERS, GERALD E. Polarized abelian varieties and the heat equations. *Compositio Math.* **49** (1983), no. 2, 173–194. [MR0704390](#) (85f:14045), [Zbl 0576.14042](#).
- [Wi] WITTEN, EDWARD. Quantum field theory and the Jones polynomial. *Comm. Math. Phys.* **121** (1989), no. 3, 351–399. [MR0990772](#) (90h:57009), [Zbl 0667.57005](#), doi: [10.1007/BF01217730](https://doi.org/10.1007/BF01217730).

(Indranil Biswas) SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, 1 HOMI BHABHA ROAD, MUMBAI 400005, INDIA
indranil@math.tifr.res.in

(Harald Upmeyer) FACHBEREICH MATHEMATIK UND INFORMATIK, PHILIPPS-UNIVERSITÄT MARBURG, LAHNBERGE, HANS-MEERWEIN-STRASSE, D-35032 MARBURG, GERMANY
upmeyer@mathematik.uni-marburg.de

This paper is available via <http://nyjm.albany.edu/j/2016/22-2.html>.