

In the following proposition we generalize the wellknown semigroup-theoretical notion of Rees congruence to universal algebras.

PROPOSITION 1.2. *Let M be a Rees subalgebra of $\mathfrak{A} = (A, \Omega)$ and ρ_M the following binary relation on A : $x \equiv y(\rho_M)$: $x, y \in M$ or $x = y$. Then ρ_M is a congruence on \mathfrak{A} .*

If ρ is a congruence on \mathfrak{A} with at most one non-singleton class M and M is a subalgebra of \mathfrak{A} , then M is a Rees subalgebra and $\rho = \rho_M$.

The proof of this proposition is obvious.

REMARK 1.3. If the algebra $\mathfrak{A} = (S, \cdot)$ is a semigroup, then every semigroup ideal I is a Rees subalgebra. But there are Rees subalgebras that are not ideals of S . Let S be a semigroup without null and $S^* = S \cup \{0\}$ with $a \cdot 0 = 0 \cdot a = 0$ for every $a \in S^*$, then S is a Rees subalgebra of S^* but not a semigroup ideal.

In Szasz (1968) Rees congruences on lattices (L, \wedge, \vee) are studied; the author proved, Theorem 1., p. 261, that an ideal I is a Rees subalgebra, if the following condition holds:

$$a \in L - I, \quad b \in I: \quad a \geq b.$$

2. General theory of Rees congruence

In this chapter we study for an arbitrary Rees subalgebra M the Rees congruence ρ_M . First we prove an isomorphism theorem that formally looks like the second isomorphism theorem in group theory.

PROPOSITION 2.1. *If $M \subseteq A$ is a Rees subalgebra and $T \subseteq A$ is a subalgebra of \mathfrak{A} such that $M \cap T \neq \emptyset$, then $M \cup T$ is a subalgebra of \mathfrak{A} , too.*

PROOF: Obviously, nullary and unary operations preserve an arbitrary union of subalgebras. Let ω be an n -ary operation, $n \geq 2$, and $\mathbf{a} = (a_1, \dots, a_n) \in (M \cup T)^n$, $b \in M \cup T \neq \emptyset$. If $\mathbf{a} \in T^n$, then $\omega(\mathbf{a}) \in T \subseteq M \cup T$. If $\mathbf{a} \notin T^n$, then let \mathbf{a}' denote the vector obtained from \mathbf{a} by substituting all its components from M with b . Then $\mathbf{a}' \equiv \mathbf{a}(\rho_M)$ componentwise, but $\mathbf{a}' \in T^n$, hence $\omega(\mathbf{a}') \in T$. Because of $\mathbf{a} \equiv \mathbf{a}'(\rho_M)$ we have $\omega(\mathbf{a}) = \omega(\mathbf{a}')$ or $\omega(\mathbf{a}), \omega(\mathbf{a}') \in M$. In the first case we obtain $\omega(\mathbf{a}) = \omega(\mathbf{a}') \in T$ and in the second case $\omega(\mathbf{a}) \in M$; and so in both cases $\omega(\mathbf{a}) \in M \cup T$ and the proposition is proved.

THEOREM 2.2. *Let M be a Rees subalgebra of $\mathfrak{A} = (A, \Omega)$ and $T \subseteq A$ a subalgebra of \mathfrak{A} such that $M \cap T \neq \emptyset$. If A/M denotes the Rees factor algebra, of \mathfrak{A} with respect to the congruence ρ_M , then the following isomorphism can be stated:*

$$T/\cup T \cong M \cup T/M.$$

PROOF: Under the condition of this theorem it is clear that $\rho_{M \cap T}$ is a congruence on T , the restriction of ρ_M on T , and similary ρ_M is a congruence on $M \cup T$. Now obviously the two algebras $T/M \cap T$ and $M \cup T/M$ are isomorphic.

DEFINITION 2.3. Let $\mathfrak{A} = (A, \Omega)$ be a universal algebra. Then a Rees series of \mathfrak{A} is a finite descending series

$$A = A_0 \supset A_1 \supset \dots \supset A_r = \emptyset$$

such that A_0, \dots, A_{r-1} are Rees subalgebras of A , $A_r = \emptyset$ and A_{i+1} is properly contained in the algebra A_i . We define the factors of the Rees series to be the Rees factor algebras A_i/A_{i+1} for $i = 0, \dots, r-1$. The length of the Rees series is the number of factors. A refinement of a Rees series is any Rees series containing every term A_i of the original series. Two Rees series are termed isomorphic if we can put their factors in 1-1 correspondence so that corresponding factors are isomorphic. Further we define a composition series to be a Rees series without proper refinements.

PROPOSITION 2.4. Let $\mathfrak{A} = (A, \Omega)$ be a universal algebra and S, T Rees subalgebras of \mathfrak{A} such that $S \cap T \neq \emptyset$. Then $S \cup T$ is a Rees subalgebra, too.

PROOF: Because of Proposition 2.1. $S \cup T$ is a subalgebra of \mathfrak{A} . Let ω be an n -ary operation of \mathfrak{A} ($n \geq 2$) and $x_1, \dots, x_{n-1} \in A$ and $(a, b) \in (S \cup T)^2$ and $d \in S \cap T \neq \emptyset$. If $a, b \in S$ we obtain

$$\begin{aligned} \omega(a, x_1, \dots, x_{n-1}), \omega(b, x_1, \dots, x_{n-1}) &\in S \text{ or} \\ \omega(a, x_1, \dots, x_{n-1}) &= \omega(b, x_1, \dots, x_{n-1}); \text{ hence} \\ \omega(a, x_1, \dots, x_{n-1}), \omega(b, x_1, \dots, x_{n-1}) &\in S \cup T \text{ or} \\ \omega(a, x_1, \dots, x_{n-1}) &= \omega(b, x_1, \dots, x_{n-1}). \end{aligned}$$

If $a, b \in T$ this results similary. Now let $a \in S, b \in T$; we obtain $a \equiv d(\rho_S)$ and $b \equiv d(\rho_T)$. ρ_S and ρ_T are congruences, $\rho_{S \cup T}$ is an equivalence relation such that $\rho_S, \rho_T \leq \rho_{S \cup T}$; therefore

$$\begin{aligned} \omega(a, x_1, \dots, x_{n-1}) &\equiv \omega(d, x_1, \dots, x_{n-1}) \quad (\rho_S) \\ \omega(b, x_1, \dots, x_{n-1}) &\equiv \omega(d, x_1, \dots, x_{n-1}) \quad (\rho_T), \end{aligned}$$

and so

$$\omega(a, x_1, \dots, x_{n-1}) \equiv \omega(b, x_1, \dots, x_{n-1}) \quad (\rho_{S \cup T}).$$

Hence

$$\begin{aligned} \omega(a, x_1, \dots, x_{n-1}), \omega(b, x_1, \dots, x_{n-1}) &\in S \cup T \text{ or} \\ \omega(a, x_1, \dots, x_{n-1}) &= \omega(b, x_1, \dots, x_{n-1}). \end{aligned}$$

If $a \in T, b \in S$ this results similary. So condition (*) of Definition 1.1. can be verified, therefore $S \cup T$ is a Rees subalgebra of \mathfrak{A} .

REMARK 2.5. We consider two groupoids G_1 and G_2 defined by the following multiplication tables:

G_1	0	a	b
0	0	0	0
a	0	a	0
b	0	0	b

G_2	0	a	e	f
0	0	0	0	0
a	0	a	0	e
e	0	0	e	e
f	0	e	e	f

The groupoid G_1 shows that we cannot omit the condition $S \cap T \neq \emptyset$ for the Rees subalgebras in Proposition 2.4., because $S = \{a\}$ and $T = \{b\}$ are Rees subalgebras of G_1 with $S \cap T \neq \emptyset$, but $S \cup T = \{a, b\}$ is not a Rees subalgebra of G_1 .

We write $S \underset{R}{<} T$, if S is a Rees subalgebra of T . Then the binary relation $\underset{R}{<}$ is not transitive in general. We take the groupoid G_2 and put $S = \{e, f\}$ and $T = \{0, e, f\}$; then $S \underset{R}{<} T$, $T \underset{R}{<} G_2$, but S is not a Rees subalgebra of G_2 .

LEMMA 2.6. Let $\mathfrak{A} = (A, \Omega)$ be a universal algebra and R, S two subalgebras and R^*, S^* be Rees subalgebras of A such that $R^* \cap S^* \neq \emptyset$. Write

$$\begin{aligned} T &= R^* \cup (R \cap S), & T^* &= R^* \cup (R \cap S^*) \\ U &= S^* \cup (R \cap S), & U^* &= S^* \cup (R^* \cap S). \end{aligned}$$

Then T^*, U^* are Rees subalgebras of T, U , respectively, and

$$T/T^* \cong U/U^*$$

PROOF: Because of the condition $R^* \cap S^* \neq \emptyset$ and Proposition 2.1. $T = R^* \cup (R \cap S)$ is a subalgebra $T^* = R^* \cup (R \cap S^*)$ is a Rees subalgebra of T ; since $R \cap S^* \underset{R}{<} A$, $R^* \underset{R}{<} A$, we obtain $R^* \cup (R \cap S^*) \underset{R}{<} A$ by Proposition 2.4.; therefore $T^* \underset{R}{<} T$, because $T \subseteq A$. Similary we obtain that U^* is a Rees subalgebra of U .

We obtain

$$T^* \cup (R \cap S) = (R^* \cup (R \cap S^*)) \cup (R \cap S) = R^* \cup (R \cap S) = T.$$

Hence by Theorem 2.2.,

$$T/T^* \cong R \cap S /_{T^* \cap (R \cap S)}.$$

But

$$(R \cap S) \cap T^* = (R^* \cap S) \cup (R \cap S^*),$$

hence

$$T/T^* \cong R \cap S /_{(R^* \cap S) \cup (R \cap S^*)}.$$

U, U^* are obtained from T, T^* by interchanging R, R^* with S, S^* . Hence as above we may prove that

$$U/U^* \cong R \cap S / (R^* \cap S) \cup (R \cap S^*);$$

and so

$$T/T^* \cong U/U^*.$$

THEOREM 2.7. *Let $\mathfrak{A} = (A, \Omega)$ be a universal algebra and*

$$A = S_0 \supset S_1 \supset \dots \supset S_r = \emptyset$$

$$A = T_0 \supset T_1 \supset \dots \supset T_s = \emptyset$$

two Rees series with $S_i \cap T_j \neq \emptyset$ for $i = 0, \dots, r-1$ and $j = 0, \dots, s-1$. Then the two series have isomorphic refinements.

PROOF: The required refinements are, respectively,

$$A = S_{00} \supset S_{01} \supset \dots \supset S_{0s} = S_{10} \supset \dots \supset S_{rs}$$

$$A = T_{00} \supset T_{10} \supset \dots \supset T_{r0} = T_{01} \supset \dots \supset T_{rs},$$

where

$$S_{ik} = S_{i+1} \cup (S_i \cap T_k), \quad T_{ik} = T_{k+1} \cup (T_k \cap S).$$

Define $R^* = S_{i+1}$, $R = S_i$, $S^* = T_{k+1}$, $S = T_k$; then R^* is a Rees subalgebra of R and S^* is a Rees subalgebra of S and $R^* \cap S^* \neq \emptyset$ ($i < r$; $k < s$). So we can apply Lemma 2.5. and obtain:

$$S_{ik}/S_{i+1} \cong T_{ik}/T_{i+1k},$$

and so the above refinements are isomorphic.

COROLLARY 2.8. *Let $\mathfrak{A} = (A, \Omega)$ be a universal algebra, such that $S \cap T \neq \emptyset$ for all subalgebras S, T of \mathfrak{A} . Then all composition series of \mathfrak{A} are isomorphic.*

REMARK 2.9. The condition $S \cap T \neq \emptyset$, for all subalgebras S, T , holds if \mathfrak{A} contains nullary operations.

In the following let $R(\mathfrak{A})$ denote the set of all Rees subalgebras of \mathfrak{A} and the empty set \emptyset . Then the inclusion \subseteq defines a lattice ordering on $R(\mathfrak{A})$ with the operations \wedge, \vee defined by

$$M \wedge N = \inf(M, N) = M \cap N$$

$$M \vee N = \sup(M, N) \quad \text{for } M, N \in R(\mathfrak{A}).$$

For every subset $T \subseteq A$ we introduce the Rees subalgebra $\langle T \rangle$ generated by T ; this is the algebra $\langle T \rangle = \bigcap_{T \subseteq M \in R(\mathfrak{A})} M$.

A Rees subalgebra M is called finetely generated in $R(\mathfrak{A})$, if a finite set $T = \{a_1, \dots, a_n\}$ exists such that $M = \langle T \rangle$.

Now every n -ary operation $\omega \in \Omega$ defines an n -ary operation ω^* on $R(\mathfrak{A})$ by

$$\omega^*(M_1, \dots, M_n) = \langle \{\omega(x_1, \dots, x_n) : x_1 \in M_1, \dots, x_n \in M_n\} \rangle$$

for $M_1, \dots, M_n \in R(\mathfrak{A})$.

Ω^* denotes the set of all such operations ω^* .

PROPOSITION 2.10. *If $\mathfrak{A} = (A, \Omega)$ is a universal algebra, then $(R(\mathfrak{A}), \Omega^*, \subseteq)$ is a lattice-ordered algebra: so*

$$\omega^*(M_1, \dots, M_n) \subseteq \omega^*(N_1, \dots, N_n)$$

for $M_1 \subseteq N_1, \dots, M_n \subseteq N_n$ and all operations $\omega^* \in \Omega^*$.

G_3	a	b	c
a	a	c	b
b	c	b	a
c	b	a	c

If $S \cap T \neq \emptyset$ for all Rees subalgebras S, T , the lattice $R(\mathfrak{A})$ is distributive; generally this does not hold (see the groupid G_3).

THEOREM 2.11. *Let $\mathfrak{A} = (A, \Omega)$ denote a universal algebra and S a proper Rees subalgebra of \mathfrak{A} . If A is finitely generated in $R(\mathfrak{A})$, then a maximal proper Rees subalgebra M of \mathfrak{A} exists such that $M \supseteq S$.*

PROOF: Let \mathfrak{M} denote the set of all proper Rees subalgebras $T \supseteq S$ and let \mathfrak{N} be a totally ordered subset of \mathfrak{M} . We prove $V = \cup_{T \in \mathfrak{N}} T \in \mathfrak{M}$. Let ω be an n -ary operation and $x_1, \dots, x_n \in V$, then a Rees subalgebra $T_0 \supseteq S$ exists such that $x_1, \dots, x_n \in T_0$, so $\omega(x_1, \dots, x_n) \in T_0 \subseteq V$ and V is a subalgebra. If $a, b \in V$ and $x_1, \dots, x_{n-1} \in A$, then a Rees subalgebra $T_1 \supseteq S$ exists, such that $a, b \in T_1$ and the condition (*) of Definition 1.1. is valid; so V is a Rees subalgebra of \mathfrak{A} .

We show $V \neq A$. A is finitely generated by the set $\{a_1, \dots, a_k\}$. If we suppose $V = A$, we obtain $a_1, \dots, a_k \in V$; therefore a Rees subalgebra $T_2 \in \mathfrak{M}$ exists with $a_1, \dots, a_k \in T_2$ and so $T_2 = A$, a contradiction to $T_2 \in \mathfrak{M}$. So we obtain $V \neq A$ and $V \in \mathfrak{M}$ is proved. Now we apply Zorn's Lemma, so \mathfrak{M} contains a maximal element and the theorem is proved.

In the following we say that the maximal chain-condition holds in $R(\mathfrak{A})$, if every increasing chain $S_0 \subset S_1 \subset S_2 \subset \dots$ of Rees subalgebras of \mathfrak{A} contains only a finite number of terms S_i . Similarly we say that the minimal chain-condition holds in $R(\mathfrak{A})$, if every descending chain $S_0 \supset S_1 \supset \dots$ of Rees subalgebras of \mathfrak{A} contains only a finite number of terms S_i .

THEOREM 1.12. *Let $\mathfrak{A} = (A, \Omega)$ be a universal algebra. Then all Rees subalgebras S of \mathfrak{A} are finitely generated in $R(\mathfrak{A})$, if and only if the maximal chain-condition holds in $R(\mathfrak{A})$.*

PROOF: First we assume that the maximal chain-condition holds. Let S be an arbitrary Rees subalgebra and $a_1 \in S$. If $S = \langle a_1 \rangle$, then S is finitely generated; if $a_2 \in S$ exists with $a_2 \notin \langle a_1 \rangle$, then $\langle a_1 \rangle \subset \langle a_1, a_2 \rangle$. So we construct an increasing chain of Rees subalgebras

$$\langle a_1 \rangle \subset \langle a_1, a_2 \rangle \subset \langle a_1, a_2, a_3 \rangle \subset \dots,$$

which must be finite because of the maximal chain-condition; so S is finitely generated in $R(\mathfrak{A})$.

Now we assume that every Rees subalgebra S is finitely generated in $R(\mathfrak{A})$ and let $S_1 \subset S_2 \subset S_3 \subset \dots$, be an increasing chain of Rees subalgebras. The union $S = \cup S_i$ of all those Rees subalgebras S_i is finitely generated because of our assumption; so $S = \langle a_1, \dots, a_k \rangle$. Therefore a Rees subalgebra S_n exists such that $a_1, \dots, a_k \in S_n$; hence $S_n = S$. So we have proved that $S = S_n = S_{n+1} = \dots$ and the maximal chain-condition is valid in $R(\mathfrak{A})$.

THEOREM 2.13. *Let $\mathfrak{A} = (A, \Omega)$ denote a universal algebra such that for all subalgebras S, T the condition $S \cap T \neq \emptyset$ holds. Then a necessary and sufficient condition for \mathfrak{A} to have a composition series is that the maximal and minimal chain-condition hold in $R(\mathfrak{A})$.*

PROOF: First we assume that a composition series with length k exists. If the maximal or minimal chain-condition does not hold we easily obtain a Rees series with length $n > k$, a contradiction, because we can apply Theorem 2.7. (for all S, T with $S \cap T \neq \emptyset$). If both chain-conditions hold, we construct a composition series as follows. Define $S_0 = A$ and S_1 a maximal Rees subalgebra with $S_1 \subset S_0$; such S_1 exists because of the maximal chain-condition. Then S_2 is a maximal Rees subalgebra of S_1 and, generally, S_n a maximal subalgebra of S_{n-1} . So we obtain a Rees series

$$A = S_0 \supset S_1 \supset S_2 \dots \supset S_{n-1} \supset S_n \supset \dots,$$

which must be finite because of the minimal chain-condition. Obviously, this Rees series is a composition series.

EXAMPLE 3.14. Let $(\mathbf{N}, +, \cdot)$ be the natural numbers with addition and multiplication; then all Rees subalgebras are of the form

$$S_\alpha = \{x \in \mathbf{N} : x \geq \alpha\}.$$

Then $S \cap T \neq \emptyset$ for all subalgebras S, T of $(\mathbf{N}, +, \cdot)$; the maximal chain-condition holds in $R(\mathbf{N})$ and the minimal chain-condition does not hold. So \mathbf{N} is finitely generated in $R(\mathbf{N})$ ($\mathbf{N} = \langle 1 \rangle$) and \mathbf{N} does not contain a composition series.

EXAMPLE 2.15. Let $\gamma = (A, \cdot, e)$ be a monoid; we define an n -ary operation ω on A by $\omega(x_1, \dots, x_n) = x_1 \cdot \dots \cdot x_n$. Then $\mathfrak{A} = (A, \omega, e)$ is an algebra with $R(\mathfrak{A}) = R(\gamma)$.

DEFINITION 2.16. Let ω be an n -ary operation of the algebra $\mathfrak{A} = (A, \Omega)$. Then an element $a \in A$ is called idempotent with respect to ω , if and only if $\omega(a, \dots, a) = a$; an element $a \in A$ is called idempotent with respect to all n -ary operations ($n \geq 1$) of \mathfrak{A} .

An algebra is called idempotent, if and only if all its elements are idempotent.

PROPOSITION 2.17. Let $\mathfrak{A} = (A, \Omega)$ be an idempotent algebra and ρ a congruence on \mathfrak{A} with at most one non-singleton class M . If M contains all nullary operations of \mathfrak{A} , ρ is a Rees congruence.

PROOF: Using Proposition 1.2, we have to prove that the non-singleton class M is a subalgebra of \mathfrak{A} . Let $a_j \in M$ ($j = 1, \dots, n$) and ω be an n -ary operation of \mathfrak{A} . Then $a_1 \equiv a_2 \equiv \dots \equiv a_n (\rho)$, because ρ is a congruence. Since a_1 is idempotent, $\omega(a_1, \dots, a_1) = a_1 \in M$ results; so we obtain $\omega(a_1, \dots, a_n) \in M$ and the proposition is proved.

REMARK 2.18. Proposition 2.17. does not hold for an arbitrary algebra \mathfrak{A} . If we take the free semigroup (A, \cdot) with $A = \{a, a^2, a^3, \dots\}$, the identity relation ρ is a congruence with at most one non-singleton class, but it is not a Rees congruence.

PROPOSITION 2.19. Let $u = (A, \Omega)$ be a universal algebra such that $S \cap T \neq \emptyset$ for all Rees subalgebras S, T of \mathfrak{A} . If \mathfrak{A} does not contain an idempotent element, then $(R(\mathfrak{A}), \wedge, \vee)$ is isomorphic to a sublattice of the congruence lattice $\mathcal{S}(\mathfrak{A})$ of \mathfrak{A} . Generally this does not hold (see the semilattice S_0).

S_0	0	e	f	p	q
0	0	0	0	0	0
e	0	e	e	0	0
f	0	e	f	0	0
p	0	0	0	p	p
q	0	0	0	p	q

PROOF: Because of Proposition 2.4. the union $S \cup T$ of two Rees subalgebras S, T is a Rees subalgebra again. So we have $S \vee T = S \cup T$ for all Rees subalgebras S, T of \mathfrak{A} .

We define the mapping $\varphi: R(\mathfrak{A}) \rightarrow \mathcal{S}(\mathfrak{A})$ by $\varphi(S) = \rho_S$ if S is a Rees subalgebra and $\varphi(\emptyset) = id$. Since \mathfrak{A} does not contain an idempotent element, φ is injective; we obtain

$$\begin{aligned} \varphi(S \wedge T) &= \varphi(S \cap T) = \rho_{(S \cap T)} = \rho_S \wedge \rho_T = \varphi(S) \wedge \varphi(T) \\ \varphi(S \vee T) &= \varphi(S \cup T) = \rho_{S \cup T} = \rho_S \vee \rho_T = \varphi(S) \vee \varphi(T); \end{aligned}$$

so φ is a lattice homomorphism; hence $\varphi(R(\mathfrak{A}))$ is a sublattice of $\mathcal{S}(\mathfrak{A})$.

3. Applications to polynomial algebras

In this chapter we consider for an algebra $\mathfrak{A} = (A, \Omega)$ the algebra $\mathcal{P}(\mathfrak{A}, n)$ of all polynomial functions in n variables over \mathfrak{A} , the operations of which are the operations of \mathfrak{A} pointwise and the $n + 1$ -ary composition o_n . If p_i ($i = 1, \dots, n + 1$) are $n + 1$ polynomial functions of $\mathcal{P}(\mathfrak{A}, n)$ represented by the polynomials $p_i(x_1, \dots, x_n)$, then the composition $o_n(p_1, \dots, p_{n+1})$ of the polynomial functions p_i is just the polynomial function represented by the polynomial

$$p_i(p_2(x_1, \dots, x_n), \dots, p_{n+1}(x_1, \dots, x_n)),$$

where the variable x_i occurring in p_1 is substituted with $p_{i+1}(x_1, \dots, x_n)$.

PROPOSITION 3.1. *Let $\mathfrak{A} = (A, \Omega)$ be an idempotent algebra such that for every polynomial function $f: A \rightarrow A$ $f \circ f = f$. Then $\mathcal{P}(\mathfrak{A}, n)$ is an idempotent algebra.*

PROOF: We proceed by induction; for $n = 1$ it is clear because of the condition of the proposition. We have to show that $\mathcal{P}(\mathfrak{A}, n + 1)$ is idempotent, if $\mathcal{P}(\mathfrak{A}, n)$ is idempotent; the only operation we must consider is o_n . Let $p = p(x_1, \dots, x_{n+1})$ be such a polynomial function; if one variable does not occur in the word representation of this function the proposition is clear by the assumption of the induction. Let \mathbf{x} be the vector (x_1, \dots, x_{n+1}) . We use the condition $f \circ f = f$ for all f defined by $f(x_{n+2}) = p(\mathbf{x}, x_1, \dots, x_n)$ fixed; therefore

$$(i) \quad p(x_1, \dots, x_n, p(\mathbf{x})) = p(\mathbf{x}).$$

If we define $g(x_1, \dots, x_n) = p(\mathbf{x})$ for fixed x_{n+1} , we obtain

$$g(g(x_1, \dots, x_n), \dots, g(x_1, \dots, x_n)) = g(x_1, \dots, x_n);$$

and so

$$(ii) \quad p(p(\mathbf{x}), \dots, p(\mathbf{x}), x_{n+1}) = p(\mathbf{x}).$$

Put $y = (p(\mathbf{x}), \dots, p(\mathbf{x}), x_{n+1})$ and combining (i) and (ii) we obtain

$$p(p(\mathbf{x}), \dots, p(\mathbf{x})) = p(p(\mathbf{x}), \dots, p(\mathbf{x}), p(y)) = p(y) = p(\mathbf{x});$$

COROLLARY 3.2. *Let $\mathfrak{A} = (A, \Omega)$ be an idempotent algebra such that $f \circ f = f$ for all polynomial functions $f: A \rightarrow A$. If ρ is a congruence on $\mathcal{P}(\mathfrak{A}, n)$ with at most one non-singleton class, then ρ is a Rees congruence.*

This is obvious; because $\mathcal{P}(\mathfrak{A}, n)$ does not contain a nullary operation. One can apply Proposition 2.17.

REMARK 3.3. Considering the lattice (L, \wedge, \vee) we obtain the following result:

L is distributive if and only if $\mathcal{P}(L, n)$ is idempotent. For $n = 1$ this is a wellknown theorem of Skhweigert (1975); for $n \in N$ it results, applying Proposition 3.1. So in the case of a distributive lattice L we can apply Corollary 3.2. to the algebra $\mathcal{P}(L, n)$. Trivially all these facts remain true, if we consider the algebra $\mathcal{P}^*(L, n)$, which also consists of all polynomial functions in n variables over L , but the only operation is o_n .

If $\mathfrak{A} = (S, \cdot)$ is a semilattice, then Proposition 3.1. and Corollary 3.2. are applicable, too.

PROPOSITION 3.4. *Let $\mathfrak{A} = (A; \Omega)$ be a universal algebra and M a Rees subalgebra of \mathfrak{A} , such that for every $p \in \mathcal{P}(\mathfrak{A}, n)$ $p(x) \in M$, if at least one coordinate of $\mathbf{x} = (x_1, \dots, x_n)$ is an element of M . If $I_M = \{p; p(\mathbf{x}) \in M \text{ for every } \mathbf{x} \in A^n\}$ and $I_M \neq \emptyset$, then I_M is a Rees subalgebra of $\mathcal{P}(\mathfrak{A}, n)$.*

PROOF: Let ω be a k -ary pointwise operation of $\mathcal{P}(\mathfrak{A}, n)$ and $p_1, \dots, p_k \in I_M$. Then $\omega(p_1, \dots, p_k)(\mathbf{x}) = \omega(p_1(\mathbf{x}), \dots, p_k(\mathbf{x})) \in M$ for every $\mathbf{x} \in A^n$ because of the definition of I_M . Similarly for every $x \in A^n$ $o_n(p_1, \dots, p_n)(\mathbf{x}) \in M$ results. So I_M is a subalgebra of $\mathcal{P}(\mathfrak{A}, n)$, if $I_M \neq \emptyset$. Trivially, for every pointwise operation the condition of Definition 1.1. holds. Let $p_1, \dots, p_{n+1} \in \mathcal{P}(\mathfrak{A}, n)$, and $p_j \in I_M$, then

$$o_n(p_1, \dots, p_{n+1})(\mathbf{x}) = p_1(p_2(\mathbf{x}), \dots, p_j(\mathbf{x}), \dots, p_{n+1}(\mathbf{x})) \in M$$

for every $x \in A$; so the proposition is proved.

REMARK 3.5. Let (S, \cdot) be a semigroup without null and M an idea of S . Then $I_M \neq \emptyset$ is a Rees subalgebra of $\mathcal{P}(S, n)$ and defines a Rees congruence ρ_M on $\mathcal{P}(S, n)$. If N is an ideal different from M , then $I_M \neq I_N$ because $\mathcal{P}(S, n)$ contains all constant functions. If we define $\rho_\emptyset = id$, the mapping $M \rightarrow \rho_M$ is a lattice isomorphism from $\mathcal{J}(S) \cup \{\emptyset\}$ into the congruence $\mathcal{C}(\mathcal{P}(S, n))$ where $\mathcal{J}(S)$ is the ideal lattice of (S, \cdot) . So $\mathcal{J}(S) \cup \{\emptyset\}$ is isomorphic to a sublattice of $\mathcal{C}(\mathcal{P}(S, n))$: S must be convergence free if $\mathcal{C}(\mathcal{P}(S, n))$ is congruence free.

REMARK 3.6. Let (L, \wedge, \vee) be a lattice with more than two elements; then a non-trivial ideal M exists; which defines I_M as in Proposition 3.4. and $I_M \neq \emptyset$. I_M defines a non-trivial Rees congruence on $\mathcal{P}^*(L, n)$; so $\mathcal{P}^*(L, n)$ cannot be congruence free. This result remains true, if we consider the algebra of all those polynomial functions, in the representation of which all variables x_1, \dots, x_n occur.

REFERENCES

- [1] H. Lausch – W. Nölower, *Algebra of Polynomials*, North – Holland Publ. (1973).
- [2] D. Rees, *On Semigroups*, Proc. Cambridge Phil. Soc. **36** (1940), 387–400.
- [3] D. Schweigert, *Über idempotente Polynomfunktionen auf Vierbänden*, Elem. d. Math. **30** (1975), 30–32.
- [4] G. Szasz, *Rees factor lattices*, Publ. Math. Debrecen **15** (1968), 259–266.

Robert F. Tichy
 Patrubangasse 5,
 A-1100, Vienna,
 Austria