

Indefinite Affine Hyperspheres Admitting a Pointwise Symmetry. Part 2*

Christine SCHARLACH

Technische Universität Berlin, Fak. II, Inst. f. Mathematik, MA 8-3, 10623 Berlin, Germany

E-mail: schar@math.tu-berlin.de

URL: <http://www.math.tu-berlin.de/~schar/>

Received May 08, 2009, in final form October 06, 2009; Published online October 19, 2009

doi:10.3842/SIGMA.2009.097

Abstract. An affine hypersurface M is said to admit a pointwise symmetry, if there exists a subgroup G of $\text{Aut}(T_p M)$ for all $p \in M$, which preserves (pointwise) the affine metric h , the difference tensor K and the affine shape operator S . Here, we consider 3-dimensional indefinite affine hyperspheres, i.e. $S = \text{HId}$ (and thus S is trivially preserved). In Part 1 we found the possible symmetry groups G and gave for each G a canonical form of K . We started a classification by showing that hyperspheres admitting a pointwise $\mathbb{Z}_2 \times \mathbb{Z}_2$ resp. \mathbb{R} -symmetry are well-known, they have constant sectional curvature and Pick invariant $J < 0$ resp. $J = 0$. Here, we continue with affine hyperspheres admitting a pointwise \mathbb{Z}_3 - or $SO(2)$ -symmetry. They turn out to be warped products of affine spheres (\mathbb{Z}_3) or quadrics ($SO(2)$) with a curve.

Key words: affine hyperspheres; indefinite affine metric; pointwise symmetry; affine differential geometry; affine spheres; warped products

2000 Mathematics Subject Classification: 53A15; 53B30

1 Introduction

Let M^n be a connected, oriented manifold. Consider an immersed hypersurface with relative normalization, i.e., an immersion $\varphi: M^n \rightarrow \mathbb{R}^{n+1}$ together with a transverse vector field ξ such that $D\xi$ has its image in $\varphi_* T_x M$. Equi-affine geometry studies the properties of such immersions under equi-affine transformations, i.e. volume-preserving linear transformations ($SL(n+1, \mathbb{R})$) and translations.

In the theory of nondegenerate equi-affine hypersurfaces there exists a canonical choice of transverse vector field ξ (unique up to sign), called the affine (Blaschke) normal, which induces a connection ∇ , a nondegenerate symmetric bilinear form h and a 1-1 tensor field S by

$$D_X Y = \nabla_X Y + h(X, Y)\xi, \quad (1)$$

$$D_X \xi = -SX, \quad (2)$$

for all $X, Y \in \mathcal{X}(M)$. The connection ∇ is called the induced affine connection, h is called the affine metric or Blaschke metric and S is called the affine shape operator. In general ∇ is not the Levi-Civita connection $\hat{\nabla}$ of h . The difference tensor K is defined as

$$K(X, Y) = \nabla_X Y - \hat{\nabla}_X Y, \quad (3)$$

for all $X, Y \in \mathcal{X}(M)$. Moreover the form $h(K(X, Y), Z)$ is a symmetric cubic form with the property that for any fixed $X \in \mathcal{X}(M)$, trace K_X vanishes. This last property is called the

*This paper is a contribution to the Special Issue "Élie Cartan and Differential Geometry". The full collection is available at <http://www.emis.de/journals/SIGMA/Cartan.html>

apolarity condition. The difference tensor K , together with the affine metric h and the affine shape operator S are the most fundamental algebraic invariants for a nondegenerate affine hypersurface (more details in Section 2). We say that M^n is indefinite, definite, etc. if the affine metric h is indefinite, definite, etc. (Because the affine metric is a multiple of the Euclidean second fundamental form, a positive definite hypersurface is locally strongly convex.) For details of the basic theory of nondegenerate affine hypersurfaces we refer to [7] and [10].

Here we will restrict ourselves to the case of affine hyperspheres, i.e. the shape operator will be a (constant) multiple of the identity ($S = HId$). Geometrically this means that all affine normals pass through a fixed point or they are parallel. There are many affine hyperspheres, even in the two-dimensional case only partial classifications are known. This is due to the fact that affine hyperspheres reduce to the study of the Monge-Ampère equations. Our question is the following: *What can we say about a three-dimensional affine hypersphere which admits a pointwise G -symmetry, i.e. there exists a non-trivial subgroup G of the isometry group such that for every $p \in M$ and every $L \in G$:*

$$K(LX_p, LY_p) = L(K(X_p, Y_p)) \quad \forall X_p, Y_p \in T_pM.$$

We have motivated this question in Part 1 [13] (see also [1, 14, 8]). A classification of 3-dimensional positive definite affine hyperspheres admitting pointwise symmetries was obtained in [14]. We continue the classification in the indefinite case. We can assume that the affine metric has index two, i.e. the corresponding isometry group is the (special) Lorentz group $SO(1, 2)$. In Part 1, it turns out that a $SO(1, 2)$ -stabilizer of a nontrivial traceless cubic form is isomorphic to either $SO(2)$, $SO(1, 1)$, \mathbb{R} , the group S_3 of order 6, $\mathbb{Z}_2 \times \mathbb{Z}_2$, \mathbb{Z}_3 , \mathbb{Z}_2 or it is trivial. We have shown that hyperspheres admitting a pointwise $\mathbb{Z}_2 \times \mathbb{Z}_2$ - resp. \mathbb{R} -symmetry are well-known, they have constant sectional curvature and Pick invariant $J < 0$ resp. $J = 0$.

In the following we classify the indefinite affine hyperspheres which admit a pointwise \mathbb{Z}_3 -, $SO(2)$ - or $SO(1, 1)$ -symmetry. They turn out to be warped products of affine spheres (\mathbb{Z}_3) or quadrics ($SO(2)$, $SO(1, 1)$) with a curve. As a result we get a new composition method. Since the methods for the proofs for $SO(1, 1)$ are similar to those for \mathbb{Z}_3 - or $SO(2)$ -symmetry (and as long) we will omit them here. Both methods and results are different in case of S_3 -symmetry and will be published elsewhere. The paper is organized as follows:

We will state the basic formulas of (equi-)affine hypersurface-theory needed in the further classification in Section 2. In Section 3, we show that in case of $SO(2)$ -, S_3 - or \mathbb{Z}_3 -symmetry we can extend the canonical form of K (cf. [13]) locally. Thus we can obtain information about the coefficients of K and ∇ from the basic equations of Gauss, Codazzi and Ricci (cf. Section 4). In Section 5 we show, that in case of \mathbb{Z}_3 - or $SO(2)$ -symmetry it follows that the hypersurface admits a warped product structure $\mathbb{R} \times_{ef} N^2$. Then we classify such hyperspheres by showing how they can be constructed starting from 2-dimensional positive definite affine spheres resp. quadrics (cf. Theorems 1–8). We end in Section 6 by stating the classification results in case of $SO(1, 1)$ -symmetry (cf. Theorems 9–16).

The classification can be seen as a generalization of the well known Calabi product of hyperbolic affine spheres [2, 6] and of the constructions for affine spheres considered in [5]. The following natural question for a (de)composition theorem, related to these constructions, gives another motivation for studying 3-dimensional hypersurfaces admitting a pointwise symmetry:

(De)composition Problem. *Let M^n be a nondegenerate affine hypersurface in \mathbb{R}^{n+1} . Under what conditions do there exist affine hyperspheres M_1^r in \mathbb{R}^{r+1} and M_2^s in \mathbb{R}^{s+1} , with $r+s = n-1$, such that $M = I \times_{f_1} M_1 \times_{f_2} M_2$, where $I \subset \mathbb{R}$ and f_1 and f_2 depend only on I (i.e. M admits a warped product structure)? How can the original immersion be recovered starting from the immersion of the affine spheres?*

Of course the first dimension in which the above problem can be considered is three and our results provide an answer in that case.

2 Basics of affine hypersphere theory

First we recall the definition of the affine normal ξ (cf. [10]). In equi-affine hypersurface theory on the ambient space \mathbb{R}^{n+1} a fixed volume form \det is given. A transverse vector field ξ induces a volume form θ on M by $\theta(X_1, \dots, X_n) = \det(\varphi_*X_1, \dots, \varphi_*X_n, \xi)$. Also the affine metric h defines a volume form ω_h on M , namely $\omega_h = |\det h|^{1/2}$. Now the affine normal ξ is uniquely determined (up to sign) by the conditions that $D\xi$ is everywhere tangential (which is equivalent to $\nabla\theta = 0$) and that

$$\theta = \omega_h. \quad (4)$$

Since we only consider 3-dimensional indefinite hyperspheres, i.e.

$$S = HId, \quad H = \text{const}, \quad (5)$$

we can fix the orientation of the affine normal ξ such that the affine metric has signature one. Then the sign of H in the definition of an affine hypersphere is an invariant.

Next we state some of the fundamental equations, which a nondegenerate hypersurface has to satisfy, see also [10] or [7]. These equations relate S and K with amongst others the curvature tensor R of the induced connection ∇ and the curvature tensor \hat{R} of the Levi-Civita connection $\widehat{\nabla}$ of the affine metric h . There are the Gauss equation for ∇ , which states that:

$$R(X, Y)Z = h(Y, Z)SX - h(X, Z)SY,$$

and the Codazzi equation

$$(\nabla_X S)Y = (\nabla_Y S)X.$$

Also we have the total symmetry of the affine cubic form

$$C(X, Y, Z) = (\nabla_X h)(Y, Z) = -2h(K(X, Y), Z). \quad (6)$$

The fundamental existence and uniqueness theorem, see [3] or [4], states that given h , ∇ and S such that the difference tensor is symmetric and traceless with respect to h , on a simply connected manifold M an affine immersion of M exists if and only if the above Gauss equation and Codazzi equation are satisfied.

From the Gauss equation and Codazzi equation above the Codazzi equation for K and the Gauss equation for $\widehat{\nabla}$ follow:

$$(\widehat{\nabla}_X K)(Y, Z) - (\widehat{\nabla}_Y K)(X, Z) = \frac{1}{2}(h(Y, Z)SX - h(X, Z)SY - h(SY, Z)X + h(SX, Z)Y),$$

and

$$\hat{R}(X, Y)Z = \frac{1}{2}(h(Y, Z)SX - h(X, Z)SY + h(SY, Z)X - h(SX, Z)Y) - [K_X, K_Y]Z$$

If we define the Ricci tensor of the Levi-Civita connection $\widehat{\nabla}$ by:

$$\widehat{\text{Ric}}(X, Y) = \text{trace}\{Z \mapsto \hat{R}(Z, X)Y\}. \quad (7)$$

and the Pick invariant by:

$$J = \frac{1}{n(n-1)}h(K, K), \quad (8)$$

then from the Gauss equation we get for the scalar curvature $\hat{\kappa} = \frac{1}{n(n-1)}(\sum_{i,j} h^{ij} \widehat{\text{Ric}}_{ij})$:

$$\hat{\kappa} = H + J. \quad (9)$$

For an affine hypersphere the Gauss and Codazzi equations have the form:

$$R(X, Y)Z = H(h(Y, Z)X - h(X, Z)Y), \quad (10)$$

$$(\nabla_X H)Y = (\nabla_Y H)X, \quad \text{i.e. } H = \text{const}, \quad (11)$$

$$(\widehat{\nabla}_X K)(Y, Z) = (\widehat{\nabla}_Y K)(X, Z), \quad (12)$$

$$\hat{R}(X, Y)Z = H(h(Y, Z)X - h(X, Z)Y) - [K_X, K_Y]Z. \quad (13)$$

Since H is constant, we can rescale φ such that $H \in \{-1, 0, 1\}$.

3 A local frame for pointwise $\text{SO}(2)$ -, S_3 - or \mathbb{Z}_3 -symmetry

Let M^3 be a hypersphere admitting a $\text{SO}(2)$ -, S_3 - or \mathbb{Z}_3 -symmetry. According to [13, Theorem 4, 2.-4.] there exists for every $p \in M^3$ an ONB $\{\mathbf{t}, \mathbf{v}, \mathbf{w}\}$ of $T_p M^3$ such that

$$\begin{aligned} K(\mathbf{t}, \mathbf{t}) &= -2a_4\mathbf{t}, & K(\mathbf{t}, \mathbf{v}) &= a_4\mathbf{v}, & K(\mathbf{t}, \mathbf{w}) &= a_4\mathbf{w}, \\ K(\mathbf{v}, \mathbf{v}) &= -a_4\mathbf{t} + a_6\mathbf{v}, & K(\mathbf{v}, \mathbf{w}) &= -a_6\mathbf{w}, & K(\mathbf{w}, \mathbf{w}) &= -a_4\mathbf{t} - a_6\mathbf{v}, \end{aligned}$$

where $a_4 > 0$ and $a_6 = 0$ in case of $\text{SO}(2)$ -symmetry, $a_4 = 0$ and $a_6 > 0$ for S_3 , and $a_4 > 0$ and $a_6 > 0$ for \mathbb{Z}_3 .

We would like to extend the ONB locally. It is well known that $\widehat{\text{Ric}}$ (cf. (7)) is a symmetric operator and we compute (some of the computations in this section are done with the CAS Mathematica¹):

Lemma 1. *Let $p \in M^3$ and $\{\mathbf{t}, \mathbf{v}, \mathbf{w}\}$ the basis constructed earlier. Then*

$$\begin{aligned} \widehat{\text{Ric}}(\mathbf{t}, \mathbf{t}) &= -2(H - 3a_4^2), & \widehat{\text{Ric}}(\mathbf{t}, \mathbf{v}) &= 0, \\ \widehat{\text{Ric}}(\mathbf{t}, \mathbf{w}) &= 0, & \widehat{\text{Ric}}(\mathbf{v}, \mathbf{v}) &= 2(H - a_4^2 + a_6^2), \\ \widehat{\text{Ric}}(\mathbf{v}, \mathbf{w}) &= 0, & \widehat{\text{Ric}}(\mathbf{w}, \mathbf{w}) &= 2(H - a_4^2 + a_6^2). \end{aligned}$$

Proof. The proof is a straight-forward computation using the Gauss equation (13). It follows e.g. that

$$\begin{aligned} \hat{R}(\mathbf{t}, \mathbf{v})\mathbf{t} &= H\mathbf{v} - K_{\mathbf{t}}(a_4\mathbf{v}) + K_{\mathbf{v}}(-2a_4\mathbf{t}) = H\mathbf{v} - a_4^2\mathbf{v} - 2a_4^2\mathbf{v} = (H - 3a_4^2)\mathbf{v}, \\ \hat{R}(\mathbf{t}, \mathbf{w})\mathbf{t} &= H\mathbf{w} - K_{\mathbf{t}}(a_4\mathbf{w}) + K_{\mathbf{w}}(-2a_4\mathbf{t}) = H\mathbf{w} - a_4^2\mathbf{w} - 2a_4^2\mathbf{w} = (H - 3a_4^2)\mathbf{w}, \\ \hat{R}(\mathbf{t}, \mathbf{v})\mathbf{w} &= -K_{\mathbf{t}}(-a_6\mathbf{w}) + K_{\mathbf{v}}(a_4\mathbf{w}) = 0. \end{aligned}$$

From this it immediately follows that

$$\widehat{\text{Ric}}(\mathbf{t}, \mathbf{t}) = -2(H - 3a_4^2)$$

and

$$\widehat{\text{Ric}}(\mathbf{t}, \mathbf{w}) = 0.$$

The other equations follow by similar computations. ■

¹See Appendix or http://www.math.tu-berlin.de/~schar/IndefSym_typ234.html.

We want to show that the basis we have constructed, at each point p , can be extended differentiably to a neighborhood of the point p such that, at every point, K with respect to the frame $\{T, V, W\}$ has the previously described form.

Lemma 2. *Let M^3 be an affine hypersphere in \mathbb{R}^4 which admits a pointwise $\text{SO}(2)$ -, S_3 - or \mathbb{Z}_3 -symmetry. Let $p \in M$. Then there exists an orthonormal frame $\{T, V, W\}$ defined in a neighborhood of the point p such that K is given by:*

$$\begin{aligned} K(T, T) &= -2a_4T, & K(T, V) &= a_4V, & K(T, W) &= a_4W, \\ K(V, V) &= -a_4T + a_6V, & K(V, W) &= -a_6W, & K(W, W) &= -a_4T - a_6V, \end{aligned}$$

where $a_4 > 0$ and $a_6 = 0$ in case of $\text{SO}(2)$ -symmetry, $a_4 = 0$ and $a_6 > 0$ in case of S_3 -symmetry, and $a_4 > 0$ and $a_6 > 0$ in case of \mathbb{Z}_3 -symmetry.

Proof. First we want to show that at every point the vector \mathbf{t} is uniquely defined (up to sign) and differentiable. We introduce a symmetric operator \hat{A} by:

$$\widehat{\text{Ric}}(Y, Z) = h(\hat{A}Y, Z).$$

Clearly \hat{A} is a differentiable operator on M . Since $2(H - 3a_4^2) \neq 2(H - a_4^2 + a_6^2)$, the operator has two distinct eigenvalues. A standard result then implies that the eigen distributions are differentiable. We take T a local unit vector field spanning the 1-dimensional eigen distribution, and local orthonormal vector fields \tilde{V} and \tilde{W} spanning the second eigen distribution. If $a_6 = 0$, we can take $V = \tilde{V}$ and $W = \tilde{W}$.

As T is (up to sign) uniquely determined, for $a_6 \neq 0$ there exist differentiable functions a_4, c_6 and $c_7, c_6^2 + c_7^2 \neq 0$, such that

$$\begin{aligned} K(T, T) &= -2a_4T, & K(\tilde{V}, \tilde{V}) &= -a_4T + c_6\tilde{V} + c_7\tilde{W}, \\ K(T, \tilde{V}) &= a_4\tilde{V}, & K(\tilde{V}, \tilde{W}) &= c_7\tilde{V} - c_6\tilde{W}, \\ K(T, \tilde{W}) &= a_4\tilde{W}, & K(\tilde{W}, \tilde{W}) &= -a_4T - c_6\tilde{V} - c_7\tilde{W}. \end{aligned}$$

As we have shown in [13], in the proof of Theorem 2 (Case 2), we can always rotate \tilde{V} and \tilde{W} such that we obtain the desired frame. \blacksquare

Remark 1. It actually follows from the proof of the previous lemma that the vector field T is (up to sign) invariantly defined on M , and therefore the function a_4 , too. Since the Pick invariant (8) $J = \frac{1}{3}(-5a_4^2 + 2a_6^2)$, the function a_6 also is invariantly defined on the affine hypersphere M^3 .

4 Gauss and Codazzi for pointwise $\text{SO}(2)$ -, S_3 - or \mathbb{Z}_3 -symmetry

In this section we always will work with the local frame constructed in the previous lemma. We denote the coefficients of the Levi-Civita connection with respect to this frame by:

$$\begin{aligned} \hat{\nabla}_T T &= a_{12}V + a_{13}W, & \hat{\nabla}_T V &= a_{12}T - b_{13}W, & \hat{\nabla}_T W &= a_{13}T + b_{13}V, \\ \hat{\nabla}_V T &= a_{22}V + a_{23}W, & \hat{\nabla}_V V &= a_{22}T - b_{23}W, & \hat{\nabla}_V W &= a_{23}T + b_{23}V, \\ \hat{\nabla}_W T &= a_{32}V + a_{33}W, & \hat{\nabla}_W V &= a_{32}T - b_{33}W, & \hat{\nabla}_W W &= a_{33}T + b_{33}V. \end{aligned}$$

We will evaluate first the Codazzi and then the Gauss equations ((12) and (13)) to obtain more information.

Lemma 3. *Let M^3 be an affine hypersphere in \mathbb{R}^4 which admits a pointwise $\text{SO}(2)$ -, S_3 - or \mathbb{Z}_3 -symmetry and $\{T, V, W\}$ the corresponding ONB. If the symmetry group is*

SO(2), then $0 = a_{12} = a_{13} = a_{23} = a_{32}$, $a_{33} = a_{22}$ and

$$T(a_4) = -4a_{22}a_4, \quad 0 = V(a_4) = W(a_4),$$

S₃, then $0 = a_{12} = a_{13}$, $a_{23} = -3b_{13} = -a_{32}$, $a_{33} = a_{22}$ and

$$T(a_6) = -a_{22}a_6, \quad V(a_6) = 3b_{33}a_6, \quad W(a_6) = -3b_{23}a_6,$$

Z₃ and $\mathbf{a}_6^2 \neq 4\mathbf{a}_4^2$, then $0 = a_{12} = a_{13} = a_{23} = a_{32}$, $a_{33} = a_{22}$, $b_{13} = 0$,

$$T(a_4) = -4a_{22}a_4, \quad 0 = V(a_4) = W(a_4), \quad \text{and}$$

$$T(a_6) = -a_{22}a_6, \quad V(a_6) = 3b_{33}a_6, \quad W(a_6) = -3b_{23}a_6,$$

Z₃ and $\mathbf{a}_6 = 2\mathbf{a}_4$, then $a_{12} = 2a_{22} = -2a_{33} = -b_{33}$,

$$a_{13} = -2a_{23} = -2a_{32} = b_{23}, \quad b_{13} = 0, \quad \text{and}$$

$$T(a_4) = 0, \quad V(a_4) = -4a_{22}a_4, \quad W(a_4) = 4a_{23}a_4.$$

Proof. An evaluation of the Codazzi equations (12) with the help of the CAS Mathematica leads to the following equations (they relate to eq1–eq6 and eq8–eq9 in the Mathematica notebook):

$$V(a_4) = -2a_{12}a_4, \quad T(a_4) = -4a_{22}a_4 + a_{12}a_6, \quad 0 = 4a_{23}a_4 + a_{13}a_6, \quad (14)$$

$$W(a_4) = -2a_{13}a_4, \quad 0 = 4a_{32}a_4 + a_{13}a_6, \quad T(a_4) = -4a_{33}a_4 - a_{12}a_6, \quad (15)$$

$$T(a_6) - V(a_4) = 3a_{12}a_4 - a_{22}a_6, \quad 0 = a_{13}a_4 + (a_{23} + 3b_{13})a_6, \quad (16)$$

$$W(a_4) = (a_{23} + a_{32})a_6, \quad W(a_6) = (-a_{23} + 3a_{32})a_4 - b_{23}a_6, \quad (17)$$

$$V(a_6) = (-a_{22} + a_{33})a_4 + 3b_{33}a_6, \quad (17)$$

$$T(a_6) = -a_{12}a_4 - a_{33}a_6, \quad W(a_4) = -3a_{13}a_4 + (-a_{32} + 3b_{13})a_6, \quad (18)$$

$$V(a_4) = (-a_{22} + a_{33})a_6, \quad W(a_6) = (3a_{23} - a_{32})a_4 - 3b_{23}a_6, \quad (19)$$

$$0 = (a_{23} - a_{32})a_4, \quad (20)$$

$$W(a_4) = -a_{13}a_4 + (a_{32} - 3b_{13})a_6. \quad (21)$$

From the first equation of (15) (we will use the notation (15).1) and (17).1 resp. (14).3 and (15).2 we get:

$$0 = 2a_{13}a_4 + (a_{23} + a_{32})a_6, \quad 0 = 2(a_{23} + a_{32})a_4 + a_{13}a_6. \quad (22)$$

From (19).1) and (14).1 resp. (14).2 and (15).3 we get:

$$0 = -2a_{12}a_4 + 2(a_{22} - a_{33})a_6, \quad 0 = 2(-a_{22} + a_{33})a_4 + a_{12}a_6. \quad (23)$$

We consider first the case, that $\mathbf{a}_6^2 \neq 4\mathbf{a}_4^2$. Then we obtain from the foregoing equations that $a_{13} = 0$, $a_{32} = -a_{23}$, $a_{12} = 0$ and $a_{33} = a_{22}$. Furthermore it follows from (14).1 that $V(a_4) = 0$, from (14).2 that $T(a_4) = -4a_{22}a_4$ and from (14).3 that $a_{23}a_4 = 0$. Equation (15).1 becomes $W(a_4) = 0$, equation (16).2 $T(a_6) = -a_{22}a_6$ and (16).3 $(a_{23} + 3b_{13})a_6 = 0$. Finally equation (17).2 resp. 3 gives $W(a_6) = -3b_{23}a_6$ and $V(a_6) = 3b_{33}a_6$.

In case of $SO(2)$ -symmetry ($a_4 > 0$ and $a_6 = 0$) it follows that $a_{23} = 0$ and thus the statement of the theorem.

In case of S_3 -symmetry ($a_4 = 0$ and $a_6 > 0$) it follows that $a_{23} = -3b_{13}$ and thus the statement of the theorem.

In case of Z_3 -symmetry ($a_4 > 0$ and $a_6 > 0$) it follows that $a_{23} = 0$ and $b_{13} = 0$ and thus the statement of the theorem.

In case that $a_6 = \pm 2a_4$ ($\neq 0$), we can choose V, W such that $\mathbf{a}_6 = 2\mathbf{a}_4$. Now equations (20), (14).3 and (16).3 lead to $a_{23} = a_{32}$, $a_{13} = -2a_{23}$ and $b_{13} = 0$. A combination of (14).2 and (15).3 gives $a_{12} = (a_{22} - a_{33})$, and then by equations (16).2, (14).1 and (14).2 that $a_{33} = -a_{22}$. Thus $T(a_4) = 0$ by (14).2, $V(a_4) = -4a_{22}a_4$ by (14).1 and $W(a_4) = 4a_{22}a_4$ by (15).1. Finally (17).2 and (15).1 resp. (17).3 and (14).1 imply that $b_{23} = -a_{23}$ resp. $b_{33} = -a_{22}$. ■

An evaluation of the Gauss equations (13) with the help of the CAS Mathematica leads to the following:

Lemma 4. *Let M^3 be an affine hypersphere in \mathbb{R}^4 which admits a pointwise $\text{SO}(2)$ -, S_3 - or \mathbb{Z}_3 -symmetry and $\{T, V, W\}$ the corresponding ONB. Then*

$$T(a_{22}) = -a_{22}^2 + a_{23}^2 + H - 3a_4^2, \quad (24)$$

$$T(a_{23}) = -2a_{22}a_{23}, \quad (25)$$

$$W(a_{22}) + V(a_{23}) = 0, \quad (26)$$

$$W(a_{23}) - V(a_{22}) = 0, \quad (27)$$

$$V(b_{13}) - T(b_{23}) = a_{22}b_{23} + (a_{23} + b_{13})b_{33}, \quad (28)$$

$$T(b_{33}) - W(b_{13}) = (a_{23} + b_{13})b_{23} - a_{22}b_{33}, \quad (29)$$

$$V(b_{33}) - W(b_{23}) = -a_{22}^2 - a_{23}^2 + 2a_{23}b_{13} + b_{23}^2 + b_{33}^2 + H + a_4^2 + 2a_6^2. \quad (30)$$

If the symmetry group is \mathbb{Z}_3 , then $a_6^2 \neq 4a_4^2$.

Proof. The equations relate to eq11–eq13 and eq16 in the Mathematica notebook. If $a_6^2 = 4a_4^2 (\neq 0)$, then we obtain by equations eq11.1 and eq12.3 resp. eq15.3 and eq12.3 that $2V(a_{22}) = -4a_{22}^2 - H + 3a_4^2$ resp. $2W(a_{23}) = 4a_{23}^2 + H - 3a_4^2$, thus $V(a_{22}) - W(a_{23}) = -2a_{22}^2 - 2a_{23}^2 - H + 3a_4^2$. This gives a contradiction to eq13.3, namely $V(a_{22}) - W(a_{23}) = -2a_{22}^2 - 2a_{23}^2 - H - 9a_4^2$. ■

5 Pointwise \mathbb{Z}_3 - or $\text{SO}(2)$ -symmetry

The following methods only work in the case of \mathbb{Z}_3 - or $\text{SO}(2)$ -symmetry, therefore the case of S_3 -symmetry will be considered elsewhere. As the vector field T is globally defined, we can define the distributions $L_1 = \text{span}\{T\}$ and $L_2 = \text{span}\{V, W\}$. In the following we will investigate these distributions. For the terminology we refer to [9].

Lemma 5. *The distribution L_1 is autoparallel (totally geodesic) with respect to $\widehat{\nabla}$.*

Proof. From $\widehat{\nabla}_T T = a_{12}V + a_{13}W = 0$ (cf. Lemmas 3 and 4) the claim follows immediately. ■

Lemma 6. *The distribution L_2 is spherical with mean curvature normal $U_2 = a_{22}T$.*

Proof. For $U_2 = a_{22}T \in L_1 = L_2^\perp$ we have $h(\widehat{\nabla}_{E_a} E_b, T) = h(E_a, E_b)h(U_2, T)$ for $E_a, E_b \in \{V, W\}$, and $h(\widehat{\nabla}_{E_a} U_2, T) = h(E_a(a_{22})T + a_{22}\widehat{\nabla}_{E_a} T, T) = 0$ (cf. Lemma 3 and (26), (27)). ■

Remark 2. a_{22} is independent of the choice of ONB $\{V, W\}$. It therefore is a globally defined function on M^3 .

We introduce a coordinate function t by $\frac{\partial}{\partial t} := T$. Using the previous lemma, according to [12], we get:

Lemma 7. *(M^3, h) admits a warped product structure $M^3 = I \times_{e^f} N^2$ with $f : I \rightarrow \mathbb{R}$ satisfying*

$$\frac{\partial f}{\partial t} = a_{22}. \quad (31)$$

Proof. Proposition 3 in [12] gives the warped product structure with warping function $\lambda_2 : I \rightarrow \mathbb{R}$. If we introduce $f = \ln \lambda_2$, following the proof we see that $a_{22}T = U_2 = -\text{grad}(\ln \lambda_2) = -\text{grad} f$. ■

Lemma 8. *The curvature of N^2 is ${}^N K(N^2) = e^{2f}(H + 2a_6^2 + a_4^2 - a_{22}^2)$.*

Proof. From Proposition 2 in [12] we get the following relation between the curvature tensor \hat{R} of the warped product M^3 and the curvature tensor \tilde{R} of the usual product of pseudo-Riemannian manifolds ($X, Y, Z \in \mathcal{X}(M)$ resp. their appropriate projections):

$$\begin{aligned}\hat{R}(X, Y)Z &= \tilde{R}(X, Y)Z + h(Y, Z)(\hat{\nabla}_X U_2 - h(X, U_2)U_2) - h(\hat{\nabla}_X U_2 - h(X, U_2)U_2, Z)Y \\ &\quad - h(X, Z)(\hat{\nabla}_Y U_2 - h(Y, U_2)U_2) + h(\hat{\nabla}_Y U_2 - h(Y, U_2)U_2, Z)X \\ &\quad + h(U_2, U_2)(h(Y, Z)X - h(X, Z)Y).\end{aligned}$$

Now $\tilde{R}(X, Y)Z = {}^N\tilde{R}(X, Y)Z$ for all $X, Y, Z \in TN^2$ and otherwise zero (cf. [11, page 89], Corollary 58) and $K(N^2) = K(V, W) = \frac{h(-\hat{R}(V, W)V, W)}{h(V, V)h(W, W) - h(V, W)^2}$ (cf. [11, page 77], the curvature tensor has the opposite sign). Since $h(X, Y) = e^{2f}h(X, Y)$ for $X, Y \in TN^2$, it follows that

$${}^N K(N^2) = e^{2f}h(-{}^N\hat{R}(V, W)V, W).$$

Finally we obtain by the Gauss equation (13) the last ingredient for the computation: $\hat{R}(V, W)V = -(H + 2a_6^2 + a_4^2)W$ (cf. the Mathematica notebook). ■

Summarized we have obtained the following structure equations (cf. (1), (2) and (3)), where $a_6 = 0$ in case of $SO(2)$ -symmetry resp. $b_{13} = 0$ in case of \mathbb{Z}_3 -symmetry:

$$D_T T = -2a_4 T - \xi, \tag{32}$$

$$D_T V = +a_4 V - b_{13} W, \tag{33}$$

$$D_T W = +b_{13} V + a_4 W, \tag{34}$$

$$D_V T = +(a_{22} + a_4) V, \tag{35}$$

$$D_W T = +(a_{22} + a_4) W, \tag{36}$$

$$D_V V = +a_6 V - b_{23} W + (a_{22} - a_4) T + \xi, \tag{37}$$

$$D_V W = +b_{23} V - a_6 W, \tag{38}$$

$$D_W V = -(b_{33} + a_6) W, \tag{39}$$

$$D_W W = +(b_{33} - a_6) V + (a_{22} - a_4) T + \xi, \tag{40}$$

$$D_X \xi = -HX. \tag{41}$$

The Codazzi and Gauss equations ((12) and (13)) have the form (cf. Lemmas 3 and 4):

$$T(a_4) = -4a_{22}a_4, \quad 0 = V(a_4) = W(a_4), \tag{42}$$

$$T(a_6) = -a_{22}a_6, \quad V(a_6) = 3b_{33}a_6, \quad W(a_6) = -3b_{23}a_6, \tag{43}$$

$$T(a_{22}) = -a_{22}^2 + H - 3a_4^2, \quad V(a_{22}) = 0, \quad W(a_{22}) = 0, \tag{44}$$

$$V(b_{13}) - T(b_{23}) = a_{22}b_{23} + b_{13}b_{33}, \tag{45}$$

$$T(b_{33}) - W(b_{13}) = b_{13}b_{23} - a_{22}b_{33}, \tag{46}$$

$$V(b_{33}) - W(b_{23}) = -a_{22}^2 + b_{23}^2 + b_{33}^2 + H + a_4^2 + 2a_6^2, \tag{47}$$

where $a_6 = 0$ in case of $SO(2)$ -symmetry resp. $b_{13} = 0$ in case of \mathbb{Z}_3 -symmetry.

Our first goal is to find out how N^2 is immersed in \mathbb{R}^4 , i.e. to find an immersion independent of t . A look at the structure equations (32)–(41) suggests to start with a linear combination of T and ξ .

We will solve the problem in two steps. First we look for a vector field X with $D_T X = \alpha X$ for some function α : We define $X := AT + \xi$ for some function A on M^3 . Then $D_T X = \alpha X$ iff $\alpha = -A$ and $\frac{\partial}{\partial t} A = -A^2 + 2a_4 A + H$, and $A := a_{22} - a_4$ solves the latter differential equation.

Next we want to multiply X with some function β such that $D_T(\beta X) = 0$: We define a positive function β on \mathbb{R} as the solution of the differential equation:

$$\frac{\partial}{\partial t}\beta = (a_{22} - a_4)\beta \quad (48)$$

with initial condition $\beta(t_0) > 0$. Then $D_T(\beta X) = 0$ and by (35), (41) and (36) we get (since β , a_{22} and a_4 only depend on t):

$$D_T(\beta((a_{22} - a_4)T + \xi)) = 0, \quad (49)$$

$$D_V(\beta((a_{22} - a_4)T + \xi)) = \beta(a_{22}^2 - a_4^2 - H)V, \quad (50)$$

$$D_W(\beta((a_{22} - a_4)T + \xi)) = \beta(a_{22}^2 - a_4^2 - H)W. \quad (51)$$

To obtain an immersion we need that $\nu := a_{22}^2 - a_4^2 - H$ vanishes nowhere, but we only get:

Lemma 9. *The function $\nu = a_{22}^2 - a_4^2 - H$ is globally defined, $\frac{\partial}{\partial t}(e^{2f}\nu) = 0$ and ν vanishes identically or nowhere on \mathbb{R} .*

Proof. Since $0 = \frac{\partial}{\partial t}^N K(N^2) = \frac{\partial}{\partial t}(e^{2f}(2a_6^2 - \nu))$ (Lemma 8) and $\frac{\partial}{\partial t}(e^{2f}2a_6^2) = 0$ (cf. (43) and (31)), we get that $\frac{\partial}{\partial t}(e^{2f}\nu) = 0$. Thus $\frac{\partial}{\partial t}\nu = -2(\frac{\partial}{\partial t}f)\nu = -2a_{22}\nu$. ■

5.1 The first case: $\nu \neq 0$ on M^3

We may, by translating f , i.e. by replacing N^2 with a homothetic copy of itself, assume that $e^{2f}\nu = \varepsilon_1$, where $\varepsilon_1 = \pm 1$.

Lemma 10. *$\Phi := \beta((a_{22} - a_4)T + \xi): M^3 \rightarrow \mathbb{R}^4$ induces a proper affine sphere structure, say $\tilde{\phi}$, mapping N^2 into a 3-dimensional linear subspace of \mathbb{R}^4 . $\tilde{\phi}$ is part of a quadric iff $a_6 = 0$.*

Proof. By (50) and (51) we have $\Phi_*(E_a) = \beta\nu E_a$ for $E_a \in \{V, W\}$. A further differentiation, using (37) (β and ν only depend on t), gives:

$$\begin{aligned} D_V\Phi_*(V) &= \beta\nu D_VV = \beta\nu((a_{22} - a_4)T + a_6V - b_{23}W + \xi) \\ &= a_6\Phi_*(V) - b_{23}\Phi_*(W) + \nu\Phi = a_6\Phi_*(V) - b_{23}\Phi_*(W) + \varepsilon_1 e^{-2f}\Phi. \end{aligned}$$

Similarly, we obtain the other derivatives, using (38)–(40), thus:

$$D_V\Phi_*(V) = a_6\Phi_*(V) - b_{23}\Phi_*(W) + e^{-2f}\varepsilon_1\Phi, \quad (52)$$

$$D_V\Phi_*(W) = b_{23}\Phi_*(V) - a_6\Phi_*(W), \quad (53)$$

$$D_W\Phi_*(V) = -(b_{33} + a_6)\Phi_*(W), \quad (54)$$

$$D_W\Phi_*(W) = (b_{33} - a_6)\Phi_*(V) + e^{-2f}\varepsilon_1\Phi, \quad (55)$$

$$D_{E_a}\Phi = \beta e^{-2f}\varepsilon_1 E_a. \quad (56)$$

The foliation at $f = f_0$ gives an immersion of N^2 to M^3 , say π_{f_0} . Therefore, we can define an immersion of N^2 to \mathbb{R}^4 by $\tilde{\phi} := \Phi \circ \pi_{f_0}$, whose structure equations are exactly the equations above when $f = f_0$. Hence, we know that $\tilde{\phi}$ maps N^2 into $\text{span}\{\Phi_*(V), \Phi_*(W), \Phi\}$, an affine hyperplane of \mathbb{R}^4 and $\frac{\partial}{\partial t}\Phi = 0$ implies $\Phi(t, v, w) = \tilde{\phi}(v, w)$.

We can read off the coefficients of the difference tensor $K^{\tilde{\phi}}$ of $\tilde{\phi}$ (cf. (1) and (3)): $K^{\tilde{\phi}}(\tilde{V}, \tilde{V}) = a_6\tilde{V}$, $K^{\tilde{\phi}}(\tilde{V}, \tilde{W}) = -a_6\tilde{W}$, $K^{\tilde{\phi}}(\tilde{W}, \tilde{W}) = -a_6\tilde{V}$, and see that $\text{trace}(K^{\tilde{\phi}})_X$ vanishes. The affine metric introduced by this immersion corresponds with the metric on N^2 . Thus $\varepsilon_1\tilde{\phi}$ is the affine normal of $\tilde{\phi}$ and $\tilde{\phi}$ is a proper affine sphere with mean curvature ε_1 . Finally the vanishing of the difference tensor characterizes quadrics. ■

Our next goal is to find another linear combination of T and ξ , this time only depending on t . (Then we can express T in terms of ϕ and some function of t .)

Lemma 11. *Define $\delta := HT + (a_{22} + a_4)\xi$. Then there exist a constant vector $C \in \mathbb{R}^4$ and a function $a(t)$ such that*

$$\delta(t) = a(t)C.$$

Proof. Using (35) resp. (36) and (41) we obtain that $D_V\delta = 0 = D_W\delta$. Hence δ depends only on the variable t . Moreover, we get by (32), (44), (42) and (41) that

$$\begin{aligned} \frac{\partial}{\partial t}\delta &= D_T(HT + (a_{22} + a_4)\xi) \\ &= H(-2a_4T - \xi) + (-a_{22}^2 + H - 3a_4^2 - 4a_{22}a_4)\xi - (a_{22} + a_4)HT \\ &= -(3a_4 + a_{22})(HT + (a_{22} + a_4)\xi) = -(3a_4 + a_{22})\delta. \end{aligned}$$

This implies that there exists a constant vector C in \mathbb{R}^4 and a function $a(t)$ such that $\delta(t) = a(t)C$. \blacksquare

Notice that for an improper affine hypersphere ($H = 0$) ξ is constant and parallel to C . Combining $\tilde{\phi}$ and δ we obtain for T (cf. Lemmas 10 and 11) that

$$T(t, v, w) = -\frac{a}{\nu}C + \frac{1}{\beta\nu}(a_{22} + a_4)\tilde{\phi}(v, w). \quad (57)$$

In the following we will use for the partial derivatives the abbreviation $\varphi_x := \frac{\partial}{\partial x}\varphi$, $x = t, v, w$.

Lemma 12.

$$\varphi_t = -\frac{a}{\nu}C + \frac{\partial}{\partial t}\left(\frac{1}{\beta\nu}\right)\tilde{\phi}, \quad \varphi_v = \frac{1}{\beta\nu}\tilde{\phi}_v, \quad \varphi_w = \frac{1}{\beta\nu}\tilde{\phi}_w.$$

Proof. As by (48) and Lemma 9 $\frac{\partial}{\partial t}\frac{1}{\beta\nu} = \frac{1}{\beta\nu}(a_{22} + a_4)$, we obtain the equation for $\varphi_t = T$ by (57). The other equations follow from (50) and (51). \blacksquare

It follows by the uniqueness theorem of first order differential equations and applying a translation that we can write

$$\varphi(t, v, w) = \tilde{a}(t)C + \frac{1}{\beta\nu}(t)\tilde{\phi}(v, w)$$

for a suitable function \tilde{a} depending only on the variable t . Since C is transversal to the image of $\tilde{\phi}$ (cf. Lemmas 10 and 11, $\nu \neq 0$), we obtain that after applying an equiaffine transformation we can write: $\varphi(t, v, w) = (\gamma_1(t), \gamma_2(t)\phi(v, w))$, in which $\tilde{\phi}(v, w) = (0, \phi(v, w))$. Thus we have proven the following:

Theorem 1. *Let M^3 be an indefinite affine hypersphere of \mathbb{R}^4 which admits a pointwise \mathbb{Z}_3 - or $SO(2)$ -symmetry. Let $a_{22}^2 - a_4^2 \neq H$ for some $p \in M^3$. Then M^3 is affine equivalent to*

$$\varphi : I \times N^2 \rightarrow \mathbb{R}^4 : (t, v, w) \mapsto (\gamma_1(t), \gamma_2(t)\phi(v, w)),$$

where $\phi : N^2 \rightarrow \mathbb{R}^3$ is a (positive definite) elliptic or hyperbolic affine sphere and $\gamma : I \rightarrow \mathbb{R}^2$ is a curve. Moreover, if M^3 admits a pointwise $SO(2)$ -symmetry then N^2 is either an ellipsoid or a two-sheeted hyperboloid.

We want to investigate the conditions imposed on the curve γ . For this we compute the derivatives of φ :

$$\begin{aligned}\varphi_t &= (\gamma'_1, \gamma'_2 \phi), & \varphi_v &= (0, \gamma_2 \phi_v), & \varphi_w &= (0, \gamma_2 \phi_w), \\ \varphi_{tt} &= (\gamma''_1, \gamma''_2 \phi), & \varphi_{tv} &= (0, \gamma'_2 \phi_v), & \varphi_{tw} &= (0, \gamma'_2 \phi_w), \\ \varphi_{vv} &= (0, \gamma_2 \phi_{vv}), & \varphi_{vw} &= (0, \gamma_2 \phi_{vw}), & \varphi_{ww} &= (0, \gamma_2 \phi_{ww}).\end{aligned}\tag{58}$$

Furthermore we have to distinguish if M^3 is proper ($H = \pm 1$) or improper ($H = 0$).

First we consider the case that M^3 is proper, i.e. $\xi = -H\varphi$. An easy computation shows that the condition that ξ is a transversal vector field, namely $0 \neq \det(\varphi_t, \varphi_v, \varphi_w, \xi) = -\gamma_2^2(\gamma_1 \gamma'_2 - \gamma'_1 \gamma_2) \det(\phi_v, \phi_w, \phi)$, is equivalent to $\gamma_2 \neq 0$ and $\gamma_1 \gamma'_2 - \gamma'_1 \gamma_2 \neq 0$. To check the condition that ξ is the Blaschke normal (cf. (4)), we need to compute the Blaschke metric h , using (1), (58), (52)–(55) and the notation $r, s \in \{v, w\}$ and g for the Blaschke metric of ϕ :

$$\begin{aligned}\varphi_{tt} &= \dots \varphi_t + \frac{\gamma'_1 \gamma''_2 - \gamma''_1 \gamma'_2}{H(\gamma_1 \gamma'_2 - \gamma'_1 \gamma_2)} \xi, & \varphi_{tr} &= \text{tang}, \\ \varphi_{rs} &= \text{tang} - \frac{\gamma'_1 \gamma_2}{H(\gamma_1 \gamma'_2 - \gamma'_1 \gamma_2)} \varepsilon_{1g} \left(\frac{\partial}{\partial r}, \frac{\partial}{\partial s} \right) \xi.\end{aligned}$$

We obtain that

$$\det h = h_{tt}(h_{vv}h_{ww} - h_{vw}^2) = \frac{\gamma'_1 \gamma''_2 - \gamma''_1 \gamma'_2}{H^3(\gamma_1 \gamma'_2 - \gamma'_1 \gamma_2)^3} (\gamma'_1)^2 \gamma_2^2 \det g.$$

Thus

$$\gamma_2^4 (\gamma_1 \gamma'_2 - \gamma'_1 \gamma_2)^2 \det(\phi_v, \phi_w, \phi)^2 = \left| \frac{\gamma'_1 \gamma''_2 - \gamma''_1 \gamma'_2}{(\gamma_1 \gamma'_2 - \gamma'_1 \gamma_2)^3} (\gamma'_1)^2 \gamma_2^2 \det g \right|$$

is equivalent to (4). Since ϕ is a definite proper affine sphere with normal $-\varepsilon_1 \phi$, we can again use (4) to obtain

$$\xi = -H\varphi \iff \gamma_2^2 |\gamma_1 \gamma'_2 - \gamma'_1 \gamma_2|^5 = |\gamma'_1 \gamma''_2 - \gamma''_1 \gamma'_2| (\gamma'_1)^2 \neq 0.$$

From the computations above (g is positive definite) also it follows that φ is indefinite iff either

$$\begin{aligned}H \operatorname{sign}(\gamma_1 \gamma'_2 - \gamma'_1 \gamma_2) &= \operatorname{sign}(\gamma'_1 \gamma''_2 - \gamma''_1 \gamma'_2) = \operatorname{sign}(\gamma'_1 \gamma_2 \varepsilon_1) & \text{or} \\ -H \operatorname{sign}(\gamma_1 \gamma'_2 - \gamma'_1 \gamma_2) &= \operatorname{sign}(\gamma'_1 \gamma''_2 - \gamma''_1 \gamma'_2) = \operatorname{sign}(\gamma'_1 \gamma_2 \varepsilon_1).\end{aligned}$$

Next we consider the case that M^3 is improper, i.e. ξ is constant. By Lemma 11 ξ is parallel to C and thus transversal to ϕ . Hence we can apply an affine transformation to obtain $\xi = (1, 0, 0, 0)$. An easy computation shows that the condition that ξ is a transversal vector field, namely $0 \neq \det(\varphi_t, \varphi_v, \varphi_w, \xi) = -\gamma_2^2 \gamma'_2 \det(\phi_v, \phi_w, \phi)$, is equivalent to $\gamma_2 \neq 0$ and $\gamma'_2 \neq 0$. To check the condition that ξ is the Blaschke normal (cf. (4)) we need to compute the Blaschke metric h , using (1), (58), (52)–(55) and the notation $r, s \in \{v, w\}$ and g for the Blaschke metric of ϕ :

$$\varphi_{tt} = \dots \varphi_t - \frac{\gamma'_1 \gamma''_2 - \gamma''_1 \gamma'_2}{\gamma'_2} \xi, \quad \varphi_{tr} = \text{tang}, \quad \varphi_{rs} = \text{tang} + \frac{\gamma'_1 \gamma_2}{\gamma'_2} \varepsilon_{1g} \left(\frac{\partial}{\partial r}, \frac{\partial}{\partial s} \right) \xi.$$

We obtain that

$$\det h = h_{tt}(h_{vv}h_{ww} - h_{vw}^2) = -\frac{\gamma'_1 \gamma''_2 - \gamma''_1 \gamma'_2}{(\gamma'_2)^3} (\gamma'_1)^2 \gamma_2^2 \det g.$$

Thus (4) is equivalent to

$$\gamma_2^4 (\gamma_2')^2 \det(\phi_v, \phi_w, \phi)^2 = \left| \frac{\gamma_1' \gamma_2'' - \gamma_1'' \gamma_2'}{(\gamma_2')^3} (\gamma_1')^2 \gamma_2^2 \det g \right|.$$

Since ϕ is a definite proper affine sphere with normal $-\varepsilon_1 \phi$, we can again use (4) to obtain

$$\xi = (1, 0, 0, 0) \iff \gamma_2^2 |\gamma_2'|^5 = |\gamma_1' \gamma_2'' - \gamma_1'' \gamma_2'| (\gamma_1')^2 \neq 0.$$

From the computations above also it follows that φ is indefinite iff either

$$\begin{aligned} -\text{sign}(\gamma_2') &= \text{sign}(\gamma_1' \gamma_2'' - \gamma_1'' \gamma_2') = \text{sign}(\gamma_1' \gamma_2 \varepsilon_1) & \text{or} \\ \text{sign}(\gamma_2') &= \text{sign}(\gamma_1' \gamma_2'' - \gamma_1'' \gamma_2') = \text{sign}(\gamma_1' \gamma_2 \varepsilon_1). \end{aligned}$$

So we have seen under which conditions we can construct a 3-dimensional indefinite affine hypersphere out of an affine sphere:

Theorem 2. *Let $\phi : N^2 \rightarrow \mathbb{R}^3$ be a positive definite elliptic or hyperbolic affine sphere (with mean curvature $\varepsilon_1 = \pm 1$), and let $\gamma = (\gamma_1, \gamma_2) : I \rightarrow \mathbb{R}^2$ be a curve. Define $\varphi : I \times N^2 \rightarrow \mathbb{R}^4$ by $\varphi(t, v, w) = (\gamma_1(t), \gamma_2(t)\phi(v, w))$.*

- (i) *If γ satisfies $\gamma_2^2 |\gamma_1' \gamma_2'' - \gamma_1'' \gamma_2'|^5 = \text{sign}(\gamma_1' \gamma_2 \varepsilon_1) (\gamma_1' \gamma_2'' - \gamma_1'' \gamma_2') (\gamma_1')^2 \neq 0$, then φ defines a 3-dimensional indefinite proper affine hypersphere.*
- (ii) *If γ satisfies $\gamma_2^2 |\gamma_2'|^5 = \text{sign}(\gamma_1' \gamma_2 \varepsilon_1) (\gamma_1' \gamma_2'' - \gamma_1'' \gamma_2') (\gamma_1')^2 \neq 0$, then φ defines a 3-dimensional indefinite improper affine hypersphere.*

Now we are ready to check the symmetries.

Theorem 3. *Let $\phi : N^2 \rightarrow \mathbb{R}^3$ be a positive definite elliptic or hyperbolic affine sphere (with mean curvature $\varepsilon_1 = \pm 1$), and let $\gamma = (\gamma_1, \gamma_2) : I \rightarrow \mathbb{R}^2$ be a curve such that $\varphi(t, v, w) = (\gamma_1(t), \gamma_2(t)\phi(v, w))$ defines a 3-dimensional indefinite affine hypersphere. Then $\varphi(N^2 \times I)$ admits a pointwise \mathbb{Z}_3 - or $SO(2)$ -symmetry.*

Proof. We already have shown that φ defines a 3-dimensional indefinite proper resp. improper affine hypersphere. To prove the symmetry we need to compute K . By assumption, ϕ is an affine sphere with Blaschke normal $\xi^\phi = -\varepsilon_1 \phi$. For the structure equations (1) we use the notation $\phi_{rs} = \phi \Gamma_{rs}^u \phi_u - g_{rs} \varepsilon_1 \phi$, $r, s, u \in \{v, w\}$. Furthermore we introduce the notation $\alpha = \gamma_1 \gamma_2' - \gamma_1' \gamma_2$. Note that $\alpha' = \gamma_1 \gamma_2'' - \gamma_1'' \gamma_2$. If φ is proper, using (58), we get the structure equations (1) for φ :

$$\begin{aligned} \varphi_{tt} &= \frac{\alpha'}{\alpha} \varphi_t + \frac{\gamma_1' \gamma_2'' - \gamma_1'' \gamma_2'}{H \alpha} \xi, & \varphi_{tr} &= \frac{\gamma_2'}{\gamma_2} \varphi_r, \\ \varphi_{rs} &= \phi \Gamma_{rs}^u \varphi_u - g_{rs} \varepsilon_1 \frac{\gamma_1 \gamma_2}{\alpha} \varphi_t - g_{rs} \varepsilon_1 \frac{\gamma_1' \gamma_2}{H \alpha} \xi. \end{aligned}$$

We compute K using (6) and obtain:

$$\begin{aligned} (\nabla_{\varphi_t} h)(\varphi_r, \varphi_s) &= \left(\left(\frac{\gamma_1 \gamma_2}{\alpha} \right)' \frac{\alpha}{\gamma_1 \gamma_2} - 2 \frac{\gamma_2'}{\gamma_2} \right) h(\varphi_r, \varphi_s), \\ (\nabla_{\varphi_r} h)(\varphi_t, \varphi_t) &= 0, \end{aligned}$$

implying that K_{φ_t} restricted to the space spanned by φ_v and φ_w is a multiple of the identity. Taking T in direction of φ_t , we see that φ_v and φ_w are orthogonal to T . Thus we can construct an ONB $\{T, V, W\}$ with V, W spanning $\text{span}\{\varphi_v, \varphi_w\}$ such that $a_1 = 2a_4$, $a_2 = a_3 = a_5 = 0$. By the considerations in [13, Section 4] we see that φ admits a pointwise \mathbb{Z}_3 - or $SO(2)$ -symmetry. If φ is improper, the proof runs completely analogous. \blacksquare

5.2 The second case: $\nu \equiv 0$ and $H \neq 0$ on M^3

Next, we consider the case that $H = a_{22}^2 - a_4^2$ and $H \neq 0$ on M^3 . It follows that $a_{22} \neq \pm a_4$ on M^3 .

We already have seen that M^3 admits a warped product structure. The map Φ we have constructed in Lemma 10 will not define an immersion (cf. (50) and (51)). Anyhow, for a fixed point t_0 , we get from (37)–(40), (50) and (51), using the notation $\tilde{\xi} = (a_{22} - a_4)T + \xi$:

$$\begin{aligned} D_V V &= a_6 V - b_{23} W + \tilde{\xi}, & D_V W &= b_{23} V - a_6 W, \\ D_W V &= -(b_{33} + a_6) W, & D_W W &= (b_{33} - a_6) V + \tilde{\xi}, & D_{E_a} \tilde{\xi} &= 0, & E_a &\in \{V, W\}. \end{aligned}$$

Thus, if v and w are local coordinates which span the second distribution L_2 , then we can interpret $\varphi(t_0, v, w)$ as a positive definite improper affine sphere in a 3-dimensional linear subspace.

Moreover, we see that this improper affine sphere is a paraboloid provided that $a_6(t_0, v, w)$ vanishes identically. From the differential equations (43) determining a_6 , we see that this is the case exactly when a_6 vanishes identically, i.e. when M^3 admits a pointwise $SO(2)$ -symmetry.

After applying a translation and a change of coordinates, we may assume that

$$\varphi(t_0, v, w) = (v, w, f(v, w), 0),$$

with affine normal $\tilde{\xi}(t_0, v, w) = (0, 0, 1, 0)$. To obtain T at t_0 , we consider (35) and (36) and get that

$$D_{E_a}(T - (a_{22} + a_4)\varphi) = 0, \quad E_a, E_b \in \{V, W\}.$$

Evaluating at $t = t_0$, this means that there exists a constant vector C , transversal to $\text{span}\{V, W, \xi\}$, such that $T(t_0, v, w) = (a_{22} + a_4)(t_0)\varphi(t_0, v, w) + C$. Since $a_{22} + a_4 \neq 0$ everywhere, we can write:

$$T(t_0, v, w) = \alpha_1(v, w, f(v, w), \alpha_2), \tag{59}$$

where $\alpha_1, \alpha_2 \neq 0$ and we applied an equiaffine transformation so that $C = (0, 0, 0, \alpha_1 \alpha_2)$. To obtain information about $D_T T$ we have that $D_T T = -2a_4 T - \xi$ (cf. (32)) and $\xi = \tilde{\xi} - (a_{22} - a_4)T$ by the definition of $\tilde{\xi}$. Also we know that $\tilde{\xi}(t_0, v, w) = (0, 0, 1, 0)$ and by (49)–(51) that $D_X(\beta\tilde{\xi}) = 0$, $X \in \mathcal{X}(M)$. Taking suitable initial conditions for the function β ($\beta(t_0) = 1$), we get that $\beta\tilde{\xi} = (0, 0, 1, 0)$ and finally the following vector valued differential equation:

$$D_T T = (a_{22} - 3a_4)T - \frac{1}{\beta}(0, 0, 1, 0).$$

Solving this differential equation, taking into account the initial conditions (59) at $t = t_0$, we get that there exist functions δ_1 and δ_2 depending only on t such that

$$T(t, u, v) = (\delta_1(t)v, \delta_1(t)w, \delta_1(t)(f(v, w) + \delta_2(t)), \alpha_2 \delta_1(t)),$$

where $\delta_1(t_0) = \alpha_1$, $\delta_2(t_0) = 0$, $\delta_1'(t) = (a_{22} - 3a_4)\delta_1(t)$ and $\delta_2'(t) = \delta_1^{-1}(t)\beta^{-1}(t)$. As $T(t, v, w) = \frac{\partial \varphi}{\partial t}(t, v, w)$ and $\varphi(t_0, v, w) = (v, w, f(v, w), 0)$ it follows by integration that

$$\varphi(t, v, w) = (\gamma_1(t)v, \gamma_1(t)w, \gamma_1(t)f(v, w) + \gamma_2(t), \alpha_2(\gamma_1(t) - 1)),$$

where $\gamma_1'(t) = \delta_1(t)$, $\gamma_1(t_0) = 1$, $\gamma_2(t_0) = 0$ and $\gamma_2'(t) = \delta_1(t)\delta_2(t)$. After applying an affine transformation we have shown:

Theorem 4. *Let M^3 be an indefinite proper affine hypersphere of \mathbb{R}^4 which admits a pointwise \mathbb{Z}_3 - or $SO(2)$ -symmetry. Let $H = a_{22}^2 - a_4^2 (\neq 0)$ on M^3 . Then M^3 is affine equivalent with*

$$\varphi : I \times N^2 \rightarrow \mathbb{R}^4 : (t, v, w) \mapsto (\gamma_1(t)v, \gamma_1(t)w, \gamma_1(t)f(v, w) + \gamma_2(t), \gamma_1(t)),$$

where $\psi : N^2 \rightarrow \mathbb{R}^3 : (v, w) \mapsto (v, w, f(v, w))$ is a positive definite improper affine sphere with affine normal $(0, 0, 1)$ and $\gamma : I \rightarrow \mathbb{R}^2$ is a curve. Moreover, if M^3 admits a pointwise $SO(2)$ -symmetry then N^2 is an elliptic paraboloid.

We want to investigate the conditions imposed on the curve γ . For this we compute the derivatives of φ :

$$\begin{aligned} \varphi_t &= (\gamma_1'v, \gamma_1'w, \gamma_1'f(v, w) + \gamma_2', \gamma_1'), \\ \varphi_v &= (\gamma_1, 0, \gamma_1f_v, 0), \quad \varphi_w = (0, \gamma_1, \gamma_1f_w, 0), \\ \varphi_{tt} &= (\gamma_1''v, \gamma_1''w, \gamma_1''f(v, w) + \gamma_2'', \gamma_1''), \\ \varphi_{tv} &= \frac{\gamma_1'}{\gamma_1}\varphi_v, \quad \varphi_{tw} = \frac{\gamma_1'}{\gamma_1}\varphi_w, \\ \varphi_{vv} &= (0, 0, f_{vv}\gamma_1, 0), \quad \varphi_{vw} = (0, 0, \gamma_1f_{vw}, 0), \quad \varphi_{ww} = (0, 0, \gamma_1f_{ww}, 0). \end{aligned} \tag{60}$$

M^3 is a proper hypersphere, i.e. $\xi = -H\varphi$. An easy computation shows that the condition that ξ is a transversal vector field, namely $0 \neq \det(\varphi_t, \varphi_v, \varphi_w, \xi) = -H\gamma_1^2(\gamma_1\gamma_2' - \gamma_1'\gamma_2)$, is equivalent to $\gamma_1 \neq 0$ and $\gamma_1\gamma_2' - \gamma_1'\gamma_2 \neq 0$. Since $(0, 0, 1, 0) = \frac{\gamma_1}{\gamma_1\gamma_2' - \gamma_1'\gamma_2}\varphi_t - \frac{\gamma_1'}{\gamma_1\gamma_2' - \gamma_1'\gamma_2}\varphi$, we have the following structure equations:

$$\begin{aligned} \varphi_{tt} &= \left(\frac{\gamma_1''}{\gamma_1} + \frac{\gamma_1'\gamma_2'' - \gamma_1''\gamma_2'}{\gamma_1} \frac{\gamma_1}{\gamma_1\gamma_2' - \gamma_1'\gamma_2} \right) \varphi_t + \frac{\gamma_1'\gamma_2'' - \gamma_1''\gamma_2'}{\gamma_1\gamma_2' - \gamma_1'\gamma_2} \frac{1}{H} \xi, \\ \varphi_{tr} &= \frac{\gamma_1'}{\gamma_1} \varphi_r, \quad \varphi_{rs} = \frac{\gamma_1^2}{\gamma_1\gamma_2' - \gamma_1'\gamma_2} f_{rs} \varphi_t + \frac{\gamma_1\gamma_1'}{\gamma_1\gamma_2' - \gamma_1'\gamma_2} f_{rs} \frac{1}{H} \xi. \end{aligned} \tag{61}$$

We obtain:

$$\det h = h_{tt}(h_{vv}h_{ww} - h_{vw}^2) = \frac{\gamma_1'\gamma_2'' - \gamma_1''\gamma_2'}{H^3(\gamma_1\gamma_2' - \gamma_1'\gamma_2)^3} \gamma_1^2(\gamma_1')^2(f_{vv}f_{ww} - f_{vw}^2).$$

Since ψ is a positive definite improper affine sphere with affine normal $(0, 0, 1)$, we get by (4) that $f_{vv}f_{ww} - f_{vw}^2 = 1$. Now (4) (for ξ) is equivalent to

$$\gamma_1^4(\gamma_1\gamma_2' - \gamma_1'\gamma_2)^2 = \left| \frac{\gamma_1'\gamma_2'' - \gamma_1''\gamma_2'}{(\gamma_1\gamma_2' - \gamma_1'\gamma_2)^3} \right| \gamma_1^2(\gamma_1')^2.$$

It follows that

$$\xi = -H\varphi \iff \gamma_1^2|\gamma_1\gamma_2' - \gamma_1'\gamma_2|^5 = |\gamma_1'\gamma_2'' - \gamma_1''\gamma_2'|(\gamma_1')^2 \neq 0.$$

From the computations above also it follows that φ is indefinite iff either

$$\begin{aligned} \text{sign}(\gamma_1'\gamma_2'' - \gamma_1''\gamma_2') &= \text{sign}(H(\gamma_1\gamma_2' - \gamma_1'\gamma_2)) = -\text{sign}(\gamma_1\gamma_1') \quad \text{or} \\ \text{sign}(\gamma_1'\gamma_2'' - \gamma_1''\gamma_2') &= -\text{sign}(H(\gamma_1\gamma_2' - \gamma_1'\gamma_2)) = -\text{sign}(\gamma_1\gamma_1'). \end{aligned}$$

So we have seen under which conditions we can construct a 3-dimensional indefinite affine hypersphere out of an affine sphere:

Theorem 5. *Let $\psi : N^2 \rightarrow \mathbb{R}^3 : (v, w) \mapsto (v, w, f(v, w))$ be a positive definite improper affine sphere with affine normal $(0, 0, 1)$, and let $\gamma : I \rightarrow \mathbb{R}^2$ be a curve. Define $\varphi : I \times N^2 \rightarrow \mathbb{R}^4$ by $\varphi(t, v, w) = (\gamma_1(t)v, \gamma_1(t)w, \gamma_1(t)f(v, w) + \gamma_2(t), \gamma_1(t))$. If $\gamma = (\gamma_1, \gamma_2)$ satisfies $\gamma_1^2|\gamma_1\gamma_2' - \gamma_1'\gamma_2|^5 = -\text{sign}(\gamma_1\gamma_1')(\gamma_1'\gamma_2'' - \gamma_1''\gamma_2')(\gamma_1')^2 \neq 0$, then φ defines a 3-dimensional indefinite proper affine hypersphere.*

Now we are ready to check the symmetries.

Theorem 6. *Let $\psi : N^2 \rightarrow \mathbb{R}^3 : (v, w) \mapsto (v, w, f(v, w))$ be a positive definite improper affine sphere with affine normal $(0, 0, 1)$, and let $\gamma : I \rightarrow \mathbb{R}^2$ be a curve such that $\varphi(t, v, w) = (\gamma_1(t)v, \gamma_1(t)w, \gamma_1(t)f(v, w) + \gamma_2(t), \gamma_1(t))$ defines a 3-dimensional indefinite proper affine hypersphere. Then $\varphi(N^2 \times I)$ admits a pointwise \mathbb{Z}_3 - or $SO(2)$ -symmetry.*

Proof. We already have shown that φ defines a 3-dimensional indefinite proper affine hypersphere with affine normal $\xi = -H\varphi$. To prove the symmetry we need to compute K . We get the induced connection and the affine metric from the structure equations (61). We compute K using (6) and obtain:

$$\begin{aligned} (\nabla_{\varphi_t} h)(\varphi_r, \varphi_s) &= \left(\frac{\partial}{\partial t} \ln \left(\frac{\gamma_1 \gamma_1'}{\gamma_1 \gamma_2' - \gamma_1' \gamma_2} \right) - 2 \frac{\gamma_1'}{\gamma_1} \right) h(\varphi_r, \varphi_s), \\ (\nabla_{\varphi_r} h)(\varphi_t, \varphi_t) &= 0, \end{aligned}$$

implying that K_{φ_t} restricted to the space spanned by φ_v and φ_w is a multiple of the identity. Taking T in direction of φ_t , we see that φ_v and φ_w are orthogonal to T . Thus we can construct an ONB $\{T, V, W\}$ with V, W spanning $\text{span}\{\varphi_v, \varphi_w\}$ such that $a_1 = 2a_4, a_2 = a_3 = a_5 = 0$. By the considerations in [13, Section 4] we see that φ admits a pointwise \mathbb{Z}_3 - or $SO(2)$ -symmetry. ■

5.3 The third case: $\nu \equiv 0$ and $H = 0$ on M^3

The final cases now are that $\nu \equiv 0$ and $H = 0$ on the whole of M^3 and hence $a_{22} = \pm a_4$.

First we consider the case that $a_{22} = a_4 =: a > 0$. Again we use that M^3 admits a warped product structure and we fix a parameter t_0 . At the point t_0 , we have by (37)–(41):

$$\begin{aligned} D_V V &= +a_6 V - b_{23} W + \xi, \\ D_V W &= +b_{23} V - a_6 W, \\ D_W V &= -(b_{33} + a_6) W, \\ D_W W &= +(b_{33} - a_6) V + \xi, \\ D_X \xi &= 0. \end{aligned}$$

Thus, if v and w are local coordinates which span the second distribution L_2 , then we can interpret $\varphi(t_0, v, w)$ as a positive definite improper affine sphere in a 3-dimensional linear subspace.

Moreover, we see that this improper affine sphere is a paraboloid provided that $a_6(t_0, v, w)$ vanishes identically. From the differential equations (43) determining a_6 , we see that this is the case exactly when a_6 vanishes identically, i.e. when M^3 admits a pointwise $SO(2)$ -symmetry.

After applying an affine transformation and a change of coordinates, we may assume that

$$\varphi(t_0, v, w) = (v, w, f(v, w), 0), \tag{62}$$

with affine normal $\xi(t_0, v, w) = (0, 0, 1, 0)$, actually

$$\xi(t, v, w) = (0, 0, 1, 0)$$

(ξ is constant on M^3 by assumption). Furthermore we obtain by (35) and (36), that $D_U T = 2aU$ for all $U \in L_2$. We define $\delta := T - 2a\varphi$, which is transversal to $\text{span}\{V, W, \xi\}$. Since a is independent of v and w (cf. (42)), $D_U \delta = 0$, and we can assume that

$$T(t_0, v, w) - 2a(t_0)\varphi(t_0, v, w) = (0, 0, 0, 1). \tag{63}$$

We can integrate (42) ($T(a) = -4a^2$) and we take $a = \frac{1}{4t}$, $t > 0$. Thus (32) becomes $D_T T = -\frac{1}{2t}T - \xi$ and we obtain the following linear second order ordinary differential equation:

$$\frac{\partial^2}{\partial t^2}\varphi + \frac{1}{2t}\frac{\partial}{\partial t}\varphi = -\xi. \quad (64)$$

The general solution is $\varphi(t, v, w) = -\frac{t^2}{3}\xi + 2\sqrt{t}A(v, w) + B(v, w)$. The initial conditions (62) and (63) imply that $A(v, w) = (\frac{v}{2\sqrt{t_0}}, \frac{w}{2\sqrt{t_0}}, \frac{f(v, w)}{2\sqrt{t_0}} + \frac{2}{3}t_0^{3/2}, \sqrt{t_0})$ and $B(v, w) = (0, 0, -t_0^2, -2t_0)$. Obviously we can translate B to zero. Furthermore we can translate the affine sphere and apply an affine transformation to obtain $A(v, w) = \frac{1}{2\sqrt{t_0}}(v, w, f(v, w), 1)$. After a change of coordinates we get:

$$\varphi(t, v, w) = (tv, tw, tf(v, w) - ct^4, t), \quad c, t > 0. \quad (65)$$

Next we consider the case that $-a_{22} = a_4 =: a > 0$. Again we use that M^3 admits a warped product structure and we fix a parameter t_0 . A look at (37)–(41) suggests to define $\tilde{\xi} = -2aT + \xi$, then we get at the point t_0 :

$$\begin{aligned} D_V V &= +a_6 V - b_{23} W + \tilde{\xi}, \\ D_V W &= +b_{23} V - a_6 W, \\ D_W V &= -(b_{33} + a_6) W, \\ D_W W &= +(b_{33} - a_6) V + \tilde{\xi}, \\ D_U \tilde{\xi} &= 0. \end{aligned}$$

Thus, if v and w are local coordinates which span the second distribution L_2 , then we can interpret $\varphi(t_0, v, w)$ as a positive definite improper affine sphere in a 3-dimensional linear subspace.

Moreover, we see that this improper affine sphere is a paraboloid provided that $a_6(t_0, v, w)$ vanishes identically. From the differential equations (43) determining a_6 , we see that this is the case exactly when a_6 vanishes identically, i.e. when M^3 admits a pointwise $SO(2)$ -symmetry.

After applying an affine transformation and a change of coordinates, we may assume that

$$\varphi(t_0, v, w) = (v, w, f(v, w), 0), \quad (66)$$

with affine normal

$$\tilde{\xi}(t_0, v, w) = (0, 0, 1, 0). \quad (67)$$

We have considered $\tilde{\xi}$ before. We can solve (48) ($\frac{\partial}{\partial t}\beta = -2a\beta$) explicitly by $\beta = c\frac{1}{\sqrt{a}}$ (cf. (42)) and get by (49)–(51) that $D_X(\frac{1}{\sqrt{a}}\tilde{\xi}) = 0$. Thus $\frac{1}{\sqrt{a}}(-2aT + \xi) =: C$ for a constant vector C , i.e. $T = -\frac{1}{2a}(\sqrt{a}C - \xi)$. Notice that by (41) ξ is a constant vector, too. We can choose $a = \frac{1}{4|t|}$, $t < 0$ (cf. (42)), and we obtain the ordinary differential equation:

$$\frac{\partial}{\partial t}\varphi = -\sqrt{|t|}C - 2t\xi, \quad t < 0. \quad (68)$$

The solution (after a translation) with respect to the initial condition (66) is $\varphi(t, v, w) = \frac{2}{3}|t|^{\frac{3}{2}}C - t^2\xi + (v, w, f(v, w), 0)$. Notice that C is a multiple of $\tilde{\xi}$ and hence by (67) a constant multiple of $(0, 0, 1, 0)$. Furthermore ξ is transversal to the space spanned by $\varphi(t_0, v, w)$. So we get after an affine transformation and a change of coordinates:

$$\varphi(t, v, w) = (v, w, f(v, w) + ct^3, t^4), \quad c, t > 0. \quad (69)$$

Combining both results (65) and (69) we have:

Theorem 7. *Let M^3 be an indefinite improper affine hypersphere of \mathbb{R}^4 which admits a pointwise \mathbb{Z}_3 - or $SO(2)$ -symmetry. Let $a_{22}^2 = a_4^2$ on M^3 . Then M^3 is affine equivalent with either*

$$\begin{aligned} \varphi : I \times N^2 \rightarrow \mathbb{R}^4 : (t, v, w) &\mapsto (tv, tw, tf(v, w) - ct^4, t), & (a_{22} = a_4) & \quad \text{or} \\ \varphi : I \times N^2 \rightarrow \mathbb{R}^4 : (t, v, w) &\mapsto (v, w, f(v, w) + ct^3, t^4), & (-a_{22} = a_4), & \end{aligned}$$

where $\psi : N^2 \rightarrow \mathbb{R}^3 : (v, w) \mapsto (v, w, f(v, w))$ is a positive definite improper affine sphere with affine normal $(0, 0, 1)$ and $c, t \in \mathbb{R}^+$. Moreover, if M^3 admits a pointwise $SO(2)$ -symmetry then N^2 is an elliptic paraboloid.

The computations for the converse statement can be done completely analogous to the previous cases, they even are simpler (the curve is given parametrized).

Theorem 8. *Let $\psi : N^2 \rightarrow \mathbb{R}^3 : (v, w) \mapsto (v, w, f(v, w))$ be a positive definite improper affine sphere with affine normal $(0, 0, 1)$. Define $\varphi(t, v, w) = (tv, tw, tf(v, w) - ct^4, t)$ or $\varphi(t, v, w) = (v, w, f(v, w) + ct^3, t^4)$, where $c, t \in \mathbb{R}^+$. Then φ defines a 3-dimensional indefinite improper affine hypersphere, which admits a pointwise \mathbb{Z}_3 - or $SO(2)$ -symmetry.*

6 Pointwise $SO(1, 1)$ -symmetry

Let M^3 be a hypersphere admitting a $SO(1, 1)$ -symmetry. We only state the classification results. The proofs are done quite similar, using a lightvector-frame instead of an orthonormal one, and will appear elsewhere. We denote a lightvector-frame by $\{E, V, F\}$, where E and F are lightvectors and V is spacelike (cf. [13]).

Lemma 13. *Let M^3 be an affine hypersphere in \mathbb{R}^4 which admits a pointwise $SO(1, 1)$ -symmetry. Let $p \in M$. Then there exists a lightvector-frame $\{E, V, F\}$ defined in a neighborhood of the point p and a positive function b_4 such that K is given by:*

$$\begin{aligned} K(V, V) &= -2b_4V, & K(V, E) &= b_4E, & K(V, F) &= b_4F, \\ K(E, E) &= 0, & K(E, F) &= b_4V, & K(F, F) &= 0. \end{aligned}$$

In the following we denote the coefficients of the Levi-Civita connection with respect to this frame by:

$$\begin{aligned} \widehat{\nabla}_E E &= a_{11}E + b_{11}V, & \widehat{\nabla}_E V &= a_{12}E - b_{11}F, & \widehat{\nabla}_E F &= -a_{12}V - a_{11}F, \\ \widehat{\nabla}_V E &= a_{21}E + b_{21}V, & \widehat{\nabla}_V V &= a_{22}E - b_{21}F, & \widehat{\nabla}_V F &= -a_{22}V - a_{21}F, \\ \widehat{\nabla}_F E &= a_{31}E + b_{31}V, & \widehat{\nabla}_F V &= a_{32}E - b_{31}F, & \widehat{\nabla}_F F &= -a_{32}V - a_{31}F. \end{aligned}$$

Similar as before, it turns out that the vector field V is globally defined, and we can define the distributions $L_1 = \text{span}\{V\}$ and $L_2 = \text{span}\{E, F\}$. Again, L_1 is autoparallel with respect to $\widehat{\nabla}$, and L_2 is spherical with mean curvature normal $-a_{12}V$. We introduce a coordinate function v by $\frac{\partial}{\partial v} := V$.

Lemma 14. *The function $\nu = b_4^2 - a_{12}^2 - H$ is globally defined, $\frac{\partial}{\partial v}(e^{2f}\nu) = 0$ and ν vanishes identically or nowhere on \mathbb{R} .*

Again we have to distinguish three cases.

6.1 The first case: $\nu \neq 0$ on M^3

Theorem 9. *Let M^3 be an indefinite affine hypersphere of \mathbb{R}^4 which admits a pointwise $SO(1, 1)$ -symmetry. Let $b_4^2 - a_{12}^2 \neq H$ for some $p \in M^3$. Then M^3 is affine equivalent to*

$$\varphi : I \times N^2 \rightarrow \mathbb{R}^4 : (v, x, y) \mapsto (\gamma_1(v), \gamma_2(v)\phi(x, y)),$$

where $\phi : N^2 \rightarrow \mathbb{R}^3$ is a one-sheeted hyperboloid and $\gamma : I \rightarrow \mathbb{R}^2$ is a curve.

Theorem 10. *Let $\phi : N^2 \rightarrow \mathbb{R}^3$ be a one-sheeted hyperboloid and let $\gamma : I \rightarrow \mathbb{R}^2$ be a curve. Define $\varphi : I \times N^2 \rightarrow \mathbb{R}^4$ by $\varphi(v, x, y) = (\gamma_1(v), \gamma_2(v)\phi(x, y))$.*

- (i) *If $\gamma = (\gamma_1, \gamma_2)$ satisfies $\gamma_2^2|\gamma_1\gamma_2' - \gamma_1'\gamma_2|^5 = |\gamma_1'\gamma_2'' - \gamma_1''\gamma_2'|(\gamma_1')^2 \neq 0$, then φ defines a 3-dimensional indefinite proper affine hypersphere.*
- (ii) *If $\gamma = (\gamma_1, \gamma_2)$ satisfies $\gamma_2^2|\gamma_2'|^5 = |\gamma_1'\gamma_2'' - \gamma_1''\gamma_2'|(\gamma_1')^2 \neq 0$, then φ defines a 3-dimensional indefinite improper affine hypersphere.*

Theorem 11. *Let $\phi : N^2 \rightarrow \mathbb{R}^3$ be a one-sheeted hyperboloid and let $\gamma : I \rightarrow \mathbb{R}^2$ be a curve, such that $\varphi(v, x, y) = (\gamma_1(v), \gamma_2(v)\phi(x, y))$ defines a 3-dimensional indefinite affine hypersphere. Then $\varphi(I \times N^2)$ admits a pointwise $SO(1, 1)$ -symmetry.*

6.2 The second case: $\nu \equiv 0$ and $H \neq 0$ on M^3

Theorem 12. *Let M^3 be an indefinite proper affine hypersphere of \mathbb{R}^4 which admits a pointwise $SO(1, 1)$ -symmetry. Let $H = b_4^2 - a_{12}^2 (\neq 0)$ on M^3 . Then M^3 is affine equivalent with*

$$\varphi : I \times N^2 \rightarrow \mathbb{R}^4 : (v, x, y) \mapsto (\gamma_1(v)x, \gamma_1(v)y, \gamma_1(v)f(x, y) + \gamma_2(v), \gamma_1(v)),$$

where $\psi : N^2 \rightarrow \mathbb{R}^3 : (x, y) \mapsto (x, y, f(x, y))$ is a hyperbolic paraboloid with affine normal $(0, 0, 1)$ and $\gamma : I \rightarrow \mathbb{R}^2$ is a curve.

Theorem 13. *Let $\psi : N^2 \rightarrow \mathbb{R}^3$ be a hyperbolic paraboloid with affine normal $(0, 0, 1)$, and let $\gamma : I \rightarrow \mathbb{R}^2$ be a curve. Define $\varphi : I \times N^2 \rightarrow \mathbb{R}^4$ by $\varphi(v, x, y) = (\gamma_1(v)x, \gamma_1(v)y, \gamma_1(v)f(x, y) + \gamma_2(v), \gamma_1(v))$. If $\gamma = (\gamma_1, \gamma_2)$ satisfies $\gamma_1^2|\gamma_1\gamma_2' - \gamma_1'\gamma_2|^5 = |\gamma_1'\gamma_2'' - \gamma_1''\gamma_2'|(\gamma_1')^2 \neq 0$, then φ defines a 3-dimensional indefinite proper affine hypersphere.*

Theorem 14. *Let $\psi : N^2 \rightarrow \mathbb{R}^3$ be a hyperbolic paraboloid with affine normal $(0, 0, 1)$, and let $\gamma : I \rightarrow \mathbb{R}^2$ be a curve, such that $\varphi(v, x, y) = (\gamma_1(v)x, \gamma_1(v)y, \gamma_1(v)f(x, y) + \gamma_2(v), \gamma_1(v))$ defines a 3-dimensional indefinite proper affine hypersphere. Then $\varphi(I \times N^2)$ admits a pointwise $SO(1, 1)$ -symmetry.*

6.3 The third case: $\nu \equiv 0$ and $H = 0$ on M^3

Theorem 15. *Let M^3 be an indefinite improper affine hypersphere of \mathbb{R}^4 which admits a pointwise $SO(1, 1)$ -symmetry. Let $a_{12}^2 = b_4^2$ on M^3 . Then M^3 is affine equivalent with either*

$$\begin{aligned} \varphi : I \times N^2 \rightarrow \mathbb{R}^4 : (v, x, y) &\mapsto (vx, vy, vf(x, y) - cv^4, v), & (a_{12} = b_4) & \quad \text{or} \\ \varphi : I \times N^2 \rightarrow \mathbb{R}^4 : (v, x, y) &\mapsto (x, y, f(x, y) + cv^3, v^4), & (-a_{12} = b_4), & \end{aligned}$$

where $\psi : N^2 \rightarrow \mathbb{R}^3 : (v, w) \mapsto (v, w, f(v, w))$ is a hyperbolic paraboloid with affine normal $(0, 0, 1)$ and $c, t \in \mathbb{R}^+$.

Theorem 16. *Let $\psi : N^2 \rightarrow \mathbb{R}^3 : (v, w) \mapsto (v, w, f(v, w))$ be a hyperbolic paraboloid with affine normal $(0, 0, 1)$. Define $\varphi(t, v, w) = (tv, tw, tf(v, w) - ct^4, t)$ or $\varphi(t, v, w) = (v, w, f(v, w) + ct^3, t^4)$, where $t \in \mathbb{R}^+$, $c \neq 0$. Then φ defines a 3-dimensional indefinite improper affine hypersphere, which admits a pointwise $SO(1, 1)$ -symmetry.*

A Computations for pointwise $SO(2)$ -, S_3 - or \mathbb{Z}_3 -symmetry

$$e[1]:=\{1, 0, 0\}; e[2]:=\{0, 1, 0\}; e[3]:=\{0, 0, 1\};$$

$$\text{ONB of } SO(1,2): e[1]=\mathbf{T}, e[2]=\mathbf{V}, e[3]=\mathbf{W}$$

Affine metric

$$h[y_{-}, z_{-}] := -y[[1]]z[[1]] + \text{Sum}[y[[i]]z[[i]], \{i, 2, 3\}];$$

Difference tensor ($r=\mathbf{a1}$, $s=\mathbf{a4}$, $u=\mathbf{a6}$)

$$\begin{aligned} K[y_{-}, z_{-}] &:= \text{Sum}[y[[i]]z[[j]]k[i, j], \{i, 1, 3\}, \{j, 1, 3\}]; \\ k[1, 1] &:= \{-r, 0, 0\}; k[1, 2] := \{0, s, 0\}; k[1, 3] := \{0, 0, r-s\}; \\ k[2, 1] &:= \{0, s, 0\}; k[2, 2] := \{-s, u, 0\}; k[2, 3] := \{0, 0, -u\}; \\ k[3, 1] &:= \{0, 0, r-s\}; k[3, 2] := \{0, 0, -u\}; k[3, 3] := \{-(r-s), -u, 0\}; \end{aligned}$$

Ricci tensor, scalar curvature and Pick invariant

$$[\mathbf{Kx}, \mathbf{Ky}]z$$

$$\begin{aligned} LK[x_{-}, y_{-}, z_{-}] &:= K[x, K[y, z]] - K[y, K[x, z]]; \\ \text{ListLK} &:= \{LK[e[1], e[2], e[1]], LK[e[1], e[2], e[2]], LK[e[1], e[2], e[3]], \\ &LK[e[1], e[3], e[1]], LK[e[1], e[3], e[2]], LK[e[1], e[3], e[3]], \\ &LK[e[2], e[3], e[1]], LK[e[2], e[3], e[2]], LK[e[2], e[3], e[3]]\}; \\ \text{FullSimplify} &[\text{ListLK}] \end{aligned}$$

Curvature tensor (of the Levi-Civita connection)

$$\begin{aligned} R[x_{-}, y_{-}, z_{-}] &:= H(h[y, z]x - h[x, z]y) - K[x, K[y, z]] + K[y, K[x, z]]; \\ \text{ListR} &:= \{R[e[1], e[2], e[1]], R[e[1], e[2], e[2]], R[e[1], e[2], e[3]], \\ &R[e[1], e[3], e[1]], R[e[1], e[3], e[2]], R[e[1], e[3], e[3]], \\ &R[e[2], e[3], e[1]], R[e[2], e[3], e[2]], R[e[2], e[3], e[3]]\}; \\ \text{Simplify} &[\text{ListR}] \end{aligned}$$

Ricci tensor (of the Levi-Civita connection)

$$\begin{aligned} \text{ric}[x_{-}, y_{-}] &:= \text{Simplify}[(-h[R[e[1], x, y], e[1]] + h[R[e[2], x, y], e[2]] + h[R[e[3], x, y], e[3]])]; \\ \text{Listric} &:= \{\text{ric}[e[1], e[1]], \text{ric}[e[1], e[2]], \text{ric}[e[1], e[3]], \text{ric}[e[2], e[2]], \text{ric}[e[2], e[3]], \text{ric}[e[3], e[3]]\}; \\ \text{Simplify} &[\text{Listric}] \end{aligned}$$

Scalar curvature (of the Levi-Civita connection)

$$\begin{aligned} \text{sc} &:= 1/6(-\text{ric}[e[1], e[1]] + \text{ric}[e[2], e[2]] + \text{ric}[e[3], e[3]]); \\ \text{Simplify} &[\text{sc}] \end{aligned}$$

Pick invariant

$$\begin{aligned} P &:= 1/6(- (k[1, 1][[1]])^2 + (k[2, 2][[2]])^2 + (k[3, 3][[3]])^2 + 3((k[1, 1][[2]])^2 + (k[1, 1][[3]])^2 - \\ &(k[2, 2][[1]])^2 - (k[3, 3][[1]])^2 + (k[2, 2][[3]])^2 + (k[3, 3][[2]])^2) - 6(k[1, 2][[3]])^2); \\ \text{Simplify} &[P] \end{aligned}$$

Lemma 1

$r = 2s$; Simplify[Listric]
Simplify[P]

Lemma 8

$R[e[2], e[3], e[2]]$

Lemma 3**Levi-Civita connection (ONB)**

$n[1, 1] := \{0, a12, a13\}$; $n[1, 2] := \{a12, 0, -b13\}$; $n[1, 3] := \{a13, b13, 0\}$;
 $n[2, 1] := \{0, a22, a23\}$; $n[2, 2] := \{a22, 0, -b23\}$; $n[2, 3] := \{a23, b23, 0\}$;
 $n[3, 1] := \{0, a32, a33\}$; $n[3, 2] := \{a32, 0, -b33\}$; $n[3, 3] := \{a33, b33, 0\}$;
 $\text{Na}[y_-, z_-] := \text{Sum}[y[[i]]z[[j]]n[i, j], \{i, 1, 3\}, \{j, 1, 3\}] + \text{Sum}[y[[1]]\text{Dt}[z[[i]], f1]e[i]$
 $+ y[[2]]\text{Dt}[z[[i]], f2]e[i] + y[[3]]\text{Dt}[z[[i]], f3]e[i], \{i, 1, 3\}]$;

Codazzi for K (affine hypersphere)

$\text{codazzi}[x_-, y_-, z_-] := \text{Na}[x, K[y, z]] - K[\text{Na}[x, y], z] - K[y, \text{Na}[x, z]] - \text{Na}[y, K[x, z]] + K[\text{Na}[y, x], z] +$
 $K[x, \text{Na}[y, z]]$;

eq1:=Simplify[codazzi[e[2], e[1], e[1]]]; eq2:=Simplify[codazzi[e[3], e[1], e[1]]];
eq3:=Simplify[codazzi[e[1], e[2], e[2]]]; eq4:=Simplify[codazzi[e[3], e[2], e[2]]];
eq5:=Simplify[codazzi[e[1], e[3], e[3]]]; eq6:=Simplify[codazzi[e[2], e[3], e[3]]];
eq7:=Simplify[codazzi[e[1], e[2], e[3]]]; eq8:=Simplify[codazzi[e[2], e[3], e[1]]];
eq9:=Simplify[codazzi[e[3], e[1], e[2]]];
eq:={eq1, eq2, eq3, eq4, eq5, eq6, eq7, eq8, eq9}; eq

1. case: $u^2 \neq 4s^2$ **conclusions from eq1,2,4:**

Simplify[eq2[[1]] - 2eq4[[1]]]
Simplify[eq1[[3]] + eq2[[2]]]
 $a13 = 0$; $a32 = -a23$;

eq

conclusions from eq1,2,6:

Simplify[-2eq6[[1]] + eq1[[1]]]
Simplify[eq1[[2]] - eq2[[3]]]
 $a12 = 0$; $a33 = a22$;

eq

Clear[a13, a12, a32, a33]

2. case: $u=2s \neq 0$

$u = 2s$; eq

conclusions from eq8, eq1:

$a32 = a23$; $a13 = -2a23$; eq

conclusions from eq3:

$b13 = 0$; eq

Simplify[eq1[[2]] - eq2[[3]]]

$a12 = -(a33 - a22)$; eq

Simplify[eq3[[2]] - 1/2eq1[[1]] + 2eq1[[2]]]
a33 = -a22; eq

It follows that $T(a_4)=0$, $V(a_4)=-4a_2^2 a_4$, $W(a_4)=4a_2^3 a_4$.

Simplify[eq4[[2]] + eq2[[1]]]

Simplify[eq4[[3]] + eq1[[1]]]

b23 = -a23; b33 = -a22;

Lemma 4

Gauss for Levi-Civita connection (affine hypersphere)

gaussLC[x_, y_, z_.]:=Na[x, Na[y, z]] - Na[y, Na[x, z]] - Na[Na[x, y] - Na[y, x], z] - Hh[y, z]x + Hh[x, z]y + K[x, K[y, z]] - K[y, K[x, z]];

eq11:=Simplify[gaussLC[e[1], e[2], e[2]]]; eq12:=Simplify[gaussLC[e[1], e[3], e[2]]];

eq13:=Simplify[gaussLC[e[2], e[3], e[2]]]; eq14:=Simplify[gaussLC[e[1], e[2], e[1]]];

eq15:=Simplify[gaussLC[e[1], e[3], e[1]]]; eq16:=Simplify[gaussLC[e[2], e[3], e[1]]];

eq17:=Simplify[gaussLC[e[1], e[2], e[3]]]; eq18:=Simplify[gaussLC[e[1], e[3], e[3]]];

eq19:=Simplify[gaussLC[e[2], e[3], e[3]]];

eqG:={eq11, eq12, eq13, eq14, eq15, eq16, eq17, eq18, eq19};

2. case: $u=2s \neq 0$

eqG

Simplify[eq11[[1]] - eq12[[3]]]

Simplify[eq15[[3]] + eq12[[3]]]

Contradiction to eq13.3

Clear[b33, b23, a33, a12, b13, a32, a13, u]

1. case: $u^2 \neq 4s^2$

a13 = 0; a32 = -a23; a12 = 0; a33 = a22; eqG

Acknowledgements

Partially supported by the DFG-Project PI 158/4-5 ‘Geometric Problems and Special PDEs’.

References

- [1] Bryant R.L., Second order families of special Lagrangian 3-folds, in *Perspectives in Riemannian Geometry, CRM Proc. Lecture Notes*, Vol. 40, Amer. Math. Soc., Providence, RI, 2006, 63–98, [math.DG/0007128](https://arxiv.org/abs/math/0607128).
- [2] Calabi E., Complete affine hyperspheres. I, in *Convegno di Geometria Differenziale (INDAM, Rome, 1971), Symposia Mathematica*, Vol. X, Academic Press, London, 1972, 19–38.
- [3] Dillen F., Equivalence theorems in affine differential geometry, *Geom. Dedicata* **32** (1989), 81–92.
- [4] Dillen F., Katsumi N., Vrancken L., Conjugate connections and Radon’s theorem in affine differential geometry, *Monatsh. Math.* **109** (1990), 221–235.
- [5] Dillen F., Vrancken L., Calabi-type composition of affine spheres, *Differential Geom. Appl.* **4** (1994), 303–328.
- [6] Hu Z., Li H.Z., Vrancken L., Characterizations of the Calabi product of hyperbolic affine hyperspheres, *Results Math.* **52** (2008), 299–314.
- [7] Li A.M., Simon U., Zhao G.S., Global affine differential geometry of hypersurfaces, *de Gruyter Expositions in Mathematics*, Vol. 11, Walter de Gruyter & Co., Berlin, 1993.

-
- [8] Lu Y., Scharlach C., Affine hypersurfaces admitting a pointwise symmetry, *Results Math.* **48** (2005), 275–300, [math.DG/0510150](#).
 - [9] Nölker S., Isometric immersions of warped products, *Differential Geom. Appl.* **6** (1996), 1–30.
 - [10] Nomizu K., Sasaki T., Affine differential geometry. Geometry of affine immersions, *Cambridge Tracts in Mathematics*, Vol. 111, Cambridge University Press, Cambridge, 1994.
 - [11] O’Neill B., Semi-Riemannian geometry. With applications to relativity, *Pure and Applied Mathematics*, Vol. 103, Academic Press, Inc. [Harcourt Brace Jovanovich Publishers], New York, 1983.
 - [12] Ponge R., Reckziegel H., Twisted products in pseudo-Riemannian geometry, *Geom. Dedicata* **48** (1993), 15–25.
 - [13] Scharlach C., Indefinite affine hyperspheres admitting a pointwise symmetry. Part 1, *Beiträge Algebra Geom.* **48** (2007), 469–491, [math.DG/0510531](#).
 - [14] Vrancken L., Special classes of three dimensional affine hyperspheres characterized by properties of their cubic form, in *Contemporary Geometry and Related Topics*, World Sci. Publ., River Edge, NJ, 2004, 431–459.