



Homology stability for outer automorphism groups of free groups

ALLEN HATCHER
KAREN VOGTMANN

Abstract We prove that the quotient map from $Aut(F_n)$ to $Out(F_n)$ induces an isomorphism on homology in dimension i for n at least $2i + 4$. This corrects an earlier proof by the first author and significantly improves the stability range. In the course of the proof, we also prove homology stability for a sequence of groups which are natural analogs of mapping class groups of surfaces with punctures. In particular, this leads to a slight improvement on the known stability range for $Aut(F_n)$, showing that its i th homology is independent of n for n at least $2i + 2$.

AMS Classification 20F65; 20F28, 57M07

Keywords Automorphisms of free groups, homology stability

1 Introduction

In this paper we present a proof that the quotient map $Aut(F_n) \rightarrow Out(F_n)$ induces an isomorphism on the integral homology groups H_i for $n \geq 2i + 4$. We also show that the natural map $Aut(F_n) \rightarrow Aut(F_{n+1})$ induces an isomorphism on H_i for $n \geq 2i + 2$, a slight improvement on the result in [6]. It follows that $H_i(Out(F_n))$ is independent of n for $n \geq 2i + 4$.

These stable homology groups have a significant amount of non-trivial torsion since they contain the homology of $\Omega^\infty S^\infty$ as a direct summand, as observed in [4]. In contrast, it is conjectured that they are rationally trivial, though a few non-trivial unstable rational classes have been found [2, 3, 6]. One of these non-trivial classes, in $H_7(Aut(F_5); \mathbf{Q})$, becomes trivial under the map to $H_7(Out(F_5); \mathbf{Q})$, showing that this map is not always an isomorphism on homology [3].

The first proof of homology stability for $Aut(F_n)$ and $Out(F_n)$, with a much weaker dimension range than we obtain here, appeared in the first author's

1995 paper [4]. However, in 2003 a mistake was discovered in the spectral sequence arguments at the end of that paper (in the last sentence on p.56, where the diffeomorphism in question may permute boundary spheres and hence not induce an inner automorphism as claimed). In the meantime an independent proof for the isomorphisms $H_i(\text{Aut}(F_n)) \cong H_i(\text{Aut}(F_{n+1}))$, with a much better dimension range, had appeared in 1998 in [6]. This left the other isomorphisms $H_i(\text{Out}(F_n)) \cong H_i(\text{Aut}(F_n))$ unproved, and the present paper fills that gap, at the same time giving a better dimension range than the one originally claimed.

The groups $\text{Out}(F_n)$ and $\text{Aut}(F_n)$ are the cases $s = 0, 1$ of a sequence of groups $\Gamma_{n,s}$ which are analogs of mapping class groups of surfaces with s punctures. We will show that $H_i(\Gamma_{n,s})$ is independent of n and s for $n \geq 2i + 4$. A more general family of groups $\Gamma_{n,s}^k$ analogous to mapping class groups of surfaces with s punctures and k boundary components is studied in [8], where homological stability for $\Gamma_{n,s}^k$ is proved in the weaker range $n \geq 3i + 3$. In particular this gives a different proof of stability for both $\text{Aut}(F_n)$ and $\text{Out}(F_n)$ in this weaker range. The groups $\Gamma_{n,1}^k$ are the groups $A_{n,k}$ of [9], and they are used in [13] to study the relation between mapping class groups of surfaces and $\text{Out}(F_n)$.

The present paper relies heavily on the geometric results of [4] concerning the connectivity of various sphere complexes, and on the Degree Theorem of [6]. The spectral sequence arguments given here are new; they completely replace those in [4] and can also be used as an alternative to the spectral sequence arguments in [6].

Acknowledgment The second author was partially supported by NSF grant DMS 0204185.

2 Tools and Outline of proof

Let $M_{n,s}$ be the compact 3-manifold obtained from the connected sum of n copies of $S^1 \times S^2$ by deleting the interiors of s disjoint balls. When $n = 0$ we take $M_{0,s}$ to be S^3 with the interiors of s disjoint balls removed. The s boundary spheres of $M_{n,s}$ are denoted $\partial_0, \dots, \partial_{s-1}$, considering ∂_0 as the base sphere. Let $\text{Diff}(M_{n,s})$ be the group of orientation-preserving diffeomorphisms of $M_{n,s}$ which fix the boundary pointwise. The mapping class group $\pi_0 \text{Diff}(M_{n,s})$ has a normal subgroup generated by Dehn twist diffeomorphisms along embedded 2-spheres, and the quotient of $\pi_0 \text{Diff}(M_{n,s})$ by this subgroup will be denoted $\Gamma_{n,s}$. A theorem of Laudenbach [10] states that $\Gamma_{n,0}$ is isomorphic to $\text{Out}(F_n)$, and $\Gamma_{n,1}$ to $\text{Aut}(F_n)$.

An inclusion $M_{n,s} \hookrightarrow M_{m,t}$ induces a homomorphism $\Gamma_{n,s} \rightarrow \Gamma_{m,t}$ by extending diffeomorphisms by the identity map outside $M_{n,s}$. We will be interested in three special cases:

- (1) *Adding a punctured handle.* The map $\alpha: \Gamma_{n,s} \rightarrow \Gamma_{n+1,s}$, $s \geq 1$, is induced by gluing a copy of $M_{1,2}$ to the base sphere ∂_0 ; the puncture in $M_{1,2}$ which is not used in the gluing is the new base sphere.
- (2) *Adding a tube.* The map $\beta: \Gamma_{n,s} \rightarrow \Gamma_{n+1,s-2}$, $s \geq 2$, is induced by gluing the boundary components of a copy of $M_{0,2}$ to the last two boundary components of $M_{n,s}$.
- (3) *Adding pants.* The map $\mu: \Gamma_{n,s} \rightarrow \Gamma_{n,s+1}$, $s \geq 1$, is induced by gluing one boundary component of a copy of $M_{0,3}$ to ∂_0 . One of the unglued boundary components of $M_{0,3}$ is the new base sphere.




Figure 1: $\alpha = \beta \circ \mu^2$

We consider the following commutative diagram of maps (see Figure 1):

$$\begin{array}{ccc}
 \Gamma_{n,s} & \xrightarrow{\alpha} & \Gamma_{n+1,s} \\
 \downarrow \mu & & \uparrow \beta \\
 \Gamma_{n,s+1} & \xrightarrow{\mu} & \Gamma_{n,s+2}
 \end{array}$$

We first show that α is an isomorphism on H_i for $n \geq 2i + 2$. This generalizes and improves the theorem in [6] that $Aut(F_n) \rightarrow Aut(F_{n+1})$ is an isomorphism on H_i for $n \geq 2i + 3$. We prove this by working with the spectral sequence associated to the action of $\Gamma_{n,s}$ on a simplicial complex $W_{n,s}$. A k -simplex of $W_{n,s}$ is a collection of $k + 1$ spheres whose complement in $M_{n,s}$ is connected, together with an additional *enveloping* sphere which separates these spheres and the base sphere from the rest of the manifold.

It follows by a formal argument that β and μ induce isomorphisms on H_i for $n \geq 2i + 2$ when $s > 0$. However, we need to show that β induces isomorphisms

on homology when $s = 0$ as well, and for this we use a similar spectral sequence argument, replacing the complex $W_{n,s}$ by a slightly different complex of spheres $X_{n,s}$ that was considered in [4]. This argument requires $n \geq 2i + 3$.




Figure 2: $\gamma \circ \alpha = \beta \circ \mu$

Finally we consider the commutative diagram

$$\begin{array}{ccc} \Gamma_{n,1} & \xrightarrow{\alpha} & \Gamma_{n+1,1} \\ \downarrow \mu & & \downarrow \gamma \\ \Gamma_{n,2} & \xrightarrow{\beta} & \Gamma_{n+1,0} \end{array}$$

where γ is induced by filling in the boundary sphere with a ball (see Figure 2). Thus γ is the quotient map $Aut(F_{n+1}) \rightarrow Out(F_{n+1})$. Since α , β , and μ induce isomorphisms on H_i for $n \geq 2i + 3$, so does γ . Replacing $n + 1$ by n , this says that the map $Aut(F_n) \rightarrow Out(F_n)$ induces an isomorphism on H_i for $n \geq 2i + 4$.

The main input needed to make the spectral sequences work is the fact that the complexes $W_{n,s}$ and $X_{n,s}$ are highly connected, as are their quotients by the action of $\Gamma_{n,s}$. For $X_{n,s}$ and its quotient this was shown in [4]. The connectivity of $W_{n,s}$ will follow from an easy generalization of the main technical result of [6], the Degree Theorem. The quotient $W_{n,s}/\Gamma_{n,s}$ is combinatorially the same as $X_{n,s}/\Gamma_{n,s}$.

3 Sphere systems and graphs

We will show that the complex $W_{n,s}$ is highly connected by comparing it to a complex of metric graphs with s distinguished points, which are allowed to coincide. The first distinguished point is the basepoint v_0 , and the rest will

be labeled v_1, \dots, v_{s-1} . The metric is determined by the lengths of the edges, and is normalized so that the sum of these lengths is 1. It is convenient also to attach a free edge of some fixed positive length to each v_i , $i > 0$, and we label the free vertex of this edge by w_i . The resulting graph is called a *thorned graph*, with thorns the $s - 1$ free edges attached at v_1, \dots, v_{s-1} (see Figure 3).




Figure 3: A thorned graph with 3 thorns and the standard thorned rose $R_{3,4}$

The *standard thorned rose* $R_{n,s}$ is a bouquet of n circles with vertex x_0 together with $s - 1$ free edges attached at x_0 , with free vertices labeled y_1, \dots, y_{s-1} . Each (oriented) petal of the rose is identified with a generator of the free group F_n . There are natural inclusions $R_{n,s} \subseteq R_{m,t}$ for all $m \geq n$ and $t \geq s$.

By a *marking* of a thorned graph $(\Gamma, v_0, \dots, v_{s-1})$ we will mean a homotopy equivalence $g: R_{n,s} \rightarrow \Gamma$ which sends x_0 to v_0 and y_i to w_i for all i . Two marked graphs $(g, \Gamma, v_0, \dots, v_{s-1})$ and $(g', \Gamma', v'_0, \dots, v'_{s-1})$ are *equivalent* if there is an isometry $h: \Gamma \rightarrow \Gamma'$ sending v_i to v'_i for all i with $g' \simeq h \circ g$ by a homotopy which is constant on the v_i . The lengths of the edges which are not thorns give barycentric coordinates for an open simplex associated to the marked thorned graph $(g, \Gamma, v_0, \dots, v_{s-1})$. Codimension one faces of this simplex correspond to marked thorned graphs obtained by collapsing an edge of Γ which is not a loop or a thorn. The space obtained from the disjoint union of all of the open simplices associated to marked thorned graphs of rank n with $s - 1$ thorns by attaching them together according to the face relations is denoted $A_{n,s}$.

Recall from [4] the complex $S(M)$ of sphere systems in $M = M_{n,s}$. A vertex of this complex is an isotopy class of embedded spheres in M that do not bound a ball and are not isotopic to a sphere of ∂M . A set of $k + 1$ vertices of $S(M)$ spans a k -simplex if these vertices are represented by disjoint spheres, no two of which are isotopic. A point of $S(M)$ can be thought of as a weighted system of spheres, the weights being given by the barycentric coordinates of the point. A system S is said to be *simple* if each component of $M - S$ is simply-connected.

We let $SS(M)$ be the subspace of $S(M)$ formed by weighted simple systems with strictly positive weights. This is a union of open simplices of $S(M)$.

As in [4, 6] which dealt with the cases $s \leq 1$, there is a homeomorphism $\Phi: SS(M) \rightarrow A_{n,s}$, defined in the following way. Given a sphere system in $SS(M)$, thicken each sphere to $S^2 \times I$, and also thicken the boundary spheres other than ∂_0 . Map M to a graph by projecting each $S^2 \times I$ onto the interval I (an edge of the graph), and each component of the closure of M minus the thickened spheres to a point (a vertex of the graph). Thus the graph is a quotient space of M . Each boundary sphere except ∂_0 will correspond to a thorn of the graph. We assign lengths to the non-thorn edges according to the weight of the corresponding sphere. To get a marking, we fix a particular simple sphere system with connected complement. Its associated graph is the standard thorned rose $R_{n,s}$, and this can be embedded in M so that the embedding induces an isomorphism on π_1 , and so that the basepoint of the image is located in ∂_0 and the thorns run out to the other boundary spheres (see Figure 4). The marking of a sphere system S is given by this embedding followed by the quotient map from M to the graph associated to S .




Figure 4: $R_{3,3}$ embedded in $M_{3,3}$

It is shown in the Appendix of [4] that Φ is a homeomorphism when $s = 0$, and the same argument works for $s > 0$ once we give an alternative description of $\Gamma_{n,s}$ in terms of homotopy equivalences of graphs. Let $G_{n,s}$ be the group of path-components of the space of homotopy equivalences of the standard thorned rose $R = R_{n,s}$ that fix the s distinguished points. Such a homotopy equivalence is actually a homotopy equivalence relative to the s distinguished points, by Proposition 0.19 of [5]. This shows in particular that $G_{n,s}$ is a group.

There is a map $\varphi: \Gamma_{n,s} \rightarrow G_{n,s}$ sending a diffeomorphism $f: M \rightarrow M$ to the composition

$$R \xrightarrow{i} M \xrightarrow{f} M \xrightarrow{r} R$$

where i is the standard inclusion and r is a retraction onto $i(R)$ inducing an isomorphism on π_1 . Since Dehn twists along 2–spheres in M have no effect on the homotopy class of $f \circ i$, the map φ is well-defined. To check that φ is a homomorphism we need to see that a composition

$$R \xrightarrow{i} M \xrightarrow{f} M \xrightarrow{r} R \xrightarrow{i} M \xrightarrow{g} M \xrightarrow{r} R$$

can be homotoped so as to eliminate the middle two maps. It is not true that $i \circ r$ is homotopic to the identity, but the composition $f \circ i$ can be homotoped, fixing the distinguished points, to have image in $i(R)$, so that $i \circ r$ is the identity on this image. This is a consequence of the fact that $\pi_1(M, i(R)) = 0$ since i induces a surjection on π_1 .

Proposition 1 *The homomorphism $\varphi: \Gamma_{n,s} \rightarrow G_{n,s}$ is an isomorphism.*

Proof When $s \leq 1$ this is a theorem of Laudenbach [10]. For the general case we do induction on s . For convenience, let us replace $Diff(M_{n,s})$ by $Diff_{n,s}$, the orientation-preserving diffeomorphisms of $M_{n,0}$ that fix s distinguished points, so $\pi_0 Diff_{n,s}$ is $\pi_0 Diff(M_{n,s})$ with Dehn twists along the boundary spheres of $M_{n,s}$ factored out. By evaluating elements of $Diff_{n,s-1}$ at the s -th distinguished point we obtain a fibration $Diff_{n,s} \rightarrow Diff_{n,s-1} \rightarrow M'$ where M' is the complement of the first $s - 1$ distinguished points in $M_{n,0}$. The last four terms of the first row of the following commutative diagram are part of the long exact sequence of homotopy groups for this fibration:

$$\begin{array}{ccccccc} 1 & \longrightarrow & F_n & \longrightarrow & \pi_0 Diff_{n,s} & \longrightarrow & \pi_0 Diff_{n,s-1} \longrightarrow 1 \\ & & \parallel & & \downarrow \psi_s & & \downarrow \psi_{s-1} \\ 1 & \longrightarrow & F_n & \longrightarrow & G_{n,s} & \longrightarrow & G_{n,s-1} \longrightarrow 1 \end{array}$$

The second row is derived in a similar way. We identify $G_{n,s-1}$ with π_0 of the space of homotopy equivalences of $R_{n,s}$ which fix the basepoint and the first $s - 2$ thorn tips, but are allowed to do anything on the last thorn. Restricting such a homotopy equivalence to the endpoint of the last thorn gives a map to $R_{n,s}$, which is a fibration with fiber consisting of homotopy equivalences fixing all the distinguished points. The long exact sequence for this fibration gives the second row. The initial 1 in this row is π_1 of the total space of the fibration, which is zero since π_2 of a graph is trivial. The maps ψ_s and ψ_{s-1} are the compositions of the quotient maps to $\Gamma_{n,s}$ and $\Gamma_{n,s-1}$ with φ . The diagram then commutes, and this yields the initial 1 in the first row.

A standard diagram chase shows that the kernels of ψ_s and ψ_{s-1} are isomorphic. By induction we may assume that the kernel of ψ_{s-1} is the subgroup generated

by twists along 2–spheres, so it follows that the same is true for ψ_s . Thus if we replace each $\pi_0 \text{Diff}$ by Γ we still have a commutative diagram of short exact sequences. The five lemma then finishes the induction step. \square

It is shown in Theorem 2.1 of [4] that $S(M)$ is contractible if $n > 0$, and we claim that when this contraction is restricted to the subspace $SS(M)$, it gives a contraction of $SS(M)$. Verifying this claim is simply a matter of following through the proof for $S(M)$ and checking that if one starts in $SS(M)$ then one never leaves it, using the observation that surgery on a simple system always produces a simple system. As the proof takes two or three pages, we will not repeat it here. The proof for $SS(M)$ is actually somewhat simpler since Lemma 2.2 in the proof for $S(M)$ is not necessary for $SS(M)$, in view of the fact that surgery takes simple systems to simple systems.

When $n = 0$, the space $S(M) = SS(M)$ is not contractible, but is instead homotopy equivalent to a wedge of spheres of dimension $s - 4$. This fact will not be needed here, so we will not give a proof.

Since $A_{n,s}$ is homeomorphic to $SS(M)$, it is also contractible when $n > 0$.

4 The degree theorem with distinguished points

We will need a variation of the Degree Theorem of [6]. In that paper, the degree of a basepointed graph of rank n was defined to be $2n - |v_0|$, where $|v_0|$ is the valence of the basepoint v_0 . The Degree Theorem states that a k –parameter family of graphs in $A_{n,1}$ can be deformed into the subspace consisting of graphs of degree at most k by a deformation during which degree is non-increasing. Since $A_{n,1}$ is contractible, this implies that the subspace of degree k graphs is $(k - 1)$ –connected.

We define the degree of a thorned graph $(\Gamma, v_0, \dots, v_{s-1})$ to be $2n + s - 1 - |v_0|$. The thorns attached at the basepoint contribute to its valence, so that the standard thorned rose has degree 0.

Lemma 1 *If $k < n/2$, then a degree k thorned graph of rank n has at least one loop at the basepoint. (Here a loop at the basepoint has no thorns attached to its interior.)*

Proof The proof is an Euler characteristic argument identical to the case of graphs with no thorns (see [6], Lemma 5.2). Here is how it goes: Suppose Γ

is a thorned graph of rank n and degree k , with basepoint v_0 . By definition, $k = 2n + s - 1 - |v_0|$. Blowing up if necessary, we may assume that all vertices other than v_0 are trivalent or univalent. For such a graph the degree equals the number of vertices other than v_0 that are trivalent since collapsing an edge from v_0 to a trivalent vertex decreases degree by one, and this can be repeated until one has the standard thorned rose.

Let Γ_1 denote the subgraph of Γ spanned by all vertices other than the basepoint. If Γ has no loops at the basepoint, then $E(\Gamma) = E(\Gamma_1) + |v_0|$. We calculate the Euler characteristic of Γ using this fact and the observation that $V(\Gamma_1) = k + s - 1$:

$$\begin{aligned} 1 - n &= \chi(\Gamma) = V(\Gamma) - E(\Gamma) \\ &= (1 + V(\Gamma_1)) - (E(\Gamma_1) + |v_0|) \\ &= (1 + k + s - 1) - (E(\Gamma_1) + (2n + s - 1 - k)) \end{aligned}$$

This simplifies to $n = 2k - E(\Gamma_1)$, hence $n \leq 2k$. This was under the assumption that Γ has no loops at the basepoint, so if $n > 2k$ there must be at least one loop at the basepoint. \square

We denote by $A_{n,s}^k$ the subspace of $A_{n,s}$ consisting of thorned graphs of degree at most k .

The version of the Degree Theorem that we will need is the following.

Theorem 1 *A map $f_0: D^k \rightarrow A_{n,s}$ is homotopic to a map $f_1: D^k \rightarrow A_{n,s}^k$ by a homotopy during which degree decreases monotonically.*

Proof The proof in [6] of the Degree Theorem applies directly to thorned graphs, and we refer the reader there for details. Here is a sketch of the proof.

We consider the natural Morse function on each graph measuring distance to the basepoint. Small initial segments of the edges below each critical point form the ϵ -cones of the graph. The thorns all point upwards with respect to the Morse function, so are never part of any ϵ -cones.

The k -parameter family of marked graphs is initially perturbed, by varying the edge-lengths of the graphs, to remove all but k critical points (counted with multiplicity) of each graph in the family. If the graphs have thorns, the remaining critical points may or may not occur at the base of thorns.

The deformation then proceeds by two operations, which may need to be repeated several times. First is *canonical splitting*, which moves non-critical vertices down towards the basepoint until they hit a critical vertex. In this process,

the non-critical bases of thorns move downward just as any other vertex would. The second operation is *sliding in the ϵ -cones*, which perturbs the attaching points of edges (including thorns) which come down into a critical vertex. The perturbation moves attaching points further down the ϵ -cones, so they can then be canonically split further down the graph. After these two processes are repeated sufficiently often, the degree of each graph in the family is at most k . \square

5 Graphs with loops at the basepoint

We will prove that $W_{n,s}$ is highly connected by comparing it to the subspace $L_{n,s}$ of $A_{n,s}$ consisting of graphs with at least one loop at the basepoint and no separating edges other than thorns.

Theorem 2 *The subspace $L_{n,s}$ of $A_{n,s}$ is $(n-3)/2$ -connected.*

Proof Note first that separating edges which are not thorns can always be eliminated by shrinking them uniformly to points, so this condition in the definition of $L_{n,s}$ is easy to achieve and we will say no more about it in the proof.

Suppose $k < n/2$, and take a map of S^{k-1} into $L_{n,s}$. Since $A_{n,s}$ is contractible, this can be extended to a map $f_0: D^k \rightarrow A_{n,s}$. By the Degree Theorem, this is homotopic to a map $f_1: D^k \rightarrow A_{n,s}^k$ by a homotopy which monotonically decreases degree. By the preceding Lemma, every graph of degree at most $n/2$ has at least one loop at the basepoint, so that $f_1(D^k) \subset L_{n,s}$.

The proof of the Degree Theorem shows that an edge which terminates at the basepoint at time t remains at the basepoint for all $t' > t$. In particular, the number of loops at the basepoint can only increase. Thus the image of S^{k-1} stays in $L_{n,s}$ for all times t and gets filled in at time $t = 1$ by a D^k in $L_{n,s}$, so $L_{n,s}$ is $(k-1)$ -connected. \square

Let $SL_{n,s}$ denote the spine of $L_{n,s}$, the simplicial complex associated to the poset of simplices of $L_{n,s}$. Thus $SL_{n,s}$ is a subcomplex of the spine of $A_{n,s}$ and is embedded in $L_{n,s}$ as a deformation retract.

We will change to the language of sphere systems in what follows, because sphere systems detect structure in $M_{n,s}$ which is invisible in terms of graphs. Let $M = M_{n,s}$, with boundary spheres $\partial_0, \dots, \partial_{s-1}$. Given a sphere system S

consisting of spheres S_1, \dots, S_k in M , the boundary of $M - S$ consists of ∂M together with pairs S_i^+, S_i^- coming from the S_i . The space $L_{n,s}$ is identified with systems S such that

- S is simple
- all spheres in S are nonseparating
- the component of $M - S$ which contains ∂_0 also contains at least one pair S_i^+, S_i^- .

6 Sphere systems with connected complement

We will be interested in sphere systems S in $M = M_{n,s}$ whose complement $M - S$ is connected. For brevity we call such systems *coconnected*. Since a subsystem of a coconnected system is coconnected, the (isotopy classes of) coconnected systems form a subcomplex $Y = Y_{n,s}$ of $S(M)$.

When $s > 0$ we will also need a refined version of Y whose definition will use the notion of an *enveloping sphere* for a coconnected system S . By this we mean a sphere S_e disjoint from S which separates $M - S$ into two pieces, one of which is a punctured ball whose boundary consists of S_e , the base boundary sphere ∂_0 , and the pairs S_i^+, S_i^- for the spheres S_i of S (Figure 5).




Figure 5: Enveloping sphere

When S has n spheres (the maximum number possible), the enveloping sphere is unique up to isotopy. If S has n spheres and $s = 1$ or $s = 2$, the enveloping sphere will be trivial, bounding a ball if $s = 1$ or parallel to ∂_1 if $s = 2$, so in these cases the enveloping sphere is not very interesting. However, when S has fewer than n spheres, the enveloping sphere is not unique, and so represents additional data not specified by S .

Let $V = V_{n,s}$ be the complex whose simplices are isotopy classes of pairs (S, S_e) where S is a coconnected system and S_e is an enveloping sphere for S . To obtain a face of the simplex (S, S_e) we pass to a subsystem S' of S , with the unique (up to isotopy) enveloping sphere S'_e disjoint from S and S_e .

Theorem 3 *The complex $Y_{n,s}$ is $(n - 2)$ -connected for $s \geq 0$, and $V_{n,s}$ is $(n - 3)/2$ -connected for $s > 0$.*

Proof For $Y_{n,s}$ this is part of Proposition 3.1 of [4]. For $V_{n,s}$ we will describe a map $f: SL_{n,s} \rightarrow V_{n,s}$ and show this is a homotopy equivalence.

For a sphere system S in $SL_{n,s}$ let M_0 denote the component of $M - S$ containing ∂_0 , and let S_1, \dots, S_k be the spheres in S which contribute two boundary components, S_i^+ and S_i^- , to M_0 . These are the spheres which represent loops at the basepoint, in the graph picture. The component M_0 is a punctured 3-sphere, as are all components of $M - S$ since S is simple. We define $f(S) = (T, T_e)$ where $T = \{S_1, \dots, S_k\}$ and T_e is the unique enveloping sphere for T which is contained in M_0 .

We claim that f is a poset map, where the poset relation on $V_{n,s}$ is also given by inclusion of faces. Note that any face S' of a sphere system S in $SL_{n,s}$ must contain all of the spheres S_1, \dots, S_k , since removing an S_i results in a system which is not simple, and therefore not in $SL_{n,s}$. Since S' is obtained by removing spheres other than the S_i , the corresponding component M'_0 is either the same or larger than M_0 . If M'_0 is larger, and $f(S') = (T', T'_e)$, then the enveloping sphere T'_e can be taken to be disjoint from T_e , so (T', T'_e) contains (T, T_e) as a proper face. Thus f is indeed a poset map, in which the poset ordering is reversed.

Now let $\tau = (T, T_e)$ be an arbitrary simplex in $V_{n,s}$ and consider the fiber

$$f_{\geq \tau} = \{S \in SL_{n,s} \mid f(S) \geq \tau\}.$$

Let k be the number of spheres in T . If $k = n$, then $f_{\geq \tau} = f^{-1}(\tau)$ is a single point (corresponding to a basic thorned rose), since all spheres in $SL_{n,s}$ are nonseparating. Now suppose $k < n$. Then $f_{\geq \tau}$ consists of all simple systems of nonseparating spheres in $M_{n,s}$ which contain T and are compatible with (disjoint from) T_e . The sphere T_e cuts $M_{n,s}$ into two pieces, one of which is homeomorphic to $M_{n-k,s}$, and $f_{\geq \tau}$ is isomorphic to the complex of simple systems of nonseparating spheres in this piece. This is the space $A_{n-k,s}$, or more precisely the subspace of $A_{n-k,s}$ obtained by shrinking separating edges. We noted earlier that $A_{n,s}$ is contractible for all $n > 0$, so it follows that the fiber $f_{\geq \tau}$ is contractible for $k < n$ as well. Quillen's Poset Lemma [11] then implies that f is a homotopy equivalence, so that $V_{n,s}$ is $(n - 3)/2$ -connected. \square

The group $\Gamma_{n,s}$ acts on Y and V , but the stabilizers of simplices are larger than we would like for the spectral sequence arguments we will be using. More specifically, we would prefer that the stabilizers not contain elements that permute the spheres in a coconnected sphere system or reverse their orientations. To deal with orientations we will use oriented versions Y^\pm and V^\pm of Y and V , in which a k -simplex is a k -simplex S or (S, S_e) with a choice of orientation for each sphere of S . Thus a k -simplex of Y or V corresponds to 2^{k+1} k -simplices of Y^\pm or V^\pm .

Proposition 2 Y^\pm is $(n - 2)$ -connected and $V_{n,s}^\pm$ is $(n - 3)/2$ -connected.

Proof For Y^\pm this was shown in Proposition 3.1 of [4], and the proof for V^\pm is essentially the same. First choose arbitrarily a positive orientation for each isotopy class of split-off handles. The subcomplex V^+ of V^\pm consisting of positively oriented systems is then a copy of V , so it will suffice to deform a given simplicial map of a triangulated i -sphere S^i into V^\pm to a map with image in V^+ , for $i \leq (n - 3)/2$. Let τ be a totally negative simplex of S^i of maximum dimension k , where *totally negative* means $f(\tau) = (T, T_e)$ and all of the spheres in T are negatively oriented. The linking sphere S^{i-k-1} is either empty (if $k = i$) or maps to V^+ , in which case f restricted to S^{i-k-1} can be viewed as a map to $V_{m,s}^+$ where $M_{m,s}$ is the piece split off by T_e . We have $m \geq n - k - 1$ since T contains at most $k + 1$ spheres. Thus $V_{m,s}^+$ is $(n - k - 4)/2$ -connected, being a copy of $V_{m,s}$. The assumption $i \leq (n - 3)/2$ implies that $i - k - 1 \leq (n - k - 4)/2$, so $f|_{S^{i-k-1}}$ extends to a map $g: D^{n-k} \rightarrow V_{m,s}^+$. (If $k = i$, g just picks a vertex of $V_{m,s}$.) Writing the star of τ in S^i in terms of joins as $\tau * S^{n-k-1} = \partial\tau * D^{n-k}$ we can therefore modify f on this star by replacing it by the map which is the join of $f|_{\partial\tau}$ with g . The modified f is homotopic to the original f via the join of $f|_\tau$ with g . The new f is an improvement on the old one in that no simplices in the star of τ of dimension k or greater are now totally negative. Outside the star the map f is unchanged, so after a finite number of such modifications we will have deformed f to have image in V^+ , as desired. \square

To eliminate permutations from the simplex stabilizers for the actions of $\Gamma_{n,s}$ on Y^\pm and V^\pm we make use of a general construction that can be applied to any simplicial complex Z . This construction associates to Z the complex $\Delta(Z)$ whose k -simplices are the simplicial maps $\Delta^k \rightarrow Z$. A k -simplex of $\Delta(Z)$ is then an ordered $(k + 1)$ -tuple of vertices of Z , all of which are vertices of a single simplex of Z . Repeated vertices are allowed in such a $(k + 1)$ -tuple, so $\Delta(Z)$ is infinite-dimensional if Z is nonempty. The complex $\Delta(Z)$ is no longer

a simplicial complex but just a Δ -complex, a CW complex with simplicial cells. There is a canonical map $\Delta(Z) \rightarrow Z$, and it is a classical fact in algebraic topology this map induces an isomorphism on homology. This is shown in Theorem 4.6.8 in [12] for example. [Here is a sketch of the proof. It is easy to construct a contraction of $\Delta(\Delta^k)$, so the result holds for Δ^k . The result for finite Z then follows by induction on the number of simplices via the five lemma and the Mayer–Vietoris sequence for simplicial homology. A direct limit argument with simplicial homology then gives the general case.]

It can be shown that the map $\Delta(Z) \rightarrow Z$ is not only a homology equivalence, but in fact a homotopy equivalence; however, we will not need this stronger statement.

We now set $X_{n,s} = \Delta(Y_{n,s}^\pm)$ and $W_{n,s} = \Delta(V_{n,s}^\pm)$.

7 The first stabilization

Using the spectral sequence associated to the action of the group $\Gamma = \Gamma_{n+1,s}$ on the complex $W = W_{n+1,s}$ we will prove:

Theorem 4 *For $s \geq 1$, the map $\alpha_*: H_i(\Gamma_{n,s}) \rightarrow H_i(\Gamma_{n+1,s})$ induced by the inclusion $\alpha: \Gamma_{n,s} \rightarrow \Gamma_{n+1,s}$ is an isomorphism for $n \geq 2i + 2$ and a surjection for $n = 2i + 1$.*

Note that for $s = 1$, the map α is the standard inclusion $Aut(F_n) \rightarrow Aut(F_{n+1})$, so in this case the theorem gives a slight improvement on Theorem 7.1 of [6] which had $2n + 3$ in place of $2n + 2$.

Proof Let C_* be the cellular chain complex of W augmented by a \mathbb{Z} in degree -1 , and let $E\Gamma_*$ be a free $\mathbb{Z}(\Gamma)$ -resolution of \mathbb{Z} . Then we have a double complex $C_* \otimes E\Gamma_*$ and hence two spectral sequences associated to the horizontal and vertical filtrations of this double complex, both converging to the same thing. The spectral sequence obtained by first taking homology with respect to the vertical boundary maps, in the $E\Gamma_*$ factor, has

$$E_{p,q}^1 = H_q(\Gamma; C_p) = \bigoplus_{\tau} H_q(\Gamma_{\tau}; \mathbb{Z}),$$

where the sum is over all orbits of p -cells τ , Γ_{τ} is the stabilizer of τ , and the second equality comes from Shapiro's lemma. Since all elements of Γ_{τ} fix τ pointwise, the coefficient module \mathbb{Z} is trivial as a Γ -module, and we will

omit this \mathbb{Z} from the notation from now on. If the image of the p -simplex $\tau: \Delta^p \rightarrow V_{n+1,s}^\pm$ is the pair (S, S_e) where the system S has k distinct spheres, then there is a natural surjection $\Gamma_{n+1-k,s} \rightarrow \Gamma_\tau$ coming from the inclusion $M_{n+1-k,s} \rightarrow M_{n+1,s}$, where $M_{n+1-k,s}$ is the part of $M_{n+1,s}$ on the side of S_e away from S . The surjection $\Gamma_{n+1-k,s} \rightarrow \Gamma_\tau$ is in fact an isomorphism by the argument for Lemma 3.3 of [4].

The fact that $W_{n+1,s}$ is m -connected for $m = (n - 2)/2$ implies that $E_{p,q}^\infty = 0$ for $p + q \leq m$. This follows by looking at the spectral sequence coming from the horizontal filtration, which first takes homology with respect to the boundary maps in C_* . This spectral sequence has $E_{p,q}^1 = 0$ for $p \leq m$, so since both spectral sequences converge to the same thing, the assertion follows.

We prove the theorem by induction on the homology dimension i . Thus we assume that for $k < i$, the map α induces an isomorphism on H_k for $n \geq 2k + 2$ and a surjection for $n = 2k + 1$.

Let us show first that α induces a surjection on H_i for $n \geq 2i + 1$. The stabilization $H_i(\Gamma_n) \rightarrow H_i(\Gamma_{n+1})$ is the differential $d^1: E_{0,i}^1 \rightarrow E_{-1,i}^1$, so to show this is surjective it suffices to show:

- (i) $E_{-1,i}^\infty = 0$.
- (ii) $E_{p,i-p}^2 = 0$ for $p > 0$.

Condition (ii) guarantees that all differentials d^r with $r > 1$ going to $E_{-1,i}^r$ must be zero, so only the d^1 differential can kill $E_{-1,i}^1$.

It is easy to see that statement (i) holds since we have already observed that $E_{p,q}^\infty = 0$ for $p + q \leq (n - 2)/2$, so in particular this holds with $(p, q) = (-1, i)$ if $i - 1 \leq (n - 2)/2$, or $n \geq 2i$. To prove (ii) we use the description of the d^1 differentials in Chapter 7 of [1]. In the q -th row these differentials compute the homology of the quotient W/Γ with coefficients in the system of groups $H_q(\Gamma_\tau)$. For $p + q \leq i$ and $q < i$, and for τ' a face of τ , the coefficient maps $H_q(\Gamma_\tau) \rightarrow H_q(\Gamma_{\tau'})$ induced by inclusion are all isomorphisms by the induction hypothesis. In addition, induction gives that the coefficient maps from the $p + q = i + 1$ terms to the $p + q = i$ terms are surjective for $q < i$. Thus for $p + q \leq i$ the groups $E_{p,q}^2$ are homology groups of W/Γ with locally constant coefficients. In fact, we claim that the coefficients are constant. The twisting of coefficients around the faces of a simplex τ is induced by diffeomorphisms of M that permute the nonseparating spheres corresponding to the vertices of τ . These diffeomorphisms can be chosen to have support in the part of M cut off by the enveloping sphere for τ . Since the stabilizer Γ_τ is supported on the other part of M , these diffeomorphisms act trivially on $H_q(\Gamma_\tau)$.

Next we need the fact that the quotient W/Γ is $(n-1)$ -connected. This was shown in Lemma 3.5 of [4] for the analogous quotient X/Γ , and the two quotients are combinatorially equivalent. This is because Γ acts transitively on k -simplices of both the complexes V^\pm and Y^\pm from which W and X are constructed, and all permutations of the vertices of a simplex of V^\pm or Y^\pm are realized by the action of Γ .

Thus if $i \leq n-1$ we conclude that (ii) holds. The condition $i \leq n-1$ is implied by $n \geq 2i+1$, so this finishes the induction step for surjectivity.

To show that $d^1: E_{0,i}^1 \rightarrow E_{-1,i}^1$ is injective, it is enough to show:

- (iii) $E_{0,i}^\infty = 0$.
- (iv) $E_{p+1,i-p}^2 = 0$ for $p > 0$.
- (v) The differential $d^1: E_{1,i}^1 \rightarrow E_{0,i}^1$ is zero.

The arguments for (iii) and (iv) are the same as for (i) and (ii), with a shift of one unit to the right, corresponding to replacing $n \geq 2i+1$ by $n \geq 2i+2$. The connectivity condition for W/Γ is now $i+1 \leq n-1$, and this is implied by $n \geq 2i+2$.

For (v) there are two orbits of 1-cells, one nondegenerate and the other degenerate. The coefficients are automatically untwisted for the degenerate 1-cell, and they are untwisted for the nondegenerate one by the argument given above. In W/Γ each 1-cell attaches at each end to the unique 0-cell, so it follows that the differential is zero. \square

If one examines the proof, one sees that in the inequalities $n \geq 2i+2$ and $n \geq 2i+1$, the coefficient 2 in the $2i$ term is forced by the induction process since the d^2 differentials go two units to the left and one unit upward. On the other hand, the constant term 2 or 1 comes from the connectivity $(n-2)/2$ of $W_{n+1,s}$. If this connectivity estimate were improved, this could yield a corresponding improvement in the constant term of the stable range.

8 The second stabilization

We turn now to the β stabilization $H_i(\Gamma_{n,s+2}) \rightarrow H_i(\Gamma_{n+1,s})$. As an easy application of the theorem in the preceding section we will show:

Theorem 5 *For $s \geq 1$, the map $\beta_*: H_i(\Gamma_{n,s+2}) \rightarrow H_i(\Gamma_{n+1,s})$ induced by β is an isomorphism for $n \geq 2i+2$ and a surjection for $n = 2i+1$.*

Proof Surjectivity of β_* on H_i follows from surjectivity of α because of the commutative diagram from Section 2:

$$\begin{array}{ccc} \Gamma_{n,s} & \xrightarrow{\alpha} & \Gamma_{n+1,s} \\ \downarrow \mu & & \uparrow \beta \\ \Gamma_{n,s+1} & \xrightarrow{\mu} & \Gamma_{n,s+2} \end{array}$$

For injectivity we will first prove the result in an unspecified range $n \gg i$ and then improve this to $n \geq 2i + 2$. We consider the relative homology groups H_i for the pairs in the following commutative diagram:

$$\begin{array}{ccc} (\Gamma_{n,s}, \Gamma_{n-1,s+2}) & \xrightarrow{\alpha} & (\Gamma_{n+1,s}, \Gamma_{n,s+2}) \\ \downarrow \mu & & \uparrow \beta \\ (\Gamma_{n,s+1}, \Gamma_{n-1,s+3}) & \xrightarrow{\mu} & (\Gamma_{n,s+2}, \Gamma_{n-1,s+4}) \end{array}$$

Since α is an isomorphism on the absolute groups H_i for $n \gg i$, it is also an isomorphism on the relative groups. Then the diagram shows that β is surjective on the relative H_i for $n \gg i$. This is the map on the left in the following commutative square:

$$\begin{array}{ccc} H_i(\Gamma_{n,s+2}, \Gamma_{n-1,s+4}) & \longrightarrow & H_{i-1}(\Gamma_{n-1,s+4}) \\ \beta_* \downarrow & & \beta_* \downarrow \\ H_i(\Gamma_{n+1,s}, \Gamma_{n,s+2}) & \longrightarrow & H_{i-1}(\Gamma_{n,s+2}) \end{array}$$

Assuming $n \gg i$ always, we can see that the lower map is injective by looking at a portion of the long exact sequence for the pair $(\Gamma_{n+1,s}, \Gamma_{n,s+2})$:

$$H_i(\Gamma_{n,s+2}) \xrightarrow{\beta_*} H_i(\Gamma_{n+1,s}) \longrightarrow H_i(\Gamma_{n+1,s}, \Gamma_{n,s+2}) \longrightarrow H_{i-1}(\Gamma_{n,s+2})$$

The first map is surjective as we saw at the beginning of the proof, so the second map is zero and the third map is injective. Returning to the commutative square, the composition from upper left to lower right passing through the upper right term is zero since it is two maps in a long exact sequence. Thus the composition through the lower left group is zero. This composition is a surjection followed by an injection, so it follows that $H_i(\Gamma_{n+1,s}, \Gamma_{n,s+2})$ is zero. Hence β induces an isomorphism on H_i for $n \gg i$.

It remains to improve the stable range to what is stated in the theorem. From the first diagram in the proof, the fact that α_* and β_* are isomorphisms for $n \gg i$ then implies that the maps μ_* are also isomorphisms for $n \gg i$, using the fact that μ_* is always injective (since $\gamma \circ \mu$ is the identity, where γ is induced

by filling in a boundary sphere with a ball). For each $s \geq 1$ and each i , we now have a diagram of maps:

$$\begin{array}{ccccccc}
 H_i(\Gamma_{n,s}) & \xrightarrow{\alpha_*} & H_i(\Gamma_{n+1,s}) & \xrightarrow{\alpha_*} & \cdots & \xrightarrow{\alpha_*} & H_i(\Gamma_{n+k,s}) \\
 \downarrow \mu_* & & \downarrow \mu_* & & & & \downarrow \mu_* \\
 H_i(\Gamma_{n,s+1}) & \xrightarrow{\alpha_*} & H_i(\Gamma_{n+1,s+1}) & \xrightarrow{\alpha_*} & \cdots & \xrightarrow{\alpha_*} & H_i(\Gamma_{n+k,s+1})
 \end{array}$$

In both rows, the horizontal maps are all isomorphisms for $n \geq 2i + 2$. For k large the map μ_* on the far right is an isomorphism. Since the diagram commutes, we can work backward to conclude that in fact all the maps μ_* in the diagram are isomorphisms, if $n \geq 2i + 2$. Now the fact that $\beta \circ \mu^2 = \alpha$ allows us to conclude that $\beta_*: H_i(\Gamma_{n,s+2}) \rightarrow H_i(\Gamma_{n+1,s})$ is an isomorphism for $n \geq 2i + 2$, proving the theorem. \square

Theorem 6 *The map $\beta_*: H_i(\Gamma_{n,2}) \rightarrow H_i(\Gamma_{n+1,0})$ induced by β is an isomorphism for $n \geq 2i + 3$ and a surjection for $n = 2i + 2$.*

Proof This is very similar to the proof of Theorem 4, using now the action of $\Gamma = \Gamma_{n+1,0}$ on the complex $X = X_{n+1,0}$. The complex X is $(n - 1)$ -connected, which is better than the connectivity $(n - 2)/2$ for W . The quotient X/Γ was shown to be $(n - 1)$ -connected in [4]. The only remaining issue to deal with in carrying the proof of Theorem 4 over to the present context is the justification for why the local coefficient system on X/Γ is untwisted in steps (ii), (iv), and (v) of the proof of Theorem 4. In the case of W/Γ , we used the enveloping spheres of simplices of W for this, but simplices of X do not have canonical enveloping spheres.

We will solve this problem by choosing somewhat arbitrary enveloping spheres, and showing that the map on homology does not depend on this choice, if n is sufficiently large. If τ is a simplex with k distinct spheres, choose an enveloping sphere which cuts $M_{n+1,0}$ into two pieces, one containing all the spheres in τ , and one diffeomorphic to $M_{n+1-k,1}$. The coefficients over τ will be untwisted if the iterated μ map $H_q(\Gamma_{n+1-k,1}) \rightarrow H_q(\Gamma_{n+1-k,2k})$ is surjective, so that $H_q(\Gamma_\tau)$ is supported on the side of the chosen enveloping sphere away from the spheres of τ . For step (ii) the worst case we have to deal with is the differential $E_{2,i-1}^1 \rightarrow E_{1,i-1}^1$ with $H_{i-1}(\Gamma_{n-2,6})$ in its domain. Here the iterated μ will be an isomorphism if $n - 2 \geq 2(i - 1) + 2$, or $n \geq 2i + 2$. Since steps (i) and (ii) give surjectivity, we conclude that β_* is surjective for $n \geq 2i + 2$.

To get injectivity, we need steps (iii)–(v). For step (iv) we repeat the argument for (ii), but shifted one unit to the right, making the inequality $n \geq 2i + 3$. For

(v) we are looking at the differential $E_{1,i}^1 \rightarrow E_{0,i}^1$ with $H_i(\Gamma_{n-1,4})$ in its domain, so we need $n - 1 \geq 2i + 2$, or $n \geq 2i + 3$. \square

Corollary 1 *The quotient map $Aut(F_n) \rightarrow Out(F_n)$ is an isomorphism on homology in dimension i for $n \geq 2i + 4$, and a surjection for $n = 2i + 3$.*

Proof The quotient map $Aut(F_n) \rightarrow Out(F_n)$ is the map γ in the following diagram, induced by filling in the boundary sphere with a ball:

$$\begin{array}{ccc} \Gamma_{n-1,1} & \xrightarrow{\alpha} & \Gamma_{n,1} \\ \downarrow \mu & & \downarrow \gamma \\ \Gamma_{n-1,2} & \xrightarrow{\beta} & \Gamma_{n,0} \end{array}$$

The map α induces an isomorphism on H_i for $n \geq 2i + 3$ by Theorem 4. It was shown in the proof of Theorem 5 that μ induces an isomorphism on H_i in the same range. The corollary follows since β induces an isomorphism for $n \geq 2i + 4$ and a surjection for $n = 2i + 3$. \square

Because the homology of $Aut(F_n) = \Gamma_{n,1}$ stabilizes, we obtain the following corollary.

Corollary 2 *The group $H_i(Out(F_n))$ is independent of n when $n \geq 2i + 4$.*

References

- [1] **KS Brown**, *Cohomology of groups*, Graduate Texts in Mathematics 87, Springer-Verlag, New York-Berlin (1982) [MathReview](#)
- [2] **J Conant, K Vogtmann**, *Morita classes in the homology of automorphism groups of free groups*, *Geom. Topol.* 8 (2004) 1471–1499
- [3] **F Gerlits**, *Invariants in chain complexes of graphs*, PhD thesis, Cornell University (2002)
- [4] **A Hatcher**, *Homological stability for automorphism groups of free groups*. *Comment. Math. Helv.* 70 (1995) 39–62 [MathReview](#)
- [5] **A Hatcher**, *Algebraic Topology*, Cambridge University Press, Cambridge (2002) [MathReview](#)
- [6] **A Hatcher, K Vogtmann**, *Cerf theory for graphs*, *J. London Math. Soc.* 58 (1998) 633–655 [MathReview](#)
- [7] **A Hatcher, K Vogtmann**, *Rational homology of $Aut(F_n)$* , *Mathematical Research Letters* 5 (1998) 759–780 [MathReview](#)

- [8] **A Hatcher, N Wahl**, *Stabilization for the automorphisms of free groups with boundaries*, preprint [arXiv:math.GT/0406277](https://arxiv.org/abs/math/0406277)
- [9] **C Jensen, N Wahl**, *Automorphisms of free groups with boundaries*, *Algebr. Geom. Topol.* 4 (2004) 543–569 [MathReview](#)
- [10] **F Laudenbach**, *Sur les 2-sphères d'une variété de dimension 3*, *Ann. of Math.* 97 (1973) 57–81 [MathReview](#)
- [11] **D Quillen**, *Homotopy properties of the poset of nontrivial p -subgroups of a group*, *Adv. in Math.* 28 (1978) 101–128 [MathReview](#)
- [12] **E H Spanier**, *Algebraic Topology*, McGraw-Hill Book Co. New York–Toronto–London (1966) [MathReview](#)
- [13] **N Wahl**, *From mapping class groups to automorphism groups of free groups*, [arXiv:math.AT/0406278](https://arxiv.org/abs/math/0406278)

*Department of Mathematics, Cornell University
Ithaca, NY 14853-4201, USA*

Email: hatcher@math.cornell.edu and vogtmann@math.cornell.edu

Received: 28 June 2004