

*Geometry & Topology Monographs*

Volume 4: Invariants of knots and 3-manifolds (Kyoto 2001)

Pages 303–311

## On the potential functions for the hyperbolic structures of a knot complement

YOSHIYUKI YOKOTA

**Abstract** We explain how to construct certain potential functions for the hyperbolic structures of a knot complement, which are closely related to the analytic functions on the deformation space of hyperbolic structures.

**AMS Classification** 57M50; 57M25, 57M27

**Keywords** Potential function, hyperbolicity equations, volume, Chern-Simons invariant

*Dedicated to Professor Mitsuyoshi Kato for his 60'th birthday*

### 1 Introduction

Let  $M$  be the complement of a hyperbolic knot  $K$  in  $S^3$ . Through the study of *Kashaev's conjecture*, we have found a complex function which gives the *volume* and the *Chern-Simons invariant* of the complete hyperbolic structure of  $M$  at the critical point corresponding to the promised solution to the hyperbolicity equations for  $M$ , see [2, 4] for details.

The purpose of this article is to explain how to construct such complex functions for the non-complete hyperbolic structures of  $M$ . Such functions are closely related to the analytic functions on the deformation space of the hyperbolic structures of  $M$ , parametrized by the eigenvalue of the holonomy representation of the meridian of  $K$ , which reveal a complex-analytic relation between the volumes and the Chern-Simons invariants of the hyperbolic structures of  $M$ , see [3, 5] for details.

In this note, we suppose  $K$  is  $5_2$  for simplicity which is represented by the diagram  $D$  depicted in Figure 1.




Figure 1

## 2 Geometry of a knot complement

### 2.1 Ideal triangulations

We first review an ideal triangulation of  $M$  due to D. Thurston. Let  $\dot{M}$  denote  $M$  with two poles  $\pm\infty$  of  $S^3$  removed. Then,  $\dot{M}$  decomposes into 5 ideal octahedra corresponding to the 5 crossings of  $D$ , each of which further decomposes into 4 ideal tetrahedra around an axis, as shown in Figure 2.




Figure 2

In fact, we can recover  $\dot{M}$  by glueing adjacent tetrahedra as shown in Figure 3.




Figure 3

As usual, we put a hyperbolic structure on each tetrahedron by assigning a complex number, called *modulus*, to the edge corresponding to the axis as shown in Figure 4. In what follows, we denote the tetrahedron with modulus  $z$  by  $T(z)$ .




Figure 4

Let  $B$  be the intersection between  $T(a_1) \cup T(b_1)$  and  $T(b_3) \cup T(c_3)$ . Then, each of

$$T(a_1), T(b_1), T(b_3), T(c_3)$$

intersects  $\partial N(B \cup K)$  in two triangles, and they are essentially one-dimensional objects in  $S^3 \setminus N(B \cup K)$ . On the other hand, each of

$$T(c_1), T(d_1), T(a_2), T(b_2), T(d_2), T(a_3), T(d_3), T(a_4), T(b_4), T(c_4), T(c_5)$$

intersects  $\partial N(B \cup K)$  in two triangles and one quadrangle, and they are essentially two-dimensional objects in  $S^3 \setminus N(B \cup K)$ . Thus, by contracting these

15 tetrahedra, we obtain an ideal triangulation  $\mathcal{S}$  of  $M$  with

$$T(c_2), T(d_4), T(a_5), T(b_5), T(d_5).$$

Figure 5 exhibits the triangulation of  $\partial N(B \cup K)$  induced by  $\mathcal{S}$ , where each couple of edges labeled with the same number are identified.




Figure 5

### 2.2 Hyperbolicity equations

If  $c_2, d_4, a_5, b_5, d_5$  above give a hyperbolic structure of  $M$ , the product of the moduli around each edge in  $\mathcal{S}$  should be 1, which is called the *hyperbolicity equations* and can be read from Figure 5 as follows.

$$\begin{aligned} d_4 b_5 &= a_5 b_5 d_5 = 1, \\ \frac{c_2 a_5 (1 - 1/d_4)}{1 - d_4} \cdot \frac{(1 - 1/d_5)(1 - 1/c_2)(1 - 1/b_5)}{(1 - a_5)(1 - b_5)} &= 1, \\ \frac{c_2 (1 - 1/a_5)}{(1 - d_5)(1 - c_2)} \cdot \frac{(1 - 1/d_5)(1 - 1/b_5)}{(1 - d_5)(1 - a_5)(1 - d_4)} &= 1, \\ \frac{d_4 (1 - 1/a_5)(1 - 1/d_4)}{1 - b_5} \cdot \frac{d_5 (1 - 1/c_2)}{1 - c_2} &= 1. \end{aligned}$$

It is easy to observe that these equations are generated by

$$d_4 b_5 = a_5 b_5 d_5 = 1, \quad \frac{c_2 a_5}{d_4} = \frac{1 - 1/a_5}{(1 - 1/c_2)(1 - d_5)} = \frac{(1 - 1/a_5)(1 - d_4)}{1 - b_5} = \frac{c_2}{d_5},$$

which suggests to put

$$c_2 = y\xi, \quad d_4 = x/\xi, \quad a_5 = x/y, \quad b_5 = \xi/x, \quad d_5 = y/\xi$$

and to rewrite the hyperbolicity equations as follows.

$$\frac{(1 - y/x)(1 - x/\xi)}{1 - \xi/x} = \frac{1 - y/x}{(1 - y/\xi)(1 - 1/y\xi)} = \xi^2.$$




Figure 6

Note that the variables  $x, y$  correspond to the interior edges of a graph depicted in Figure 6, which is  $D$  with some edges deleted.

A solution to the equations above determines a hyperbolic structure of  $M$ , where  $\xi$  is nothing but the eigenvalue of the holonomy representation of the meridian of  $K$ . The set  $\mathcal{D}$  of such solutions is called the *deformation space* of the hyperbolic structures of  $M$  and can be parametrized by  $\xi$  or the eigenvalue  $\eta$  of the holonomy representation of the longitude of  $K$ . In our example,  $\eta$  is given by

$$\eta = \frac{y\xi^6}{x} \cdot (1 - 1/y\xi) = \frac{y\xi^6}{x} \cdot \frac{1 - \xi/x}{(1 - x/\xi)(1 - y/\xi)}.$$

Note that the factors  $1 - x/\xi, 1 - y/\xi, 1 - \xi/x$  and  $1 - 1/y\xi$  correspond to the corners of  $D$  which touch the unbounded regions.

### 3 Potential functions

Curious to say, we can always construct a *potential function* for the hyperbolicity equations and  $\eta$  combinatorially by using Euler's dilogarithm function

$$\text{Li}_2(z) = - \int_0^z \frac{\log(1-w)}{w} dw,$$

where we remark that the volume of a tetrahedron with modulus  $z$  is given by

$$D(z) = \text{Im Li}_2(z) + \log |z| \arg(1 - z).$$

### 3.1 Neumann-Zagier's functions

In fact, we define  $V(x, y, \xi)$  by

$$-\text{Li}_2(1/y\xi) + \text{Li}_2(y/\xi) - \text{Li}_2(y/x) + \text{Li}_2(\xi/x) + \text{Li}_2(x/\xi) + \log \xi \log \frac{x^2}{y^2\xi^6} - \frac{\pi^2}{6},$$

the principal part of which is nothing but the sum of dilogarithm functions associated to the corners of the graph as shown in Figure 7.




Figure 7

Then, we have

$$x \frac{\partial V}{\partial x} = \log \frac{\xi^2(1 - \xi/x)}{(1 - y/x)(1 - x/\xi)}, \quad y \frac{\partial V}{\partial y} = \log \frac{1 - y/x}{\xi^2(1 - y/\xi)(1 - 1/y\xi)},$$

both of which vanish on  $\mathcal{D}$ , and

$$\begin{aligned} \xi \frac{\partial V}{\partial \xi} &= \log \frac{x^2(1 - x/\xi)(1 - y/\xi)}{y^2\xi^{12}(1 - \xi/x)(1 - 1/y\xi)} \\ &= \log \left\{ \frac{x}{y\xi^6} \cdot \frac{1}{1 - 1/y\xi} \right\}^2 - x \frac{\partial V}{\partial x} - y \frac{\partial V}{\partial y} \\ &= \log \left\{ \frac{x}{y\xi^6} \cdot \frac{(1 - x/\xi)(1 - y/\xi)}{1 - \xi/x} \right\}^2 + x \frac{\partial V}{\partial x} + y \frac{\partial V}{\partial y}, \end{aligned}$$

that is,

$$\xi \frac{\partial V}{\partial \xi} = -\log \eta^2$$

on  $\mathcal{D}$ , which shows  $V(x, y, e^u)$  coincides with  $\Phi(u)$  given in [3, Theorem 3].

### 3.2 Dehn fillings

Furthermore, for a slope  $\alpha \in \mathbb{Q}$ , we put

$$V_\alpha(x, y, \xi) = V(x, y, \xi) + \frac{\log \xi(2\pi\sqrt{-1} - p \log \xi)}{q},$$

where  $p, q \in \mathbb{Z}$  denote the numerator and the denominator of  $\alpha$ . Then, we have

$$\xi \frac{\partial V_\alpha}{\partial \xi} = \xi \frac{\partial V}{\partial \xi} + \frac{2\pi\sqrt{-1} - p \log \xi^2}{q} = \frac{2\pi\sqrt{-1} - p \log \xi^2 - q \log \eta^2}{q},$$

and so a solution  $(x_\alpha, y_\alpha, \xi_\alpha)$  to the equations

$$dV_\alpha(x, y, \xi) = 0$$

determines the complete hyperbolic structure of the closed 3-manifold  $M_\alpha$  obtained from  $M$  by  $\alpha$  Dehn filling. Note that, by choosing  $r, s \in \mathbb{Z}$  such that  $ps - qr = 1$ , we can compute the logarithm of the eigenvalue of the holonomy representation of the core geodesic  $\gamma_\alpha$  of  $M_\alpha$  which is related to the *length* and the *torsion* of  $\gamma_\alpha$  as follows, see [3, Lemma 4.2].

$$\log \xi^r \eta^s = \frac{s\pi\sqrt{-1} - \log \xi}{q} = \frac{\text{length}(\gamma_\alpha) + \sqrt{-1} \cdot \text{torsion}(\gamma_\alpha)}{2}.$$

### Volumes and Chern-Simons invariants

#### 3.3 Yoshida's functions

As in [4], we can observe

$$\begin{aligned} \text{Im } V_\alpha(x, y, \xi) &= -D(1/y\xi) + D(y/\xi) - D(y/x) + D(\xi/x) + D(x/\xi) \\ &\quad + \log |x| \cdot \text{Im } x \frac{\partial V_\alpha}{\partial x} + \log |y| \cdot \text{Im } y \frac{\partial V_\alpha}{\partial y} + \log |\xi| \cdot \text{Im } \xi \frac{\partial V_\alpha}{\partial \xi}, \end{aligned}$$

and so

$$\text{Im } V_\alpha(x_\alpha, y_\alpha, \xi_\alpha) = \text{vol}(M_\alpha).$$

To detect  $\text{Re } V_\alpha(x_\alpha, y_\alpha, \xi_\alpha)$ , we shall consider

$$R(x, y, \xi) = -R(1/y\xi) + R(y/\xi) - R(y/x) + R(\xi/x) + R(x/\xi) - \frac{\pi^2}{6},$$

where  $R(z)$  denotes Roger's dilogarithm function defined by

$$R(z) = \text{Li}_2(z) + \log z \log(1 - z)/2.$$

Then,  $R(x, y, \xi)$  can be expressed as

$$\begin{aligned} &- \text{Li}_2(1/y\xi) + \text{Li}_2(y/\xi) - \text{Li}_2(y/x) + \text{Li}_2(\xi/x) + \text{Li}_2(x/\xi) \\ &- \frac{\log x}{2} \left( x \frac{\partial V}{\partial x} - \log \xi^2 \right) - \frac{\log y}{2} \left( y \frac{\partial V}{\partial y} + \log \xi^2 \right) - \frac{\log \xi}{2} \left( \xi \frac{\partial V}{\partial \xi} - \log \frac{x^2}{y^2 \xi^{12}} \right), \end{aligned}$$

and so  $R(x, y, \xi)$  agrees with

$$V(x, y; \xi) + \log \xi \log \eta$$

on  $\mathcal{D}$  and with

$$V_\alpha(x, y, \xi) - \frac{\log \xi (2\pi\sqrt{-1} - p \log \xi)}{q} + \log \xi \log \eta = V_\alpha(x, y, \xi) - \frac{\pi\sqrt{-1} \cdot \log \xi}{q}$$

at  $(x_\alpha, y_\alpha, \xi_\alpha) \in \mathcal{D}$ . Therefore, we have

$$\begin{aligned} R(x_\alpha, y_\alpha, \xi_\alpha) &= V_\alpha(x_\alpha, y_\alpha, \xi_\alpha) - \frac{s\pi^2 + \pi\sqrt{-1} \cdot \log \xi_\alpha}{q} + \frac{s\pi^2}{q} \\ &= V_\alpha(x_\alpha, y_\alpha, \xi_\alpha) + \frac{\pi\sqrt{-1}}{2} \cdot \{\text{length}(\gamma_\alpha) + \sqrt{-1} \cdot \text{torsion}(\gamma_\alpha)\} + \frac{s\pi^2}{q}. \end{aligned}$$

In particular,

$$\begin{aligned} \text{Im} \frac{2}{\pi} \cdot R(x_\alpha, y_\alpha, \xi_\alpha) &= \text{Im} \frac{2}{\pi} \cdot V_\alpha(x_\alpha, y_\alpha, \xi_\alpha) + \frac{2 \log |\xi_\alpha|}{q} \\ &= \frac{2}{\pi} \cdot \text{vol}(M_\alpha) + \text{length}(\gamma_\alpha), \end{aligned}$$

which shows that, up to a pure imaginary constant,

$$\frac{2}{\pi\sqrt{-1}} R(x, y, e^u)$$

must coincide with  $2\pi f(u)$  of [3, Theorem 2], and that

$$\text{Re} \frac{2}{\pi} \cdot R(x_\alpha, y_\alpha, \xi_\alpha) = \text{Re} \frac{2}{\pi} \cdot \left\{ V_\alpha(x_\alpha, y_\alpha, \xi_\alpha) + \frac{s\pi^2}{q} \right\} - \text{torsion}(\gamma_\alpha)$$

must coincide with  $-4\pi CS(M_\alpha) - \text{torsion}(\gamma_\alpha)$ . Consequently, up to some constant which is independent of  $\alpha$ , we have

$$\text{Re} \left\{ V_\alpha(x_\alpha, y_\alpha, \xi_\alpha) + \frac{s\pi^2}{q} \right\} = -2\pi^2 CS(M_\alpha).$$

## 4 Concluding remarks

We redefine  $V_\alpha(x, y, \xi)$  as follows.

$$V_\alpha(x, y, \xi) = V(x, y, \xi) + \frac{\log \xi (2\pi\sqrt{-1} - p \log \xi) + s\pi^2}{q}.$$

Then,  $dV_\alpha(x, y, \xi) = 0$  gives the hyperbolicity equations for  $M_\alpha$ , and

$$V_\alpha(x_\alpha, y_\alpha, \xi_\alpha) = -2\pi^2 CS(M_\alpha) + \text{vol}(M_\alpha)\sqrt{-1}$$

up to a real constant, where  $(x_\alpha, y_\alpha, \xi_\alpha)$  is a solution to the equations above.

We finally remark that such a construction always works, even for a *link*, and the analytic functions in [3, 5] are now combinatorially constructed up to a constant. For the figure-eight knot and  $\alpha \in \mathbb{Z}$ , our potential function coincides with the function in [1] which appears in the “optimistic” limit of the quantum  $SU(2)$  invariants of  $M_\alpha$ .



## References

- [1] **H Murakami**, *Optimistic calculations about the Witten-Reshetikhin-Turaev invariants of closed three-manifolds obtained from the figure-eight knot by integral Dehn surgeries*, “Recent Progress Toward the Volume Conjecture”, RIMS Kokyuroku 1172 (2000) 70–79
- [2] **H Murakami, J Murakami, M Okamoto, T Takata, Y Yokota**, *Experiment. Math.* 11 (2002) 427–435
- [3] **W D Neumann, D Zagier**, *Volumes of hyperbolic 3-manifolds*, *Topology* 24 (1985) 307–332
- [4] **Y Yokota**, *On the volume conjecture for hyperbolic knots*, preprint available at <http://www.comp.metro-u.ac.jp/~jojo/volume-conjecture.ps>
- [5] **T Yoshida**, *The  $\eta$ -invariant of hyperbolic 3-manifolds*, *Invent. Math.* 81 (1985) 473–514

*Department of Mathematics, Tokyo Metropolitan University  
Tokyo, 192-0397, Japan*

Email: [jojo@math.metro-u.ac.jp](mailto:jojo@math.metro-u.ac.jp)

Received: 5 December 2001      Revised: 19 February 2002