

# Some Remarks on the Poisson-Nijenhuis and the Jacobi Structures

J. M. Nunes da Costa<sup>1</sup>

*Departamento de Matemática, Universidade de Coimbra,  
Apartado 3008, 3000 Coimbra, Portugal  
e-mail: jmcosta@mat.uc.pt*

**Abstract.** We study the relationship between two (compatible) Jacobi structures on a manifold  $M$ , using their associated homogeneous Poisson structures on  $\mathbb{R} \times M$ , in the case where these Poisson tensors are related by a Nijenhuis tensor.

M. S. Classification (1991): 58F05.

Keywords: Poisson-Nijenhuis manifold, Jacobi manifold.

## Introduction

All objects considered in this paper (manifolds, maps, differential forms, vector and tensor fields) are assumed to be differentiable of class  $C^\infty$ .

Let  $M$  be a smooth manifold equipped with a *Poisson tensor*  $\Lambda$  and a *Nijenhuis tensor*  $N$ , that is, a tensor field of type  $(1, 1)$  whose *Nijenhuis torsion*  $\tau(N)$  vanishes everywhere. The Nijenhuis torsion  $\tau(N)$  of  $N$  is given by the following formula, where  $X$  and  $Y$  are vector fields on  $M$ ,

$$\tau(N)(X, Y) = [NX, NY] - N([NX, Y] + [X, NY] - N[X, Y]).$$

We denote by  $\Lambda^\# : T^*M \rightarrow TM$  the vector bundle map such that, for any  $x \in M$ ,  $\alpha, \beta \in T_x^*M$ ,

$$\langle \beta, \Lambda^\#(\alpha) \rangle = \Lambda_x(\alpha, \beta).$$

---

<sup>1</sup>This work was partially supported by CMUC-FCT and PRAXIS. The paper is in final form and no other version has been submitted for publication elsewhere.

With  $\Lambda$  and  $N$  we can define a tensor field  $\mathcal{R}(\Lambda, N)$  of type  $(2, 1)$ , called the *Magri-Morosi concomitant* [Ma-Mo 84] of  $\Lambda$  and  $N$ , that is defined, for any pair of a 1-form  $\alpha$  and a vector field  $X$ , by

$$\mathcal{R}(\Lambda, N)(\alpha, X) = (\mathcal{L}_{\Lambda^\#(\alpha)}N)X - \Lambda^\#(\mathcal{L}_X({}^tN\alpha)) + \Lambda^\#(\mathcal{L}_{NX}\alpha), \tag{1}$$

where  ${}^tN$  is the transpose of  $N : TM \rightarrow TM$ .

Let  $\Lambda$  and  $N$  be, respectively, a Poisson tensor and a Nijenhuis tensor on  $M$ . The triple  $(M, \Lambda, N)$  is called a *Poisson-Nijenhuis manifold* [KSch-Ma 90] if

$$N\Lambda^\# = \Lambda^\#{}^tN \quad \text{and} \quad \mathcal{R}(\Lambda, N) = 0.$$

If  $(M, \Lambda, N)$  is a Poisson-Nijenhuis manifold, there exists a sequence  $(\Lambda_k)_{k \in \mathbb{N}}$  of Poisson tensors on  $M$ , with  $\Lambda_k = N^k\Lambda$ . Moreover, these Poisson tensors are pairwise compatible, that is,  $[\Lambda_l, \Lambda_k] = 0$ , for all  $l, k \in \mathbb{N}$ .

Besides the Poisson-Nijenhuis structure, we are going to use, in what follows, the notion of Jacobi manifold.

A *Jacobi manifold* [Lich 78] is a triple  $(M, C, E)$  where  $C$  and  $E$  are respectively a bivector and a vector field on  $M$ , such that

$$[E, C] = 0 \quad \text{and} \quad [C, C] = 2E \wedge C.$$

When  $E = 0$ , the Jacobi manifold is a Poisson manifold. The *Jacobi bracket* of  $f, g \in C^\infty(M, \mathbb{R})$  is given by

$$\{f, g\} = C(df, dg) + f(E.g) - g(E.f),$$

and it defines a *local Lie algebra structure* on  $C^\infty(M, \mathbb{R})$ .

If  $(M, C, E)$  is a Jacobi manifold and  $h$  is a nowhere vanishing function on  $M$ , then the pair  $(hC, C^\#(dh) + hE) = (C_h, E_h)$  defines a new Jacobi structure on  $M$ , which is said to be *conformally equivalent* to  $(C, E)$ .

With each Jacobi manifold  $(M, C, E)$ , we may associate a homogeneous Poisson structure  $(\Lambda, \frac{\partial}{\partial t})$  on  $\mathbb{R} \times M$ , with  $\Lambda$  given by

$$\Lambda = \exp(-t)(C + \frac{\partial}{\partial t} \wedge E), \tag{2}$$

where  $t$  is the canonical coordinate on  $\mathbb{R}$ .

This paper is divided into two sections. Section 1 is devoted to the subject of compatible Jacobi manifolds. We show how the compatibility is related with the Lichnerowicz-Jacobi cohomology and we present a way of generating compatible Jacobi structures on a manifold. In Section 2, we establish the conditions on the Poisson-Nijenhuis structure of  $\mathbb{R} \times M$  to ensure the compatibility of the corresponding Jacobi structures on  $M$ .

### 1. Compatible Jacobi manifolds

The notion of compatibility of two Jacobi structures on a manifold was introduced in [NdC 98]. We recall that two Jacobi structures  $(C_1, E_1)$  and  $(C_2, E_2)$  on a manifold  $M$  are said to be *compatible* if  $(C_1 + C_2, E_1 + E_2)$  is again a Jacobi structure on  $M$ .

**Proposition 1.1.** [NdC 98] *Let  $M$  be a manifold endowed with two Jacobi structures  $(C_1, E_1)$  and  $(C_2, E_2)$ . Then,  $(C_1, E_1)$  and  $(C_2, E_2)$  are compatible if and only if*

$$[E_1, C_2] + [E_2, C_1] = 0 \quad \text{and} \quad [C_1, C_2] = E_1 \wedge C_2 + E_2 \wedge C_1.$$

There are some equivalent ways of expressing the compatibility of two Jacobi structures on a manifold [NdC 98]. But the study of the compatibility of two Jacobi structures on a manifold, can also be done using the Lichnerowicz-Jacobi cohomology. If  $(M, C, E)$  is a Jacobi manifold, let us denote by  $A^k(M)$  the space of skew-symmetric contravariant tensor fields of order  $k$  ( $k$ -tensors) on  $M$  and define the differential operator

$$\sigma : A^k(M) \rightarrow A^{k+1}(M), \quad \sigma(P) = -[C, P] + kE \wedge P. \quad (3)$$

The restriction of  $\sigma$  to the subspace

$$A_I^k(M) = \{P \in A^k(M) : \mathcal{L}_E P = 0\}$$

of the invariant  $k$ -tensors with respect to the vector field  $E$  is a *cohomology operator* on the Jacobi manifold and the resultant cohomology is called the *Lichnerowicz-Jacobi cohomology* of  $M$  [Le-Ma-Pa 97].

**Proposition 1.2.** *Two Jacobi structures  $(C_1, E_1)$  and  $(C_2, E_2)$  on a manifold  $M$  are compatible if and only if*

$$\sigma_1(C_2) = -\sigma_2(C_1) \quad \text{and} \quad \sigma_1(E_2) = -\sigma_2(E_1),$$

where  $\sigma_i, i = 1, 2$ , are the cohomology operators of the Lichnerowicz-Jacobi cohomology of  $M$ , with respect to both Jacobi structures.

*Proof.* A direct computation using Proposition 1.1 and the definition (3) of the cohomology operators  $\sigma_i, i = 1, 2$ , gives the desired result.  $\square$

In [NdC 98] it was proved that two conformally equivalent Jacobi structures on  $M$  are compatible. Another way of obtaining compatible Jacobi structures uses the Lie derivative on the direction of some vector field.

**Proposition 1.3.** *Let  $X$  be a vector field on the Jacobi manifold  $(M, C, E)$  such that*

$$\mathcal{L}_X(\mathcal{L}_X C) = 0 \quad \text{and} \quad \mathcal{L}_X(\mathcal{L}_X E) = 0.$$

*Then the pair  $(C_1, E_1) = (\mathcal{L}_X C, \mathcal{L}_X E)$  defines a new Jacobi structure on  $M$  which is compatible with  $(C, E)$ .*

*Proof.* With  $E_1 = \mathcal{L}_X E = [X, E]$ , we have  $\mathcal{L}_{E_1} C = -\mathcal{L}_E(\mathcal{L}_X C)$ , that is

$$\mathcal{L}_{E_1} C + \mathcal{L}_E C_1 = 0. \quad (4)$$

So,

$$\begin{aligned}\mathcal{L}_{E_1}C_1 &= \mathcal{L}_X(\mathcal{L}_{E_1}C) \\ &= -\mathcal{L}_X(\mathcal{L}_E C_1) \\ &= -\mathcal{L}_{[X,E]}(\mathcal{L}_X C) \\ &= -\mathcal{L}_{E_1}C_1,\end{aligned}$$

and

$$\mathcal{L}_{E_1}C_1 = 0. \quad (5)$$

On the other hand,

$$\begin{aligned}\mathcal{L}_X([C, C]) &= [\mathcal{L}_X C, C] + [C, \mathcal{L}_X C] \\ &= 2[C_1, C]\end{aligned}$$

and

$$\mathcal{L}_X(E \wedge C) = E_1 \wedge C + E \wedge C_1;$$

so,

$$[C_1, C] = E_1 \wedge C + E \wedge C_1. \quad (6)$$

Because  $\mathcal{L}_X E_1 = \mathcal{L}_X C_1 = 0$ , if we take the Lie derivative on the direction of  $X$  of both members of (6), we obtain

$$[C_1, C_1] = 2E_1 \wedge C_1. \quad (7)$$

The equalities (5) and (7) prove that  $(C_1, E_1)$  is a Jacobi structure, while (4) and (6) show the compatibility.  $\square$

We can also prove the following result.

**Proposition 1.4.** *Let  $X_1$  and  $X_2$  be two vector fields on the Jacobi manifold  $(M, C, E)$  such that*

$$[X_1, X_2] = 0, \quad [X_1, \mathcal{L}_{X_2}C] = 0, \quad [X_1, \mathcal{L}_{X_2}E] = 0$$

and

$$\mathcal{L}_{X_i}(\mathcal{L}_{X_i}C) = 0, \quad \mathcal{L}_{X_i}(\mathcal{L}_{X_i}E) = 0, \quad i = 1, 2.$$

Then

$$(C_1, E_1) = (\mathcal{L}_{X_1}C, \mathcal{L}_{X_1}E) \quad \text{and} \quad (C_2, E_2) = (\mathcal{L}_{X_2}C, \mathcal{L}_{X_2}E)$$

are compatible Jacobi structures on  $M$ .

*Proof.* We only have to check the compatibility. By Proposition 1.3, we know that the Jacobi structures  $(C, E)$  and  $(C_2, E_2)$  are compatible. Then,

$$[C, C_2] = E \wedge C_2 + E_2 \wedge C \quad \text{and} \quad [E, C_2] + [E_2, C] = 0. \quad (8)$$

If we take the Lie derivative, on the direction of  $X_1$ , of both members of equalities (8), we obtain

$$[C_1, C_2] = E_1 \wedge C_2 + E_2 \wedge C_1 \quad \text{and} \quad [E_1, C_2] + [E_2, C_1] = 0. \quad (9)$$

$\square$

## 2. Homogeneous Poisson-Nijenhuis manifolds

Let  $M$  be a manifold endowed with two Jacobi structures  $(C_1, E_1)$  and  $(C_2, E_2)$ . Take the corresponding homogeneous Poisson structures on  $\mathbb{R} \times M$ ,  $(\Lambda_i = \exp(-t)(C_i + \frac{\partial}{\partial t} \wedge E_i), \frac{\partial}{\partial t})$ ,  $i = 1, 2$ .

**Proposition 2.1.** [NdC 98] *The Jacobi structures  $(C_1, E_1)$  and  $(C_2, E_2)$  on  $M$  are compatible if and only if  $\Lambda_1$  and  $\Lambda_2$  are compatible Poisson tensors on  $\mathbb{R} \times M$ .*

Using this result, we want to study the relationship between two compatible Jacobi structures on  $M$  and their associated (compatible) homogeneous Poisson structures on  $\mathbb{R} \times M$ , in the case where these Poisson structures are related by a Nijenhuis tensor.

Let  $\bar{N}$  be a  $(1, 1)$ -tensor on  $\mathbb{R} \times M$  such that  $\mathcal{L}_{\frac{\partial}{\partial t}} \bar{N} = 0$ . Then,  $\bar{N}$  is given by

$$\bar{N} = N + Y \otimes dt + \frac{\partial}{\partial t} \otimes \gamma + g \frac{\partial}{\partial t} \otimes dt, \tag{10}$$

where  $N$  is a  $(1, 1)$ -tensor on  $M$ ,  $Y$  is a vector field on  $M$ ,  $\gamma$  is a 1-form on  $M$  and  $g \in C^\infty(M, \mathbb{R})$ . Reciprocally, if  $\bar{N}$  is given by (10), then  $\mathcal{L}_{\frac{\partial}{\partial t}} \bar{N} = 0$ .

### Remarks

1. The image of a vector field on  $\mathbb{R} \times M$  of the form  $\bar{X} = \exp(-t)(f \frac{\partial}{\partial t} + X)$ , where  $X$  is a vector field on  $M$  and  $f \in C^\infty(M, \mathbb{R})$ , by the  $(1, 1)$ -tensor given by (10) is a vector field on  $\mathbb{R} \times M$  of the same type of  $\bar{X}$ .
2. If  $\Lambda$  is the homogeneous Poisson tensor on  $\mathbb{R} \times M$ , given by (2), then  $\bar{N}\Lambda$  is a homogeneous bivector on  $\mathbb{R} \times M$ :  $\mathcal{L}_{\frac{\partial}{\partial t}}(\bar{N}\Lambda) = -\bar{N}\Lambda$

**Proposition 2.2.** *Let  $\bar{N}$  be a  $(1, 1)$ -tensor on  $\mathbb{R} \times M$  given by (10). Then  $\bar{N}$  is a Nijenhuis tensor on  $\mathbb{R} \times M$  if and only if*

- i)  $\tau(N) = Y \otimes d\gamma$ ;
- ii)  $\mathcal{L}_N \gamma = g d\gamma$ ;
- iii)  $\mathcal{L}_Y N = -Y \otimes dg$ ;
- iv)  ${}^t N(dg) = \mathcal{L}_Y \gamma + g dg$ .

*Proof.* First, remark that if  $X$  is a vector field on  $M$ , then  $\bar{N}X = NX + \langle \gamma, X \rangle \frac{\partial}{\partial t}$  and that  $\bar{N}(\frac{\partial}{\partial t}) = Y + g \frac{\partial}{\partial t}$ . If  $X_1$  and  $X_2$  are vector fields on  $M$ , then

$$\begin{aligned} \tau(\bar{N})(X_1, X_2) &= \tau(N)(X_1, X_2) + (X_2 \cdot \langle \gamma, X_1 \rangle - X_1 \cdot \langle \gamma, X_2 \rangle \\ &\quad + \langle \gamma, [X_1, X_2] \rangle) \bar{N}(\frac{\partial}{\partial t}) + ((NX_1) \cdot \langle \gamma, X_2 \rangle - (NX_2) \cdot \langle \gamma, X_1 \rangle \\ &\quad - \langle \gamma, [NX_1, X_2] \rangle - \langle \gamma, [X_1, NX_2] \rangle - \langle \gamma, N[X_1, X_2] \rangle) \frac{\partial}{\partial t} \\ &= \tau(N)(X_1, X_2) - d\gamma(X_1, X_2)Y - (gd\gamma(X_1, X_2) - \mathcal{L}_N \gamma(X_1, X_2)) \frac{\partial}{\partial t}. \end{aligned}$$

So,  $\tau(\bar{N})(X_1, X_2) = 0$  if and only if

$$\tau(N)(X_1, X_2) = (Y \otimes d\gamma)(X_1, X_2) \quad (11)$$

and

$$\mathcal{L}_N\gamma(X_1, X_2) = gd\gamma(X_1, X_2). \quad (12)$$

Let  $X$  be a vector field on  $M$ . Then,

$$\begin{aligned} \tau(\bar{N})(X, \frac{\partial}{\partial t}) &= [NX, Y] + ((NX).g)\frac{\partial}{\partial t} - (Y. \langle \gamma, X \rangle)\frac{\partial}{\partial t} \\ &\quad - N[X, Y] - \langle \gamma, [X, Y] \rangle \frac{\partial}{\partial t} - (X.g)(Y + g\frac{\partial}{\partial t}) \\ &= -(\mathcal{L}_Y N)(X) - (Y \otimes dg)(X) \\ &\quad + (\langle {}^t Ndg, X \rangle - \langle \mathcal{L}_Y \gamma, X \rangle - \langle gdg, X \rangle)\frac{\partial}{\partial t}. \end{aligned}$$

So,  $\tau(\bar{N})(X, \frac{\partial}{\partial t}) = 0$  if and only if

$$(\mathcal{L}_Y N)(X) = -(Y \otimes dg)(X) \quad (13)$$

and

$$\langle {}^t Ndg - \mathcal{L}_Y \gamma - gdg, X \rangle = 0. \quad (14)$$

Equalities (11), (12), (13) and (14) end the proof.  $\square$

Let us take the homogeneous Poisson tensor  $\Lambda = \exp(-t)(C + \frac{\partial}{\partial t} \wedge E)$  on  $(\mathbb{R} \times M)$ .

**Lemma 2.1.** *With the notations of Proposition 2.2,  $\bar{N}\Lambda = \Lambda {}^t \bar{N}$  if and only if*

- i)  $NE = C^\#(\gamma) + gE$ ;
- ii)  $NC - C {}^t N = E \otimes Y + Y \otimes E$ ;
- iii)  $\langle \gamma, E \rangle = 0$ .

*Proof.* Let  $\alpha$  be any 1-form on  $M$ . Then,

$$\begin{aligned} \bar{N}(\Lambda^\#(\alpha)) &= \exp(-t)(\bar{N}(C^\#(\alpha) - \langle \alpha, E \rangle \frac{\partial}{\partial t})) \\ &= \exp(-t)(N(C^\#(\alpha)) - \langle \alpha, E \rangle Y - (\langle \alpha, C^\#(\gamma) \rangle + \langle \alpha, E \rangle g)\frac{\partial}{\partial t}) \end{aligned}$$

and, taking account that

$${}^t \bar{N}(\alpha) = {}^t N\alpha + \langle \alpha, Y \rangle dt,$$

we compute

$$\Lambda^\#({}^t \bar{N}\alpha) = \exp(-t)(C^\#({}^t N\alpha) + \langle \alpha, Y \rangle E - \langle \alpha, NE \rangle \frac{\partial}{\partial t})$$

and we conclude that  $\bar{N}\Lambda^\#(\alpha) = \Lambda^\#({}^tN\alpha)$  if and only if

$$\langle \alpha, NE \rangle = \langle \alpha, C^\#(\gamma) + gE \rangle \quad (15)$$

and

$$(NC^\# - Y \otimes E)\alpha = (C^\#{}^tN + E \otimes Y)\alpha. \quad (16)$$

Equalities (15) and (16) give conditions i) and ii).

On the other hand,

$$\bar{N}(\Lambda^\#(dt)) = \exp(-t)(NE + \langle \gamma, E \rangle \frac{\partial}{\partial t}) \quad (17)$$

and, because  ${}^t\bar{N}(dt) = \gamma + gdt$ ,

$$\Lambda^\#({}^t\bar{N}(dt)) = \exp(-t)(C^\#(\gamma) + gE). \quad (18)$$

Once the right members of (17) and (18) are equal, we obtain condition iii).  $\square$

Let us take the Magri-Morosi concomitant  $\mathcal{R}(\Lambda, \bar{N})$  of  $\Lambda$  and  $\bar{N}$ , (1).

**Lemma 2.2.** *With the notations of Proposition 2.2,  $\mathcal{R}(\Lambda, \bar{N}) = 0$  if and only if*

- i)  $NE + \mathcal{L}_Y E = gE - C^\#(dg)$
- ii)  $\mathcal{L}_Y C = gC - NC + Y \otimes E$
- iii)  $(\mathcal{L}_E N)X + \langle \gamma, X \rangle E = C^\#(i_X d\gamma) + (X.g)E$
- iv)  $\mathcal{R}(C, N)(df, X) = (X.(Y.f))E - (X.(E.f))Y - \langle \gamma, X \rangle C^\#(df)$ ,

where  $X$  is any vector field on  $M$  and  $f$  is any function of  $C^\infty(M, \mathbb{R})$ .

*Proof.* Let  $f$  be any function of  $C^\infty(M, \mathbb{R})$ . The component of the vector field  $\mathcal{R}(\Lambda, \bar{N})(df, \frac{\partial}{\partial t})$  on  $\frac{\partial}{\partial t}$  is

$$\exp(-t)(\langle df, C^\#(dg) + \mathcal{L}_Y E + NE - gE \rangle). \quad (19)$$

The other components (without  $\frac{\partial}{\partial t}$ ) of  $\mathcal{R}(\Lambda, \bar{N})(df, \frac{\partial}{\partial t})$ , obtained from the computation of  $\langle dh, \mathcal{R}(\Lambda, \bar{N})(df, \frac{\partial}{\partial t}) \rangle$ , for any  $h \in C^\infty(M, \mathbb{R})$ , are

$$\exp(-t)(\langle dh, (-\mathcal{L}_Y C)^\#(df) + gC^\#(df) - N(C^\#(df)) + (E.f)Y \rangle). \quad (20)$$

From (19) and (20), we conclude that  $\mathcal{R}(\Lambda, \bar{N})(df, \frac{\partial}{\partial t}) = 0$  gives conditions i) and ii) of the Lemma.

On the other hand, if  $X$  is any vector field on  $M$ , the component of the vector field  $\mathcal{R}(\Lambda, \bar{N})(df, X)$  on  $\frac{\partial}{\partial t}$  is

$$\exp(-t)(\langle df, -(\mathcal{L}_E N)X - \langle \gamma, X \rangle E + C^\#(i_X d\gamma) + (X.g)E \rangle, \quad (21)$$

while the components without  $\frac{\partial}{\partial t}$ , obtained from the computation of  $\langle dh, \mathcal{R}(\Lambda, \bar{N})(df, X) \rangle$ , for any  $h \in C^\infty(M, \mathbb{R})$ , are

$$\exp(-t)(\langle dh, \mathcal{R}(C, N)(df, X) + \langle \gamma, X \rangle C^\#(df) + (X.(E.f))Y - (X.(Y.f))E \rangle. \quad (22)$$

From (21) and (22), we conclude that  $\mathcal{R}(\Lambda, \bar{N})(df, X) = 0$  gives conditions iii) and iv) of the Lemma.  $\square$

**Theorem 2.1.** *Let  $\Lambda = \exp(-t)(C + \frac{\partial}{\partial t} \wedge E)$  and  $\bar{N} = N + Y \otimes dt + \frac{\partial}{\partial t} \otimes \gamma + g \frac{\partial}{\partial t} \otimes dt$ , be respectively the homogeneous Poisson tensor and the Nijenhuis tensor on  $\mathbb{R} \times M$ . Then, the triple  $(\mathbb{R} \times M, \Lambda, \bar{N})$  is a Poisson-Nijenhuis manifold if and only if conditions i), ii) and iii) of Lemma 2.1 and i) – iv) of Lemma 2.2 hold. Moreover, the homogeneous Poisson tensor  $\bar{N}\Lambda$  is given by  $\exp(-t)(C_1 + \frac{\partial}{\partial t} \wedge E_1)$ , where*

$$(C_1, E_1) = (gC - \mathcal{L}_Y C, gE - C^\#(dg) - \mathcal{L}_Y E) \quad (23)$$

is a Jacobi structure on  $M$ , compatible with  $(C, E)$ .

*Proof.* We only have to find the expressions of  $C_1$  and  $E_1$ . For any pair  $(f, h)$  of functions on  $M$ ,

$$\langle dh, (\bar{N}\Lambda)^\#(df) \rangle = \exp(-t) \langle dh, NC^\#(df) - (Y \otimes E)(df) \rangle \quad (24)$$

and

$$\langle dh, (\Lambda^\#({}^t\bar{N}))^\#(df) \rangle = \exp(-t) \langle dh, C^\#({}^tN(df)) + (E \otimes Y)(df) \rangle. \quad (25)$$

Since  $\bar{N}\Lambda = \frac{1}{2}(\bar{N}\Lambda + \Lambda {}^t\bar{N})$ , we obtain from (24) and (25), using conditions ii) of Lemma 2.1 and ii) of Lemma 2.2,

$$\begin{aligned} (\bar{N}\Lambda)(df, dh) &= \exp(-t)(NC - Y \otimes E)(df, dh) \\ &= \exp(-t)(gC - \mathcal{L}_Y C)(df, dh). \end{aligned} \quad (26)$$

Also,

$$\langle dh, \bar{N}(\Lambda^\#(dt)) \rangle = \exp(-t) \langle dh, NE \rangle \quad (27)$$

and

$$\langle dh, \Lambda^\#({}^t\bar{N}(dt)) \rangle = \exp(-t) \langle dh, C^\#(\gamma) + gE \rangle. \quad (28)$$

From (27) and (28), and taking into account conditions i) of Lemma 2.1 and i) of Lemma 2.2, we conclude that

$$\begin{aligned} (\bar{N}\Lambda)(dt, dh) &= \exp(-t)NE(dt, dh) \\ &= \exp(-t)(gE - C^\#(dg) - \mathcal{L}_Y E)(dt, dh). \end{aligned} \quad (29)$$

$\square$

**Acknowledgements.** The author wishes to thank Professors C.-M. Marle and J. Monterde for helpful discussions.



## References

- [KSch-Ma 90] Kosmann-Schwarzbach Y. and Magri F.: *Poisson-Nijenhuis structures*. Ann. Inst. Henri Poincaré **53** (1990), 35.
- [Le-Ma-Pa 97] León M. de, Marrero J. C. and Padrón E.: *Lichnerowicz-Jacobi cohomology*. J. Phys. A: Math. Gen. **30** (1997), 6029.
- [Lich 78] Lichnerowicz A.: *Les variétés de Jacobi et leurs algèbres de Lie associées*. J. Math. pures et appl. **57** (1990), 453.
- [Ma-Mo 84] Magri F. and Morosi C.: *A Geometrical characterization of integrable Hamiltonian systems through the theory of Poisson-Nijenhuis manifolds*. Quaderno S19, Univ. of Milan, 1984.
- [NdC 98] Nunes da Costa J. M.: *Compatible Jacobi manifolds: geometry and reduction*. J. Phys. A: Math. Gen. **31** (1998), 1025.