



NEW RESULTS ON THE GEOMETRY OF TRANSLATION SURFACES*

MARIAN I. MUNTEANU and ANA I. NISTOR

*Department of Mathematics, University Al. I. Cuza of Iași, Bd. Carol I, n. 11
Iași 700506, Romania*

Abstract. In this paper we study the second mean curvature for different hypersurfaces in space forms. We furnish some examples and we remind some connections between II -minimality and biharmonicity. The main result consists in proving that there are no II -minimal translation surfaces in the Euclidean three-space.

1. Introduction

The study of the second fundamental form II was initiated in the early papers of Weingarten [16], Darboux [5] and Cartan [3] where appeared for the first time notions like connection or curvature associated to II . Later on, Erard [7] introduced the second fundamental form as metric on the surface. This is possible only when II is non-degenerate and hence it can be regarded as a (pseudo)-Riemannian metric on the surface. At this point one can consider a connected smooth surface M endowed with II as metric in order to study new characteristics associated to (M, II) . In the classical case when the metric on the surface is given by the first fundamental form I , i.e., for (M, I) , there are well known formulae to compute the **Gaussian curvature** K and the **mean curvature** H in order to analyze the properties of M that arise from this “measures”. In a similar manner, the **second Gaussian curvature** denoted by K_{II} and the **second mean curvature**, denoted by H_{II} , were considered. In [3], K_{II} was introduced for the first time by Cartan, as the analogous of the Gaussian curvature. Concerning H_{II} , it was defined by Glässner in [8]. An overview over the literature dedicated to the second fundamental form and the associated curvatures for different type of submanifolds in different ambient spaces can be found in [15] and its references. Regarding the second

*Reprinted from JGSP 18 (2010) 49–62.

mean curvature, the critical points of the area functional of the second fundamental form are those surfaces for which the mean curvature of the second fundamental form vanishes. A non-developable surface is said to be **II-flat** if $K_{II} = 0$ and respectively **II-minimal** if $H_{II} = 0$. Consequently, the well known result that there are no compact minimal submanifolds in the Euclidean space was discussed also in the case of the second mean curvature and in [8] it is proved that *do not exist compact II-minimal surfaces in Euclidean space*. Despite this non-existence result confirmed also in some other particular ambient spaces, in [15] it is proved that compact II-minimal surfaces may exist in some general ambient spaces.

In the present paper we are interested in the study of the second mean curvature for different examples of submanifolds. More exactly we study the II-minimality property, equivalently, the condition $H_{II} = 0$. The classical examples in the theory of harmonic and biharmonic maps are proved to be also interesting examples concerning the II-minimality property. Let us remind the following situations of classical biharmonic maps which are also II-minimal, namely the standard embedding $\mathbb{S}^n(\frac{1}{\sqrt{2}}) \subset \mathbb{S}^{n+1}$ and the hypersurface $\mathbb{S}^k(\frac{1}{\sqrt{2}}) \times \mathbb{S}^{n-k}(\frac{1}{\sqrt{2}}) \subset \mathbb{S}^{n+1}$ for $k = 1, \dots, n-1$ (see e.g., [4], [10] and [15]). In the sequel we formulate some generalizations of these results.

Returning to the theory of surfaces, in [11] it is proved that a ruled surface with nowhere vanishing Gaussian curvature is II-minimal if and only if it is a piece of helicoid. A study on II-minimal affine translation surfaces written as a sum of two curves is contained in [12]. Here it is stated that *there are no affine II-minimal translation surfaces of this type*. An interesting situation occurs when the two curves are situated in orthogonal planes. This is our case of translation surfaces. One can retrieve the same non-existence result concerning II-minimal translation surfaces in the Euclidean three-space. The aim of this article is to give a proof of this statement. It is contained in Section 3, after we acquaint the reader with the basic notions about translation surfaces and II-minimality in the Preliminaries of this article. Moreover, we provide also interesting examples of II-minimal hypersurfaces in different space forms.

2. Preliminaries

In the general theory of surfaces and hypersurfaces generically denoted by M and isometrically immersed in some ambient space $(\widetilde{M}, \widetilde{g})$, one can associate different “measures”. Naturally, a way to describe a **metric** g on M is taking the restriction of the metric \widetilde{g} from the ambient space. If the immersion $(M, g) \hookrightarrow (\widetilde{M}, \widetilde{g})$ has codimension one, we write the Gauss and Weingarten formulas

$$\begin{aligned}\tilde{\nabla}_X Y &= \nabla_X Y + II(X, Y) N \\ \tilde{\nabla}_X N &= -AX\end{aligned}$$

for every X, Y tangent to M . The corresponding Levi-Civita connections on the ambient space and on the surface are denoted by $\tilde{\nabla}$ and ∇ , respectively. Moreover II is a symmetric $(1, 2)$ -tensor field called *the second fundamental form* of the surface M and A is a symmetric $(1, 1)$ -tensor field known as *the shape operator* associated to the unit normal to the surface N . The following relation holds $II(X, Y) = g(X, AY)$, where X, Y are vector fields tangents to M .

Concerning the curvature tensor R on the surface and using the previous notations, recall that we use the following sign convention $R(X, Y) = [\nabla_X, \nabla_Y] - \nabla_{[X, Y]}$ for any $X, Y \in T(M)$.

Having now some basic working tools on M we can construct its intrinsic and extrinsic geometry by means of the characterization of the curvatures. The most used metrics on a surface is given by the first fundamental form I associated to the immersion which gives the parametrization. But, one can think of rebuilding all the geometry corresponding to I by taking the second fundamental form as a new metric on the surface. One elementary condition that II must satisfy consists of non-degeneracy, namely the surface must be non-developable.

2.1. Translation Surfaces

Let us consider a surface having the Cartesian parametrization given by

$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto A(x) \begin{pmatrix} \mathbf{f}(y) \\ \mathbf{g}(y) \\ \mathbf{h}(y) \end{pmatrix} + \begin{pmatrix} a(x) \\ b(x) \\ c(x) \end{pmatrix}$$

where $A(x) \in \text{SO}(3)$ (sometimes in $\text{O}(3)$). This surface represents a union of “equal” curves i.e., it is the image of one curve, called **generatrix**, obtained by isometries of the space. Some authors call this kind of surface a *surface of Darboux*. Some known examples are to be mentioned, namely

1. $A = I_3$: **translation surfaces**
2. $A =$ matrix of rotation (the axe and the angle are fixed), $a = b = c = 0$: **rotation surfaces**
3. $A =$ matrix of rotation (the axe \bar{n} and the angle are fixed), $(a, b, c) = x \bar{n}$: **helicoidal surfaces**

If the generatrix is

- a. a straight line : **ruled surfaces**

- b. a circle : **circled surfaces** including, e.g. tubes. For a smooth curve γ , the tube of unit radius around it is given by

$$r(s, t) = \gamma(t) + \cos s N(t) + \sin s B(t)$$

where s is the arclength parameter, $N(s)$ and $B(s)$ are respectively the normal and the binormal of the curve. As a Darboux surface, a tube can be written as

$$r(s, t) = \gamma(t) + A(t) \mathbb{S}^1$$

where by \mathbb{S}^1 we mean the unit circle.

The special Euclidean group of the n -dimensional space or the Euclidean motion group $\text{SE}(n)$ is the semi-direct product of \mathbb{R}^n with the special orthogonal group $\text{SO}(n)$. In the three-dimensional case $\text{SE}(3) = \mathbb{R}^3 \times \text{SO}(3)$. To be more precise, for two elements $h = (a, A)$ and $h' = (b, B)$ in $\text{SE}(3)$, the group multiplication and the inverse are given by $h \circ h' = (a + Ab, AB)$, respectively $h^{-1} = (-A^T a, A^T)$. It is also possible to represent any element of $\text{SE}(n)$ as an $(n + 1) \times (n + 1)$ homogeneous transformation matrix of the form $\begin{pmatrix} A & a \\ O & 1 \end{pmatrix}$ (see e.g. [14], p.45).

A surface of Darboux can be thought as the action of the one-parameter family of matrices in $\text{SE}(3)$ to a given curve.

As we have seen, a translation surface is a “sum” of two curves. If the two curves are situated in orthogonal planes the surface can be represented as

$$(x, y) \longmapsto (x, y, f(x) + g(y)). \quad (1)$$

Examples: planes, cylinders, hyperbolic and elliptic paraboloids, the egg box surface, Scherk surface (the only minimal translation surface in \mathbb{E}^3).

2.2. II-minimality

We dedicate this section to some very nice examples of II -minimal hypersurfaces immersed in the sphere \mathbb{S}^n and we give a nonexistence result in the case of II -minimal hypersurfaces in the hyperbolic space \mathbb{H}^n . The motivation comes from another interesting property of these immersions intensively studied in recent years, namely the biharmonicity. Similar to the variational characterization of the mean curvature H , the curvature of the second fundamental form, denoted by H_{II} is introduced as a measure for the rate of change of the II -area under a normal deformation. Let us denote by M a m -dimensional hypersurface in a semi-Riemannian manifold $(\widetilde{M}, \tilde{g})$ with the second fundamental form as semi-Riemannian metric. Accordingly (see [15]), the mean curvature associated to the second mean curvature H_{II} is given by

$$H_{II} = \frac{1}{2} \left(mH - \sum_{i=1}^m \tilde{g}(\tilde{R}(V_i, N)V_i, N)\kappa_i + \frac{\alpha}{2} \Delta_{II} \log |\det A| - \alpha \operatorname{div}_{II} \mathcal{Z} \right).$$

Let us explain in few words the notations used above. The classical mean curvature of the first fundamental form is denoted by H , the unit normal is called N and by A we mean the shape operator of M . Moreover, V_i , $i = 1, \dots, m$ form an orthonormal basis on M with respect to II and let $\kappa_i = II(V_i, V_i) = \pm 1$, $i = 1, \dots, m$. The vector field \mathcal{Z} in $T(M)$ is constructed as $\mathcal{Z} = \operatorname{tr}_{II} B = \sum_{i=1}^m B(V_i, V_i)\kappa_i$, where the tensor $B : T(M) \times T(M) \rightarrow T(M)$ is defined by $(V, W) \mapsto A^{-1} \left\{ \left(\tilde{R}(V, N)W \right)^{\mathfrak{t}} \right\}$. Here, we denote by \mathfrak{t} the tangential component of the corresponding vector field. The shape operator is thought here as a field of endomorphisms of each tangent spaces in points of M , namely $A : T(M) \rightarrow T(M)$, $V \mapsto -\tilde{\nabla}_V N$.

If the ambient space is a space form (its sectional curvature is constant), then the tensor B vanishes and hence $\mathcal{Z} = 0$. Moreover, if in addition the shape operator has constant determinant (and this often happens) the second mean curvature can be computed by using an easier formula, namely

$$H_{II} = \frac{1}{2} \left(mH - \sum_{i=1}^m \tilde{g}(\tilde{R}(V_i, N)V_i, N)\kappa_i \right). \quad (2)$$

In the sequel we present some examples of II -minimal hypersurfaces in spheres.

Example 1. The standard embedding $\mathbb{S}^{n-1}(r) \hookrightarrow \mathbb{S}^n(1)$ is II -minimal if and only if $r = \frac{1}{\sqrt{2}}$.

Proof: Let (x^0, x^1, \dots, x^n) be global coordinates in \mathbb{E}^{n+1} and denote by p either the point or the position vector in \mathbb{E}^{n+1} . Without loss of the generality one can think the sphere $\mathbb{S}^{n-1}(r)$ as obtained by cutting the unit sphere \mathbb{S}^n by the hyperplane $x^0 = \sqrt{1-r^2}$. Thus, an arbitrary vector field tangent to $\mathbb{S}^{n-1}(r)$ can be expressed as $X = (0, X^1, \dots, X^n)$, where X^i are smooth functions depending on x^1, \dots, x^n such that $\sum_{i=1}^n X^i x^i = 0$. In order to compute the second mean curvature, one can express first the unit normal of the embedding as $\nu = \pm \left(-r, \frac{\sqrt{1-r^2}}{r} (x^1, \dots, x^n) \right)$. Fixing an orientation we choose, e.g. the “plus” sign. The second fundamental form is given by $II(X, Y) = -\frac{\sqrt{1-r^2}}{r} \langle X, Y \rangle$ for all $X, Y \in T(\mathbb{S}^{n-1}(r))$, where $\langle \cdot, \cdot \rangle$ denotes the usual Euclidean scalar product. Note that II is negatively defined. We are able now to obtain the second mean

curvature. After straightforward computations in (2) one gets

$$H_{II} = \frac{1}{2} \left(\frac{r}{\sqrt{1-r^2}} - \frac{\sqrt{1-r^2}}{r} \right).$$

Under the assumption of II -minimality, $H_{II} = 0$ is equivalent with $r = \frac{1}{\sqrt{2}}$. Hence the conclusion. □

More generally we have

Example 2. The embedding $\mathbb{S}^{n-1}(r) \hookrightarrow \mathbb{S}^n(R)$ is II -minimal if and only if $r = \frac{R}{\sqrt{R^2+1}}$.

Proof: Similar computations as in previous example. □

Remark 1. The following chain of embeddings

$$\mathbb{S}^1\left(\frac{1}{\sqrt{n}}\right) \hookrightarrow \mathbb{S}^2\left(\frac{1}{\sqrt{n-1}}\right) \hookrightarrow \dots \hookrightarrow \mathbb{S}^{n-1}\left(\frac{1}{\sqrt{2}}\right) \hookrightarrow \mathbb{S}^n(1)$$

is such that each embedding $\mathbb{S}^k\left(\frac{1}{\sqrt{n-k+1}}\right) \hookrightarrow \mathbb{S}^{k+1}\left(\frac{1}{\sqrt{n-k}}\right)$ is II -minimal for any $k \in \{1, \dots, n-1\}$.

Let us give another example of a II -minimal surface in the unit sphere \mathbb{S}^3 .

Example 3. Let us consider the following parametrization

$$r : M \longrightarrow \mathbb{S}^3(1), \quad r(s, t) = (\cos s \cos t, \sin s \cos t, \cos s \sin t, \sin s \sin t).$$

Then second mean curvature of the surface M vanishes identically.

Proof: The proof of this statement is straightforward. □

The next example ends the series of II -minimal surfaces in spheres presented in this paper.

Example 4. Let $M = \mathbb{S}^{m_1}(r_1) \times \mathbb{S}^{m_2}(r_2) \hookrightarrow \mathbb{S}^{m+1}(r) = \widetilde{M}$ be the usual embedding with $r_1^2 + r_2^2 = r^2$ and $m_1 + m_2 = m$. Then, M is II -minimal in \widetilde{M} if and only if $r_1 = r \sqrt{\frac{m_1+m_2r^2}{m(r^2+1)}}$ and $r_2 = r \sqrt{\frac{m_2+m_1r^2}{m(r^2+1)}}$.

Proof: Consider $(x^0, x^1, \dots, x^{m_1})$ and $(y^0, y^1, \dots, y^{m_2})$ global coordinates on \mathbb{R}^{m_1+1} , respectively on \mathbb{R}^{m_2+1} . The unit normal is $\nu = \frac{1}{r} \left(-\frac{r_2}{r_1} x, \frac{r_1}{r_2} y \right)$ where

$x = (x^0, x^1, \dots, x^{m_1})$ and $y = (y^0, y^1, \dots, y^{m_2})$. Hence the shape operator can be expressed as

$$A = \begin{pmatrix} \frac{r_2}{rr_1} I_{m_1} & O \\ O & -\frac{r_1}{rr_2} I_{m_2} \end{pmatrix}$$

whose determinant is constant. By I_k we denoted the identity $k \times k$ matrix. After straightforward computations we get from (2) that the second mean curvature is given by

$$H_{II} = \frac{1}{2rr_1r_2} (m_1r_2^2 - m_2r_1^2 - m_1r^2r_1^2 + m_2r^2r_2^2).$$

Hence the conclusion. \square

Remark 2. In particular, if $m = 2$ and $r = 1$ we get that the Clifford torus

$$\mathbb{S}^1 \left(\frac{1}{\sqrt{2}} \right) \times \mathbb{S}^1 \left(\frac{1}{\sqrt{2}} \right) \hookrightarrow \mathbb{S}^3(1)$$

is II -minimal.

Proof: See for example [9]. \square

Having in mind these results, there is another interesting property involving the curvatures of a surface that we can study, the Weingarten property. If A, B are two different type curvatures of a (non-developable) surface, and if there is a non-trivial functional relation between A and B , then the surface is called an $\{A, B\}$ -generalized Weingarten surface. See for details [6].

Remark 3. The generalized Clifford torus $\mathbb{S}^1(r_1) \times \mathbb{S}^1(r_2) \hookrightarrow \mathbb{S}^3(1)$, with $r_1^2 + r_2^2 = 1$ is a $\{H, H_{II}\}$ -generalized Weingarten surface.

Proof: Easy computations yield the following relation: $H_{II} = 2H$, namely, the two mean curvatures corresponding to the first and to the second fundamental forms respectively, are proportional. \square

Seeing all these nice examples for spheres, we wonder what happens when the ambient is the hyperbolic space. At this point, if we consider similar problems, e.g. II -minimality, in the hyperbolic spaces, we get the following non-existence result

Proposition 1. There is no $r > 0$ such that $\mathbb{H}^{n-1}(-r) \hookrightarrow \mathbb{H}^n(-1)$ is II -minimal. Here $\mathbb{H}_R^k := \mathbb{H}^k(-R) = \{x \in \mathbb{R}_1^{k+1}; \langle x, x \rangle_1 = -R^2, x_0 > 0\}$ ($R > 0$) where $\langle \cdot, \cdot \rangle_1$ denotes the usual **Lorentzian scalar product** with signature $(- + \dots +)$.

Proof: After similar computations as in the previous examples we find the expression of the second mean curvature, but under the restriction of II -minimality we reach a contradiction! \square

Having in mind all these examples in spheres and in hyperbolic spaces, let us recall now another interesting property for surfaces and hypersurfaces, the *biharmonicity*. As the aim of our article does not consist in the study of biharmonicity, the reader is invited to check [1, 2] for more details on the subject.

In the end of this section we would like to bring into attention some classical results concerning the biharmonicity of the surfaces and hypersurfaces studied above from the II -minimality point of view. Concerning the spheres, it is known that the proper biharmonic surfaces in \mathbb{S}^3 are also II -minimal surfaces. Moreover, the hyperspheres $\mathbb{S}^m\left(\frac{1}{\sqrt{2}}\right)$ and the generalized Clifford tori $\mathbb{S}^{m_1}\left(\frac{1}{\sqrt{2}}\right) \times \mathbb{S}^{m_2}\left(\frac{1}{\sqrt{2}}\right)$, $m_1 \neq m_2$ are the only known examples of proper biharmonic hypersurfaces in \mathbb{S}^{m+1} .

If the problem is considered in hyperbolic spaces, only few results are obtained. For example (see [1]), there exist no proper biharmonic hypersurfaces in \mathbb{H}^4 .

3. II -minimal Translation Surfaces

In this section we analyze II -minimal translation surfaces with a Riemannian second fundamental form, namely we study under which conditions the second mean curvature vanishes, i.e., $H_{II} = 0$. Having in mind the usual technique for computing the second mean curvature by using the normal variation of the area functional one gets for surfaces in \mathbb{E}^3

$$H_{II} = H + \frac{1}{4}\Delta_{II} \log(K)$$

where K and H denote the usual Gaussian, respectively mean curvatures of our surface and Δ_{II} is the Laplacian for functions computed with respect to the second fundamental form as metric. H_{II} can be equivalently expressed as

$$H_{II} = H + \frac{1}{2\sqrt{\det II}} \sum_{i,j} \frac{\partial}{\partial u^i} \left(\sqrt{\det II} h^{ij} \frac{\partial}{\partial u^j} (\ln \sqrt{K}) \right). \quad (3)$$

Here II denotes the second fundamental form, (h_{ij}) is the associated matrix with its inverse (h^{ij}) , the indices i, j belong to the set $\{1, 2\}$ and the parameters u^1, u^2 are x , respectively y from the parametrization (1). Moreover, II becomes a metric on the surface if it is positive definite (or, more generally, if it is non-degenerated). Sometimes, the second mean curvature is taken with the opposite sign (see [15]).

Using the parametrization (1) of a translation surface and denoting by r the corresponding immersion in the Euclidean three-space endowed with the Euclidean

scalar product \langle , \rangle , namely $r : M \rightarrow \mathbb{E}^3, (x, y) \mapsto r(x, y) = (x, y, f(x) + g(y))$ one easily computes its **first fundamental form**

$$I = E dx^2 + 2F dx dy + G dy^2$$

where E, F, G - the coefficients of I - are given by $E = \langle r_x, r_x \rangle, F = \langle r_x, r_y \rangle, G = \langle r_y, r_y \rangle$ and the **second fundamental form**

$$II = L dx^2 + 2M dx dy + N dy^2$$

with the coefficients given by $L = \frac{(r_x, r_y, r_{xx})}{\sqrt{EG-F^2}}, M = \frac{(r_x, r_y, r_{xy})}{\sqrt{EG-F^2}}$ and $N = \frac{(r_x, r_y, r_{yy})}{\sqrt{EG-F^2}}$. Here $(r_x, r_y, r_{xx}) = \langle r_x \times r_y, r_{xx} \rangle$ and the same definition is valid for the other expressions.

Denoting $f' = \alpha$ and $g' = \beta$, we get

$$I = (1 + \alpha(x)^2) dx^2 + 2\alpha(x)\beta(y) dx dy + (1 + \beta(y)^2) dy^2$$

$$II = \frac{1}{\sqrt{\Delta}} (\alpha'(x) dx^2 + \beta'(y) dy^2)$$

where $\Delta = 1 + \alpha(x)^2 + \beta(y)^2$.

The inverse matrix (h^{ij}) of the second fundamental form of a translation surface has the following expression

$$(h^{ij})_{i,j} = \begin{pmatrix} \frac{\sqrt{1+\alpha^2+\beta^2}}{\alpha'} & 0 \\ 0 & \frac{\sqrt{1+\alpha^2+\beta^2}}{\beta'} \end{pmatrix}.$$

The curvatures corresponding to the first fundamental form, the Gaussian curvature $K = \frac{LN-M^2}{EG-F^2}$ and the mean curvature $H = \frac{EN-2FM+GL}{EG-F^2}$ become in this case

$$K = \frac{\alpha'(x)\beta'(y)}{\Delta^2}$$

and

$$H = \frac{(1 + \beta^2(y))\alpha'(x) + (1 + \alpha^2(x))\beta'(y)}{2\Delta^{3/2}}.$$

After straightforward computations, the sum in (3) has the following expression

$$\begin{aligned} \sum_{i,j} &= \frac{1}{4\Delta^2} \sqrt{\frac{\beta'}{\alpha'}} \left(\frac{2\alpha'\alpha'''' - 3\alpha''^2}{\alpha'^2} \Delta^2 + (-4\alpha\alpha'' - 8\alpha'^2)\Delta + 16\alpha^2\alpha'^2 \right) \\ &+ \frac{1}{4\Delta^2} \sqrt{\frac{\alpha'}{\beta'}} \left(\frac{2\beta'\beta'''' - 3\beta''^2}{\beta'^2} \Delta^2 + (-4\beta\beta'' - 8\beta'^2)\Delta + 16\beta^2\beta'^2 \right). \end{aligned}$$

Notice that $\alpha'\beta' > 0$ since the second fundamental form is positive definite, so the square roots are well defined.

We are interested to find II -minimal translation surfaces in the Euclidean three-space. Having now all the necessary tools, the condition $H_{II} = 0$ for a translation surface is equivalent to

$$\frac{2\alpha'\alpha'''' - 3\alpha''^2}{2\alpha'^3} + \frac{2\beta'\beta'''' - 3\beta''^2}{2\beta'^3} - \frac{2}{\Delta} \left(\frac{\alpha'^2 + \alpha\alpha''}{\alpha'} + \frac{\beta'^2 + \beta\beta''}{\beta'} \right) + \frac{6}{\Delta^2} (\alpha^2\alpha' + \beta^2\beta') = 0. \tag{4}$$

The first two terms in (4) are functions only of x respectively of y , hence we take the derivatives in the previous equation successively with respect to x and y .

Denoting by

$$\phi(x) = \frac{\alpha'^2 + \alpha\alpha''}{\alpha'}, \quad \psi(y) = \frac{\beta'^2 + \beta\beta''}{\beta'}, \quad p(x) = \alpha^2\alpha' \quad \text{and} \quad q(y) = \beta^2\beta'$$

we get

$$\frac{\partial}{\partial y} \frac{\partial}{\partial x} \left(-\frac{2}{\Delta} (\phi + \psi) + \frac{6}{\Delta^2} (p + q) \right) = 0.$$

After straightforward computations and multiplying with $\frac{\Delta^3}{8\alpha\alpha'\beta\beta'}$ it follows

$$(F + G)\Delta^2 - 2(P + Q)\Delta + 18(p + q) = 0 \tag{5}$$

where $F(x) = \frac{\phi'}{2\alpha\alpha'}$, $G(y) = \frac{\psi'}{2\beta\beta'}$, $P(x) = \phi + \frac{3p'}{2\alpha\alpha'}$ and $Q(y) = \psi + \frac{3q'}{2\beta\beta'}$. Repeating similar operations, namely taking the two partial derivatives and dividing by $4\alpha\alpha'\beta\beta'$ one gets

$$(A + B)\Delta + a + b = 0 \tag{6}$$

where $A(x) = \frac{F'}{2\alpha\alpha'}$, $B(y) = \frac{G'}{2\beta\beta'}$, $a(x) = F - \frac{P'}{2\alpha\alpha'}$ and $b(y) = G - \frac{Q'}{2\beta\beta'}$.

Finally, using the same technique, we should have

$$\frac{A'}{2\alpha\alpha'} = c, \quad \frac{B'}{2\beta\beta'} = -c, \quad c \in \mathbb{R}.$$

Solving the above equations we obtain $A(x) = c\alpha^2 + d_1$ and $B(y) = -c\beta^2 + d_2$. Replacing these expressions in the previous ODEs we find that

$$F(x) = \frac{c}{2}\alpha^4 + d_1\alpha^2 + \mu_1$$

$$G(y) = -\frac{c}{2}\beta^4 + d_2\beta^2 + \mu_2$$

$$\phi(x) = \frac{c}{6}\alpha^6 + \frac{d_1}{2}\alpha^4 + \mu_1\alpha^2 + \tau_1$$

$$\psi(y) = -\frac{c}{6}\beta^6 + \frac{d_2}{2}\beta^4 + \mu_2\beta^2 + \tau_2$$

$$\alpha'(x) = \frac{c}{42}\alpha^6 + \frac{d_1}{10}\alpha^4 + \frac{\mu_1}{3}\alpha^2 + \tau_1 + \frac{\rho_1}{\alpha}$$

$$\beta'(y) = -\frac{c}{42}\beta^6 + \frac{d_2}{10}\beta^4 + \frac{\mu_2}{3}\beta^2 + \tau_2 + \frac{\rho_2}{\beta}$$

$$p(x) = \frac{c}{42}\alpha^8 + \frac{d_1}{10}\alpha^6 + \frac{\mu_1}{3}\alpha^4 + \tau_1\alpha^2 + \rho_1\alpha$$

$$q(y) = -\frac{c}{42}\beta^8 + \frac{d_2}{10}\beta^6 + \frac{\mu_2}{3}\beta^4 + \tau_2\beta^2 + \rho_2\beta$$

$$P(x) = \frac{19c}{42}\alpha^6 + \frac{7d_1}{5}\alpha^4 + 3\mu_1\alpha^2 + 4\tau_1 + \frac{3\rho_1}{2\alpha}$$

$$Q(y) = -\frac{19c}{42}\beta^6 + \frac{7d_2}{5}\beta^4 + 3\mu_2\beta^2 + 4\tau_2 + \frac{3\rho_2}{2\beta}$$

$$a(x) = -\frac{6c}{7}\alpha^4 - \frac{9d_1}{5}\alpha^2 - 2\mu_1 + \frac{3\rho_1}{4\alpha^3}$$

$$b(y) = \frac{6c}{7}\beta^4 - \frac{9d_2}{5}\beta^2 - 2\mu_2 + \frac{3\rho_2}{4\beta^3}$$

where $d_1, d_2, \mu_1, \mu_2, \tau_1, \tau_2, \rho_1, \rho_2 \in \mathbb{R}$. In order to determine all these integration constants, we substitute the corresponding expressions in (6), obtaining a sum of polynomials in α and β which are equal to zero. This means that there exists $\xi \in \mathbb{R}$ such that

$$\frac{c}{7}\alpha^4 + \left(c - \frac{4}{5}d_1 + d_2\right)\alpha^2 + \frac{3\rho_1}{4\alpha^3} + d_1 - 2\mu_1 - \xi = 0$$

$$-\frac{c}{7}\beta^4 + \left(-c + d_1 - \frac{4}{5}d_2\right)\beta^2 + \frac{3\rho_2}{4\beta^3} + d_2 - 2\mu_2 + \xi = 0.$$

At this point, by the same argument as in the previous section, all the coefficients in the above (algebraic) expressions must be zero and consequently we get $c = 0$, $d_1 = d_2 = 0$, $\rho_1 = \rho_2 = 0$, $\mu_1 = -\frac{\xi}{2}$ and $\mu_2 = \frac{\xi}{2}$. Thus, the previous expressions

can be expressed in a simpler form

$$\begin{aligned}
 F(x) &= -\frac{\xi}{2}, & G(y) &= \frac{\xi}{2} \\
 \phi(x) &= -\frac{\xi}{2}\alpha^2 + \tau_1, & \psi(y) &= \frac{\xi}{2}\beta^2 + \tau_2 \\
 \alpha'(x) &= -\frac{\xi}{6}\alpha^2 + \tau_1, & \beta'(y) &= \frac{\xi}{6}\beta^2 + \tau_2 \\
 p(x) &= -\frac{\xi}{6}\alpha^4 + \tau_1\alpha^2, & q(y) &= \frac{\xi}{6}\beta^4 + \tau_2\beta^2 \\
 P(x) &= -\frac{3\xi}{2}\alpha^2 + 4\tau_1, & Q(y) &= \frac{3\xi}{2}\beta^2 + 4\tau_2 \\
 a(x) &= \xi, & b(y) &= -\xi.
 \end{aligned} \tag{7}$$

Let us look now at (5). By the same reasoning as above we deduce

$$(3\xi + 10\tau_1 - 8\tau_2)\alpha^2 - 8\tau_1 = \eta$$

$$(-3\xi - 8\tau_1 + 10\tau_2)\beta^2 - 8\tau_2 = -\eta$$

for an arbitrary $\eta \in \mathbb{R}$. Moreover, the constants τ_1 , τ_2 and ξ are given by $\tau_1 = -\frac{\eta}{8}$, $\tau_2 = \frac{\eta}{8}$, $\xi = \frac{3\eta}{4}$. We conclude that $\alpha' = -\frac{\eta}{8}(\alpha^2 + 1)$ and $\beta' = \frac{\eta}{8}(\beta^2 + 1)$. Finally α and β must satisfy also the condition (4). After straightforward computations it follows that $\eta = \xi = 0$.

It follows that $\alpha' = \beta' = 0$, which cannot occur since if this happened, the second fundamental form would vanish identically. Hence, the second mean curvature is not defined and we end this section with the following non-existence theorem.

Theorem 1. *There are no II-minimal translation surfaces in Euclidean three-space.*

Acknowledgments

The first author was supported by grant PN-II ID 398/2007-2010, CNCSIS, Romania. The second author was partially supported by grant PN-II ID 457/2009-2011, CNCSIS, Romania.

References

- [1] Balmuş A., *Biharmonic Maps and Submanifolds*, PhD. Thesis, Università degli Studi di Cagliari, Italy, 2007.
- [2] Balmuş A., Montalo S. and Oniciuc C., *Properties of Biharmonic Submanifolds in Spheres*, JGSP **17** (2010) 87-102.

- [3] Cartan E., *Les surfaces qui admettent une seconde forme fondamentale donnée*, Bull. Sci. Math. **67** (1943) 28–32.
- [4] Caddeo R., Montaldo S. and Oniciuc O., *Biharmonic Submanifolds of \mathbb{S}^3* , Internat. J. Math. **12** (2001) 867–876.
- [5] Darboux G., *Leçons sur la théorie générale des surfaces*, Gauthiers-Villars, Paris 1915.
- [6] Dillen F. and Sodsiri W., *Ruled Surfaces of Weingarten Type in Minkowski 3-space*, J. Geom. **83** (2005) 10–21.
- [7] Erard P., *Über die zweite Fundamentalform von Flächen im Raum*, Abhandlung zur Erlangung der Würde eines Doktors der Mathematik der Eidgenössischen Technischen Hochschule Zürich, Diss. # 4234, ETH Zürich, 1968.
- [8] Glässner E., *Über die Minimalflächen der Zweiten Fundamentalform*, Monatsh. Math. **78** (1974) 193–214.
- [9] Haesen S., Verpoort S. and Verstraelen L., *The Mean Curvature of the Second Fundamental Form*, Houston J. Math. **34** (2008) 703–719.
- [10] Jiang G., *2-harmonic Maps and Their First and Second Variation Formulas* (in Chinese), Ann. Math. Ser. A **7** (1986) 389–402.
- [11] Manhart F., *Die II-Minimalregelflächen*, Anzeiger der Österreichische Akademie der Wissenschaften, Mathematisch-Naturwissenschaftliche Klasse, **119** (1982) 157–160.
- [12] Manhart F., *Die Affinminimalrückungsflächen*, Arch. Math. **44** (1985) 547–556.
- [13] Munteanu M. and Nistor A., *On the Geometry of the Second Fundamental Form of Translation Surfaces in \mathbb{E}^3* , to appear in Houston J. Math., arXiv:0812.3166v1 [math.DG].
- [14] Selig J., *Geometric Fundamentals of Robotics*, Second Edition, Springer, Berlin, 2005.
- [15] Verpoort S., *The Geometry of the Second Fundamental Form: Curvature Properties and Variational Aspects*, PhD. Thesis, Katholieke Universiteit Leuven, Belgium, 2008.
- [16] Weingarten J., *Über die Deformationen einer biegsamen unausdehnbaren Fläche*, J. für die Reine und Angew. Math. (Crelle's Journal) **100** (1887) 296–310.